


ARTICLE

DOI: 10.1038/s41467-017-01855-z

OPEN

$G\alpha_i$ is required for carvedilol-induced β_1 adrenergic receptor β -arrestin biased signaling

Jialu Wang^{1,2}, Kenji Hanada², Dean P. Staus², Michael A. Makara², Giri Raj Dahal², Qiang Chen², Andrea Ahles³, Stefan Engelhardt^{3,4}  & Howard A. Rockman^{1,2,5}

The β_1 adrenergic receptor (β_1 AR) is recognized as a classical $G\alpha_s$ -coupled receptor. Agonist binding not only initiates G protein-mediated signaling but also signaling through the multifunctional adapter protein β -arrestin. Some β AR ligands, such as carvedilol, stimulate β AR signaling preferentially through β -arrestin, a concept known as β -arrestin-biased agonism. Here, we identify a signaling mechanism, unlike that previously known for any $G\alpha_s$ -coupled receptor, whereby carvedilol induces the transition of the β_1 AR from a classical $G\alpha_s$ -coupled receptor to a $G\alpha_i$ -coupled receptor stabilizing a distinct receptor conformation to initiate β -arrestin-mediated signaling. Recruitment of $G\alpha_i$ is not induced by any other β AR ligand screened, nor is it required for β -arrestin-bias activated by the β_2 AR subtype of the β AR family. Our findings demonstrate a previously unrecognized role for $G\alpha_i$ in β_1 AR signaling and suggest that the concept of β -arrestin-bias may need to be refined to incorporate the selective bias of receptors towards distinct G protein subtypes.

¹Department of Cell Biology, Duke University Medical Center, Durham, NC 27710, USA. ²Department of Medicine, Duke University Medical Center, Durham, NC 27710, USA. ³Institute of Pharmacology and Toxicology, Technical University of Munich, Munich 80802, Germany. ⁴German Center for Cardiovascular Research (DZHK), Partner Site Munich Heart Alliance, Munich 80802, Germany. ⁵Departments of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC 27710, USA. Correspondence and requests for materials should be addressed to H.A.R. (email: h.rockman@duke.edu)

G protein-coupled receptors (GPCRs) represent the largest and the most versatile family of cell surface receptors¹. Members of this receptor family translate diverse extracellular cues to intracellular responses, and are commonly targeted for medicinal therapeutics^{2,3}. One of the most commonly used therapeutic agents in medicine are ligands that target β adrenergic receptors (β ARs) because they regulate many important physiological processes involved in the regulation of cardiovascular and pulmonary function⁴.

GPCRs selectively couple to different heterotrimeric G protein complexes ($G\alpha\beta\gamma$) that are classified into four families based on their α -subunits: $G\alpha_{\text{stimulatory}}$ ($G\alpha_s$), $G\alpha_{\text{inhibitory/olfactory}}$ ($G\alpha_i/o$), $G\alpha_{q/11}$, and $G\alpha_{12/13}$ ⁵. Among the different G protein subtypes, β ARs primarily transmit signals through $G\alpha_s$ ⁶. In the classical paradigm of β AR signaling, receptors exist in two distinct conformational states: active or inactive. Agonist binding stabilizes an active β AR conformation that promotes coupling with heterotrimeric G proteins, triggering guanine nucleotide exchange of $G\alpha_s$ and its dissociation from the $G\beta\gamma$ subunits, leading to the activation of adenylyl cyclase and triggering second messenger cyclic AMP signaling^{7,8}. Subsequent to agonist binding, activated β ARs are phosphorylated by G protein-coupled receptor kinases (GRKs) leading to recruitment of the multifunctional β -arrestins (β -arrestin1 and β -arrestin2) and inhibition of further G protein coupling, a process termed desensitization⁸. It is now appreciated that β -arrestins also act as signal transducers in their own right⁷ to stimulate a distinct array of signaling and cellular responses, such as transactivation of the epidermal growth factor receptor (EGFR)^{9,10}, induction of extracellular signal-regulated kinase (ERK)^{10–13}, and activation of Ca^{2+} /calmodulin kinase II (CaMKII)¹⁴. Current data suggest a much greater complexity of GPCR signaling than the two-state (active or inactive) model whereby multiple receptor conformations can exist, each with a different affinity for its transducer, resulting in the activation of distinct cellular signaling pathways^{15–17}. Whereas balanced ligands, such as isoproterenol, stabilize β AR conformations signal with equal efficacy through G proteins and β -arrestins, some ligands stabilize conformations that selectively recruit only one of the transducers to stimulate a specific subset of cellular signals, a process termed “biased agonism”^{18,19}. As biased ligands may be capable of selectively activating beneficial signaling while simultaneously blocking untoward receptor activated pathways²⁰, understanding mechanisms of biased agonism can have important implications for drug discovery targeting GPCRs.

The β -blocker carvedilol is a β -arrestin-biased β AR ligand that preferentially activates β -arrestin-mediated pathways while having inverse agonism towards $G\alpha_s$ signaling^{7,10,19,21}. To date, the prevailing mechanistic concept of β -arrestin-bias for the $G\alpha_s$ -coupled β_1 AR is ligand-stimulated activation of β -arrestin in the absence of G protein coupling. However, recently it has been demonstrated for the angiotensin II type 1 receptor that the weak β -arrestin-biased agonist, [¹Sar⁴Ile⁸Ile]-angiotensin II, is capable of activating both $G\alpha_q$ and $G\alpha_i$ ²², indicating a possible role of G proteins in β -arrestin-mediated signaling. Moreover, recent biophysical work suggests that both G protein and β -arrestin can simultaneously interact with an activated GPCR to form super complexes²³, raising the possibility that association of β -arrestin with the receptor may not preclude interaction with a G protein. Here, we set out to test whether G protein coupling is a critical component of β -arrestin-biased β AR signaling. Our findings show that carvedilol, unique among other β AR agonists or antagonists tested, selectively promotes the recruitment of $G\alpha_i$ to β_1 ARs to initiate β -arrestin-biased signaling. These data underscore the complexity of β -arrestin-biased agonism and have important implications for identifying new therapeutic agents to selectively target β -arrestin-biased signaling.

Results

$G\alpha_i$ is required for carvedilol-induced β_1 AR-mediated ERK. Previous studies have demonstrated that carvedilol induces β AR-mediated ERK phosphorylation in a $G\alpha_s$ -independent, β -arrestin-dependent manner^{10,21}. To determine whether $G\alpha_i$ is required for carvedilol-stimulated β AR signaling, we tested the effect of the $G\alpha_i$ inhibitor pertussis toxin (PTX) on carvedilol-stimulated ERK phosphorylation in HEK293 cells stably expressing FLAG-tagged β_1 AR or β_2 AR. PTX catalyzes the ADP-ribosylation of $G\alpha_i$ and prevents $G\alpha_i$ coupling to ligand bound receptors. In β_1 AR stable cells, carvedilol dose dependently increased ERK phosphorylation, which was significantly diminished by pretreatment with the $G\alpha_i$ inhibitor PTX (Fig. 1a, Supplementary Fig. 1a). In contrast, PTX had no effect on the carvedilol-induced β_2 AR-mediated ERK phosphorylation (Fig. 1a, Supplementary Fig. 1a). These observations suggest that $G\alpha_i$ is needed for carvedilol-induced β_1 AR, but not β_2 AR signaling.

To further delineate the role of $G\alpha_i$ in carvedilol-induced β AR signaling, we measured the level of ERK activation in β_1 AR or β_2 AR stable cells after removing $G\alpha_i$ using CRISPR/Cas9 gene editing. All three subtypes of $G\alpha_i$ ($G\alpha_{i1}$, $G\alpha_{i2}$, and $G\alpha_{i3}$) were depleted with their specifically targeted guide RNAs (Supplementary Fig. 1b). $G\alpha_i$ depletion markedly blocked carvedilol-induced ERK phosphorylation in β_1 AR stable cells, while it had no effect in β_2 AR stable cells (Fig. 1b). The absence of $G\alpha_i$ was considerably more robust in abrogating carvedilol-stimulated ERK phosphorylation compared to that observed with PTX treatment (Fig. 1a).

We then determined if a similar signaling mechanism is involved in heart tissue by measuring ERK phosphorylation in Langendorff perfused mouse hearts following carvedilol stimulation. To study the specific effect of carvedilol on the β_1 AR, we used previously generated β_2 AR knockout mice²⁴. Carvedilol perfusion robustly stimulated ERK phosphorylation in hearts of β_2 AR knockout mice, which was entirely abrogated in hearts of PTX-pretreated mice (Fig. 1c). In contrast, in β_1 AR knockout mice²⁵ while carvedilol robustly induced ERK phosphorylation by activating the β_2 AR, PTX pretreatment was unable to block ERK activation (Fig. 1c). These data are consistent with our in vitro data and indicate a previously unrecognized, β AR subtype specific, requirement for $G\alpha_i$ in carvedilol-induced β_1 AR signaling.


Carvedilol-induced β_1 AR conformational change requires $G\alpha_i$.

Different ligands for the same receptor stabilize unique conformational states promoting coupling to selective signal transducers and activation of distinct downstream signaling pathways^{7,20}. Since we showed that $G\alpha_i$ is required for carvedilol-induced β_1 AR signaling, we tested whether it allosterically stabilizes a unique carvedilol-bound β_1 AR conformation. We utilized a fluorescence resonance energy transfer (FRET)-based β_1 AR conformational sensor in which Cerulean (Cer) and YFP are inserted in the C-terminus and third intracellular loop of the receptor, respectively (Fig. 2a)²⁶. Agonist-induced β_1 AR activation is represented by the loss of FRET, i.e., decrease of YFP/Cer ratio²⁶. To test whether $G\alpha_i$ stabilizes a carvedilol-induced β_1 AR conformation, HEK293 cells stably expressing the β_1 AR FRET sensor were pretreated with vehicle or PTX, then stimulated with the balanced agonist isoproterenol or the β -arrestin-biased agonist carvedilol while monitoring the FRET ratio in real time. We found that compared to isoproterenol which caused a decrease in the FRET ratio, carvedilol induced a directional opposite response to the FRET signal, whereas the β_1 AR antagonist metoprolol showed no effect (Fig. 2b). Importantly, pretreatment with PTX significantly diminished the carvedilol-induced FRET ratio without any effect on the isoproterenol stimulated FRET-based receptor biosensor (Fig. 2c). Lastly, PTX alone did not affect the

FRET ratio (Supplementary Fig. 2). These data demonstrate the β_1 AR adopts a distinct conformational state when bound to isoproterenol compared to carvedilol and that $G\alpha_i$ is needed to stabilize the carvedilol-bound β_1 AR conformation.

Carvedilol selectively promotes $G\alpha_i$ recruitment to β_1 ARs. To determine the mechanism of how $G\alpha_i$ is involved in carvedilol-induced β_1 AR signaling, we measured ligand-promoted $G\alpha_i$ recruitment to β ARs with an in situ proximity ligation assay

(PLA), a confocal-microscopy based assay that allows direct visualization and quantification of protein-protein interactions. Using HEK293 cells stably expressing β_1 ARs, we show an over twofold increase in the PLA signal after carvedilol treatment, indicating recruitment of $G\alpha_i$ to the β_1 AR (Fig. 3a). In contrast, carvedilol had no effect on the recruitment of $G\alpha_i$ to β_2 ARs, but $G\alpha_i$ was robustly recruited by isoproterenol consistent with the known process of G protein switching for β_2 ARs²⁷ (Fig. 3a). Importantly, pretreatment with the β AR antagonist propranolol blocked the carvedilol-induced $G\alpha_i$ recruitment to β_1 ARs



(Fig. 3b), indicating the recruitment is dependent on ligand interaction with the β_1 AR orthosteric binding pocket. To further demonstrate recruitment of $G\alpha_i$ to carvedilol-stimulated β_1 AR, we also performed co-immunoprecipitation experiments. Carvedilol stimulation increased the amount of $G\alpha_i$ bound to β_1 ARs in a dose-dependent manner, whereas it resulted in a decrease of $G\alpha_i$ that could be co-immunoprecipitated with β_2 ARs (Fig. 3c, Supplementary Fig. 3a). As a control for the effect of detergent on protein interaction during the co-immunoprecipitation, experiments were also performed with 1% n-Dodecyl β -D-maltoside (DDM) lysis buffer and showed similar results (Supplementary Fig. 3b). The amount of $G\alpha_i$ bound to β_2 ARs was increased by the balanced agonist isoproterenol (Fig. 3c), as we observed with the PLA experiments and again consistent with the previously identified process of $G\alpha_s/G\alpha_i$ switching²⁷.

We next determined whether carvedilol could induce $G\alpha_i$ protein activation using an antibody that specifically recognizes the active GTP-bound $G\alpha_i$. Carvedilol stimulation promoted the activation of $G\alpha_i$ in β_1 AR stable cells, but not in β_2 AR stable cells (Fig. 3d), which was blocked by PTX (Supplementary Fig. 3c).

To determine whether $G\alpha_i$ recruitment is specifically stimulated by carvedilol, we tested a number of β AR agonists and antagonists with PLA (Fig. 4a) and co-immunoprecipitation (Fig. 4b, Supplementary Fig. 4). Remarkably, no other ligand tested induced $G\alpha_i$ recruitment to β_1 ARs, suggesting that this process may be a unique property of the β -arrestin-biased ligand carvedilol.

Collectively, these data support a concept that carvedilol selectively promotes the recruitment and activation of $G\alpha_i$ to the β_1 AR subtype triggering β -arrestin-mediated signaling.

Signaling dependence on both $G\alpha_i$ and β -arrestins. Previous studies have shown that carvedilol stimulation of β_1 ARs promotes the internalization and activation of EGFRs, which in turn activates downstream signaling such as ERK phosphorylation¹⁰. To dissect the mechanism of carvedilol-induced $G\alpha_i$ -dependent signaling, we tested the effect of PTX on β_1 AR-mediated EGFR internalization. We transfected HEK293 cells stably expressing β_1 ARs with GFP-tagged EGFR, and monitored internalization by confocal microscopy. When stimulated with isoproterenol or carvedilol, GFP-EGFR redistributed from the plasma membrane into endosomes, similar to that observed after EGF treatment (Fig. 5a). Pretreatment with PTX significantly blocked the carvedilol-induced EGFR internalization, while without any effect on the isoproterenol response, indicating a requirement for $G\alpha_i$ for carvedilol-induced response (Fig. 5a).

Consistent with previous study showing that carvedilol-induced β_1 AR-mediated EGFR transactivation is β -arrestin-dependent¹⁰, siRNA knockdown of β -arrestin1 and β -arrestin2 abrogated both isoproterenol- and carvedilol-induced EGFR internalization (Fig. 5b, Supplementary Fig. 5a). While

transactivation triggered EGFR internalization induced by both ligands are β -arrestin dependent, the precise molecular mechanism appears to have distinct features. Whereas the carvedilol-induced response requires both $G\alpha_i$ and β -arrestin, the isoproterenol-induced response is PTX insensitive.

To more robustly determine the role of $G\alpha_i$ and β -arrestin in carvedilol-stimulated EGFR transactivation, we generated β -arrestin or $G\alpha_i$ deficient cells using CRISPR-Cas9 gene editing (Supplementary Fig. 5b). The wild type, $G\alpha_i$ knockout or β -arrestin1/2 knockout cells were transfected with CFP-tagged β_1 AR. After ligand stimulation, the level of cell surface EGFRs was analyzed by flow cytometry (Fig. 5c). In wild-type cells, EGFRs were internalized following the treatment with EGF, isoproterenol or carvedilol. The depletion of $G\alpha_i$ blocked carvedilol-induced EGFR internalization, whereas absence of $G\alpha_i$ had no effect on EGF- or isoproterenol-induced responses. In contrast, β -arrestin1/2 knockout cells showed impaired EGFR internalization in response to both isoproterenol and carvedilol. Taken together, these results suggest that the carvedilol-induced EGFR internalization are dependent on both $G\alpha_i$ and β -arrestins.

Consistent with our observation for EGFR internalization, carvedilol-induced ERK phosphorylation required both $G\alpha_i$ and β -arrestins (Fig. 6a, b). Either $G\alpha_i$ inhibition by PTX or β -arrestin knockdown with siRNA diminished carvedilol-induced ERK phosphorylation (Fig. 6a). Moreover, in HEK293 cells transfected with FLAG- β_1 ARs but depleted of either $G\alpha_i$ or β -arrestin, carvedilol stimulated ERK activation was completely abrogated (Fig. 6b). Interestingly, removing either β -arrestin1 or β -arrestin2 prevented carvedilol-stimulated ERK phosphorylation, suggesting that both isoforms are required for carvedilol-stimulated signaling (Fig. 6b).

When β ARs are stimulated by the balanced agonist isoproterenol, protein kinase A (PKA) activated by $G\alpha_s$ -dependent cyclic AMP phosphorylates the receptor leading to a switch of β_2 AR G protein coupling from $G\alpha_s$ to $G\alpha_i$. The now $G\alpha_i$ coupled β_2 AR acts as a negative regulator of $G\alpha_s$ signaling and activates ERK signaling via dissociated $G\beta\gamma$ subunits from heterotrimeric $G\alpha_i$ ^{27–29}. Here, we sought to determine if $G\beta\gamma$ subunits are required for carvedilol-stimulated $G\alpha_i$ -dependent ERK phosphorylation. $G\beta\gamma$ inhibition was achieved by transfection of T8- β ARKct, a chimeric molecule consisting of two components: the C-terminus of the β adrenergic receptor kinase (β ARKct) that competitively binds $G\beta\gamma$, therefore acting as an inhibitor of $G\beta\gamma$ ³⁰; and the extracellular and transmembrane domain of CD8 receptor, which anchors the chimeric protein to the plasma membrane and potentiates its inhibitory effect³¹. The $G\beta\gamma$ blockade efficiency of T8- β ARKct was confirmed by testing its effect on lysophosphatidic acid (LPA)-induced phosphorylation of cyclic AMP-responsive element-binding protein (CREB) (Fig. 7a). We show that the inhibition of $G\beta\gamma$ by T8- β ARKct did not affect the carvedilol-induced ERK activation (Fig. 7b). This suggests that unlike isoproterenol stimulated $G\alpha_i$ -signaling achieved by G

Fig. 1 $G\alpha_i$ is required for the carvedilol-induced β_1 AR-mediated ERK phosphorylation both in vitro and in vivo. **a** Effect of PTX on carvedilol-induced β AR-mediated ERK phosphorylation in HEK293 cells. HEK293 cells stably expressing FLAG-tagged β_1 ARs or β_2 ARs were pretreated with vehicle or 200 ng/ml PTX for 16 h, then stimulated with indicated concentration of carvedilol for 5 min. Carvedilol induced ERK phosphorylation in both β_1 AR or β_2 AR stable cells in dose-dependent manner. The response in β_1 AR stable cells was blocked by PTX, whereas that in β_2 AR stable cells was PTX insensitive. **b** Effect of $G\alpha_i$ knockout on β AR-mediated ERK phosphorylation in HEK293 cells. The $G\alpha_i$ expression in β_1 AR or β_2 AR stable cells was depleted with CRISPR-Cas9 gene editing. Compared with wild-type β_1 AR stable cells, the carvedilol-induced ERK phosphorylation in $G\alpha_i$ knockout β_1 AR stable cells was diminished. In comparison, the response in β_2 AR stable cells was not affected. **c** Effect of PTX on carvedilol-stimulated ERK phosphorylation in Langendorff perfused hearts from β_2 AR knockout mice or β_1 AR knockout mice. Mice were pretreated with vehicle or 25 μ g per kg PTX through intraperitoneal injection. 48 h after injection, mice hearts were excised and perfused with vehicle or 10 μ M carvedilol for 10 min. PTX diminished the carvedilol-induced ERK phosphorylation in hearts from β_2 AR knockout mice, but not β_1 AR knockout mice. Data represent the mean \pm SEM for *n* independent experiments (**a**, **b**) or *n* animals (**c**) as marked on the figure. Statistical significance vs. control was assessed using two-way ANOVA (**a**, **b**) or one-way ANOVA (**c**) with Bonferroni correction. NS no stimulation; Carv carvedilol; Iso isoproterenol; p-ERK phosphorylated ERK; t-ERK total ERK; WT wild type; KO knockout




Fig. 2 $G\alpha_i$ is required for carvedilol-induced β_1 AR conformation change. **a** In the FRET-based β_1 AR conformation sensor, YFP and Cerulean (Cer) are inserted in the third intracellular loop and the C-tail of β_1 AR respectively. **b** Ligand-induced changes of the FRET ratio in HEK293 cells stably expressing β_1 AR FRET sensor. The stable cells were stimulated with 100 μ M carvedilol, 10 μ M isoproterenol, 10 μ M or 100 μ M β_1 AR antagonist metoprolol, while FRET was monitored in real-time as the emission ratio of YFP to Cer. Carvedilol stimulation increased the FRET ratio, while isoproterenol decreased it, demonstrating the distinct β_1 AR conformations induced by these two ligands. Metoprolol had no effect on the FRET ratio. **c** Effect of PTX on ligand-induced FRET ratio change. Cells were pretreated with vehicle or 200 ng per ml PTX for 16 h before ligand stimulation. PTX blocked carvedilol-induced change, while having no effect on the isoproterenol response, suggesting that $G\alpha_i$ is required to stabilize the carvedilol-induced β_1 AR conformation. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells was assessed using one-way ANOVA with Bonferroni correction (**b**); statistical significance between PTX-pretreated and non-pretreated cells was assessed using two-way ANOVA with Bonferroni correction (**c**, left panel), or two-tailed paired Student's t -test (**c**, right panel)

protein switching, carvedilol-induced $G\alpha_i$ -dependent β_1 AR signaling does not require $G\beta\gamma$.

Collectively, these data demonstrate that both $G\alpha_i$ and β -arrestins are required for carvedilol-induced β_1 AR signaling. Notably, either PTX pretreatment or β -arrestin knockdown was able to significantly block the carvedilol-induced β_1 AR-mediated EGFR internalization and ERK phosphorylation, and these responses were completely abrogated when either $G\alpha_i$ or β -arrestin was depleted by gene editing. Taken together these data suggest that $G\alpha_i$ and β -arrestins are likely involved in the same signaling cascade, rather than acting in parallel pathways downstream of β_1 AR.

Phosphorylation of β_1 AR is not required for $G\alpha_i$ recruitment.

GRK-mediated receptor phosphorylation plays a critical role in β -arrestin-dependent signaling of β ARs³². When β_1 ARs are stimulated by balanced agonists, such as isoproterenol or dobutamine, GRK-mediated β_1 AR phosphorylation of the carboxyl-terminal tail occurs and is required for agonist mediated β -arrestin recruitment^{9,33}. For the β_2 AR, a similar process has been shown to occur whereby stimulation with the β -arrestin-biased agonist carvedilol promotes β_2 AR phosphorylation at specific


GRK sensitive amino acid residues 355 and 356 of the c-terminal tail^{21,32}. Here, we sought to determine whether GRK-mediated phosphorylation of the β_1 AR is a necessary step in the carvedilol-induced $G\alpha_i$ recruitment to the receptor. To address this question, we used a mutant β_1 AR that lacks the putative GRK phosphorylation sites within the receptor carboxyl-terminal tail (GRK- β_1 AR) and therefore unable to be phosphorylated by GRKs^{9,33}. We show that carvedilol stimulation increased $G\alpha_i$ recruitment to a similar extent between wild type and GRK- β_1 ARs, as assessed by co-immunoprecipitation (Fig. 8a) and suggests that GRK-mediated β_1 AR phosphorylation is not required for carvedilol-induced $G\alpha_i$ recruitment to the β_1 AR.

β ARs can switch coupling from $G\alpha_s$ to $G\alpha_i$ when stimulated with a balanced agonist^{27,29}. In the $G\alpha_s$ - $G\alpha_i$ switching model, agonist stimulated β_2 AR- $G\alpha_i$ coupling is dependent on PKA-mediated receptor phosphorylation^{27,29}. To determine whether a similar mechanism is involved in the carvedilol-induced $G\alpha_i$ recruitment to β_1 ARs, we used a β_1 AR mutant lacking the putative PKA phosphorylation sites (PKA- β_1 AR) or the PKA inhibitor H89. In our experiments, carvedilol stimulation promotes the $G\alpha_i$ recruitment to PKA- β_1 ARs, similar as to wild-type receptors (Fig. 8a), whereas PKA inhibition with H89 did not have a significant effect (Fig. 8b).

Taken together, these data suggest that neither GRK- nor PKA-mediated receptor phosphorylation is required for carvedilol-induced $G\alpha_i$ recruitment to β_1 ARs.

β_1 AR C-tail is required but not sufficient for $G\alpha_i$ coupling. Since the C-terminus of the β ARs play vital roles in recruiting signal effectors and regulating downstream signaling³⁴, we postulated that specific amino acid residues within the β_1 AR C-tail

are critical for receptor subtype specificity of $G\alpha_i$ recruitment. To test this hypothesis, we transfected HEK293 cells with β AR chimera mutants in which the C-tail of β_1 ARs was exchanged with that of β_2 ARs¹⁴, and assessed $G\alpha_i$ recruitment to chimera β ARs with co-immunoprecipitation. Carvedilol stimulation promoted the recruitment of $G\alpha_i$ to the wild-type β_1 ARs, but was abrogated when the β_1 AR contained the C-tail from the β_2 AR ($\beta_{1/2}$ AR) (Fig. 8c). In contrast, the effect of carvedilol on $G\alpha_i$ recruitment to the β_2 AR with the β_1 AR C-tail ($\beta_{2/1}$ AR) was similar to that of



wild-type β_2 ARs. These data suggest that the C-tail of the β_1 AR is required for $G\alpha_i$ recruitment, but alone is insufficient for this process to occur and is consistent with the crystal structure of the β_2 AR and G protein complex showing multiple receptor-G protein contact points³⁵.

Discussion

In this study, we provide new insight into the molecular mechanism of biased agonism at the β_1 AR. Carvedilol, a ligand classically known as a β AR antagonist, activates β -arrestin signaling by switching the uniquely $G\alpha_s$ -coupled β_1 AR to a $G\alpha_i$ -coupled receptor. We show that carvedilol is unique among a number of agonists and antagonists tested to promote the recruitment and activation of $G\alpha_i$ to β_1 ARs. The recruited $G\alpha_i$ in turn stabilizes a carvedilol-bound β_1 AR conformation that is required for β -arrestin-biased β_1 AR signaling as measured by EGFR internalization and ERK phosphorylation. These results indicate that the previously defined G protein bias vs. β -arrestin-bias may be attributed to ligand-induced selective coupling of receptors to specific G protein subtypes, i.e., G protein subtype bias. In our conceptual model for β_1 AR biased signaling, we speculate that binding of carvedilol to the β_1 AR stabilizes a unique receptor conformation that recruits and activates $G\alpha_i$ to promote β -arrestin-mediated signaling (Fig. 9). While carvedilol is also known to stimulate β_2 AR signaling, $G\alpha_i$ recruitment was not required for β_2 AR-mediated β -arrestin-biased signaling and suggests that different mechanisms for bias may be operative between β AR subtypes.

In the classical view of GPCR signaling, agonist stabilization of specific active conformational states promotes coupling of heterotrimeric G proteins and stimulation of downstream signaling³⁶. Receptor signaling is then terminated by a process involving receptor phosphorylation, β -arrestin recruitment and receptor internalization. However, recent studies suggest that the classical “on-off” (active and inactive) model is oversimplified²⁰, as GPCRs transmit signaling through multiple transducers to regulate diverse arrays of pathways. First, some GPCRs can couple to multiple G proteins. For example, the isoproterenol-activated β_2 AR switches coupling from $G\alpha_s$ to $G\alpha_i$ ²⁷. In this study, we show that carvedilol switches the classical $G\alpha_s$ -coupled receptor β_1 AR to a $G\alpha_i$ -coupled receptor. However, in contrast to the $G\alpha_s$ - $G\alpha_i$ switching of the β_2 AR, the carvedilol-induced β_1 AR- $G\alpha_i$ coupling does not involve $G\alpha_s$ activation and PKA-mediated receptor phosphorylation. The carvedilol-induced β_1 AR- $G\alpha_i$ signaling is also different from the actions of classical $G\alpha_i$ -coupled receptors such as the muscarinic M_2 receptor and the α_2 adrenergic receptor³⁷, as its activation of ERK is not mediated through $G\beta\gamma$ subunits. Second, in addition to their role as signal terminators for G protein signaling, β -arrestins can act as signal transducers in their own right. Current conceptual

models support the idea that ligands may differentially stabilize distinct receptor conformations that recruit divergent portfolio of signaling transducers and effector proteins to active a select suite of cellular signaling pathways, a concept termed functional selectivity or biased agonism¹⁵.

The β -arrestin-biased ligand carvedilol has three unique features at the β_1 AR: (1) it has inverse efficacy for $G\alpha_s$ -dependent adenylyl cyclase activity; (2) it promotes the recruitment of $G\alpha_i$, not $G\alpha_s$, to the β_1 AR; (3) it activates the classical β -arrestin signaling using a $G\alpha_i$ paradigm. These unique signaling properties of carvedilol may be attributed to its ability to stabilize a distinct receptor active conformation¹⁵. For the β_2 AR, carvedilol uniquely induces significant conformational rearrangement around residue Lys263 and Cys265 in the third intracellular loop of the receptor, which may expose the loop toward intracellular surface and facilitate the receptor interaction with β -arrestins¹⁵. Though a previous study suggests the crystal structure of carvedilol-bound β_1 AR is similar to that of the cyanopindolol-bound inactive state structure³⁸, additional conformations stabilized by carvedilol may require the binding of transducers such as $G\alpha_i$ or β -arrestin. This requirement of transducer binding for receptor conformational stability is supported by the structural study of the β_2 AR showing that the interaction of a G protein, or a G protein-like-protein nanobody, is required to stabilize the agonist-induced receptor active conformation³⁹. In our study, using a FRET-based β_1 AR conformation sensor, we show that carvedilol induces a change of FRET ratio, representing a receptor conformational change. Notably, the β_1 AR conformation induced by carvedilol is distinct from the one induced by the balanced agonist isoproterenol, as carvedilol increased the FRET ratio while isoproterenol decreased it. This further supports a concept that receptors can adopt distinct conformations when stimulated by different ligands. As our results show that carvedilol promotes the recruitment of $G\alpha_i$ to β_1 ARs, while a wide range of other β AR ligands tested do not, it is possible that carvedilol induces a β_1 AR conformational change that exposes allosteric binding sites on the receptor to allow for receptor- $G\alpha_i$ interaction. In turn, the bound $G\alpha_i$ stabilizes the carvedilol-induced active receptor conformation and is consistent with our data where pretreatment with the $G\alpha_i$ inhibitor PTX impairs the carvedilol activated β_1 AR conformation. Together these data support the concept that carvedilol-induced $G\alpha_i$ is a positive allosteric modulator of the β -arrestin-biased β_1 AR active conformation.

While we have not determined the precise mechanism of how $G\alpha_i$ binding to the carvedilol-occupied β_1 AR triggers β -arrestin signaling, we postulate that it may involve subsequent receptor phosphorylation in a process known as the “barcode” hypothesis¹⁹. Upon ligands stimulation, GPCRs can be phosphorylated by distinct GRK subtypes at specific sites. Previous study identified β_2 AR sites that are specifically phosphorylated by GRK2 and GRK6³². While the balanced agonist isoproterenol stimulates

Fig. 3 Carvedilol promotes $G\alpha_i$ recruitment and activation in β_1 AR stable cells, but not in β_2 AR stable cells. HEK293 cells stably expressing FLAG-tagged β_1 ARs or β_2 ARs were stimulated with 10 μ M carvedilol or 10 μ M isoproterenol for 5 min. **a** In proximity ligation assay (PLA), cells were immuno-stained with $G\alpha_i$ antibody raised in mouse and β_1 AR (or β_2 AR) antibody raised in rabbit. The red PLA signal represents the protein interactions of $G\alpha_i$ and β_1 AR (or β_2 AR). The area in yellow squares are enlarged for better view. Carvedilol promoted $G\alpha_i$ recruitment to β_1 ARs, but not to β_2 ARs. Scale bar = 20 μ m. **b** β_1 AR stable cells were pretreated with vehicle or 10 μ M propranolol for 30 min. The β AR antagonist propranolol blocked the carvedilol response, suggesting that β_1 AR- $G\alpha_i$ coupling is induced by the binding of carvedilol to the β_1 AR orthosteric binding pocket. Scale bar = 20 μ m. **c** The effect of carvedilol and isoproterenol on $G\alpha_i$ recruitment was confirmed with co-immunoprecipitation assays. FLAG-tagged β_1 ARs or β_2 ARs were immunoprecipitated with anti-FLAG M2 beads, and $G\alpha_{i3}$ was detected with its specific antibody by western blot. **d** Carvedilol specifically activated $G\alpha_i$ in β_1 AR stable cells. Cells were transfected with $G\alpha_{i2}$. 48 h after transfection, cells were treated with 10 μ M carvedilol for 5 min or 30 min, or 10 μ M isoproterenol for 5 min. Activated $G\alpha_i$ was immunoprecipitated with an antibody specifically recognizing the active form of $G\alpha_i$, and immunoblotted with an $G\alpha_i$ antibody. In the middle lane marked as act $G\alpha_i$, cells were transfected with constitutively active $G\alpha_{i2}$, serving as positive control. Carvedilol stimulation activated $G\alpha_i$ in β_1 AR stable cells, but not in β_2 AR stable cells. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells (NS) was assessed using one-way ANOVA with Bonferroni correction




Fig. 4 A number of β AR agonists or antagonists tested do not have significant effect on $G\alpha_i$ recruitment. **a** β_1 AR stable cells were stimulated with vehicle or 10 μM indicated ligands for 5 min. Interaction of β_1 AR and $G\alpha_i$ were detected by PLA. Scale bar = 20 μm . **b** β_1 AR or β_2 AR stable cells were stimulated with ligands at indicated concentration for 5 min. $G\alpha_i$ recruitment was detected by co-immunoprecipitation. Both assays suggested that none of the ligands tested had similar effect of carvedilol on $G\alpha_i$ recruitment. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells was assessed using one-way ANOVA with Bonferroni correction

β_2 AR phosphorylation at both GRK2- and GRK6-specific sites, carvedilol only stimulates receptor phosphorylation at the GRK6-specific sites. This “barcode” phosphorylation pattern of receptors plays essential roles in regulating the recruitment and functionality of signaling transducers¹⁹. For instance, β_2 AR phosphorylation mediated by GRK2 and GRK6 induces distinct β -arrestin conformations, and differentially regulates receptor internalization and ERK activation³². Similarly for the β_1 AR, GRK2-mediated and GRK5/6-mediated receptor phosphorylation leads to distinct cellular responses^{9, 40}, suggesting that a phosphorylation barcode for the β_1 AR may also direct β -arrestin signaling. To dissect the mechanism of how $G\alpha_i$ regulates β_1 AR signaling,

future studies will need to compare the isoproterenol- or carvedilol-induced barcode phosphorylation patterns of the β_1 AR, as well as the effect of $G\alpha_i$ inhibitor PTX on it.

While our data show that both $G\alpha_i$ and β -arrestins are required for carvedilol-induced biased signaling of the β_1 AR, whether β -arrestin is recruited to the carvedilol occupied β_1 AR remains to be determined. Using a number of methodologies, such as co-immunoprecipitation, confocal- or bioluminescence resonance energy transfer-based assays, we were unable to detect carvedilol-induced β -arrestin recruitment to the β_1 AR. This may be due to a number of reasons: (1) ligand-induced β -arrestin recruitment and activation is rapid, within 2 s after stimulation, and reversible⁴¹,




Fig. 5 Both $G\alpha_i$ and β -arrestins are required for carvedilol-induced β_1 AR-mediated EGFR internalization. **a** The effect of PTX on ligand-stimulated EGFR internalization. β_1 AR stable cells with transient transfection of GFP-EGFR were pretreated with vehicle or 200 ng per ml PTX for 16 h. Then the cells were stimulated with 10 μ M carvedilol, 10 μ M isoproterenol or 10 ng per ml EGF for 5 min. Both carvedilol and isoproterenol promoted EGFR internalization, but only the carvedilol-induced response was PTX sensitive. Scale bar = 10 μ m. **b** Either PTX pretreatment or β -arrestins knockdown blocked carvedilol-induced EGFR internalization. β_1 AR stable cells were transfected with GFP-EGFR together with scrambled control siRNA or β -arrestin1/2 siRNA. 48 h after transfection, cells were pretreated with vehicle or 200 ng per ml PTX for 16 h before stimulation. Scale bar = 10 μ m. **c** Carvedilol-induced EGFR internalization was abrogated in β -arrestins or $G\alpha_i$ knockout cells. Wild type, β -arrestin1/2 knockout or $G\alpha_i$ knockout cells were transfected with CFP-tagged β_1 ARs. Cells were stimulated with 10 μ M carvedilol, 10 μ M isoproterenol or 10 ng per ml EGF for 5 min. The EGFR level on cell surface was assessed by flow cytometry. Both carvedilol- and isoproterenol-induced EGFR internalization were impaired in the β -arrestin knockout cells, whereas only the carvedilol-induced response was blocked in the $G\alpha_i$ knockout cells. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance was assessed using two-tailed paired Student's t -test (**a**) or one-way ANOVA with Bonferroni correction (**b**, **c**)

(2) the affinity of the β_1 AR- β -arrestin interaction is low. Both the β_1 AR and the β_2 AR are known as class A receptors, since they are characterized by transient and weak interaction with β -arrestins along with a rapid recycling to the plasma membrane after internalization. To demonstrate carvedilol triggered β -arrestin recruitment to the β_2 AR, previous studies used a chimeric receptor consisting of the β_2 AR fused to vasopressin V_2 receptor

cytoplasmic tail (β_2 AR- V_2 R) to increase the affinity of β -arrestin to the ligand occupied receptor²¹. However, as we have shown (Fig. 8c), the C-tail of the β_1 AR is required for $G\alpha_i$ recruitment. Therefore substituting the β_1 AR C-tail with the V_2 R tail would not provide a chimeric receptor suitable to study the role of $G\alpha_i$ in carvedilol stimulated β -arrestin recruitment. Importantly, we cannot exclude that the carvedilol-stimulated β_1 AR signaling is




Fig. 6 Carvedilol-induced β_1 AR-mediated ERK phosphorylation is dependent on both $G\alpha_i$ and β -arrestins. **a** The effect of PTX and β -arrestin knockdown on carvedilol-stimulated ERK phosphorylation. β_1 AR stable cells with transfection of control siRNA or β -arrestin1/2 siRNA were pretreated with vehicle or PTX, then stimulated with 10 μ M carvedilol for 5 min or 30 min. Carvedilol-stimulated ERK phosphorylation was diminished by either PTX pretreatment or β -arrestins siRNA, suggesting the requirement of $G\alpha_i$ and β -arrestins for this signaling. **b** The β_1 AR-mediated ERK phosphorylation in β -arrestin or $G\alpha_i$ knockout cells. Wild type, β -arrestin knockout or $G\alpha_i$ knockout HEK293 cells were transfected with FLAG-tagged β_1 ARs. Cells were stimulated with 10 μ M carvedilol for 5 or 30 min, 10 μ M isoproterenol or 10 ng per ml EGF for 5 min. The depletion of either β -arrestins or $G\alpha_i$ impaired carvedilol-induced ERK phosphorylation. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells was assessed using one-way ANOVA with Bonferroni correction

mediated by β -arrestin by an indirect mechanism that does not require direct binding of β -arrestin to the β_1 AR. A recent study identified unique features for the β_1 AR with respect to β -arrestin interaction and activation⁴², where a brief interaction with the activated β_1 AR is sufficient to target β -arrestin2 to clathrin-coated structures and trigger ERK signaling even in the absence of receptor association⁴². This β -arrestin “activation at a distance” mechanism suggests that a β_1 AR- β -arrestin complex may not be essential for the activation of β -arrestin-dependent signaling and could explain our findings for a role of β -arrestin in carvedilol-induced signaling without a direct β_1 AR- β -arrestin interaction. Lastly, it is also possible that instead of directly engagement with the β_1 AR, β -arrestins could associate with other components of the signaling cascade such as the transactivated EGFR. This has recently been shown for the vasopressin V_2 receptor signaling, where β -arrestins are recruited to, and act downstream of, the transactivated insulin-like growth factor receptor⁴³.

Carvedilol is a β AR antagonist (β -blocker), a family of drugs that are widely used in the therapeutic treatment of cardiovascular diseases such as hypertension and heart failure, as β_1 ARs and β_2 ARs are predominant GPCR subtypes expressed in

mammalian heart and play vital roles in the regulation of cardiac function⁴. In heart failure, treatment with β -blockers improves left ventricle function, reverses the pathological cardiac remodeling, and reduces mortality and morbidity^{44, 45}. However, β -blockers have different clinical efficacies. Some evidence suggests that carvedilol has a superior effect on cardiovascular survival to other β -blockers⁴⁶. The molecular basis for this remains to be elucidated, but has been attributed to the additional properties of carvedilol other than β -blockers, such as the antioxidant, anti-proliferative effects and α_1 adrenergic receptor blockade⁴⁷. Interestingly, carvedilol appears to be unique among β AR blockers in that it can activate β -arrestin-dependent signaling that confers cardioprotection^{10, 21}. Given the possible cardioprotective role of $G\alpha_i$ during cardiac stress⁴⁸ and the ability of carvedilol to promote β_1 AR- $G\alpha_i$ coupling, it is possible that this unique property of carvedilol is also important for its therapeutic efficacy.

In conclusion, we identify a new signaling mechanism of GPCR biased agonism. To date, the β_1 AR was considered to be predominantly coupled to $G\alpha_s$, and β -arrestin-dependent β_1 AR signaling to be independent of G proteins. However, our data supports a concept where carvedilol has three unique properties




Fig. 7 $G\beta\gamma$ subunits are not required for carvedilol-induced β_1 AR-mediated ERK phosphorylation. **a** Validation of the $G\beta\gamma$ inhibition by T8- β ARKct. HEK293 cells with or without transient transfection of T8- β ARKct were stimulated with 10 μ M LPA for 5 min. T8- β ARKct diminished the LPA-induced CREB phosphorylation, a known $G\beta\gamma$ -dependent process, confirming the inhibition of $G\beta\gamma$ subunits by T8- β ARKct. **b** The $G\beta\gamma$ subunits are not required for carvedilol-induced β_1 AR-mediated ERK phosphorylation. β_1 AR stable cells with or without T8- β ARKct transfection was pretreated with vehicle or 200 ng per ml PTX for 16 h. The cells were then stimulated with 10 μ M carvedilol or 10 μ M isoproterenol for 5 min. T8- β ARKct did not have significant effect on ERK phosphorylation, suggesting that $G\beta\gamma$ subunits were not required. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells was assessed using one-way ANOVA with Bonferroni correction

at the β_1 AR: (1) it is inert with respect to $G\alpha_s$; (2) it recruits $G\alpha_i$ and converts the β_1 AR from a $G\alpha_s$ -coupled receptor to one that couples to $G\alpha_i$; and (3) it activates classical β -arrestin-dependent signaling in a $G\alpha_i$ paradigm. These data suggest a greater complexity for receptor signaling bias than previously appreciated in that coupling of distinct G protein subtypes to the activated receptor are needed for β -arrestin-biased agonism. These data also have important implications when considering the development of new therapeutic ligands designed to selectively target β -arrestin-biased signaling pathways.

Methods

Cell culture. HEK293 cells (American Type Culture Collection) stably expressing FLAG-tagged β_1 AR or β_2 AR are maintained and transfected as previously described^{33, 49}. Cells were periodically treated with BMCyclin (Roche) to avoid mycoplasma contamination. Cells were incubated overnight in serum-free medium supplemented with 0.1% BSA, 10 mM HEPES and 1% penicillin-streptomycin and pretreated with pertussis toxin (200 ng per ml, overnight), H89 (10 μ M, 30 min) or propranolol (10 μ M, 30 min) before ligand stimulation. HEK293 cells stably expressing β_1 AR-FRET sensor were used for the FRET experiments.

Generation of β -arrestin or $G\alpha_i$ knockout cell line. Plasmids carrying *S. pyogenes* Cas9 (SpCas9) next to a cloning site for guide RNA (gRNA) with EGFP (pSpCas9 (BB)-2A-GFP, Addgene 48138) or puromycin resistant gene (pSpCas9(BB)-2A-Puro, Addgene 48139) were obtained from Addgene (deposited by the laboratory of Dr. F. Zhang⁵⁰). Designing of the guide RNAs for $G\alpha_i$ or β -arrestins and cloning the guide RNAs into the Cas9 plasmids were performed as previously described⁵⁰.

For β -arrestin knockout cells, β -arrestin1 was targeted using guide sequence oligos (top: CACCGCATCGACCTCGTGACCCTG; bottom: AACCCAGGGTCCACGAGGTCGATGC). β -arrestin2 was targeted using guide sequence oligos (top: CACCGGTAGATCACCTGGACAAAAG; bottom: AAACCTTTGTCCAGGTGATCTACGC). The guide sequence oligos were cloned into pSpCas9(BB)-2A-Puro. After confirming the cloning by sequencing, plasmids were transfected into HEK293 cells using Fugene 6 transfection reagent (Promega). 72 h after transfection, cells were harvested to check INDEL (insertion deletion) in the genome by surveyor's assay. Puromycin (2.5 μ g per ml) was added into the medium of surveyor positive cells to select cells with the plasmid containing puromycin resistant gene along with guide RNA and Cas9. The knockout of β -arrestins were confirmed by western blot.

For $G\alpha_i$ knockout cells, $G\alpha_{i1}$ was targeted using guide sequence oligos (top: CACCGCGCCGTCTCAGGAGGTTG; bottom: AAACCAACCTCCGTGAGGACGGCGC). $G\alpha_{i2}$ was targeted using guide sequence oligos (top: CACCGAGACAACCGCCCGTACTGC; bottom: AAACGCAGTACCGGGCGGTTGTCTC), and $G\alpha_{i3}$ was targeted using guide sequence oligos (top: CACGGGACGGCTAAAGATTGACTT; bottom: AAACAAGTCAATCTTTAGCCGTCCC). The guide sequence oligos were cloned into pSpCas9 (BB)-2A-GFP. Plasmids targeting the three $G\alpha_i$ subtypes were co-transfected into HEK293 cells. GFP positive cells were selected by fluorescence-activated cell sorting, diluted for growth and single cell colonies were obtained. The $G\alpha_i$ knockout were confirmed by western blot.

Immunoblotting and immunoprecipitation. Following stimulation, cells were scraped in 1% NP-40 lysis buffer (20 mM Tris, pH 7.4, 137 mM NaCl, 20% glycerol, 1% Nonidet P-40, 2 mM sodium orthovanadate, 1 mM PMSF, 10 mM sodium fluoride, 10 μ g per ml aprotinin, 5 μ g per ml leupeptin and phosphatase inhibitors) or 1% DDM lysis buffer (20 mM HEPES, 150 mM NaCl, 1% n-Dodecyl β -D-maltoside, protease inhibitors and phosphatase inhibitors). For immunoprecipitation of FLAG-tagged β_1 AR or β_2 AR, 1–2 mg of protein was incubated overnight with 30 μ l of anti-FLAG M2 magnetic beads (Sigma). For immunoprecipitation of active $G\alpha_i$, protein was incubated for 2 h with anti-active $G\alpha_i$ antibody (New East Biosciences) and Protein A/G beads (EMD Millipore). Immunoprecipitates or cell lysate samples were separated by SDS-PAGE, transferred to PVDF membrane (Bio-Rad) and subjected to immunoblotting with various primary antibodies. Immunoblots were detected using enhanced chemiluminescence (Thermo Fisher Scientific) and analyzed with ImageJ software. Uncropped blots are shown in Supplementary Fig. 6.

Antibodies. Please refer the information of antibodies to Supplementary Table 1.

ERK phosphorylation in mice heart. Eight to 12-week-old gender-matched β_1 AR knockout (β_1 AR KO) mice and β_2 AR KO mice⁵¹ were used for this study. Three to six animals were used for each experimental group based on previous experiments. Randomization and blinding were not performed. Mice were pretreated with vehicle or 25 μ g per kg pertussis toxin (PTX) via intraperitoneal injection. After 48 h, mice were anesthetized with ketamine (100 mg per kg) and xylazine (2.5 mg per kg) for 10 min. Heart was then excised and, with aorta cannulated to needle, perfused with perfusion buffer (118 mM NaCl, 4.7 mM KCl, 1.2 mM $MgSO_4$, 1.2 mM KH_2PO_4 , 2.5 mM $CaCl_2$, 25 mM $NaHCO_3$, 0.5 mM Na-EDTA, 5.5 mM glucose) with O_2 bubbling through Langendorff apparatus (Hugo Sachs Harvard Apparatus) set at 37 $^\circ$ C. After 10 min perfusion, buffer was changed to perfusion buffer with vehicle or 10 μ M carvedilol, and perfused for another 10 min. Heart was




Fig. 8 Neither PKA-mediated nor GRK-mediated β_1 AR phosphorylation is required for carvedilol-induced $G\alpha_i$ recruitment. C-tail of β_1 AR is required but not sufficient for $G\alpha_i$ recruitment. **a** HEK293 cells were transfected with FLAG-tagged wild-type, PKA- or GRK- β_1 ARs. Carvedilol promoted $G\alpha_i$ recruitment to mutant β_1 ARs lacking the putative PKA- or GRK-mediated phosphorylation sites, to a similar extent as to the wild-type β_1 ARs. **b** HEK293 cells stably expressing FLAG-tagged β_1 ARs were pretreated with vehicle or 10 μM H89 for 30 min. The PKA inhibitor H89 did not have a significant effect on carvedilol-induced β_1 AR- $G\alpha_i$ coupling. **c** HEK293 cells were transfected with FLAG-tagged β_1 AR, β_2 AR or chimeric β AR constructs in which the receptor C-tails were exchanged between the two receptor subtypes. Carvedilol did not promote $G\alpha_i$ recruitment to the β_1 AR with C-tail from β_2 AR. On the other hand, the β_1 AR C-tail did not make β_2 AR capable of recruiting $G\alpha_i$ with carvedilol stimulation. Data represent the mean \pm SEM for n independent experiments as marked on the figure. Statistical significance vs. unstimulated cells was assessed using one-way ANOVA with Bonferroni correction




Fig. 9 Schematic model of the carvedilol-induced $G\alpha_i$ - β -arrestin-biased signaling of β_1 ARs. Binding of carvedilol to the orthosteric site of the β_1 AR stabilizes a distinct intermediate conformation that then promotes the recruitment of $G\alpha_i$. The carvedilol- and $G\alpha_i$ -bound receptor in turn stabilizes a unique β_1 AR conformation that mediates β -arrestin-biased signaling

then removed from the system and left ventricle was excised and snap frozen in liquid nitrogen. Animal experiments carried out for this study were handled according to approved protocols and animal welfare regulations the Animal Care and Use Committee of Duke University Medical Center.

Fluorescence resonance energy transfer measurement. FRET measurement was performed as previously described²⁶. Briefly, HEK293 cells stably expressing β_1 AR-FRET sensor were cultured in glass-bottomed confocal dish. Cells were pretreated with vehicle or 200 ng per ml PTX for 16 h before experiment. On the day of experiment, cells were maintained in FRET buffer (10 mM HEPES, 0.2% BSA, 140 mM NaCl, 4.5 mM KCl, 2 mM $CaCl_2$, 2 mM $MgCl_2$, pH 7.4). FRET experiments were performed using an Olympus IX-71 microscope. FRET was monitored as the emission ratio of YFP to Cerulean. Images were taken at 10 s interval and analyzed with ImageJ software.

In situ proximity ligation assay. β_1 AR or β_2 AR stable cells were cultured in 35 mm poly-D-lysine coated glass-bottom confocal dish (MatTek). Following stimulation, cells were fixed in 4% paraformaldehyde for 15 min and permeabilized with 0.2% Triton-X-100 for 10 min. After blocked with blocking buffer from Duolink Detection Kit (Sigma) for 30 min, cells were incubated overnight at 4 °C with anti- β_1 AR (or β_2 AR) antibody from rabbit (Santa Cruz) in conjunction with anti- $G\alpha_i$ antibody from mouse (New East Biosciences). The proximity ligation reaction was performed according to the manufacturer's protocol using the Duolink Detection Kit (Sigma). Cells were mounted with DAPI Fluoromount-G (Southern Biotech). Images were recorded with Zeiss Axio Observer Z1 confocal microscope with $\times 40$ objective. Data analysis was performed with ImageJ software. To quantify the mean PLA signal per cell, the red PLA fluorescence intensity was divided by the number of cells. The mean PLA signal of each data set was corrected by subtracting the background staining determined as the mean PLA signal of HEK293 cells without receptor overexpression. The relative fold over non-stimulation was normalized to the mean PLA signal of the unstimulated cells. In each experiment, 20–40 cells from three images were quantified for each condition.

EGFR internalization assessed by confocal microscopy. HEK293 cells stably expressing FLAG-tagged β_1 AR were transfected with EGFR-GFP together with control siRNA or β -arrestin siRNA as described below. After 24 h, the transfected cells were plated into glass-bottomed confocal dish and kept in culture for additional 24 h. Following pretreatment with PTX and stimulation with ligands, cells were washed with ice-cold PBS and fixed with 4% paraformaldehyde for 15 min. EGFR internalization was visualized with Zeiss Axio Observer Z1 confocal microscope with $\times 63$ objective. In each experiment, 50 cells of each condition were counted under microscope. The percentage of cells showing EGFR internalization was determined by the number of cells showing the intracellular aggregate of EGFR-GFP.

EGFR internalization assessed by flow cytometry. HEK293 cells (wildtype, β -arrestin1/2 knockout or $G\alpha_i$ knockout) were transfected with CFP-tagged β_1 AR. 48 h after transfection, cells were serum starved for 4 h before ligand stimulation. Following stimulation, cells were dissociated with accutase, washed with PBS and fixed in 4% formaldehyde for 15 min at room temperature. Fixed cells were enumerated, washed twice with staining buffer (PBS, 0.5% BSA, 2 mM EDTA) and blocked with 5% rat serum (Sigma) in staining buffer for 15 min. 1×10^6 cells for each sample were stained with equal concentrations of either PE-conjugated EGFR antibody (R&D systems) or isotype control (R&D systems); PE-conjugated rat IgG2A) for 30 min at room temperature. Following staining, cells were washed twice with staining buffer and resuspended in PBS for analysis utilizing a BD LSRII flow cytometer (BD Biosciences). Data analysis was performed with FlowJo

software. Following doublet exclusion, single cells were gated for CFP positivity. To quantify relative EGFR internalization following ligand stimulation, the following formula was utilized: geometric mean fluorescence intensity of the PE-EGFR signal for each data set minus MFI of the isotype control. The resultant value was normalized to the MFI of the unstimulated cells to assess the relative percentage of EGFR internalization.

β -arrestin siRNA knockdown. siRNAs targeting β -arrestin have been described previously¹³. A nonsilencing RNA duplex (5'-AAUUCUCCGAACGUGU-CACGU-3') was used as a control. HEK293 cells stably expressing FLAG-tagged β_1 AR were seeded into 10 cm dish on the day before to reach 30–40% confluence at the time of transfection. siRNA were transfected using GeneSilencer Transfection Reagent (Genlantis) according to the manufacturer's protocol. In brief, 20 μ g siRNA and 240 μ l siRNA dilution buffer were added into 180 μ l serum-free medium, whereas 51 μ l of transfection reagent was mixed with 300 μ l serum-free medium. Both solutions were allowed to stand for 5 min at room temperature, then combined and incubated for additional 20 min. The mixture was then added to cells in the 10 cm dish with 4 ml serum-free medium. After 4 h incubation at 37 °C and 5% CO_2 , 5.5 ml of MEM containing 20% FBS and 2% penicillin-streptomycin were added into the dish. All assays were performed 3 d after siRNA transfection.

Statistical analysis. Data are expressed as mean \pm SEM. Statistical comparisons were performed using two-tailed Student's *t*-test or ANOVA with Bonferroni correction for multiple comparisons in Graphpad Prism. Normality test was performed with Shapiro-Wilk test. Outlier data points more than two standard deviations from the mean were excluded from analysis. Differences were considered statistically significant at $P < 0.05$.

Data availability. All data supporting the findings of this study are available from the authors upon request.

Received: 6 October 2016 Accepted: 19 October 2017

Published online: 22 November 2017

References

- Lagerstrom, M. C. & Schiöth, H. B. Structural diversity of G protein-coupled receptors and significance for drug discovery. *Nat. Rev. Drug Discov.* **7**, 339–357 (2008).
- Ma, P. & Zimmel, R. Value of novelty? *Nat. Rev. Drug Discov.* **1**, 571–572 (2002).
- Overington, J. P., Al-Lazikani, B. & Hopkins, A. L. How many drug targets are there? *Nat. Rev. Drug Discov.* **5**, 993–996 (2006).
- Woo, A. Y. & Xiao, R. P. beta-Adrenergic receptor subtype signaling in heart: from bench to bedside. *Acta Pharmacol. Sin.* **33**, 335–341 (2012).
- Wettschureck, N. & Offermanns, S. Mammalian G proteins and their cell type specific functions. *Physiol. Rev.* **85**, 1159–1204 (2005).
- Salazar, N. C., Chen, J. & Rockman, H. A. Cardiac GPCRs: GPCR signaling in healthy and failing hearts. *Biochim. Biophys. Acta* **1768**, 1006–1018 (2007).
- Rajagopal, S., Rajagopal, K. & Lefkowitz, R. J. Teaching old receptors new tricks: biasing seven-transmembrane receptors. *Nat. Rev. Drug Discov.* **9**, 373–386 (2010).
- Rockman, H. A., Koch, W. J. & Lefkowitz, R. J. Seven-transmembrane-spanning receptors and heart function. *Nature* **415**, 206–212 (2002).
- Noma, T. et al. Beta-arrestin-mediated beta1-adrenergic receptor transactivation of the EGFR confers cardioprotection. *J. Clin. Invest.* **117**, 2445–2458 (2007).
- Kim, I. M. et al. Beta-blockers alprenolol and carvedilol stimulate beta-arrestin-mediated EGFR transactivation. *Proc. Natl Acad. Sci. USA* **105**, 14555–14560 (2008).
- Luttrell, L. M. et al. Beta-arrestin-dependent formation of beta2 adrenergic receptor-Src protein kinase complexes. *Science* **283**, 655–661 (1999).
- Shenoy, S. K. et al. Beta-arrestin-dependent, G protein-independent ERK1/2 activation by the beta2 adrenergic receptor. *J. Biol. Chem.* **281**, 1261–1273 (2006).
- Ahn, S., Wei, H., Garrison, T. R. & Lefkowitz, R. J. Reciprocal regulation of angiotensin receptor-activated extracellular signal-regulated kinases by beta-arrestins 1 and 2. *J. Biol. Chem.* **279**, 7807–7811 (2004).
- Mangmool, S., Shukla, A. K. & Rockman, H. A. Beta-arrestin-dependent activation of Ca(2+)/calmodulin kinase II after beta(1)-adrenergic receptor stimulation. *J. Cell. Biol.* **189**, 573–587 (2010).
- Kahsai, A. W. et al. Multiple ligand-specific conformations of the beta2-adrenergic receptor. *Nat. Chem. Biol.* **7**, 692–700 (2011).
- Manglik, A. et al. Structural insights into the dynamic process of beta2-adrenergic receptor signaling. *Cell* **161**, 1101–1111 (2015).

17. Ye, L., Van Eps, N., Zimmer, M., Ernst, O. P. & Prosser, R. S. Activation of the A2A adenosine G-protein-coupled receptor by conformational selection. *Nature* **533**, 265–268 (2016).
18. Rajagopal, S. et al. Quantifying ligand bias at seven-transmembrane receptors. *Mol. Pharmacol.* **80**, 367–377 (2011).
19. Wisler, J. W., Xiao, K., Thomsen, A. R. & Lefkowitz, R. J. Recent developments in biased agonism. *Curr. Opin. Cell Biol.* **27**, 18–24 (2014).
20. Staus, D. P. et al. Allosteric nanobodies reveal the dynamic range and diverse mechanisms of G-protein-coupled receptor activation. *Nature* **535**, 448–452 (2016).
21. Wisler, J. W. et al. A unique mechanism of beta-blocker action: carvedilol stimulates beta-arrestin signaling. *Proc. Natl Acad. Sci. USA* **104**, 16657–16662 (2007).
22. Sauliere, A. et al. Deciphering biased-agonism complexity reveals a new active AT1 receptor entity. *Nat. Chem. Biol.* **8**, 622–630 (2012).
23. Thomsen, A. R. et al. GPCR-G protein-beta-arrestin super-complex mediates sustained G protein signaling. *Cell* **166**, 907–919 (2016).
24. Chruscinski, A. J. et al. Targeted disruption of the beta2 adrenergic receptor gene. *J. Biol. Chem.* **274**, 16694–16700 (1999).
25. Rohrer, D. K. et al. Targeted disruption of the mouse beta1-adrenergic receptor gene: developmental and cardiovascular effects. *Proc. Natl Acad. Sci. USA* **93**, 7375–7380 (1996).
26. Rochais, F. et al. Real-time optical recording of beta1-adrenergic receptor activation reveals supersensitivity of the Arg389 variant to carvedilol. *J. Clin. Invest.* **117**, 229–235 (2007).
27. Daaka, Y., Luttrell, L. M. & Lefkowitz, R. J. Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. *Nature* **390**, 88–91 (1997).
28. Xiao, R. P., Ji, X. & Lakatta, E. G. Functional coupling of the beta 2-adrenoceptor to a pertussis toxin-sensitive G protein in cardiac myocytes. *Mol. Pharmacol.* **47**, 322–329 (1995).
29. Martin, N. P., Whalen, E. J., Zamah, M. A., Pierce, K. L. & Lefkowitz, R. J. PKA-mediated phosphorylation of the beta1-adrenergic receptor promotes Gs/Gi switching. *Cell. Signal.* **16**, 1397–1403 (2004).
30. Koch, W. J., Inglese, J., Stone, W. C. & Lefkowitz, R. J. The binding site for the beta gamma subunits of heterotrimeric G proteins on the beta-adrenergic receptor kinase. *J. Biol. Chem.* **268**, 8256–8260 (1993).
31. Crespo, P., Cachero, T. G., Xu, N. & Gutkind, J. S. Dual effect of beta-adrenergic receptors on mitogen-activated protein kinase. Evidence for a beta gamma-dependent activation and a G alpha s-cAMP-mediated inhibition. *J. Biol. Chem.* **270**, 25259–25265 (1995).
32. Nobles, K. N. et al. Distinct phosphorylation sites on the beta(2)-adrenergic receptor establish a barcode that encodes differential functions of beta-arrestin. *Sci. Signal.* **4**, ra51 (2011).
33. Rapacciuolo, A. et al. Protein kinase A and G protein-coupled receptor kinase phosphorylation mediates beta-1 adrenergic receptor endocytosis through different pathways. *J. Biol. Chem.* **278**, 35403–35411 (2003).
34. Ritter, S. L. & Hall, R. A. Fine-tuning of GPCR activity by receptor-interacting proteins. *Nat. Rev. Mol. Cell Biol.* **10**, 819–830 (2009).
35. Rasmussen, S. G. et al. Crystal structure of the beta2 adrenergic receptor-Gs protein complex. *Nature* **477**, 549–555 (2011).
36. Dohlman, H. G., Thorner, J., Caron, M. G. & Lefkowitz, R. J. Model systems for the study of seven-transmembrane-segment receptors. *Annu. Rev. Biochem.* **60**, 653–688 (1991).
37. van Biesen, T., Luttrell, L. M., Hawes, B. E. & Lefkowitz, R. J. Mitogenic signaling via G protein-coupled receptors. *Endocr. Rev.* **17**, 698–714 (1996).
38. Warne, T., Edwards, P. C., Leslie, A. G. & Tate, C. G. Crystal structures of a stabilized beta1-adrenoceptor bound to the biased agonists bucindolol and carvedilol. *Structure* **20**, 841–849 (2012).
39. Rosenbaum, D. M. et al. Structure and function of an irreversible agonist-beta (2) adrenoceptor complex. *Nature* **469**, 236–240 (2011).
40. Tilley, D. G., Kim, I. M., Patel, P. A., Violin, J. D. & Rockman, H. A. beta-Arrestin mediates beta1-adrenergic receptor-epidermal growth factor receptor interaction and downstream signaling. *J. Biol. Chem.* **284**, 20375–20386 (2009).
41. Nuber, S. et al. beta-Arrestin biosensors reveal a rapid, receptor-dependent activation/deactivation cycle. *Nature* **531**, 661–664 (2016).
42. Eichel, K., Jullie, D. & von Zastrow, M. beta-Arrestin drives MAP kinase signalling from clathrin-coated structures after GPCR dissociation. *Nat. Cell Biol.* **18**, 303–310 (2016).
43. Oligny-Longpre, G. et al. Engagement of beta-arrestin by transactivated insulin-like growth factor receptor is needed for V2 vasopressin receptor-stimulated ERK1/2 activation. *Proc. Natl Acad. Sci. USA* **109**, E1028–E1037 (2012).
44. Packer, M. et al. Effect of carvedilol on survival in severe chronic heart failure. *N. Engl. J. Med.* **344**, 1651–1658 (2001).
45. McMurray, J. J. & Pfeffer, M. A. Heart failure. *Lancet* **365**, 1877–1889 (2005).
46. Poole-Wilson, P. A. et al. Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the Carvedilol Or Metoprolol European Trial (COMET): randomised controlled trial. *Lancet* **362**, 7–13 (2003).
47. Ohlstein, E. H. et al. Carvedilol, a cardiovascular drug, prevents vascular smooth muscle cell proliferation, migration, and neointimal formation following vascular injury. *Proc. Natl Acad. Sci. USA* **90**, 6189–6193 (1993).
48. DeGeorge, B. R. Jr et al. Targeted inhibition of cardiomyocyte Gi signaling enhances susceptibility to apoptotic cell death in response to ischemic stress. *Circulation* **117**, 1378–1387 (2008).
49. Naga Prasad, S. V. et al. Phosphoinositide 3-kinase regulates beta2-adrenergic receptor endocytosis by AP-2 recruitment to the receptor/beta-arrestin complex. *J. Cell Biol.* **158**, 563–575 (2002).
50. Ran, F. A. et al. Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**, 2281–2308 (2013).
51. Rohrer, D. K., Chruscinski, A., Schauble, E. H., Bernstein, D. & Kobilka, B. K. Cardiovascular and metabolic alterations in mice lacking both beta1- and beta2-adrenergic receptors. *J. Biol. Chem.* **274**, 16701–16708 (1999).

Acknowledgements

We thank Dr K. Xiao (University of Pittsburgh, USA) for intellectual input. We thank Dr R.J. Lefkowitz (Duke University, USA) for providing the beta-arrestin antibodies. We thank Dr M. Bouvier (University of Montreal, Canada) and Dr J.S. Gutkind (University of California, San Diego, USA) for providing the T8-betaARKct construct. This work was supported by National Institutes of Health grants HL056687 and HL075443 to H.A.R. and AHA Predoctoral Fellowship 14PRE20480352 to J.W.

Author contributions

H.A.R. supervised the entire study; H.A.R., J.W. and D.P.S. wrote the manuscript with comments from all co-authors; J.W. designed and performed most of the experiments and analyzed data; K.H. performed the ERK phosphorylation assay in mice heart; M.A.M. assisted with flow cytometry assay; G.R.D. generated the Gα_s and β-arrestin knockout cells; Q.C. assisted with co-immunoprecipitation experiments; A.A. and S.E. provided material and performed preliminary experiments for β₁AR FRET-sensor.

Additional information

Supplementary Information accompanies this paper at doi:10.1038/s41467-017-01855-z.

Competing interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017