

Comparative Evaluation of Several Gene Targets for Designing a Multiplex-PCR for an Early Diagnosis of Extrapulmonary Tuberculosis

Ankush Raj^{1*}, Netrapal Singh^{1*}, Krishna B. Gupta², Dhruva Chaudhary³, Aparna Yadav⁴, Anil Chaudhary⁵, Kshitij Agarwal⁵, Mandira Varma-Basil⁶, Rajendra Prasad⁶, Gopal K. Khuller⁷, and Promod K. Mehta¹

¹Centre for Biotechnology, Maharshi Dayanand University, Rohtak;

Departments of ²TB & Respiratory Medicine, ³Pulmonary and Critical Care Medicine, and ⁴Microbiology, Postgraduate Institute of Medical Sciences (PGIMS), Rohtak;

⁵Rajan Babu Institute of Pulmonary Medicine and Tuberculosis (RBIPMT), Delhi;

⁶Vallabhbai Patel Chest Institute, University of Delhi, Delhi;

⁷Department of Biochemistry, Postgraduate Institute of Medical Education and Research, Chandigarh, India.

Purpose: Diagnosis of extrapulmonary tuberculosis (EPTB) poses serious challenges. A careful selection of appropriate gene targets is essential for designing a multiplex-polymerase chain reaction (M-PCR) assay.

Materials and Methods: We compared several gene targets of *Mycobacterium tuberculosis*, including *IS6110*, *devR*, and genes encoding MPB-64 (*mpb64*), 38kDa (*pstS1*), 65kDa (*hsp65*), 30kDa (*fbpB*), ESAT-6 (*esat6*), and CFP-10 (*cfp10*) proteins, using PCR assays on 105 EPTB specimens. From these data, we chose the two best gene targets to design an M-PCR.

Results: Among all gene targets tested, *mpb64* showed the highest sensitivity (84% in confirmed cases and 77.5% in clinically suspected cases), followed by *IS6110*, *hsp65*, *38kDa*, *30kDa*, *esat6*, *cfp10*, and *devR*. We used *mpb64+IS6110* for designing an M-PCR assay. Our M-PCR assay demonstrated a high sensitivity of 96% in confirmed EPTB cases and 88.75% in clinically suspected EPTB cases with a high specificity of 100%, taking clinical diagnosis as the gold standard.

Conclusion: These M-PCR results along with the clinical findings may facilitate an early diagnosis of EPTB patients and clinical management of disease.

Key Words: *Mycobacterium tuberculosis*, extrapulmonary tuberculosis, PCR, multiplex-PCR, diagnosis

INTRODUCTION

Tuberculosis (TB) remains one of the foremost infectious diseases throughout the world, with an estimated 9.0 million incident cases in 2013, and India ranks first (24% of cases) in total

TB incident cases throughout the world.¹ According to the National Tuberculosis Control Programme, 2.6 million new cases of sputum smear-positive pulmonary TB (PTB), 2.0 million new cases of sputum smear-negative PTB, and 0.8 million new cases of extrapulmonary tuberculosis (EPTB) were observed in 2013 worldwide.¹ EPTB comprises about 15–20% of TB cases and can comprise up to 50% of TB cases in human immunodeficiency virus (HIV)-infected individuals.^{1,2} According to the Revised National Tuberculosis Control Programme, 0.22 million new cases of EPTB were documented in 2013 in India.³ Beyond the Indian subcontinent, EPTB remains a significant health problem in other developing and under-developed countries such as China, Korea, Vietnam, Brazil, Tunisia, Burkina Faso, etc.¹

Diagnosis of smear-positive PTB has been somewhat established; however, the diagnosis of smear-negative PTB, TB-HIV

Received: January 16, 2015 **Revised:** March 21, 2015

Accepted: May 2, 2015

Corresponding author: Dr. Promod K. Mehta, Centre for Biotechnology, Maharshi Dayanand University, Rohtak-124001 (Haryana), India.
Tel: 91-9896504193, Fax: 91-1262-274640, E-mail: pkmehta3@hotmail.com

*Ankush Raj and Netrapal Singh contributed equally to this work.

•The authors have no financial conflicts of interest.

© Copyright: Yonsei University College of Medicine 2016

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

co-infection, and EPTB exhibits serious challenges.^{4,5} Diagnosis of EPTB can be elusive due to paucibacillary load in the biological specimens, lack of adequate clinical sample volume, and localization of disease at sites that are difficult to access.^{2,6} Smear microscopy is the most extensively used method for the diagnosis of EPTB as it is inexpensive and rapid; however, the method has drawbacks due to low and variable sensitivity values (0–40%).⁷ Culture identification, though considered as a gold standard, has drawbacks including low and variable sensitivities (0–80%) in various clinical forms of EPTB,^{2,4,8} a slow turnaround time, and a labor-intensive process. The BACTEC system, histopathological/cytological examination, and interferon- γ release assays are also employed for the diagnosis of EPTB; however, these assays have their own limitations.^{2,7} Detection of adenine deaminase is utilized in diagnosing EPTB specimens, although there is no established cut-off value, and this test has been shown to be positive in other diseases such as lymphomas and collagen vascular diseases.^{2,9} Recently, nucleic acid amplification (NAA) tests such as PCR have emerged as potentially important tools for the diagnosis of EPTB specimens. Various mycobacterial gene targets such as insertion sequence (*IS*)6110; *IS1081*; 16S rRNA DNA; *devR* (transcriptional regulator, Rv3133c); genes encoding 65kDa (heat shock protein 65, *hsp65*; Rv0440), MPB-64/MPT-64 (mycobacterial protein from species tuberculosis, *mpb64*; Rv1980c), 38kDa (phosphate-binding lipoprotein, *pstSI*; Rv0934), and MTP-40 (membrane associated phospholipase C1; Rv2351c) proteins; *TRC4* (conserved repetitive element); *GCRS* (guanine-cytosine-rich repetitive sequence); *fbp*, encoding fibronectin-binding protein B (30kDa, Ag85B protein; Rv1886c); *hupB*, encoding histone-like DNA-binding protein (Rv2986c); and *dnaJ*, encoding chaperone protein (Rv0352), have been exploited for such tests.^{2,5,7,10,11} However, limited information is available for the utilization of the *M. tuberculosis*-specific region of difference (RD) 1 *esat6*, encoding early secretory antigenic target-6 (ESAT-6, Rv3875), and *cfp10*, encoding culture filtrate protein-10 (CFP-10, Rv3874), gene targets. Moreover, there is a high variation in these tests, owing to different gene targets employed as well as the different gold standards adopted in various laboratories. Recently, the introduction of the MTB/RIF GeneXpert (Cepheid, Sunnyvale, CA, USA) assay, which targets conserved *rpoB* gene encoding RNA β polymerase subunit, has been a major breakthrough in the diagnosis of EPTB;¹² however, its use is limited in resource-poor settings due to its high cost. Therefore, we planned a study to compare several gene targets (*IS6110*, *devR*, *mpb64*, *hsp65*, *30kDa*, *38kDa*, *esat6*, and *cfp10*) for PCR assays on EPTB specimens to identify the two best gene targets exhibiting high sensitivities, using clinical diagnosis as the gold standard, and to design a simple, reliable, and cost-effective multiplex-polymerase chain reaction (M-PCR) assay with the two resulting targets for an early diagnosis of EPTB.

MATERIALS AND METHODS

Clinical specimens

The clinical specimens were collected from the patients visiting the clinical departments of Rajan Babu Institute for Pulmonary Medicine & Tuberculosis, Delhi; Vallabhbai Patel Chest Institute, University of Delhi; and the Postgraduate Institute of Medical Sciences, Rohtak during the period of June, 2013 to September, 2014. Written informed consent was obtained from all patients participating in this study. This work was part of a project approved by the Maharshi Dayanand University Institute Ethical Committee under the protocol number MDU/CBT/11/793. This study was double blinded, as the laboratory was not aware of the clinical data and the clinicians were not aware of the laboratory data until all of the analyses were complete.

The sample size (n=155) was determined to be statistically significant with a 95% level of confidence, taking into account the prevalence rate of the study population (Delhi/Rohtak). Samples were divided into two groups. Group 1, the EPTB group (n=105) was subdivided into 1) confirmed EPTB cases [n=25; smear positive (n=21) or culture positive (n=4)] and 2) clinically suspected EPTB cases (n=80) that were smear negative and culture negative yet suspected on the basis of imaging, clinical findings, histological/cytological observations, and response to anti-tubercular therapy (ATT). Group 2 was the control group (n=50), which comprised non-TB individuals.

Among 105 EPTB specimens, there were 64 pleural fluids, six ascitic fluids, 20 pus specimens from different origins, two lymph node aspirates, ten urine specimens, two pleural biopsies, and one synovial fluid specimen. Similarly, pleural fluids (n=23), pus (n=5), ascitic fluids (n=16), and urine (n=6) were collected from non-TB individuals with renal failure, trauma, or cancer as controls. All EPTB patients were in the age group of 15–60 years, and both sexes were represented. The inclusion criteria in the selection of TB pleuritis and TB lymphadenitis patients were 1) fever, cough, pleuritic pain, malaise, anorexia, and exudative pleural effusion with lymphocytic predominance on diagnostic tap and 2) untreated patients with peripheral (superficial) TB lymphadenitis of cervical or axillary regions. Similarly, the inclusion criteria for the selection of abdominal TB patients were fever and weight loss with one or more of the following: diarrhea persisting >1 month; ascites; abdominal lymphadenopathy or mesenteric masses on ultrasound; or hepatomegaly, splenomegaly, generalized pain, or tenderness persisting over 7 days. The exclusion criteria were 1) patients with a history of TB in the preceding 2 years, those already receiving ATT, those with other coexisting medical illnesses (e.g., HIV-positive individuals), pregnant women, and diabetes mellitus patients and 2) smear- and culture-negative suspected EPTB patients who did not respond to ATT even up to 8 weeks of treatment. All EPTB and non-TB specimens except urine were collected in the minor operation theater, fol-

lowing all standard aseptic surgical precautions of the participating institutes. All specimens were stored at 4°C and processed within 24 h.

Sample processing and DNA extraction

All clinical specimens including pleural biopsies (after homogenization) were digested and decontaminated using 1% N-acetyl-L-cysteine combined with 4% NaOH (NALC-NaOH). Briefly, an equal volume of NALC-NaOH solution was added to a clinical sample in a 15-mL centrifuge tube, mixed by inversion and incubated at room temperature for 20 minutes. Later, 0.067 M phosphate buffer (pH 6.8) was added, and the reaction mixture was subjected to centrifugation at 3000× g for 15 min. The supernatant was discarded, and the pellet was used for DNA extraction, smear microscopy, and the culture. The decontamination and centrifugation steps were performed in a biosafety hood. The mycobacterial genomic DNA was extracted according to the method of van Helden, et al.¹³ with few modifications using the cetyl-*N,N,N*-trimethyl ammonium bromide (CTAB)-phenol-chloroform method. Briefly, the pellet was suspended in 500 µL of Tris-EDTA buffer, heated at 100°C for 10 minutes, and incubated with 20 µL of lysozyme (10 mg/mL) at 37°C for 2 h. Then, 30 µL of 10% SDS and 10 µL of proteinase K (20 mg/mL) were added, briefly vortexed, and incubated at 37°C for 1 hour, followed by incubation with 5 M NaCl (100 µL)+10% CTAB (80 µL) at 65°C for 20 min. Later, an equal volume of phenol-chloroform-isoamyl alcohol (25:24:1) mixture was added, mixed by inversion and centrifuged at 14000× g for 10 minutes. This step was repeated twice for the removal of DNA contaminants. The supernatant was removed, followed by the addition of 3 M sodium acetate and isopropanol for DNA precipitation. The precipitate was washed with 70% ethanol and air dried. The DNA pellet was suspended in

100 µL of Tris-EDTA and stored at -20°C until further analysis.

PCR

We designed primers for *IS6110*, *mpb64*, *esat6*, *cfp10*, *38kDa*, *30kDa*, *devR*, and *hsp65* (targets of *M. tuberculosis* H37Rv) using Primer 3 software for PCR and validated the designs with a primer-designing software tool (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). All primers were obtained from Eurofins Genomic India Pvt. Ltd., Bengaluru. Table 1 shows the primer sequence as well as the annealing temperature used for different targets. For PCR amplification of individual gene targets, 1 µL of purified *M. tuberculosis* H37Rv DNA (1 ng/mL) was added to 25 µL of PCR master mix containing 2.5 µL 10× PCR buffer, 1.5 µL of 25 mM MgCl₂, 0.5 µL of 200 µM (each) of the four deoxyribonucleoside triphosphates, 10 pM of each forward and reverse primer, PCR grade water, and 0.625 U of *Taq* DNA polymerase (Bangalore Genei Pvt. Ltd., Bengaluru, India). The DNA amplification was performed in a thermal cycler (T100™ BIO-RAD). After an initial denaturation at 95°C for 3 min, each cycle was composed of three steps: denaturation at 95°C for 30 s followed by annealing of respective primers at an appropriate temperature (Table 1) for 30 s, extension at 72°C for 45 s for 30 cycles, and a final extension at 72°C for 5 min. The PCR-amplified products were resolved by gel electrophoresis using 1.5% agarose in 1X TAE buffer containing ethidium bromide (0.5 µg/mL) at 80 volts for about 40 min and viewed under the Gel Documentation System (MiniLumi, DNR Bio-Imaging Systems Ltd., Jerusalem, Israel) to determine the presence of a specific size band for each gene target. The PCR amplification for DNA isolated from the clinical specimens was performed in a similar manner. All samples were run in duplicate. In each experiment, a positive control (*M. tuberculosis* DNA at 1 ng/mL) and two negative controls (no template DNA, only PCR grade water) were included.

Table 1. Primer Sequences Used for Different Gene Targets of *M. tuberculosis* H37Rv

Gene target	Primers sequence (5'→3')	Annealing conditions (°C/30 s)	Expected amplicon size (bp)
<i>IS6110</i>	Forward GAAGAATCCGCTGAGATAAAGC Reverse GGTTGATGTGGTCGTAGTAGGT	53	258
<i>mpb64</i>	Forward GACTTCTGGTCGGGGTAGTAAC Reverse CTGTCTTTTTGCTCTGTTGTTC	54	163
<i>38kDa (pstS1)</i>	Forward ACACCTTCTTGTTCACCCAGTA Reverse GATGGCGTACTCGTAGTTGATG	53	383
<i>30kDa (fbpB)</i>	Forward TGTACCAGTCGCTGTAGAAG Reverse GACATCAAGGTTCAAGTTCC	55	190
<i>hsp65</i>	Forward GGGCTACATCTCGGGGTA Reverse GGTCTCGTCCTTGGTGAC	52	400
<i>devR</i>	Forward CGTAGTTCTTCACCGTCTT Reverse GACATCAAGGGAATGGAGTT	49	241
<i>esat6</i>	Forward GTCCATTCATTCCCTCCT Reverse CTATGCGAACATCCCAGT	53	220
<i>cfp10</i>	Forward CAGAGATGAAGACCGATG Reverse GAGTTCCTGCTTCTGCTTA	54	203

Table 2. Combination of Different Concentrations of Two Primer Pairs Used for Optimization of M-PCR

Primer pairs	Concentration of primers (μM)			
<i>mpb64</i>	0.2	0.2	0.4	0.4
<i>IS6110</i>	0.2	0.4	0.4	0.2

M-PCR, multiplex-polymerase chain reaction.

M-PCR

M-PCR (*mpb64*+*IS6110*) amplification was performed with the same concentrations of master mix components as for multiplex PCR, with the exception of the different primer pair concentrations. DNA amplification was performed for 30 cycles as described above, although the annealing step was performed at 53°C for 30 s. For the optimization of M-PCR, the combinations of different concentrations of two primer pairs were evaluated (Table 2) using 1 μL of purified *M. tuberculosis* DNA (1 ng/mL) to choose an appropriate ratio for *mpb64*:*IS6110*, which showed the clear amplicons of specific sizes for the two gene targets. PCR amplification for DNA isolated from the clinical specimens was performed in a similar manner and determined the presence or absence of a single band or two bands of specific size(s) under the Gel documentation system. Both positive and negative controls were run as explained above. The PCR/M-PCR results were compared using clinical diagnosis as the gold standard (comprising smear, culture, clinical findings, imaging, histological and cytological observations, response to ATT, and all together).

Statistical methods

The sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) were calculated as detailed by Srivastava, et al.¹⁴

RESULTS

Prior to taking up this study in EPTB samples, validation of the PCR tests using different gene targets was carried out with few sputum samples (n=6) from PTB patients, which were found to be smear-positive and culture-positive for *M. tuberculosis*.

Comparison of PCR results among EPTB specimens with different gene targets

Fig. 1 shows the amplicons of different size(s) using different gene targets (*IS6110*, *mpb64*, *esat6*, *cfp10*, *38kDa*, *30kDa*, *devR*, and *hsp65*) with the purified *M. tuberculosis* DNA via PCR. After optimization of PCR with the purified DNA, we performed PCR using the clinical EPTB as well as non-TB samples. Representatives of positive clinical samples using different gene targets have been depicted (Fig. 1). The order of sensitivity with different targets in 105 EPTB cases (both confirmed and suspected cases) in descending order was *mpb64*>*IS6110*>*hsp65*>*38kDa*>*30kDa*>*esat6*>*cfp10*>*devR* (Table 3). However, the specificity of




Fig. 1. PCR gel picture of several gene targets tested on the same clinical EPTB specimens. L1, L18, L19, and L36 represent 100 bp molecular marker; L2, L6, L10, L14, L20, L24, L28, and L32 were positive controls with the purified *M. tuberculosis* H37RvDNA; L3, L7, L11, L15, L21, L25, L29, and L33 were negative controls without template DNA; L4, L5, L8, L9, L12, L13, L16, L17, L22, L23, L26, L27, L30, L31, L34, and L35 represent clinical EPTB specimens. EPTB, extrapulmonary tuberculosis.

the PCR assay was high (88–100%) with all gene targets employed. The PPV, NPV, and accuracy observed with the individual gene targets have been summarized as percentages in Table 3.

Optimizing the primer-pair concentrations for M-PCR

Considering that *mpb64* and *IS6110* showed the highest positivity among all eight targets examined, we chose these two targets for M-PCR. We optimized the primer concentrations for *mpb64*+*IS6110* (Table 2) to develop an M-PCR. Among the different combinations of primer concentrations, a combination of 0.2 μM of *mpb64* and 0.4 μM of *IS6110* showed sharp amplicons with the purified *M. tuberculosis* DNA (Fig. 2). Therefore, the same combination of primers was incorporated in the PCR master mixture to evaluate the clinical EPTB samples.

M-PCR

The M-PCR targeting *mpb64*+*IS6110* showed better % sensitivity (90.5%) than multiplex PCR (Table 3 and 4) with individual *mpb64* (79%) and *IS6110* (75%) with a high specificity (100%) in 105 EPTB and 50 non-TB specimens (controls). Among the 25 confirmed EPTB cases, M-PCR revealed 24 positive cases, thus leading to 96% positivity. Out of 24 positive cases, 17 were positive with both *IS6110* and *mpb64*, three were positive with *IS6110* only, and four were positive with *mpb64* only. Out of 80 clinically suspected EPTB cases, M-PCR showed 71 positive cases, thus leading to 88.75% positivity. Out of 71 positive cases, 50 were positive with both *IS6110* and *mpb64*, nine were positive with *IS6110* only, and 12 were positive with *mpb64* only (Fig. 3). The M-PCR test demonstrated 100% PPV, 83.33% NPV, and

Table 3. PCR Results Using Various Gene Targets

Results (+/-)	Gene targets															
	mpb64		IS6110		38kDa		hsp65		30kDa		esat6		cfp10		devR	
	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-
n=25	21	4	20	5	18	7	19	6	16	9	17	8	17	8	15	10
Sensitivity (%)	84		80		72		76		64		68		68		60	
PPV (%)	100		100		81.81		82.62		80		77.27		77.27		71.42	
NPV (%)	92.59		90.90		86.79		88.46		83.63		84.90		84.90		81.48	
ACC (%)	86.15		94.66		93.33		85.33		86.66		82.66		82.66		78.66	
n=80	62	18	59	21	51	29	54	26	47	33	45	35	43	37	42	38
Sensitivity (%)	77.5		73.75		63.75		67.50		58.75		56.25		53.75		52.50	
PPV (%)	100		100		92.72		93.10		92.15		90		89.58		87.50	
NPV (%)	73.52		70.42		61.33		63.88		58.22		56.25		54.87		53.65	
ACC (%)	86.15		83.84		74.61		76.92		71.53		69.23		67.69		66.15	
n=105	83	22	79	26	69	36	73	32	63	42	62	43	60	45	57	48
Sensitivity (%)	79.04		75.23		65.71		69.52		60.0		59.04		57.14		54.28	
PPV (%)	100		100		94.52		94.80		94.02		92.53		92.30		90.47	
NPV (%)	69.44		65.78		56.09		58.97		52.27		51.13		50.0		47.82	
ACC (%)	85.16		83.22		74.19		74.19		70.32		69.03		67.74		65.16	
n=50	0	50	0	50	4	46	4	46	4	46	5	45	5	45	6	44
Specificity (%)	100		100		92		92		92		90		90		88	

n, number of specimens; +, PCR positive cases; -, PCR negative cases; PPV, positive predictive value; NPV, negative predictive value; ACC, accuracy; EPTB, extrapulmonary tuberculosis; Total EPTB cases, confirmed+clinically suspected cases.

93.54% accuracy in 105 EPTB cases. The M-PCR results of individual EPTB specimens, including pleural fluids, pleural biopsies, pus, and urine, have been summarized in Table 5. Overall, the sensitivity with microscopy, culture, and M-PCR in 105 EPTB cases was 20%, 3.8%, and 90.47%, respectively, although high specificity (100%) was observed for all these tests in 50 non-TB specimens (controls).

DISCUSSION

The diagnosis of EPTB remains inconclusive in most cases, which can lead to serious consequences. Due to the inadequate sensitivities of conventional bacteriological methods, an unprecedented interest has been stimulated regarding the development of NAA tests such as PCR to facilitate an early diagnosis of EPTB. However, meta-analysis studies have revealed that heterogeneous sensitivities and specificities are observed with in-house PCR tests, while commercial PCR tests yield high specificities yet variable sensitivities.^{2,15} The major drawback of commercial tests is their high cost, which makes them unaffordable in most developing countries with a high TB burden.^{2,15} Furthermore, the true accuracy of PCR tests may actually be different than the reported one if using an imperfect gold or reference standard.^{6,12} Culture is the most widely used gold standard, though it is suboptimal for validating EPTB specimens with varying sensitivities, which can lead to inaccurate PCR results.^{2,16} Several researchers^{6,12,17} used clinical diagnosis and composite reference standard (CRS; combination of smear, culture, histology and cytology, clinical findings, response to ATT, etc.) to validate PCR results when diagnosing EPTB specimens, and this study also used this validation method.

We chose *mpb64* and *IS6110* to develop an M-PCR, which revealed a high sensitivity both in confirmed (96%) and clinically suspected (88.75%) EPTB cases (Table 4). Our M-PCR test was able to detect 16 cases in which *IS6110* was missing yet *mpb64* was present. Similarly, there were 12 cases in which *mpb64* was missing yet *IS6110* was present (Table 4). No false positive results were observed with the 50 non-TB individuals, thus leading to 100% specificity. This was likely due to the utilization of highly specific *M. tuberculosis*-specific gene targets (*mpb64* and *IS6110*) and also good laboratory practices, such as when handling the samples, as well as the inclusion of proper negative controls. Similar to our studies, high sensitivity and specificity has been documented for an M-PCR with *mpb64+IS6110*, designed for the diagnosis of TB lymphadenitis, osteoarticular TB, and gastrointestinal TB,^{10,11,18} however, no report is available on TB pleuritis. We demonstrated high sensitivity (95.3%) and specificity (100%) for M-PCR using *mpb64+IS6110* from pleural fluids (64 from TB patients and 23 from non-TB individuals) for the diagnosis of TB pleuritis (Table 5). Based on a meta-analysis of six studies (598 samples) on the Xpert assay, lower sensitivity (21.4%) has been reported for the diagnosis of TB pleuritis from

pleural fluids, using CRS as the reference standard.¹⁷ Therefore, the World Health Organization (WHO) has not recommended the use of the Xpert assay for the diagnosis of TB pleuritis, though the same assay has been recommended for the diagnosis of TB lymphadenitis and TB meningitis.¹⁷ Interestingly, our M-PCR test was positive in six of ten urine samples collected from the suspected TB pleuritis patients (Table 5). Among those six positive cases, two were missed by *IS6110* yet were detected by *mpb64*. The detection of *M. tuberculosis* in urine samples by PCR can be a useful method for the diagnosis of TB pleuritis and other clinical EPTB forms in which sample collection is difficult and requires aggressive techniques,¹⁹ though it needs further validation in a large number of specimens. The PCR/M-PCR assay is also limited in that it does not discriminate between alive and dead or degraded bacilli. However, this is likely a theoretical limitation, and its relevance must be judged in light of the overall picture of EPTB cases.

Genotype studies of *M. tuberculosis* strains in the Indian pop-

ulation using spoligotyping and *IS6110*-restriction fragment length polymorphism reveal the East African Indian family to be the major clade in the south India, whereas the Central Asian family (CAS) predominates in the northern part of India,^{20,21} and the largest clade among the CAS belongs to the CAS1-Delhi lineage, which possesses variable *IS6110* copy numbers. *IS6110*, which belongs to a family of IS of the IS3 category has been widely employed in PCR tests due to the presence of its multiple copies in the *M. tuberculosis* complex genome, which is believed to confer higher sensitivity.²² However, in recent years, several clinical investigations raised concerns over *IS6110* specificity as well as sensitivity in the diagnosis of TB owing to false-positive (due to homology with other target DNA besides *M. tuberculosis*) or false-negative (due to a single copy or absence of copies of *IS6110*) results, particularly from Southeast Asian countries,^{2,6} thereby suggesting the use of other potential gene targets for monoplex or M-PCR assays. Our PCR results with individual *IS6110* and *mpb64* (Table 3) are similar to previ-

Table 4. Sensitivity and Specificity of M-PCR

Type	Specimens	Results	M-PCR (<i>mpb64</i> + <i>IS6110</i>)	Both <i>IS6110</i> & <i>mpb64</i>	Only <i>IS6110</i>	Only <i>mpb64</i>	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	ACC (%)
Group I	Confirmed EPTB cases n=25	+	24	17	3	4	96		100	98.03	98.66
		-	1								
	Suspected EPTB cases n=80	+	71	50	9	12	88.75		100	84.74	93.07
-		9									
Total EPTB cases n=105	+	95	67	12	16	90.47		100	83.33	93.54	
	-	10									
Group II Negative control n=50		+	0	0	0	0		100			
		-	50								

n, number of specimens; M-PCR, multiplex-polymerase chain reaction; +, PCR positive cases; -, PCR negative cases; PPV, positive predictive value; NPV, negative predictive value; ACC, accuracy; EPTB, extrapulmonary tuberculosis; Total EPTB cases, confirmed+clinically suspected cases.

Table 5. Sensitivity of M-PCR in Individual EPTB Specimens

Groups	Type of EPTB specimens	Results	M-PCR (<i>mpb64</i> + <i>IS6110</i>)	Both <i>IS6110</i> & <i>mpb64</i>	Only <i>IS6110</i>	Only <i>mpb64</i>	Sensitivity (%)
Confirmed EPTB cases (n=25)	Pleural fluids (21)	+	20	13	3	4	95.23
		-	1				
	Lymph node aspirates (2)	+	2	2			100
		-					
	Pleural biopsies (2)	+	2	2			100
		-					
Suspected EPTB cases (n=80)	Pleural fluids (43)	+	41	25	7	9	95.34
		-	2				
	Ascitic fluids (6)	+	5	3	1	1	83.33
		-	1				
	Pus (20)	+	18	15	1	2	90
		-	2				
	Urine (10)	+	6	4			60
		-	4				
	Synovial fluid (1)	+	1	1			100
		-					

n, number of specimens; +, PCR positive cases; -, PCR negative cases; M-PCR, multiplex-polymerase chain reaction; EPTB, extrapulmonary tuberculosis.




Fig. 2. M-PCR: amplification of 163 bp region of *mpb64* gene and 258 bp region of *IS6110* of *M. tuberculosis* H37RvDNA in the same tube with different ratios of primers. L1 represents 100 bp molecular marker; L2, L3, L4, and L5 represent *mpb64* and *IS6110* primer concentrations (μ M) in ratios of 0.2:0.2, 0.2:0.4, 0.4:0.4, and 0.4:0.2 with *M. tuberculosis* H37Rv DNA; L6, negative control (no template DNA). M-PCR, multiplex-polymerase chain reaction.

ous studies^{10,16} yet differ from the other studies^{23,24} reporting a sensitivity of <50%. Balne, et al.²⁵ recently reported a very low sensitivity of 9% only with *IS6110*-based PCR in ocular TB specimens, whereas 70.2% and 40% sensitivities were observed with individual *mpb64* and *38kDa*, respectively. The reason for low sensitivity values with *IS6110* in many studies could be due to south Indian isolates, 40% of which are shown to possess zero or low copy numbers of *IS6110*, while only 10–15% of North Indian isolates are shown to possess zero or low copy numbers of *IS6110*.^{2,22} Other reasons for low sensitivity with *IS6110* or other targets could also relate to insufficient lysis of bacterial cells, loss of DNA during purification, or the presence of PCR inhibitors in the EPTB samples.

The *fbpB* gene (Rv1886c), encoding 30 kDa protein, was included in these assays, as this protein constitutes up to 41% of total mycobacterial proteins in log-phase culture supernatants. The mRNA coding for 30 kDa protein has been utilized as a target for reverse transcriptase-PCR²⁶ in order to diagnose EPTB, and it can also detect viable mycobacteria; however, it is cumbersome to work with mRNA routinely. We performed PCR targeting the *30kDa* gene (Table 3), which revealed a slightly lower sensitivity (70%) than earlier observations (87.5% sensitivity) made by Kidane, et al.²⁷ when diagnosing TB lymphadenitis. The PCR results with the individual *hsp65* and *38kDa* gene targets (65.7–69.5% sensitivity) (Table 3) are almost similar to previous studies with EPTB specimens, such as pleural fluids, pus, and skin or abdomen biopsies.^{9,26,28} When performed individually, our combined *IS6110* PCR and *devR* PCR tests showed a sensitivity of 80.9% and a specificity of 88% in 105 EPTB specimens (Table 3), which is comparable to such combined PCR tests reported by Chakravorty, et al.⁶ in pleural fluids and lymph




Fig. 3. M-PCR: amplification of 163 bp region of *mpb64* gene and 258 bp region of *IS6110* in the same tube; L1 and L11 represent 100 bp molecular marker; L2, positive control (*M. tuberculosis* H37Rv DNA); L3, negative control (no template DNA); L4, negative control (only PCR grade water); L5–8, representative positive clinical EPTB samples; L9–10, representative negative EPTB samples. M-PCR, multiplex-polymerase chain reaction; EPTB, extrapulmonary tuberculosis.

node biopsies using the universal sample processing method. However, M-PCR in a single tube has advantages, as it reduces errors of cross contamination as well as cost and increases the sensitivity of the test.

The simultaneous identification and differentiation of *M. tuberculosis* and *M. avium* complexes and the non-tuberculous mycobacteria directly from PTB and EPTB specimens have been documented for an M-PCR using genus-specific primers targeting *hsp65*, *esat6/cfp10*, and *M. avium* complex-specific primer pairs targeting 16S-23S rDNA internal transcribed spacer-1 sequences.^{29,30} Similarly, Kim, et al.³¹ devised an M-PCR for the identification of *M. tuberculosis* complex to the species level in 37 strains and 178 clinical isolates targeting *rpoB* as well as RD1 and RD8 sequences. While performing PCR with the individual RD1 *esat-6* and *cfp10* gene targets for the diagnosis of EPTB specimens, lesser sensitivities (57–59%) (Table 3) were observed as compared to *mpb64* or *IS6110*, likely due to the genetic variability of RD1 genes or low copies of ESAT-6 gene clusters in the genomes of certain *M. tuberculosis* isolates.^{32,33} We recently detected RD1 and RD2 proteins (ESAT-6, MPB-64, etc.) from both PTB and EPTB specimens using a novel ultra-sensitive immuno-PCR assay (PCR amplified immunoassay) with a good diagnostic efficacy,³⁴ though it needs to be better optimized for routine use.

Similar to our studies, Dubey, et al.³⁵ used *IS6110+mpb64* in designing an M-nested PCR for the detection of *M. tuberculosis* directly from the blood samples of PTB as well as EPTB patients, using a Blacklight Card Disc with high sensitivity (95.7%) and specificity (100%). However, Elnifro, et al.³⁶ earlier suggested avoiding the use of nested primers for the M-PCR assay as this could lead to false positive results. Other researchers developed M-PCR using an *IS6110+38kDa*, *IS6110+devR* or *IS6110+IS*-like element *B9*, for the diagnosis of PTB and EPTB.^{37–39} The combination of three gene targets (*IS6110+dnaJ+hsp65* or *IS6110+16S rRNA+devR*) for designing an M-PCR has also been document-

ed as a diagnosis tool for EPTB.^{9,40} However, we were unable to further raise the sensitivity of M-PCR using three gene targets, such as *IS6110+mpb64+38kDa* or *IS6110+mpb64+hsp65* (data not shown), as our M-PCR exhibited substantially high sensitivity and specificity with two gene targets.

In conclusion, this study demonstrates the superiority of *mpb64*, followed by *IS6110*, among the several gene targets tested and further indicates the effectiveness of *mpb64+IS6110* in the design of an efficient M-PCR test that facilitates an early diagnosis of smear-negative and culture-negative paucibacillary EPTB specimens, which are difficult to diagnose with the available standard methods. To the best of our knowledge, this is the first report to evaluate M-PCR with *mpb64+IS6110* for the efficient diagnosis of TB pleuritis. Presently, we are validating these results in a large number of samples from varied clinical EPTB types in order for this simple and cost-effective test to be included in the diagnostic panel on a routine basis, which may prove to be a better alternative than the *rpoB*-based Xpert assay in resource-poor settings. This test might perform well in varied clinical EPTB types from various countries that harbor *M. tuberculosis* isolates with high copy numbers of *IS6110* and also countries that harbor *M. tuberculosis* isolates with zero or low copy numbers of *IS6110* (e.g., South Indian isolates), as most can be detected by *mpb64*.

ACKNOWLEDGEMENTS

The financial assistance provided by the University Grant Commission, New Delhi, the Department of Biotechnology (Builder-Programme), New Delhi and the Department of Science and Technology, New Delhi to carry out this work is duly acknowledged. We sincerely thank Prof. Jaya Tyagi, AIIMS, New Delhi for providing *M. tuberculosis* H37Rv DNA and Dr. Sunil Kumar, RBIPMT, Delhi for collecting EPTB samples.

REFERENCES

- World health organization. Global tuberculosis report. Geneva, Switzerland: World Health Organization; 2014.
- Mehta PK, Raj A, Singh N, Khuller GK. Diagnosis of extrapulmonary tuberculosis by PCR. *FEMS Immunol Med Microbiol* 2012;66:20-36.
- RNTCP 2014 TB India. Revised national TB control program annual status report. New Delhi: Central TB Division; 2014.
- Mehta PK, Kalra M, Khuller GK, Behera D, Verma I. Development of an ultrasensitive polymerase chain reaction-amplified immunoassay based on Mycobacterial RD antigens: implications for the serodiagnosis of tuberculosis. *Diagn Microbiol Infect Dis* 2012;72:166-74.
- Makeshkumar V, Madhavan R, Narayanan S. Polymerase chain reaction targeting insertion sequence for the diagnosis of extrapulmonary tuberculosis. *Indian J Med Res* 2014;139:161-6.
- Chakravorty S, Sen MK, Tyagi JS. Diagnosis of extrapulmonary tuberculosis by smear, culture, and PCR using universal sample processing technology. *J Clin Microbiol* 2005;43:4357-62.
- Haldar S, Bose M, Chakrabarti P, Dagainawala HF, Harinath BC, Kashyap RS, et al. Improved laboratory diagnosis of tuberculosis—the Indian experience. *Tuberculosis (Edinb)* 2011;91:414-26.
- Negi SS, Anand R, Basir SF, Pasha ST, Gupta S, Khare S, et al. Protein antigen b (Pab) based PCR test in diagnosis of pulmonary and extra-pulmonary tuberculosis. *Indian J Med Res* 2006;124:81-8.
- Bandyopadhyay D, Gupta S, Banerjee S, Gupta S, Ray D, Bhat-tacharya S, et al. Adenosine deaminase estimation and multiplex polymerase chain reaction in diagnosis of extra-pulmonary tuberculosis. *Int J Tuberc Lung Dis* 2008;12:1203-8.
- Sharma K, Sinha SK, Sharma A, Nada R, Prasad KK, Goyal K, et al. Multiplex PCR for rapid diagnosis of gastrointestinal tuberculosis. *J Glob Infect Dis* 2013;5:49-53.
- Sharma K, Gupta N, Sharma A, Singh G, Gupta PK, Rajwanshi A, et al. Multiplex polymerase chain reaction using insertion sequence 6110 (IS6110) and Mycobacterial protein fraction from BCG of Rm 0.64 in electrophoresis target genes for diagnosis of tuberculous lymphadenitis. *Indian J Med Microbiol* 2013;31:24-8.
- Tortoli E, Russo C, Piersimoni C, Mazzola E, Dal Monte P, Pascarella M, et al. Clinical validation of Xpert MTB/RIF for the diagnosis of extrapulmonary tuberculosis. *Eur Respir J* 2012;40:442-7.
- van Helden PD, Victor TC, Warren RM, van Helden EG. Isolation of DNA from Mycobacterium tuberculosis. *Methods Mol Med* 2001;54:19-30.
- Srivastava R, Kumar D, Waskar MN, Sharma M, Katoch VM, Srivastava BS. Identification of a repetitive sequence belonging to a PPE gene of Mycobacterium tuberculosis and its use in diagnosis of tuberculosis. *J Med Microbiol* 2006;55(Pt 8):1071-7.
- Pai M, Flores LL, Hubbard A, Riley LW, Colford JM Jr. Nucleic acid amplification tests in the diagnosis of tuberculous pleuritis: a systematic review and meta-analysis. *BMC Infect Dis* 2004;4:6.
- Vadwai V, Shetty A, Rodrigues C. Using likelihood ratios to estimate diagnostic accuracy of a novel multiplex nested PCR in extrapulmonary tuberculosis. *Int J Tuberc Lung Dis* 2012;16:240-7.
- Denkinger CM, Schumacher SG, Boehme CC, Dendukuri N, Pai M, Steingart KR. Xpert MTB/RIF assay for the diagnosis of extrapulmonary tuberculosis: a systematic review and meta-analysis. *Eur Respir J* 2014;44:435-46.
- Sharma K, Sharma A, Sharma SK, Sen RK, Dhillon MS, Sharma M. Does multiplex polymerase chain reaction increase the diagnostic percentage in osteoarticular tuberculosis? A prospective evaluation of 80 cases. *Int Orthop* 2012;36:255-9.
- Rebollo MJ, San Juan Garrido R, Folguedra D, Palenque E, Díaz-Pedroche C, Lumbleras C, et al. Blood and urine samples as useful sources for the direct detection of tuberculosis by polymerase chain reaction. *Diagn Microbiol Infect Dis* 2006;56:141-6.
- Varma-Basil M, Kumar S, Arora J, Angrup A, Zozio T, Banavaliker JN, et al. Comparison of spoligotyping, Mycobacterial interspersed repetitive units typing and IS6110-RFLP in a study of genotypic diversity of Mycobacterium tuberculosis in Delhi, North India. *Mem Inst Oswaldo Cruz* 2011;106:524-35.
- Gupta A, Kulkarni S, Rastogi N, Anupurba S. A study of Mycobacterium tuberculosis genotypic diversity & drug resistance mutations in Varanasi, North India. *Indian J Med Res* 2014;139:892-902.
- Sankar S, Kuppanan S, Balakrishnan B, Nandagopal B. Analysis of sequence diversity among IS6110 sequence of Mycobacterium tuberculosis: possible implications for PCR based detection. *Bioinformation* 2011;6:283-5.
- Jin XJ, Kim JM, Kim HK, Kim L, Choi SJ, Park IS, et al. Histopathology and TB-PCR kit analysis in differentiating the diagnosis of intestinal tuberculosis and Crohn's disease. *World J Gastroenterol* 2010;16:2496-503.
- Rana T, Singh UB, Kulshrestha V, Kaushik A, Porwal C, Agarwal N, et al. Utility of reverse transcriptase PCR and DNA-PCR in the di-

- agnosis of female genital tuberculosis. *J Med Microbiol* 2011;60(Pt 4):486-91.
25. Balne PK, Modi RR, Choudhury N, Mohan N, Barik MR, Padhi TR, et al. Factors influencing polymerase chain reaction outcomes in patients with clinically suspected ocular tuberculosis. *J Ophthalmic Inflamm Infect* 2014;4:10.
 26. Negi SS, Anand R, Pasha ST, Gupta S, Basir SF, Khare S, et al. Diagnostic potential of IS6110, 38kDa, 65kDa and 85B sequence-based polymerase chain reaction in the diagnosis of *Mycobacterium tuberculosis* in clinical samples. *Indian J Med Microbiol* 2007; 25:43-9.
 27. Kidane D, Olobo JO, Habte A, Negesse Y, Aseffa A, Abate G, et al. Identification of the causative organism of tuberculous lymphadenitis in ethiopia by PCR. *J Clin Microbiol* 2002;40:4230-4.
 28. Kulkarni S, Vyas S, Supe A, Kadival G. Use of polymerase chain reaction in the diagnosis of abdominal tuberculosis. *J Gastroenterol Hepatol* 2006;21:819-23.
 29. Gopinath K, Singh S. Multiplex PCR assay for simultaneous detection and differentiation of *Mycobacterium tuberculosis*, *Mycobacterium avium* complexes and other *Mycobacterial* species directly from clinical specimens. *J Appl Microbiol* 2009;107:425-35.
 30. Chia JH, Wu TL, Su LH, Kuo AJ, Lai HC. Direct identification of *Mycobacteria* from smear-positive sputum samples using an improved multiplex polymerase chain reaction assay. *Diagn Microbiol Infect Dis* 2012;72:340-9.
 31. Kim Y, Choi Y, Jeon BY, Jin H, Cho SN, Lee H. A simple and efficient multiplex PCR assay for the identification of *Mycobacterium* genus and *Mycobacterium tuberculosis* complex to the species level. *Yonsei Med J* 2013;54:1220-6.
 32. Gey Van Pittius NC, Gamielidien J, Hide W, Brown GD, Siezen RJ, Beyers AD. The ESAT-6 gene cluster of *Mycobacterium tuberculosis* and other high G+C gram-positive bacteria. *Genome Biol* 2001; 2:RESEARCH0044.
 33. Uplekar S, Heym B, Friocourt V, Rougemont J, Cole ST. Comparative genomics of *esx* genes from clinical isolates of *Mycobacterium tuberculosis* provides evidence for gene conversion and epitope variation. *Infect Immun* 2011;79:4042-9.
 34. Mehta PK, Raj A, Singh NP, Khuller GK. Detection of potential microbial antigens by immuno-PCR (PCR-amplified immunoassay). *J Med Microbiol* 2014;63(Pt 5):627-41.
 35. Dubey A, Gwal R, Agrawal S. *Mycobacterium tuberculosis* detection in blood using multiplex nested polymerase chain reaction. *Int J Tuberc Lung Dis* 2013;17:1341-5.
 36. Elnifro EM, Ashshi AM, Cooper RJ, Klapper PE. Multiplex PCR: optimization and application in diagnostic virology. *Clin Microbiol Rev* 2000;13:559-70.
 37. Kulkarni S, Singh P, Memon A, Nataraj G, Kanade S, Kelkar R, et al. An in-house multiplex PCR test for the detection of *Mycobacterium tuberculosis*, its validation & comparison with a single target TB-PCR kit. *Indian J Med Res* 2012;135:788-94.
 38. Sharma SK, Sethi S, Sharma M, Meharwal SK, Katoch VM, Jindal SK, et al. Development and evaluation of a multiplex polymerase chain reaction for the detection of *Mycobacterium tuberculosis* from pulmonary specimens. *Scand J Infect Dis* 2012;44:739-44.
 39. Tang TH, Ahmed SA, Musa M, Zainuddin ZF. Rapid detection of *Mycobacterium tuberculosis* in clinical samples by multiplex polymerase chain reaction (mPCR). *World J Microbiol Biotechnol* 2013; 29:2389-95.
 40. Hallur V, Sharma M, Sethi S, Sharma K, Mewara A, Dhatwalia S, et al. Development and evaluation of multiplex PCR in rapid diagnosis of abdominal tuberculosis. *Diagn Microbiol Infect Dis* 2013; 76:51-5.