


# Evaluation of carbohydrate counting knowledge among individuals with type 1 diabetes mellitus in Saudi Arabia: a cross-sectional study

Nahla Mohammed Bawazeer <sup>1</sup>, Leena Hamdan Alshehri,<sup>1</sup>  
Nouf Mohammed Alharbi,<sup>1</sup> Noha Abdulaziz Alhazmi,<sup>1</sup> Alhanouf Fahad Alrubaysh,<sup>1</sup>  
Alia Riad Alkasser,<sup>1</sup> Khaled Hani Aburishah<sup>2</sup>

**To cite:** Bawazeer NM, Alshehri LH, Alharbi NM, *et al*. Evaluation of carbohydrate counting knowledge among individuals with type 1 diabetes mellitus in Saudi Arabia: a cross-sectional study. *BMJ Nutrition, Prevention & Health* 2022;5:e000553. doi:10.1136/bmjnph-2022-000553

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjnph-2022-000553>).

<sup>1</sup>Clinical Nutrition Program, Department of Health Sciences, College of Health and Rehabilitation Sciences, Princess Nourah Bint Abdul Rahman University, Riyadh, Saudi Arabia

<sup>2</sup>University Diabetes Center, King Saud University Medical City, King Saud University College of Medicine, Riyadh, Saudi Arabia

## Correspondence to

Dr Nahla Mohammed Bawazeer; [NMBawazeer@pnu.edu.sa](mailto:NMBawazeer@pnu.edu.sa)

Received 12 September 2022  
Accepted 17 November 2022  
Published Online First  
28 November 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

## ABSTRACT

**Introduction** Carbohydrate counting (CC) is an important nutritional strategy to improve glycaemic outcomes among patients with diabetes. Few studies have investigated CC knowledge among individuals with type 1 diabetes mellitus (T1DM) in Saudi Arabia. Therefore, we aimed to evaluate CC knowledge in Saudi adults with T1DM.

**Study design and methods** A cross-sectional study was conducted between December 2021 and February 2022, including 224 patients with T1DM from the University Diabetes Center, Riyadh. Adults aged  $\geq 18$  years, diagnosed with T1DM for  $>1$  year, and residing in Saudi Arabia were included. CC knowledge was assessed using a previously well-studied tool (AdultCarbQuiz), which was translated into Arabic and tested for validity by a group of dietitians. Descriptive statistics were used for data analysis, and bivariate and regression analyses were conducted.

**Results** The AdultCarbQuiz questionnaire-Arabic version had good validity and reliability (Cronbach's  $\alpha$ : 0.87). The CC method was used by 54% of the participants. The mean CC knowledge score was  $23.01 \pm 7.31$ . A significant negative linear relationship between the participants' CC knowledge scores, and age and glycated haemoglobin (HbA1c) levels, was revealed by simple regression analysis. Furthermore, significant independent variables related to CC knowledge scores were CC use, HbA1c levels, being taught about CC ( $>5$  times), insulin pump usage and DM duration ( $\leq 15$  years).

**Conclusions** Approximately half of the patients used the CC method. The mean CC knowledge scores were better in patients who used the CC method, were more frequently taught about CC, were treated using an insulin pump, and had a shorter DM duration than their counterparts. Therefore, designing and implementing a well-structured nutrition education programme tailored to individuals with diabetes is crucial to provide them with up-to-date dietary information, as well as the necessary knowledge and skills, to improve their outcomes and manage their condition.

## INTRODUCTION

Type 1 diabetes mellitus (T1DM) is an autoimmune disease with hyperglycaemia due to insulin deficiency resulting from the loss of

## WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Carbohydrate counting (CC) is a meal planning strategy that aids in tracking carbohydrate intake and prandial glycaemia and facilitates flexible food selection without adversely affecting the metabolic outcomes of patients. Previous studies on the efficacy and safety of the CC method in patients with type 1 diabetes mellitus (T1DM) reported a significant reduction in glycated haemoglobin (HbA1c) levels.
- ⇒ CC is a complex skill that is subject to error among children and young adults with T1DM, as well as their families, and efforts are therefore needed to develop skills and achieve accurate CC by the patients and glycaemic management.

## WHAT THIS STUDY ADDS

- ⇒ The current study is among the first to evaluate CC knowledge in adults with T1DM in Saudi Arabia. The Arabic version of the AdultCarbQuiz questionnaire showed good reliability and validity in evaluating the CC knowledge of Saudi adults with T1DM, and approximately half of the participants intended to use the CC method for diabetes management.
- ⇒ The CC knowledge scores were significantly higher among participants who used the CC method, were taught about CC  $\geq 5$  times, used an insulin pump, had lower HbA1c levels and had a DM duration  $\leq 15$  years.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ A well-structured nutrition education programme is needed to provide patients with up-to-date information, as well as the necessary knowledge and skills, to improve their outcomes and manage their condition. A prospective and/or randomised controlled study, using a similar questionnaire, is recommended to assess CC knowledge before and after the nutrition education programme.

pancreatic islet  $\beta$ -cells.<sup>1 2</sup> Approximately 10% of all diabetes mellitus (DM) cases are T1DM, which can affect individuals of all ages but

usually develops in children or young adults. The causes of T1DM remain unclear but are linked to genetic and environmental factors.<sup>1</sup> Over the past 30 years, the incidence of T1DM has increased in Saudi Arabia (SA).<sup>3</sup> In 2017, more than 35 000 children and adolescents in SA had T1DM, with the highest incidence of 3900 cases.<sup>3,4</sup>

Poor glycaemic management among patients with T1DM can lead to acute hypoglycaemia and long-term macrovascular and microvascular complications.<sup>5</sup> Nutrition therapy is associated with improved glycaemic outcomes.<sup>6,7</sup> One of the nutritional approaches to T1DM emphasises the connection between the number of consumed carbohydrates and insulin dosage. Carbohydrate counting (CC) is a meal planning strategy that aids in tracking carbohydrate intake and prandial glycaemia<sup>8</sup> and facilitates flexible food selection without adversely affecting the metabolic outcomes of patients.<sup>9</sup> It is becoming a more widely accepted method after being used as one of four meal-planning approaches with intensive insulin therapy in the Diabetes Control and Complications Trial in the early 1990s.<sup>10</sup> Consequently, many diabetes centres in SA currently consider intensive insulin therapy part of the routine management of T1DM.<sup>11,12</sup> Therefore, nutrition education programmes were tailored to train dietitians in the use of CC as an education modality for patients with diabetes.

Previous studies have evaluated the efficacy and safety of the CC method in patients with T1DM.<sup>8,13–15</sup> Scavone *et al* reported a significant reduction in glycated haemoglobin (HbA1c) levels, fewer hypoglycaemic events and decreased rapid insulin analogue doses after participants were taught about CC.<sup>8</sup> A meta-analysis of randomised controlled trials showed evidence favouring the use of the CC method and a significant reduction in HbA1c levels compared with that of controls.<sup>13</sup> However, CC is a complex skill that is subject to error among children and young adults with T1DM and their families.<sup>16</sup> Therefore, identifying opportunities to develop CC skills can improve interventions and achieve accurate CC by patients and improved glycaemic management.<sup>16</sup>

Few studies have investigated CC knowledge among individuals with T1DM in SA. A pilot study conducted with 94 Emirati and Omani adults with diabetes showed low scores for knowledge of the carbohydrate content of foods.<sup>17</sup> In addition, a cross-sectional study involving 178 Saudi individuals with DM reported a low level of knowledge about carbohydrate foods.<sup>18</sup> Therefore, this study aimed to evaluate CC knowledge among Saudi adults and to assess the associations between CC knowledge and HbA1c levels and other factors, such as educational level, DM duration, type of insulin therapy, regular follow-up with a dietitian and learning and practising the CC method, because these factors may affect carbohydrate estimation knowledge and accuracy.<sup>18–20</sup> We hypothesised that patients with good CC knowledge would have a higher educational level, receive insulin pump therapy, use the CC method and have lower HbA1c levels. An extensive review was conducted to identify an easy, quick

and previously tested tool that could assess CC knowledge levels. According to the literature, the AdultCarbQuiz is a self-administered questionnaire with good validity and reliability that was developed to aid healthcare practitioners in rapidly assessing CC knowledge among adults with T1DM.<sup>21</sup> This tool was translated into Arabic and reviewed by expert dietitians for information adequacy and validity prior to its application in this study.<sup>22</sup>

## METHODS

### Study design and participants

A cross-sectional study was conducted with adults with T1DM who were recruited during routine clinical visits to the University Diabetes Center (UDC) in King Saud University Medical City (Riyadh, SA) between December 2021 and February 2022. This study included adult participants aged  $\geq 18$  years who were diagnosed with T1DM for  $>1$  year. Patients who could not read or understand Arabic and those with cognitive impairment were excluded.

### Sample size estimation and sampling method

The sample size was estimated as follows: approximately 1000 patients from the UDC had T1DM, and approximately 25% of these patients used the CC method as a meal plan to control DM. The following sample size equation was used:  $(n) = Z^2 pq/d^2$  where the Z value for the 95% CI was 1.96, p was the prevalence of CC method usage (25%), q=1 – p=75%, and d was the margin of error (0.06 (6%)). The required sample size was 200. Furthermore, considering a 15% non-response rate, the target sample size was 230. Six patients with incomplete responses were excluded; therefore, the study included 224 patients. Participants were selected by convenience sampling.

### Research instrument

The study questionnaire included (1) sociodemographic information, such as age, sex, educational level and marital status; (2) disease information, such as DM duration, last HbA1c level, type of insulin therapy and DM complications or other diseases and (3) nutrition education questions, such as CC method usage, reasons for using CC, frequency of visits to a dietitian in the past 2 years, and the number of times wherein CC was taught. HbA1c levels were obtained from the patients' records for the last 6 months or less.

In this study, a validated AdultCarbQuiz questionnaire was used to evaluate CC knowledge,<sup>22</sup> with permission from the author. The questionnaire was translated into Arabic and reviewed by Arabic language experts. A panel of nutrition experts, including clinical diabetes dietitian experts in CC and academic nutrition faculties, reviewed the translated questionnaire for information adequacy and validity using a scoring sheet that listed each question and required the experts to score each item out of 10 and add further comments where necessary. Questions

that scored  $\leq 5$  were reviewed and modified following the comments, which were mostly associated with linguistic issues. Subsequently, the questionnaire was verified for readability with 20 patients. The participants in this pilot study confirmed that the questionnaire's instructions, layout, length, simplicity in completion and time for completion were appropriate, with no reported issues in understanding the questions.

The AdultCarbQuiz questionnaire comprises 43 items in 6 domains: identifying carbohydrates in commonly consumed foods (19 items), ability to count the carbohydrate content in typical portions of simple foods (6 items), ability to read a nutrition label for carbohydrate content (4 items), understanding the glycaemic targets (4 items), knowledge on hypoglycaemia prevention and treatment using carbohydrate foods (5 items) and ability to sum up the carbohydrate content of a meal (4 items). Scores were 1, 0 and 43 for a correct response, incorrect response and overall score, respectively. The 'don't know' answers were scored as incorrect.

### Data analysis

Data were analysed using IBM SPSS for Windows V.26.0 (IBM). Descriptive statistics (frequencies, percentages, means and SD) were used to define categorical and quantitative variables. Bivariate analysis was performed using Student's t-test for independent samples and one-way analysis of variance followed by Tukey's test to compare the mean CC knowledge scores of categorical variables with two or more categories. Simple and multiple regression analyses were performed to measure the linear relationships between quantitative dependent variables (CC knowledge scores) and a set of independent categorical and quantitative variables. For categorical independent variables with more than two categories, (k-1) dummy variables were included in the model. Regression coefficients of the model were used to assess how changes in each independent variable affected the CC knowledge scores. In addition, tolerance and variance inflation factor (VIF) criteria were used to evaluate the model's multicollinearity of independent variables. The coefficient of variation ( $R^2$ ) value was used to quantify the change in CC knowledge scores that were explained by the significant independent variables in the model. The reliability of the tool's Arabic version was assessed by split-half reliability using odd-numbered and even-numbered items. The Spearman-Brown correction was applied to correct the reliability coefficient. Internal consistency was used for each of the domains and all the instrument items, and validity was tested by comparing the mean CC knowledge scores regarding the three independent variables. A  $p < 0.05$  indicated statistical significance.

## RESULTS

### Sociodemographic and clinical characteristics

Of the 224 patients with T1DM, 40.6% were men and 68.8% were single. The mean patient age was  $28.2 \pm 7.8$

years. More than 60% had a graduate degree, and DM duration was relatively evenly distributed, with 48.7% and 51.3% having  $\leq 15$  and  $> 15$  years of DM, respectively. The mean HbA1c level was  $8.3\% \pm 1.4\%$ , with 61.2% showing no associated complications of DM. Notably, most patients (88.4%) used multiple daily insulin injections, and CC was prevalent in 54%. Only 6.3% of the participants had not visited a dietitian in the past 2 years, while 27.2% had visited a dietitian  $> 5$  times. Responses to the number of times wherein CC was taught were  $> 5$ , 3–5 and 1–2 times in 23.7%, 23.2% and 49.6% of the participants, respectively (table 1).

### CC knowledge

The CC knowledge questionnaire showed that the frequency of correct responses for food containing carbohydrates was the highest for bread, rice, spaghetti and baked potato, whereas cheese and butter had the lowest correct responses. The highest and lowest percentages of correct responses concerning the grams of carbohydrates present in each serving of food were for a cup of milk and a cup of cooked rice, respectively. Moreover, 67% and 61.2% of the participants could identify the serving size, and grams of carbohydrates in one serving, from the nutrition label, respectively (online supplemental table 1).

The highest and lowest frequencies of correct responses were provided for the questions related to '2 hours post-prandial blood glucose level' and 'one carb choice will raise your blood glucose level by how many points?', respectively. Lastly, regarding a meal's carbohydrate content, 25% of the participants identified the grams of carbohydrates in the provided breakfast example, followed by the snack and supper; however, only 4% identified the grams of carbohydrates in the lunch example (online supplemental table 1).

### Relationship between CC knowledge and related variables

The responses to the 43 items of the CC knowledge questionnaire were converted into scores, and the mean values were compared across the participants' demographic and clinical characteristics. The bivariate analysis showed significant differences in the mean CC knowledge scores across educational levels, DM duration, type of insulin therapy, CC use, frequency of visits to a dietitian in the past 2 years, and the number of times wherein CC was taught. The mean CC knowledge scores were significantly higher in participants with postgraduate and graduate qualifications than in those with high school qualifications or lower but were similar between participants with  $\leq 15$  and  $> 15$  years of DM. The scores were significantly higher in participants who used an insulin pump than in those who used multiple daily insulin injections. In addition, the participants who used the CC method had significantly higher mean CC knowledge scores than those who had never used CC. The mean CC knowledge scores were significantly higher in patients who had visited a dietitian  $> 5$  and 3–5 times than in those who had visited a dietitian

**Table 1** Distribution of sociodemographic and clinical characteristics of patients with T1DM (n=224)

Characteristics	No	(%)
Age (years)*	28.2*	(7.8)
Sex, male	91	(40.6)
Marital status		
Single	154	(68.8)
Married	60	(26.8)
Divorce	10	(4.5)
Educational level		
High school or lower	51	(22.8)
Graduate	149	(66.5)
Postgraduate	24	(10.7)
DM duration (years)		
≤15	109	(48.7)
>15	115	(51.3)
HbA1c levels (%)*	8.3*	(1.4)
DM complications		
No disease	137	(61.2)
Hypertension	22	(9.8)
Dyslipidaemia	42	(18.7)
Diabetic retinopathy	17	(7.6)
Diabetic foot	2	(0.9)
Diabetic nephropathy	13	(5.8)
Diabetic neuropathy	6	(2.7)
Other diseases	42	(18.8)
Type of intensive insulin therapy		
Multiple daily insulin injections	198	(88.4)
Insulin pump	26	(11.6)
CC use		
Yes	121	(54.0)
No	103	(46.0)
Frequency of visits to a dietitian in the past 2 years		
Has not visited a dietitian	14	(6.3)
1–2 times	93	(41.5)
3–5 times	56	(25.0)
>5 times	61	(27.2)
No of times taught about CC		
Never	8	(3.6)
1–2 times	111	(49.6)
3–5 times	52	(23.2)
>5 times	53	(23.7)

\*Mean (SD).  
CC, carbohydrate counting; DM, diabetes mellitus; HbA1c, glycated haemoglobin; T1DM, type 1 diabetes mellitus.

1–2 times and those who had never visited. Similar results were observed among participants who had been taught about CC >5 and 3–5 times, compared with those who had

**Table 2** Comparison of the mean CC knowledge scores with the demographic and clinical characteristics of the patients with T1DM

Characteristics	Mean	(SD)	t-value/ F-value	P value
Sex				
Male	23.02	(7.0)	0.015	0.988
Female	23.01	(7.5)		
Marital status				
Single	23.51	(7.0)	2.457	0.088
Married	22.48	(7.7)		
Divorce	18.50	(8.0)		
Educational level				
High school or lower	20.53	(8.2)	4.091	0.018
Graduate	23.62	(7.0)		
Postgraduate	24.54	(5.8)		
DM duration (years)				
≤15	24.36	(7.2)	2.719	0.007
>15	21.74	(7.2)		
Type of insulin therapy				
Multiple daily insulin injections	22.24	(7.3)	–4.578	< 0.0001
Insulin pump	28.92	(3.9)		
CC use				
Yes	25.98	(6.8)	7.332	< 0.0001
No	19.52	(6.3)		
Frequency of visits to a dietitian in the past 2 years				
Has not visited a dietitian	20.0	(6.9)	3.356	0.020
1–2 times	21.72	(6.8)		
3–5 times	24.05	(7.7)		
>5 times	24.72	(7.3)		
No of times taught about CC				
Never taught	17.25	(4.9)	7.938	< 0.0001
1–2 times	21.50	(7.8)		
3–5 times	23.65	(5.6)		
>5 times	26.43	(6.7)		

CC, carbohydrate counting; DM, diabetes mellitus; T1DM, type 1 diabetes mellitus.

been taught 1–2 times and those who had never learnt about CC. However, there was no significant difference in the mean CC knowledge scores associated with sex and marital status (table 2).

The simple regression analysis between CC knowledge scores and participants' age showed a significant negative linear relationship. Conversely, the regression coefficient of age was –0.159, implying that for every 1-year increase in age, the participants' CC knowledge scores decreased



**Table 3** Relationship between CC knowledge scores and the age and HbA1c levels of patients with T1DM by simple regression analysis

Independent variables	Unstandardised coefficients		Standard coefficient			Collinearity Statistics		Model summary	
	B	SE	Beta	t-value	P value	Tolerance	VIF	R <sup>2</sup>	P value
Constant	27.485	1.804		15.239	<0.0001				
Age	-0.159	0.062	-0.170	-2.573	0.011	1.00	1.00	0.029	0.011
Constant	35.967	2.885		12.465	<0.0001				
HbA1c levels	-1.561	0.343	-0.292	-4.549	<0.0001	1.00	1.00	0.085	<0.0001

B, unstandardised beta (rate of change per unit time); CC, carbohydrate counting; HbA1c, glycated haemoglobin; R<sup>2</sup>, coefficient of variation value; T1DM, type 1 diabetes mellitus; VIF, variance inflation factor.

by -0.159 units on average, which was significant. The constant coefficient value of 27.485 indicated that the mean CC knowledge score when age was zero was also significant. The VIF (1.00) and tolerance values (1.00) of this model indicated no collinearity, as both values were below 4 (VIF) and above 0.25 (tolerance). The R<sup>2</sup> value of 0.029 revealed that only 2.9% of the change in CC knowledge scores was explained by the participants' age, which was significant (p=0.011) (table 3).

In addition, the participants' CC knowledge scores and HbA1c levels showed a significant negative linear relationship. The regression coefficient of -1.561 for HbA1c levels indicated that for every one-unit increase in HbA1c levels, the participants' CC knowledge scores decreased by -1.561 units on average, which was significant. Conversely, a constant coefficient value of 35.967 suggested a significant mean CC knowledge score when the HbA1c value was zero. Furthermore, the VIF (1.00) and tolerance values (1.00) of this model indicated no collinearity, as both values were below 4 (VIF) and above 0.25 (tolerance). The R<sup>2</sup> value of 0.085 indicated that approximately 8.5% of the change in CC knowledge scores could be explained by the participants' HbA1c levels, and was significant (p<0.0001) (table 3).

Multiple regression analysis between the quantitative outcome variable (CC knowledge scores) and a

set of independent variables was performed. The independent variables included in the model were: age (in years), educational levels (graduate and postgraduate), frequency of visits to a dietitian, being taught about CC, CC use (yes), HbA1c levels, type of insulin therapy (insulin pump usage) and DM duration (≤15 years).

Among the variables included in the model, the significant independent variables were CC use (yes), HbA1c levels (high), being taught about CC (>5 times), type of insulin therapy (insulin pump usage) and DM duration (≤15 years) (table 4). The regression coefficients of the three variables (CC use, being taught about CC (>5 times) and DM duration (≤15 years)) indicated a significant positive relationship with CC knowledge scores. Conversely, the regression coefficients of the other two variables (HbA1c levels and type of insulin therapy (insulin pump usage)) showed a significant negative association with CC knowledge scores. Moreover, the CC knowledge scores increased by 4.641 units, on average, in participants who used CC, compared with those who had never used CC. The CC knowledge scores decreased by -1.039 units, on average, for every 1-unit increase in HbA1c levels. Conversely, the CC knowledge scores increased by 2.555 units, on average, in participants who were taught about CC >5 times, compared with those who had never been taught about CC. CC knowledge increased

**Table 4** Independent factors related to the CC knowledge scores of patients with T1DM by multiple linear regression analysis

Independent variables	Unstandardised coefficients		Standard coefficient			Collinearity Statistics		Model summary	
	B	SE	Beta	t-value	P value	Tolerance	VIF	R <sup>2</sup>	P value
Constant	27.225	2.800		9.724	<0.0001			0.301	<0.0001
CC use (yes)	4.641	0.898	0.317	5.168	<0.0001	0.851	1.175		
HbA1c levels (high)	-1.039	0.311	-0.194	-3.338	0.001	0.945	1.058		
Being taught about CC (> 5 times)	2.555	1.014	0.149	2.518	0.013	0.917	1.090		
Type of insulin therapy (insulin pump)	3.239	1.388	0.142	2.334	0.021	0.863	1.159		
DM duration (≤15 years)	1.901	0.832	0.130	2.285	0.023	0.986	1.014		

B, unstandardised beta (rate of change per unit time); CC, carbohydrate counting; DM, diabetes mellitus; HbA1c, glycated haemoglobin; R<sup>2</sup>, coefficient of variation value; T1DM, type 1 diabetes mellitus; VIF, variance inflation factor.

**Table 5** Reliability of each of the six domains and all the items of the AdultCarbQuiz tool

Name of domain	No of items	Cronbach's $\alpha$ (95% CI)	Split-half reliability coefficient with Spearman-Brown correction
Carbohydrate food recognition	19	0.864 (0.836 to 0.888)	
Counting carbohydrates in each of the foods	7	0.899 (0.878 to 0.918)	
Interpreting nutrition labels for carbohydrates	4	0.794 (0.747 to 0.835)	
Glycaemic targets	4	0.488 (0.369 to 0.589)	
Hypoglycaemia prevention and treatment	5	0.577 (0.483 to 0.659)	
Counting carbohydrates in a meal	4	0.815 (0.772 to 0.852)	
All items	43	0.874 (0.849 to 0.896)	
Odd-numbered and even-numbered items			0.865

by 3.239 units, on average, in participants using an insulin pump compared with those using multiple daily insulin injections. For the independent variable, DM duration ( $\leq 15$  years), the CC knowledge scores increased by 1.901 units, on average, in participants with DM duration  $\leq 15$  years compared with those with DM duration  $> 15$  years. The collinearity statistics (tolerance and VIF) showed no multicollinearity, as the tolerance and VIF values were not below and above 0.25 and 4, respectively. The  $R^2$  value of 0.301 implied that the five significant independent variables in the model accounted for 30.1% of the change in the CC knowledge scores, which was significant ( $p < 0.0001$ ) (table 4).

#### Reliability and validity of the CC knowledge questionnaire

The Cronbach's  $\alpha$  values for the 6 domains and 43 items indicated an acceptable reliability of 0.874 (0.849, 0.896). The 'glycaemic target' and 'hypoglycaemia prevention and treatment' domains had lower reliabilities, of 0.488 (0.369, 0.589) and 0.577 (0.483, 0.659), respectively. In addition, with Spearman-Brown prediction correction, the split-half reliability coefficient comparing the odd-numbered to even-numbered items was 0.865, indicating good internal consistency (table 5).

Regarding the questionnaire's validity, the mean CC knowledge score of the 224 participants was 23.01 (SD: 7.31; maximum score: 38). The mean CC knowledge scores were significantly higher in participants who had visited a dietitian  $> 5$  and 3–5 times than in those who had visited a dietitian 1–2 times and those who had never visited. In addition, the mean values were significant for participants who had been taught about CC  $> 5$  and 3–5 times compared with those who had been taught 1–2 times or never taught about CC. Furthermore, the CC knowledge scores were inversely associated with HbA1c levels, wherein they decreased as the HbA1c levels increased, which was significant. These analyses demonstrated that the CC knowledge questionnaire had adequate validity.

#### DISCUSSION

This study evaluated CC knowledge in Saudi adults with T1DM. The main findings were (1) approximately half of the participants used the CC method; (2) participants had acceptable mean CC knowledge scores; and (3) the mean CC knowledge scores were significantly higher in participants who used the CC method, were taught about CC  $\geq 5$  times, used an insulin pump, had lower HbA1c levels and had DM duration  $\leq 15$  years. Furthermore, this study used a previously well-studied tool (AdultCarbQuiz) that was translated into Arabic, reviewed by expert dietitians, and demonstrated good validity and reliability.

The results showed that 54% of the participants intended to use the CC method for DM management, visited a dietitian and were frequently taught about the CC method. However, this trend contrasts with other SA studies, in which 18% of participants used the CC method<sup>18</sup> and 19.8% reported visiting a dietitian.<sup>23</sup> The reason for these differences may be that the UDC, where the study was conducted, is a long-established, specialised diabetes centre in Riyadh that provides different services and treatment options for DM management, including intensive insulin therapy with multidisciplinary teams. Furthermore, the mean knowledge score was  $23.01 \pm 7.3$ , which is consistent with the mean knowledge score of the original questionnaire ( $23.9 \pm 8.3$ ).<sup>21</sup>

Patients with T1DM should have good knowledge about foods containing carbohydrates and be able to identify the carbohydrate content in each serving size to adjust the meal insulin dose and manage glycaemia. However, accurate CC requires highly developed literacy and numeracy skills and broad nutritional knowledge to estimate portion sizes correctly, read food labels, weigh and measure foods, and determine carbohydrate content.<sup>19</sup> In this study, participants' knowledge was assessed, with correct answers recorded in the carbohydrate food recognition domain for bread, cooked rice and pasta, and baked potatoes. This positive result could be because these food items are among the most consumed in Saudi culture.<sup>24</sup> Conversely, participants could not identify butter and

cheese, possibly because they were considered dairy products containing carbohydrates, such as milk. In addition, participants could not respond correctly to the grams of carbohydrates per serving. The scale of carbohydrates was measured in grams per cup; however, some dietitians may have used different measurement scales, during the teaching process, for certain foods (such as rice) using spoons, which is easier and more practical for patients. Therefore, patients using CC may need more information on different resources to identify carbohydrate content to answer questions correctly.<sup>25</sup> Overall, 60% of the patients in this study could interpret the carbohydrate nutrition label. A previous study reported that individuals with DM and other chronic diseases were more likely to read labels regularly to control their medical condition.<sup>26 27</sup>

In addition, patients responded correctly to different glycaemic targets and hypoglycaemia prevention and treatment scenarios. This result could be attributed to the availability of a multidisciplinary team at the UDC, which helped improve their knowledge of DM management. However, 20.5% of the participants answered correctly about how many points of blood glucose would be produced after eating a carbohydrate choice. This result may be due to the participants who had never learnt about CC or had difficulty identifying carbohydrate choices. Only a few participants correctly answered about CC per meal, which could be explained by the knowledge and practice gap in teaching the CC method. This drawback should be resolved before committing the patients to CC<sup>21</sup> and could be feasible by engaging the patients in practical, real-life, teaching scenarios.<sup>19</sup> Therefore, ongoing DM education offered by qualified health professionals, particularly for patients receiving intensive insulin therapy, is necessary to achieve optimal results.<sup>14</sup>

In this study, approximately 30% of the change in the CC knowledge scores was explained by the five independent variables in the regression model ( $p < 0.0001$ ). Being taught about CC  $> 5$  times throughout patients' lives was significantly positively correlated with CC knowledge. Notably, with the use of the CC method, the knowledge scores increased by 4.641 units on average, as compared with not using CC. Therefore, patients with T1DM who intend to perform CC should have more frequent access to qualified dietitians for information on CC and clarification on important issues.<sup>20 28</sup> In addition, the CC knowledge scores decreased by  $-1.039$  units on average for every one-unit increase in HbA1c levels. Consequently, patients with higher CC knowledge scores had better glycaemic control, which is consistent with the findings of previous studies, in which HbA1c levels decreased in patients who used the CC method.<sup>8 14 29</sup> A 2014 meta-analysis showed that 24 of the 27 included studies reported a reduction of 0.2%–1.2% in HbA1c levels after commencing CC.<sup>30</sup> Furthermore, a randomised controlled trial in children with T1DM reported a significant difference in the mean HbA1c levels between the control and CC groups at follow-up, suggesting that the CC method may improve metabolic control in patients with T1DM.<sup>31</sup> CC knowledge

increased, on average, by 3.239 units in participants using an insulin pump compared with those using multiple daily dose injections. This trend is expected because patients on insulin pump therapy receive extensive education, advanced skills and frequent follow-up with the health-care team, including dietitians, to achieve strict glycaemic control while minimising hypoglycaemia risk.<sup>32</sup> Lastly, the analysis revealed that CC knowledge scores increased by 1.901 units, on average, in participants with  $\leq 15$  years, compared with those with  $> 15$  years, of DM. This finding is validated by previous studies, which reported an inverse relationship between CC and time since DM diagnosis.<sup>33 34</sup> These studies were conducted with youth and children. Contrarily, no association between DM duration and CC was found in adults<sup>20</sup> or adolescents<sup>35</sup> who received intensive insulin therapy.

The current study provides a comprehensive assessment of CC knowledge in Saudi adults with T1DM. Of note, multiple variables were included to support the scores achieved, and a valid and reliable tool suitable for assessing CC knowledge was used. The AdultCarbQuiz questionnaire's Arabic version had good internal reliability (Cronbach's  $\alpha$ : 0.87), similar to that of the original questionnaire (0.90).<sup>21</sup> However, this study was limited to adult patients recruited from a single diabetes centre. Therefore, this study should be replicated on a larger scale, which includes a multicentre scenario. Further research involving children and their parents could reveal interesting findings to design early education interventions to promote a better quality of life. The questionnaire was administered once; therefore, additional studies are warranted to establish the stability of CC knowledge. Furthermore, a prospective and/or randomised controlled study using a similar questionnaire is recommended to assess CC knowledge before and after a nutrition education programme.

In conclusion, approximately half of the participants used the CC method. Specifically, the mean CC knowledge scores were higher in participants who used the CC method, were taught about CC  $\geq 5$  times, used an insulin pump, had lower HbA1c levels and had DM duration  $\leq 15$  years.

Dietitians have an important role to play in promoting the use of CC and assessing knowledge and skills before allowing patients with T1DM to take responsibility for their own care. Therefore, designing a well-structured nutrition education programme tailored to individuals with diabetes is crucial to provide them with up-to-date dietary information, and the necessary knowledge and skills to improve their outcomes and manage their condition.

**Twitter** Nahla Mohammed Bawazeer @BawazeerNahla

**Acknowledgements** The authors thank all participants in this study and the expert professionals for their feedback on the questionnaire.

**Contributors** All the authors were involved in the study. NMB contributed to the conception and design of the study and wrote the manuscript. LHA, NMA, NAA, AFA, ARA and KHA participated in data collection and entry. NMB interpreted the data.



KHA had overall responsibility for the study. All the authors read and approved the final manuscript for publication. NMB acts as guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not applicable.

**Ethics approval** The study procedure was approved by the Institutional Review Board of King Saud University College of Medicine (IRB no.: 21/01127; approval date: 05 December 2021). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed by Dr. Nada Benajiba, Ibn Tofail University Joint Research Unit in Nutrition and Food Regional Designated Center of Nutrition AFRA/IAEA, UK of Great Britain and Northern Ireland.

**Data availability statement** All data relevant to the study are included in the article or uploaded as online supplemental information.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iD

Nahla Mohammed Bawazeer <http://orcid.org/0000-0001-5087-7802>

## REFERENCES

- International Diabetes Federation. Type 1 diabetes 2020, 2020. Available: <https://www.idf.org/aboutdiabetes/type-1-diabetes.html> [Accessed 26 June 2022].
- Katsarou A, Gudbjörnsdóttir S, Rawshani A, et al. Type 1 diabetes mellitus. *Nat Rev Dis Primers* 2017;3:17016.
- Robert AA, Al-Dawish A, Mujammami M, et al. Type 1 diabetes mellitus in Saudi Arabia: a soaring epidemic. *Int J Pediatr* 2018;2018:1–9.
- International Diabetes Federation. IDF diabetes atlas, 2017. Available: <https://www.diabetesatlas.org> [Accessed 23 Jun 2022].
- Diabetes Control and Complications Trial Research Group, Nathan DM, Genuth S, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med* 1993;329:977–86.
- Franz MJ, Boucher JL, Evert AB. Evidence-Based diabetes nutrition therapy recommendations are effective: the key is individualization. *Diabetes Metab Syndr Obes* 2014;7:65–72.
- Atik Diabetes Association. 6. Glycemic Targets: *Standards of Medical Care in Diabetes-2021*. *Diabetes Care* 2021;44:S73–84.
- Scavone G, Manto A, Pitocco D, et al. Effect of carbohydrate counting and medical nutritional therapy on glycaemic control in type 1 diabetic subjects: a pilot study. *Diabet Med* 2010;27:477–9.
- Bayram S, Kızıltan G, Akın O. Effect of adherence to carbohydrate counting on metabolic control in children and adolescents with type 1 diabetes mellitus. *Ann Pediatr Endocrinol Metab* 2020;25:156–62.
- Gillespie SJ, Kulkarni KD, Daly AE. Using carbohydrate counting in diabetes clinical practice. *J Am Diet Assoc* 1998;98:897–905.
- Al-Qahtani SM, Shati AA, Alqahtani YA, et al. Factors affecting glycaemic control among Saudi children with type 1 diabetes mellitus in Aseer region, southwestern Saudi Arabia. *Int J Environ Res Public Health* 2022;19:11558.
- Al-Saleh Y, Al Motairi F, Hassan E, et al. Insulin pump therapy issues among adults with type 1 diabetes mellitus in Saudi Arabia: a retrospective study. *Diabetes Ther* 2020;11:2993–3001.
- Builes-Montaño CE, Ortiz-Cano NA, Ramirez-Rincón A, et al. Efficacy and safety of carbohydrate counting versus other forms of dietary advice in patients with type 1 diabetes mellitus: a systematic review and meta-analysis of randomised clinical trials. *J Hum Nutr Diet* 2022;35:1030–42.
- Laurenzi A, Bolla AM, Panigoni G, et al. Effects of carbohydrate counting on glucose control and quality of life over 24 weeks in adult patients with type 1 diabetes on continuous subcutaneous insulin infusion: a randomized, prospective clinical trial (GIOCAR). *Diabetes Care* 2011;34:823–7.
- Sämann A, Mühlhauser I, Bender R, et al. Glycaemic control and severe hypoglycaemia following training in flexible, intensive insulin therapy to enable dietary freedom in people with type 1 diabetes: a prospective implementation study. *Diabetologia* 2005;48:1965–70.
- Khoiriani IN, Dhiyanti AY, Firmansyah RF, et al. The Benefits of Carbohydrate Counting in Type 1 Diabetes Mellitus: A Scoping Review. *Amerta Nutrition* 2021;5:395.
- Ali H, Bernsen R, Taleb S. Carbohydrate food knowledge of Emirati and Omani adults with diabetes: results of a pilot study. *Int J Diabetes Metab* 2008;16:25–8.
- Alanazi B, Alshammari M, Alhawas F, et al. Reliability and validity of an Arabic version of a questionnaire to assess carbohydrate counting skills, knowledge of heart-healthy foods, and nutrition label-reading skills in adults with diabetes. *IJMDC* 2020;4:461–8.
- Kawamura T, Takamura C, Hirose M, et al. The factors affecting on estimation of carbohydrate content of meals in carbohydrate counting. *Clin Pediatr Endocrinol* 2015;24:153–65.
- Meade LT, Rushton WE. Accuracy of carbohydrate counting in adults. *Clin Diabetes* 2016;34:142–7.
- Watts SA, Anselmo JM, Kern E. Validating the AdultCarbQuiz: a test of carbohydrate-counting knowledge for adults with diabetes. *Diabetes Spectrum* 2011;24:154–60.
- Alhaji JH, Alshuwaier GO, Alharbi NS, et al. Translation and validation of the Arabic version of the capability assessment for diet and activity (CADA) questionnaire in Saudi university employed women. *Int J Environ Res Public Health* 2021;18:6246.
- Alshareef SM, Alkhatlan MA, Alwabel AA, et al. How does the utilization of diabetes dietitian and educator service in Saudi Arabia affect glycemic outcomes? *J Family Community Med* 2018;25:108–13.
- Bawazeer NM, Al-Qahtani SJ, Alzaben AS. The association between dietary patterns and socio-demographic and lifestyle characteristics: a sample of Saudi Arabia. *Curr Res Nutr Food Sci* 2021;9.
- Arslan M. Assessment of carbohydrate count method knowledge levels and insulin types of individuals with type 1 DM. *Clin Exp Health Sci* 2019;9:345–9.
- Kessler H, Wunderlich SM. Relationship between use of food labels and nutrition knowledge of people with diabetes. *Diabetes Educ* 1999;25:549–59.
- Post RE, Mainous AG, Diaz VA, et al. Use of the nutrition facts label in chronic disease management: results from the National health and nutrition examination survey. *J Am Diet Assoc* 2010;110:628–32.
- O’Gorman JR, O’Leary O, Finner N, et al. Knowledge of carbohydrate counting and insulin dose calculations among hospital staff in a regional General paediatrics unit. *Springerplus* 2015;4:727.
- Koontz MB, Cuttler L, Palmert MR, et al. Development and validation of a questionnaire to assess carbohydrate and insulin-dosing knowledge in youth with type 1 diabetes. *Diabetes Care* 2010;33:457–62.
- Schmidt S, Schelde B, Nørgaard K. Effects of advanced carbohydrate counting in patients with type 1 diabetes: a systematic review. *Diabet Med* 2014;31:886–96.
- Gökşen D, Atik Altınok Y, Ozen S, et al. Effects of carbohydrate counting method on metabolic control in children with type 1 diabetes mellitus. *J Clin Res Pediatr Endocrinol* 2014;6:74–8.
- Berget C, Messer LH, Forlenza GP. A clinical overview of insulin pump therapy for the management of diabetes: past, present, and future of intensive therapy. *Diabetes Spectr* 2019;32:194–204.
- Marker AM, Noser AE, Knecht N, et al. A time-friendly, feasible measure of nutrition knowledge in type 1 diabetes: the electronic Nutrition and Carbohydrate Counting Quiz (eNCQ). *J Diabetes Sci Technol* 2019;13:68–74.
- Smart CE, Ross K, Edge JA, et al. Can children with type 1 diabetes and their caregivers estimate the carbohydrate content of meals and snacks? *Diabet Med* 2010;27:348–53.
- Spiegel G, Bortsov A, Bishop FK, et al. Randomized nutrition education intervention to improve carbohydrate counting in adolescents with type 1 diabetes study: is more intensive education needed? *J Acad Nutr Diet* 2012;112:1736–46.