



Body-specific motor imagery of hand actions: neural evidence from right- and left-handers

Roel M. Willems^{1,2*}, Ivan Toni¹, Peter Hagoort^{1,3} and Daniel Casasanto³

¹ Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, Nijmegen, The Netherlands

² Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley, CA, USA

³ Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

Edited by:

Olaf Blanke, Ecole Polytechnique
Fédérale de Lausanne, Switzerland;
University of Geneva, Switzerland

Reviewed by:

Thomas F. Münte, University of
Magdeburg, Germany
Douglas O. Cheyne, Hospital for Sick
Children Research Institute, Canada

*Correspondence:

Roel M. Willems, Helen Wills Institute
for Neuroscience, University of
California Berkeley, 132 Barker Hall,
Berkeley, CA 94720-3190, USA.
e-mail: roelwillems@berkeley.edu

If motor imagery uses neural structures involved in action execution, then the neural correlates of imagining an action should differ between individuals who tend to execute the action differently. Here we report fMRI data showing that motor imagery is influenced by the way people habitually perform motor actions with their particular bodies; that is, motor imagery is ‘body-specific’ (Casasanto, 2009). During mental imagery for complex hand actions, activation of cortical areas involved in motor planning and execution was left-lateralized in right-handers but right-lateralized in left-handers. We conclude that motor imagery involves the generation of an action plan that is grounded in the participant’s motor habits, not just an abstract representation at the level of the action’s goal. People with different patterns of motor experience form correspondingly different neurocognitive representations of imagined actions.

Keywords: imagery, motor, handedness, fMRI, neuroimaging, embodied

INTRODUCTION

Studies employing behavioral, electrophysiological, as well as neuroimaging techniques indicate that motor imagery involves the generation of an action plan (Decety et al., 1989, 1994; Jeannerod, 1994, 2001; Parsons, 1994; Beisteiner et al., 1995; Lang et al., 1996; Porro et al., 1996; Bonnet et al., 1997; Schnitzler et al., 1997; Parsons et al., 1998; Neuper et al., 1999, 2005; Pfurtscheller et al., 1999; Gerardin et al., 2000; de Lange et al., 2005; Helmich et al., 2007; Szameitat et al., 2007a,b; Munzert et al., 2009). As such, motor imagery can be thought of as covert motor execution (Jeannerod, 1994, 2001). A remaining question is whether motor imagery is *body-specific* (Casasanto, 2008, 2009). That is, does the way one performs an action in the real world influence neural activation during motor imagery? Alternatively, it may be that the motor plan generated during motor imagery is abstracted away from individual motor experience or specific effectors and occurs at the level of *goal* of the imagined action (Rijntjes et al., 1999). Here we aimed to distinguish between these possibilities by measuring cerebral activity in left- and right-handed participants while they imagined performing everyday motor activities.

Previous research on this issue is inconclusive. Consistent with a body-specific view of mental imagery, there is some work showing different lateralization when imagining actions with the right hand as compared to actions with the left hand (Szameitat et al., 2007a), and decreased motor imagery performance specifically for the affected hand in Parkinson’s disease patients (Helmich et al., 2007). Yet, other work suggests motor planning may occur at the level of an action’s goal instead of at a more specific level such as preferred hand (Rijntjes et al., 1999), in which case the neural correlates of motor imagery should not necessarily vary with handedness.

In our own previous research, we have conducted fMRI experiments in left- and right-handers to investigate the influence of hand preference on the neural representation of action execution

and action observation (Willems and Hagoort, 2009), and on the neural basis of action semantics in language (Willems et al., in press). In one study we showed that differences in motor cortex activation between left- and right-handers in terms of action execution were similarly observed when participants viewed movie clips of these hand actions (Willems and Hagoort, 2009). Thus, differences in the way we act upon the world are reflected not only in the brain areas that subserve action production, but also in the neural substrates of action perception. In another study, we found that the motor component of action verb semantics is differently lateralized in right- and left-handers. When right-handers understand a verb that names a hand action (e.g. ‘grasp’, ‘throw’) they preferentially activate left premotor areas, whereas left-handers preferentially activate right premotor areas. At least by default, right- and left-handers represent action verb meanings from an egocentric perspective, which reflects the way they perform these actions with their dominant hands (Willems et al., in press).

Here we investigate effects of handedness on motor imagery, which lies in between the concrete domains of action execution and perception and the more abstract domain of language. If the way one performs an action in the real world is reflected in neural activation during motor imagery (i.e., if motor imagery is body-specific), then left- and right-handed participants should show differently lateralized activity in brain areas involved in the planning and execution of hand actions. Alternatively, if motor imagery only involves generating a motor plan that is abstracted away from the participant’s own motor experience, we expect to see no lateralization differences between the two groups. Such a result would be expected for instance if motor imagery is mainly based on visual experience, provided that both left- and right-handers mainly observe right-handers perform actions in the world.

MATERIALS AND METHODS

PARTICIPANTS

We tested 32 healthy participants with no known history of neurological problems, dyslexia or other language-related problems, and with normal or corrected-to-normal vision, all of whom gave informed consent. Half of the participants were left-handed ($N = 16$, 12 female, mean age 23.4 years, range 19–32 years, adapted Dutch version of Edinburgh Handedness Inventory (EHI) score (Oldfield, 1971; Van Strien, 1992): mean = -94.3 , SD = 8.7, range -82 to -100 , mode = -100), and half were right handed ($N = 16$, 10 female, mean age 23.2, range 20–29 years, EHI score: mean = 96.6, SD = 7.3, range 82–100, mode = 100). The groups did not differ in age [$|t(30)| < 1$], or in absolute EHI value [$t(30) < 1$]. The local ethics committee approved the study.

MATERIALS

Stimuli were 96 Dutch verbs expressing concrete actions. Half of these were related to manual actions (MAN, e.g. *to throw*), half of them were not related to manual actions (NONMAN, e.g. *to kneel*, see the Appendix for the complete list of stimuli). The distinction between MAN and NONMAN was pretested in a group of raters who did not participate in the fMRI experiment ($N = 16$), who scored for each verb how much they associated that action with their hand(s) on a 1- to 7-point scale. MAN words were significantly more associated with hand actions than NONMAN words [$t(94) = 23.60$, $p < 0.001$; mean MAN = 5.55, SD = 0.53; mean NONMAN = 2.04, SD = 0.83]. Raters also indicated whether they preferentially acted out the hand actions with the left, right, or with both hands. Materials were selected to ensure that the number of raters indicating to use both hands in that particular action, was low (out of 16 raters: mean = 3.36, SD = 1.89, median = 3, mode = 2). MAN and NONMAN word lists did not differ in imageability (assessed by the same group of raters) [$t(94) < 1$], number of phonemes [$t(94) < 1$] or lexical frequency [$t(94) < 1$]; defined using the CELEX database; Baayen et al., 1993).

EXPERIMENTAL PROCEDURE

Stimuli were presented using Presentation software (version 10.2¹). Each trial started with a fixation cross (250 ms) followed by presentation of a written verb (1500 ms) in the middle of the screen. Participants were instructed to read the word, close their eyes, imagine performing this action and open their eyes to indicate that they had finished motor imagery after which the next trial would start (after a variable intertrial interval between 2 and 6 s in steps of 250 ms (mean = 4 s) (Dale, 1999)). Participants were instructed to ‘vividly imagine performing this action several times and open your eyes when done’. This means that there was no constraint on the amount or duration of motor imagery, this was left to each individual participant. In terms of experimental design, we were hence vulnerable to the possible confound that participants would take longer in imagining one of the action types (MAN or NONMAN). This however turned out not to be the case (see Results).

Closing and opening of the eyes was monitored by an infrared IviewX eyetracker² and coded on-line by one of the experimenters. We used opening and closing of the eyes to signal start and finish of imagery instead of button presses since using the eyes as a response measure enables to measure imaging times and does not contaminate hand motor cortex activation due to button presses. Previous work shows that motor imagery with eyes closed entails similar processes as motor imagery with eyes open (Heremans et al., 2008) and has been successfully used before in neuroimaging studies (Szameitat et al., 2007a,b; Bakker et al., 2008). It is possible that opening and closing of the eyes leads to differential motor cortex lateralization in left- and right-handers. However, this is not problematic for the present study since opening and closing of the eyes was required for MAN and NONMAN trials alike and should hence cancel out when comparing these conditions with each other.

Stimuli were presented in pseudo-randomized order such that a condition was repeated maximally three times in a row. A mirrored presentation order was employed in half of the participants. Participants were familiarized with the procedure by 10 practice items containing different words than those used in the experiment.

DATA ACQUISITION AND ANALYSIS

Whole-brain cho-Planar Images were acquired with a 8-channel head coil on a Siemens MR system with 3T magnetic field strength (TR = 2060 ms; TE = 30 ms; flip angle 85°, 31 transversal slices; voxel size 3.5 mm × 3.5 mm × 3 mm, 0.5 mm gap between slices). Data analysis was done using SPM5³. Preprocessing involved realignment through rigid body registration, slice timing correction to the onset of the first slice, normalization to Montreal Neurological Institute (MNI) space, interpolation of voxel sizes to 2 mm × 2 mm × 2 mm, and spatial smoothing (8 mm FWHM kernel). First-level analysis involved a multiple regression analysis with regressors describing the expected hemodynamic responses during imagery of MAN words and NONMAN words. Each trial was modeled as the actual duration of the trial, convolved with a canonical hemodynamic response function (Friston et al., 1998). MR disturbances due to small head movements were accounted for by a series of nuisance regressors, namely the linear and exponential changes in the scan-by-scan estimated head motion, scan-by-scan average signals from outside the brain, white matter, and cerebrospinal fluid (Verhagen et al., 2006).

A second-level whole brain group analysis with subjects as a random factor (‘random effects analysis’) involved a model with factors VERB TYPE (MAN, NONMAN) and GROUP (left-handers, right-handers). Correction for multiple comparisons was applied by thresholding group maps at $p < 0.005$ uncorrected and subsequently taking the cluster extent into account by using the theory of Gaussian Random Fields to correct maps at $p < 0.05$ corrected for multiple comparisons (Poline et al., 1997). Differential lateralization differences between the groups was tested by means of repeated measures analysis of variance to the mean contrast

¹www.nbs.com

²www.smi.de

³http://www.fil.ion.ucl.ac.uk/spm/software/spm5/

estimates from the regions sensitive to the MAN > NONMAN comparison with factors HEMISPHERE (left, right) and GROUP (left-handers, right-handers). We circumvented a bias in regions of interest selection, since the regions of interest were based on the overall contrast across the two groups (Kriegeskorte et al., 2009). Follow-up one-sided planned comparisons of within group hemispheric differences (Right-handers_{left hem} > Right-handers_{right hem} and Left-handers_{right hem} > Left-handers_{left hem}) and between-group comparisons (Right-handers_{left hem} > Left-handers_{left hem} and Left-handers_{right hem} > Right-handers_{right hem}) were performed.

We also looked at common activations across the two groups. We implemented this as a conjunction analysis (Nichols et al., 2005) testing for areas activated to MAN > NONMAN in left-handers as well as to MAN > NONMAN in right-handers ($\text{MAN} > \text{NONMAN}_{\text{left-handers}} \cap \text{MAN} > \text{NONMAN}_{\text{right-handers}}$). For general interest, we also conducted a conjunction analysis to investigate overlapping areas during imagery of MAN actions in both groups ($\text{MAN}_{\text{left-handers}} \cap \text{MAN}_{\text{right-handers}}$).

Given the heterogeneity of effectors to which the NONMAN verbs refer, we never compared NONMAN > MAN directly.

RESULTS

BEHAVIORAL

It took participants on average 5.63 s (SD = 2.17) to imagine the MAN verbs and 5.55 s (SD = 1.95) to imagine the NONMAN verbs (Table 1). Right- and left-handers did not differ in imagining times, neither in overall times, nor in MAN or NONMAN times separately [MAN + NONMAN: $t(30) = 1.15$, $p = 0.26$; MAN: $t(30) = 1.20$, $p = 0.24$; NONMAN: $t(30) = 1.08$, $p = 0.28$]. There were no statistically significant differences within groups between MAN and NONMAN times [Left-handers: $t(15) = 1.64$, $p = 0.12$; Right-handers: $t < 1$].

NEURAL

There were no areas sensitive to the main effect of GROUP. No areas were sensitive to the VERB TYPE \times GROUP interaction at a whole-brain corrected statistical threshold. However, informal inspection at $p < 0.005$ uncorrected revealed such effects in bilateral postcentral sulcus. A wide-spread network of areas, including bilateral postcentral sulcus (mainly encompassing Brodmann Area 2;

Eickhoff et al., 2005), bilateral precentral sulcus (BA6) and bilateral inferior temporal cortex, was sensitive to the MAN > NONMAN comparison (Figure 1; Table 2).

Second, we determined which regions showed sensitivity to the MAN > NONMAN contrast in each group in isolation. In the left-handers there was stronger activation to MAN as compared to NONMAN imagery in right postcentral sulcus (BA2) extending into intraparietal sulcus, right precentral sulcus (BA6) and right inferior temporal sulcus (Figure 2; Table 3). Conversely, for the right-handers there was stronger activation for MAN as compared to NONMAN imagery in left postcentral sulcus (BA2), left precentral sulcus (BA6) extending into intraparietal sulcus, and left inferior temporal sulcus (Figure 2; Table 3). Informal inspection at a liberal statistical threshold ($p < 0.01$ uncorrected) showed that in both groups, there was no activation in primary motor cortex (cytoarchitectonically defined BA4a and BA4p; Eickhoff et al., 2005) to the MAN > NONMAN comparison. In previous work we observed that employing subject-specific regions of interest substantially improves sensitivity in detecting between-group differences (Willems et al., in press; see also Aziz-Zadeh et al., 2006). As an additional check on the involvement of primary

Table 1 | Behavioral results. Mean (and standard deviation) imagining times in seconds for left-handers (left column) and right-handers (middle column), and the results of an independent samples *t*-test to test for a difference in imagining times between the groups and within-group differences between MAN and NONMAN durations (right column). Data are displayed separately for MAN alone, NONMAN alone and MAN + NONMAN.

	Left-handers Mean (SD)	Right-handers Mean (SD)	Left-handers > right-handers
MAN	6.09 (2.68)	5.18 (1.44)	$t(30) = 1.20$, $p = 0.24$
NONMAN	5.93 (2.42)	5.18 (1.32)	$t(30) = 1.08$, $p = 0.28$
MAN + NONMAN	5.63 (2.17)	5.55 (1.95)	$t(30) = 1.15$, $p = 0.26$
MAN > NONMAN	$t(16) = 1.64$, $p = 0.12$	$t(16) < 1$	




FIGURE 1 | Result from whole brain analysis for both groups combined.

Displayed are the results for comparing MAN > NONMAN across the two groups ($\text{MAN}_{\text{left-handers}} + \text{MAN}_{\text{right-handers}} > \text{NONMAN}_{\text{left-handers}} + \text{NONMAN}_{\text{right-handers}}$).

Motor imagery of manual actions activated dorsal precentral sulcus, postcentral sulcus and inferior temporal sulcus bilaterally. Results are corrected for multiple comparisons at $p < 0.05$ corrected.




FIGURE 2 | Results from whole brain analysis for each group separately. Displayed are the results for the MAN > NONMAN comparison for left-handers (yellow) and right-handers (blue). Note the strong lateralization of responses in precentral, postcentral and inferior temporal sulcus. Results are corrected for multiple comparisons at $p < 0.05$ corrected.

Table 2 | Results of overall MAN > NONMAN comparison (including both groups). Displayed are an anatomical description of activated areas, the coordinates of local maxima within a region in MNI space and the number of voxels per area (2 mm × 2 mm × 2 mm voxels). Results are corrected for multiple comparisons by thresholding at $p < 0.005$ and taking cluster extent into account to arrive at a $p < 0.05$ corrected p -value.

Region	MNI coordinates	nr Voxels
R dorsal precentral/middle frontal sulcus	22 -8 42 28 -8 58	1097
R postcentral sulcus (extending into intraparietal sulcus)	38 -38 48 34 -46 58	5724
R inferior/middle temporal sulcus	58 -62 -2 48 -52 -12	
R thalamus	16 -28 -8	
L dorsal precentral/middle frontal sulcus	-24 -12 52 -24 -2 68	772
L postcentral sulcus (extending into intraparietal sulcus)	-38 -38 44 -16 -64 52	716
L inferior/middle temporal sulcus	-54 -64 4	1292
L thalamus	-14 -30 4	
L fusiform gyrus	-38 -26 -18	

motor cortex in body-specific motor imagery, we therefore determined subject-specific regions of interest as 4-mm spheres around local maxima to imagery of MAN trials in BA4 (again, using cytoarchitecturally defined BA4a and BA4p), and extracted the MAN > NONMAN contrast values for each subject separately. These were analyzed with repeated measures analysis of variance with factors HEMISPHERE (left, right) and GROUP (left-handers, right-handers). We did observe a HEMISPHERE × GROUP interaction to the MAN > NONMAN contrast values [$F(1, 30) = 4.41$, $MSE = 0.06$, $p = 0.044$], with each group showing greater activation in the hemisphere contralateral to the preferred hand.

Third, to determine whether there were lateralization differences between the groups, we extracted MAN > NONMAN contrast values from precentral, postcentral and inferior/middle temporal regions sensitive to the overall MAN > NONMAN contrast (Figure 1; Table 2), and analyzed these in an ANOVA with factors HEMISPHERE (left, right) and GROUP (left-handers, right-handers). In all these regions, there was a significant

HEMISPHERE × GROUP interaction (Table 4; Figure 3). Moreover, in postcentral and precentral structures, each group showed stronger activation in the hemisphere contralateral to the dominant hand than in the ipsilateral region (Table 4; Figures 3A,B), albeit that this difference was only a trend in precentral sulcus in left-handers ($p = 0.08$).

There were no areas commonly activated to MAN > NONMAN in both left and right-handers ($MAN > NONMAN_{\text{left-handers}} \cap MAN > NONMAN_{\text{right-handers}}$). However, at a lower $p < 0.005$ uncorrected statistical threshold, such overlap was observed in left dorsal premotor cortex. An extensive set of areas were commonly activated in left- as well as in right-handers during imagery of MAN actions ($MAN_{\text{left-handers}} \cap MAN_{\text{right-handers}}$). Activated areas include bilateral dorsal premotor cortex, bilateral inferior frontal gyrus, bilateral postcentral sulcus, bilateral inferior/middle temporal gyrus and bilateral middle occipital gyrus (Figure S1 and Table S1 in Supplementary Material).

To summarize, there were three bilateral pairs of regions (in precentral, postcentral and inferior/middle temporal sulci) which showed sensitivity to the MAN > NONMAN comparison either across the two groups (left-handers + right-handers), in left-handers (right-hemisphere regions), or in right-handers (left-hemisphere regions). Left-handers activated postcentral and precentral motor cortex more strongly in the right as compared to the left hemisphere, whereas the opposite pattern was observed in right-handers (left > right). A similar effect was observed in primary motor cortex (BA4) when employing subject-specific region of interest analysis, but not in the whole brain analysis.

DISCUSSION

In this study we investigated whether motor imagery involves the generation of a motor plan that is grounded in the way an individual typically performs the imagined action in the real world. Our results indicate that explicit motor imagery of everyday hand actions is body-specific (Casasanto, 2008, 2009). Left- and right-handers showed differential and opposite lateralization of activity in premotor and postcentral motor regions when they imagined performing one-handed manual actions, as compared to nonmanual actions. The hemisphere that primarily controls the dominant hand also subserves mental imagery for actions that people usually perform with this hand (see Szameitat et al., 2007a for a compatible finding in right-handers).

Table 3 | Results from whole brain analysis comparing MAN > NONMAN motor imagery in left-handers (top) and right-handers (bottom). Displayed are an anatomical description of activated areas, the coordinates of local maxima within a region in MNI space and the number of voxels per area (2 mm × 2 mm × 2 mm voxels). Results are corrected for multiple comparisons by thresholding at $p < 0.005$ and taking the cluster extent into account to arrive at a $p < 0.05$ corrected p -value.

	Region	MNI coordinates	nr Voxels	
Left-handers MAN > NONMAN	R precentral/middle frontal sulcus	32 -10 58 24 -8 40 22 -4 44 20 -10 48	619	
	R postcentral sulcus (extending into intraparietal sulcus)	36 -44 62 40 -38 50 36 -44 54 54 -20 36 24 -52 68	1052	
	R inferior/middle temporal sulcus	46 -50 -12 56 -60 0 54 -54 -4	364	
	Right-handers MAN > NONMAN	L dorsal precentral/middle frontal sulcus	-26 -8 54 -28 -12 52 -24 -16 50 -18 -6 46 -24 -10 70 -20 14 66	1022
		L postcentral sulcus (extending into intraparietal sulcus)	-38 -40 46	404
		L inferior/middle temporal sulcus	-56 -66 2 -42 -52 -12 -48 -60 -14	1213

Our results are in line with earlier work showing that present body posture influences motor imagery (de Lange et al., 2006). Here we extend this by showing that long term motor history (i.e. a preference to execute an action with one hand) also influences motor imagery. In addition to finding body-specific laterality effects in dorsal premotor cortex (BA6), we also observed these effects in primary somatosensory cortex (S1, roughly corresponding with BA2), and, when using subject-specific regions of interest, in BA4. S1 activation during motor imagery has been argued to reflect the predicted somatosensory consequences of the imagined actions (i.e., a forward model, see Wolpert and Ghahramani, 2000), but is observed only in some of the relevant neuroimaging studies (see Munzert et al., 2009 for review). Szameitat et al. (2007b) also reported S1 activation during motor imagery when participants were required to perform imagery of everyday actions, just as in the present study (see also Sacco et al., 2006). Some studies that did not observe S1 activation for instance employed the Parsons' hand laterality judgment task, which arguably requires less elaborate motor imagery (e.g. Parsons et al., 1998; de Lange et al., 2005, 2006). It is possible that the verbal instruction to imagine a relatively complex action for a more extended period of time (as in the present study and in




Table 4 | Results in regions of interest taken from the whole brain analysis (Figure 1; Table 2). The whole brain analysis revealed three pairs of cortical regions that were sensitive to the overall MAN > NONMAN contrast (i.e. across groups, **Figure 1**): bilateral precentral sulcus, bilateral central sulcus, and bilateral inferior temporal sulcus. Results from these regions were analysed in an ANOVA with factors HEMISPHERE (left, right) and GROUP (left-handers, right-handers). In all regions there was a significant GROUP \times HEMISPHERE interaction. To subsequently test the direction of this effect, within group hemisphere differences were tested (i.e. Right handers_{left hem} > Right handers_{right hem} and Left handers_{right hem} > Left handers_{left hem}). *t*-tests were one-sided. Differences significant at the $p < 0.05$ level are indicated in bold typeface.

	Precentral sulcus		Postcentral sulcus		Inferior temporal sulcus	
GROUP \times HEM	$F(1,30) = 15.21$, MSe = 0.003	$p = 0.001$	$F(1,30) = 23.67$, MSe = 0.006	$p < 0.001$	$F(1,30) = 9.09$, MSe = 0.003	$p = 0.005$
HEMISPHERE	$F(1,30) = 2.76$, MSe = 0.003	$p = 0.11$	$F < 1$, MSe = 0.006	n.s.	$F(1,30) = 5.47$, MSe = 0.003	$p = 0.026$
GROUP	$F < 1$, MSe = 0.019	n.s.	$F(1,30) = 1.28$, MSe = 0.027	0.27	$F < 1$, MSe = 0.011	n.s.
Right handers _{left hem} > Right handers _{right hem}	$t(15) = 4.45$	$p < 0.001$	$t(15) = 3.28$	$p = 0.003$	$t(15) = 2.86$	$p = 0.006$
Left handers _{right hem} > Left handers _{left hem}	$t(15) = 1.45$	$p = 0.084$	$t(15) = 3.61$	$p = 0.001$	$t(15) < 1$	n.s.

Szameitat et al., 2007b) leads to a more elaborate forward model and hence to stronger S1 activation⁴. A similar reasoning could be applied to the effects we observed in primary motor cortex (BA4). Involvement of primary motor cortex is observed in some (e.g. Tomasino et al., 2007, 2008), but not in other investigations of motor imagery (see Munzert et al., 2009 for overview). The findings in BA4 should be interpreted with caution since we did observe a Hemisphere \times Group interaction in subject-specific regions of interest, but not in within-group comparisons in the whole brain analysis, even when assessed at liberal and uncorrected statistical thresholds.

It should be acknowledged that we did not measure EMG to ensure that there was no supra-threshold muscle activity during motor imagery. It is possible that the effects we observed in primary motor cortex/somatosensory cortex could be driven by supra-threshold muscle activation. It should be stressed however that participants were in no way encouraged to move and were explicitly instructed only to imagine performing the actions. Moreover, we checked visually that participants were not actually acting out the movements they were required to imagine. Hence primary motor cortex and somatosensory cortex activation cannot be due to actual acting out of these actions, which does not preclude subthreshold motor activation.

Interestingly, a similar lateralization difference was observed in inferior/middle temporal cortex. This suggests that the influence of hand preference extends beyond the cortical motor and language systems. Indeed in previous work we observed that lateralization of

extrastriate regions involved in observation of faces (fusiform face area) and bodies (extrastriate body area) is influenced by handedness (Willems et al., in press). However in that study extrastriate body area was right-lateralized in both groups, but to a lesser extent in left- as compared to right-handers. In the present study we observe a different lateralization pattern with a strong left-lateralization in inferior/middle temporal cortex for right-handers (**Figure 3**). It is unclear what cognitive process drives these differences in temporal cortex.

The present findings suggest that when participants were asked to ‘vividly imagine performing an action’, they did so from their own (egocentric) perspective. While this may not be surprising, it was by no means a foregone conclusion. Although left-handers tend to perform actions like *throwing* with their left hands, they probably observe a far greater number of throws performed with the right hand, since the majority of throwers are right-handed. In principle, right- and left-handers could all generate motor images that reflect the statistics of observed actions rather than performed actions, which would result in similar patterns of motor activity across groups. The fact that participants generated body-specific images from their own perspective is consistent with our previous studies showing that people perceive actions and understand action language egocentrically, at least by default (Willems and Hagoort, 2009; Willems et al., in press). These findings should not be interpreted as indicating that people are *only* capable of imagining actions from their own perspective. It remains a question for future research how actions imagined from another’s perspective are instantiated in right- and left-handers’ motor systems (see Szameitat et al., 2007b; Tomasino et al., 2007).

On a methodological note, this study validates the use of differential lateralization in the motor system between left- and right-handers as an experimental tool (Longcamp et al., 2003, 2005; Lewis et al., 2006). Earlier studies of motor execution differences in left- and right-handers did not observe such lateralization differences (Kim et al., 1993; Kloppel et al., 2007). However, a crucial difference with the present study is that in these previous studies very simple hand actions were used. For instance in Kim et al., the hand actions

⁴This distinction is reminiscent of that between visual versus kinesthetic motor imagery (e.g. Guillot et al., 2009). In kinesthetic motor imagery participants are explicitly trained to focus on the kinesthetic consequences of the actions that they imagine as compared to visual motor imagery in which participants are instructed to focus on the visual aspects of motor imagery. Guillot et al. (2009) observed increased activations during kinesthetic motor imagery compared to a low-level control condition in somatosensory cortex. Such an effect was not observed when comparing visual motor imagery to the same control condition. A direct comparison between kinesthetic and visual motor imagery, however, did not reveal increased activation during kinesthetic motor imagery in S1. In our experiment we did not specifically instruct participants to focus/direct attention on one of these aspects of motor imagery and this interpretation is hence speculative.

were simple finger-thumb oppositions. In Kloppel et al. (2007) the hand actions involved pressing a button. Indeed we previously also did not observe the clear-cut difference in lateralization between left- and right-handers when they performed simple contractions and extension of the fingers (Willems and Hagoort, 2009). Future research should more systematically investigate how the complexity of performed/imagined actions influences the amount of lateralization in left- and right-handers. For present purposes it is important to note that our finding of differently lateralized motor cortex activation in left- and right-handed participants lends support to using handedness-related lateralization to study the body-specificity of other cognitive processes as well (e.g. Casasanto, 2009; Willems et al., in press; Willems and Hagoort, 2007). Moreover, this finding could have implications for clinical practice, in which motor imagery is sometimes used as a therapeutic tool (see Munzert et al., 2009 for comprehensive review).

Previous studies have suggested that the generation of an action plan occurs at the level of an action's overall goal (Rijntjes et al., 1999). Although some components of motor imagery may be abstracted away from motor experience, the present data show that motor imagery also involves generating an action plan consistent with the kinematics of actions as we tend to perform them with our particular bodies.

APPENDIX

Stimuli used. Shown are the original stimuli in Dutch, with a translation in English. These verbs were matched and pretested on several measures, please see main text for details. Due to the literal translation in English it may be less obvious that some of the words are MAN or NONMAN verbs. It should be noted that the original Dutch stimulus set was carefully pretested to ensure that all MAN words were associated with hand actions and that the NONMAN words were not. This may not be fully captured by the translation in English and hence some of the verbs may not be obviously related to the hand(s) in the English translation. Moreover, the English translation sometimes has the same word appear twice, this was not the case in the Dutch originals.

Dutch	English	Dutch	English
MAN words			
aaien	to stroke	kloppen	to knock
bonken	to bonk	knijpen	to squeeze
bonzen	to pound	malen	to grind
borstelen	to brush	meppen	to smack
deppen	to dab	noteren	to jot
dobbelen	to dice	peuteren	to pick
epileren	to pluck	plamuren	to fill
gooien	to throw	plukken	to pick
grabbelen	to scramble	porren	to prod
graveren	to engrave	prakken	to mash
grijpen	to catch	prikken	to pierce
grissen	to snatch	reiken	to reach
hameren	to hammer	roeren	to stir
kerven	to notch	salueren	to salute

Dutch	English	Dutch	English
MAN words			
scheren	to shave	stoten	to push
schieten	to shoot	strelen	to stroke
schminken	to put make up on	strijken	to iron
schrapen	to scrape	strooien	to strew
slaan	to beat	tekenen	to draw
smeren	to lubricate	tikken	to tap
smijten	to fling	verven	to paint
snijden	to cut	werpen	to throw
steken	to stab	wijzen	to point
stompen	to stomp	zwaaien	to wave

Dutch	English	Dutch	English
NONMAN words			
ademen	to breathe	mopperen	to grumble
beven	to shake	neuriën	to hum
bijten	to bite	nippen	to sip
blazen	to blow	observeren	to observe
brommen	to hum	ontwijken	to dodge
brullen	to roar	rillen	to shiver
dicteren	to dictate	ruiken	to smell
drinken	to drink	schouderophalen	to shrug
fluisteren	to whisper	schrikken	to scare up
giechelen	to giggle	slikken	to swallow
gillen	to scream	smakken	to smack lips
grijnzen	to smirk	snuiven	to sniff
grinniken	to chortle	snurken	to snore
hoesten	to cough	speuren	to trace
inademen	to breathe	spugen	to spit
knabbelen	to nibble	staren	to stare
knagen	to gnaw	stotteren	to stutter
knielen	to kneel	turen	to gaze
knikken	to nod	uitademen	to exhale
knipperen	to flash	waarnemen	to perceive
knorren	to snore	zingen	to sing
krijzen	to screech	zuigen	to suck
leunen	to lean	zweten	to sweat
likken	to lick	zwijgen	to keep silent

ACKNOWLEDGMENTS

Supported by grants from European Union Joint-Action Science and Technology Project (IST-FP6-003747), the Dutch Organisation for Scientific research (NWO Rubicon 446-08-008) and the Niels Stensen foundation. We thank Martin Laverman, Jacqueline de Nooijer, Daan van Rooij and Paul Gaalman for assistance.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://www.frontiersin.org/humanneuroscience/paper/10.3389/neuro.09/039.2009/>

REFERENCES

- Aziz-Zadeh, L., Wilson, S. M., Rizzolatti, G., and Iacoboni, M. (2006). Congruent embodied representations for visually presented actions and linguistic phrases describing actions. *Curr. Biol.* 16, 1818–1823.
- Baayen, R. H., Piepenbrock, R., and van Rijn, H. (1993). The CELEX Lexical Database. Philadelphia, PA, Linguistic Data Consortium, University of Pennsylvania.
- Bakker, M., De Lange, F. P., Helmich, R. C., Scheeringa, R., Bloem, B. R., and Toni, I. (2008). Cerebral correlates of motor imagery of normal and precision gait. *Neuroimage* 41, 998–1010.
- Beisteiner, R., Höllinger, P., Lindinger, G., Lang, W., and Berthoz, A. (1995). Mental representations of movements. Brain potentials associated with imagination of hand movements. *Electroencephalogr. Clin. Neurophysiol.* 96, 183–193.
- Bonnet, M., Decety, J., Jeannerod, M., and Requin, J. (1997). Mental simulation of an action modulates the excitability of spinal reflex pathways in man. *Brain Res. Cogn. Brain Res.* 5, 221–228.
- Casasanto, D. (2008). The Body-Specificity Hypothesis. Proceedings of the 49th Annual Meeting of the Psychonomic Society, Chicago, IL.
- Casasanto, D. (2009). Embodiment of abstract concepts: good and bad in right and left-handers. *J. Exp. Psych. Gen.* 138, 351–367.
- Dale, A. M. (1999). Optimal experimental design for event-related fMRI. *Hum. Brain Mapp.* 8, 109–114.
- de Lange, F. P., Hagoort, P., and Toni, I. (2005). Neural topography and content of movement representations. *J. Cogn. Neurosci.* 17, 97–112.
- de Lange, F. P., Helmich, R. C., and Toni, I. (2006). Posture influences motor imagery: an fMRI study. *Neuroimage* 33, 609–617.
- Decety, J., Jeannerod, M., and Prablanc, C. (1989). The timing of mentally represented actions. *Behav. Brain Res.* 34, 35–42.
- Decety, J., Perani, D., Jeannerod, M., Bettinardi, V., Tadary, B., Woods, R., Mazziotta, J. C., and Fazio, F. (1994). Mapping motor representations with positron emission tomography. *Nature* 371, 600–602.
- Eickhoff, S. B., Stephan, K. E., Mohlberg, H., Grefkes, C., Fink, G. R., Amunts, K., and Zilles, K. (2005). A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage* 25, 1325–1335.
- Friston, K. J., Fletcher, P., Josephs, O., Holmes, A., Rugg, M. D., and Turner, R. (1998). Event-related fMRI: characterizing differential responses. *Neuroimage* 7, 30–40.
- Gerardin, E., Sirigu, A., Lehericy, S., Poline, J. B., Gaymard, B., Marsault, C., Agid, Y., and Le Bihan, D. (2000). Partially overlapping neural networks for real and imagined hand movements. *Cereb. Cortex* 10, 1093–1104.
- Guillot, A., Collet, C., Nguyen, V. A., Malouin, F., Richards, C., and Doyon, J. (2009). Brain activity during visual versus kinesthetic imagery: an fMRI study. *Hum. Brain Mapp.* 30, 2157–2172.
- Helmich, R. C., de Lange, F. P., Bloem, B. R., and Toni, I. (2007). Cerebral compensation during motor imagery in Parkinson's disease. *Neuropsychologia* 45, 2201–2215.
- Heremans, E., Helsen, W. F., and Feys, P. (2008). The eyes as a mirror of our thoughts: quantification of motor imagery of goal-directed movements through eye movement registration. *Behav. Brain Res.* 187, 351–360.
- Jeannerod, M. (1994). The representing brain: neural correlates of motor intention and imagery. *Behav. Brain Sci.* 17, 187–245.
- Jeannerod, M. (2001). Neural simulation of action: a unifying mechanism for motor cognition. *Neuroimage* 14, S103–S109.
- Kim, S. G., Ashe, J., Hendrich, K., Ellermann, J. M., Merkle, H., Ugurbil, K., and Georgopoulos, A. P. (1993). Functional magnetic resonance imaging of motor cortex: hemispheric asymmetry and handedness. *Science* 261, 615–617.
- Kloppel, S., van Eimeren, T., Glauche, V., Vongersichten, A., Munchau, A., Frackowiak, R. S., Buchel, C., Weiller, C., and Siebner, H. R. (2007). The effect of handedness on cortical motor activation during simple bilateral movements. *Neuroimage* 34, 274–280.
- Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S., and Baker, C. I. (2009). Circular analysis in systems neuroscience: the dangers of double dipping. *Nature Neuroscience*, 12, 535–540.
- Lang, W., Cheyne, D., Hollinger, P., Gerschlag, W., and Lindinger, G. (1996). Electric and magnetic fields of the brain accompanying internal simulation of movement. *Brain Res. Cogn. Brain Res.* 3, 125–129.
- Lewis, J. W., Phinney, R. E., Brefszynski-Lewis, J. A., and DeYoe, E. A. (2006). Lefties get it “right” when hearing tool sounds. *J. Cogn. Neurosci.* 18, 1314–1330.
- Longcamp, M., Anton, J. L., Roth, M., and Velay, J. L. (2003). Visual presentation of single letters activates a premotor area involved in writing. *Neuroimage* 19, 1492–1500.
- Longcamp, M., Anton, J. L., Roth, M., and Velay, J. L. (2005). Premotor activations in response to visually presented single letters depend on the hand used to write: a study on left-handers. *Neuropsychologia* 43, 1801–1809.
- Munzert, J., Lorey, B., and Zentgraf, K. (2009). Cognitive motor processes: the role of motor imagery in the study of motor representations. *Brain Res. Rev.* 60, 306–326.
- Neuper, C., Scher, R., Reiner, M., and Pfurtscheller, G. (2005). Imagery of motor actions: differential effects of kinesthetic and visual-motor mode of imagery in single-trial EEG. *Brain Res. Cogn. Brain Res.* 25, 668–677.
- Neuper, C., Schlogl, A., and Pfurtscheller, G. (1999). Enhancement of left-right sensorimotor EEG differences during feedback-regulated motor imagery. *J. Clin. Neurophysiol.* 16, 373–382.
- Nichols, T., Brett, M., Andersson, J., Wager, T., and Poline, J. B. (2005). Valid conjunction inference with the minimum statistic. *Neuroimage* 25, 653–660.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9, 97–113.
- Parsons, L. M. (1994). Temporal and kinematic properties of motor behavior reflected in mentally simulated action. *J. Exp. Psychol. Hum. Percept. Perform.* 20, 709–730.
- Parsons, L. M., Gabrieli, J. D., Phelps, E. A., and Gazzaniga, M. S. (1998). Cerebrally lateralized mental representations of hand shape and movement. *J. Neurosci.* 18, 6539–6548.
- Pfurtscheller, G., Neuper, C., Ramoser, H., and Müller-Gerking, J. (1999). Visually guided motor imagery activates sensorimotor areas in humans. *Neurosci. Lett.* 269, 153–156.
- Poline, J. B., Worsley, K. J., Evans, A. C., and Friston, K. J. (1997). Combining spatial extent and peak intensity to test for activations in functional imaging. *Neuroimage*, 5, 83–96.
- Porro, C. A., Francescato, M. P., Cettolo, V., Diamond, M. E., Baraldi, P., Zuiani, C., Bazzocchi, M., and di Prampero, P. E. (1996). Primary motor and sensory cortex activation during motor performance and motor imagery: a functional magnetic resonance imaging study. *J. Neurosci.* 16, 7688–7698.
- Rijntjes, M., Dettmers, C., Buchel, C., Kiebel, S., Frackowiak, R. S., and Weiller, C. (1999). A blueprint for movement: functional and anatomical representations in the human motor system. *J. Neurosci.* 19, 8043–8048.
- Sacco, K., Cauda, F., Cerliani, L., Mate, D., Duca, S., and Geminiani, G. C. (2006). Motor imagery of walking following training in locomotor attention. The effect of “the tango lesson”. *Neuroimage* 32, 1441–1449.
- Schnitzler, A., Salenius, S., Salmelin, R., Jousmaki, V., and Hari, R. (1997). Involvement of primary motor cortex in motor imagery: a neuromagnetic study. *Neuroimage* 6, 201–208.
- Szameitat, A. J., Shen, S., and Sterr, A. (2007a). Effector-dependent activity in the left dorsal premotor cortex in motor imagery. *Eur. J. Neurosci.* 26, 3303–3308.
- Szameitat, A. J., Shen, S., and Sterr, A. (2007b). Motor imagery of complex everyday movements. An fMRI study. *Neuroimage* 34, 702–713.
- Tomasino, B., Fink, G. R., Sparing, R., Dafotakis, M., and Weiss, P. H. (2008). Action verbs and the primary motor cortex: a comparative TMS study of silent reading, frequency judgments, and motor imagery. *Neuropsychologia* 46, 1915–1926.
- Tomasino, B., Werner, C. J., Weiss, P. H., and Fink, G. R. (2007). Stimulus properties matter more than perspective: an fMRI study of mental imagery and silent reading of action phrases. *Neuroimage* 36(Suppl. 2), T128–T141.
- Van Strien, J. W. (1992). Classificatie van links – en rechtshandige proefpersonen/Classification of left-handed and right-handed test subjects. *Ned. Tijdschr. Psychol.* 47, 88–92.
- Verhagen, L., Grol, M., Dijkerman, H., and Toni, I. (2006). Studying visually-guided reach-to-grasp movements in an MR-environment. *Neuroimage* 31, S45.
- Willems, R. M., and Hagoort, P. (2007). Neural evidence for the interplay between language, gesture, and action: a review. *Brain Lang.* 101(3), 278–289.
- Willems, R. M., and Hagoort, P. (2009). Hand preference influences neural correlates of action observation. *Brain Res.* 1269, 90–104.
- Willems, R. M., Hagoort, P., and Casasanto, D. (in press). Body-specific representations of action verbs: Neural

evidence from right- and left-handers. *Psychol. Sci.*

Willems, R. M., Peelen, M. V., and Hagoort, P. (in press). Cerebral lateralization of face-selective and body-selective visual areas depends upon handedness. *Cereb. Cortex.*

Wolpert, D. M., and Ghahramani, Z. (2000). Computational principles

of movement neuroscience. *Nat. Neurosci.* 3, 1212–1217.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 20 June 2009; paper pending published: 28 August 2009; accepted: 09 October 2009; published online: 10 November 2009.

Citation: Willems RM, Toni I, Hagoort P and Casasanto D (2009) Body-specific motor imagery of hand actions: neural evidence from right- and left-handers. *Front. Hum. Neurosci.* 3:39. doi: 10.3389/neuro.09.039.2009

Copyright © 2009 Willems, Toni, Hagoort and Casasanto. This is an open-access article subject to an exclusive license agreement between the authors and the Frontiers Research Foundation, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are credited.