

# “I Felt I Was Reaching a Point with My Health”: Understanding Reasons for Engagement and Acceptability of Treatment Services for Unhealthy Alcohol Use in Primary Care

Journal of Primary Care & Community Health  
Volume 12: 1–10  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/21501327211003005  
journals.sagepub.com/home/jpc



Megan A. O’Grady<sup>1</sup> , Joseph Conigliaro<sup>2,3</sup>, Svetlana Levak<sup>2</sup>, Jeanne Morley<sup>2,3</sup>, Sandeep Kapoor<sup>2,3</sup>, Melanie Ritter<sup>2</sup>, Christina Marini<sup>4</sup>, and Jon Morgenstern<sup>2,3</sup>

## Abstract

**Introduction/Objectives:** Despite increasing need, there are large gaps in provision of care for unhealthy alcohol use. Primary care practices have become increasingly important in providing services for unhealthy alcohol use, yet little is known about the reasons patients engage in these services and their views on acceptability of such programs. The purpose of this study was to examine primary care patients’ reasons for engagement, experiences with, and acceptability of a primary care practice-based program for treating unhealthy alcohol use. **Methods:** This qualitative study was conducted in a primary care practice that was developing a collaborative care model for treating unhealthy alcohol use in primary care. Semi-structured interviews were conducted with 24 primary care patients. Data were analyzed using conventional qualitative content analysis. **Results:** Findings suggest that patients engaged for both internal (concerns about drinking and health) and external reasons (family or provider concern). Patient experiences in the program were shaped by their affective responses (enjoyable, enlightening), as well as therapeutic benefits (gaining new insights about drinking; staff/provider support). Acceptability was driven by core program elements (medication, therapy, integration) as well as positive impacts on drinking cognition and behavior and flexible, patient-centered approaches. **Conclusions:** Offering flexible and comprehensive programs with multiple elements and both abstinence and moderation goals could also improve patient engagement and views on acceptability. Primary care practices will need to be thoughtful about the resources needed to implement these programs in terms of staffing, training, and program support.

## Keywords

hazardous drinking, primary care, qualitative methods, collaborative care, screening and intervention

Dates received: 23 February 2021; revised: 23 February 2021; accepted: 24 February 2021.

Alcohol consumption and use of alcohol-related acute care have increased in the United States over the past 2 decades and the number of alcohol-related deaths per year has doubled during this time.<sup>1</sup> National data suggests that of the over 140 million current alcohol users in the United States, 47% exceeded recommended per occasion drinking amounts and one-quarter of those individuals did so on 5 or more days in the past month.<sup>2</sup> Approximately 14 million adults in the United States meet criteria for alcohol use disorder (AUD).<sup>3</sup>

Despite increasing need, large gaps in providing care for unhealthy drinking persist. Most individuals with unhealthy alcohol use (UAU) receive no clinical care.<sup>4</sup> UAU is defined

as a continuum of behaviors from risky or harmful use (exceeding recommended daily, weekly, or per occasion amounts) to AUD.<sup>5</sup> Only about 1 in 6 binge drinkers are

<sup>1</sup>University of Connecticut School of Medicine, Farmington, CT, USA

<sup>2</sup>Northwell Health, Manhasset, NY, USA

<sup>3</sup>Zucker School of Medicine at Northwell/Hofstra, Hempstead, NY, USA

<sup>4</sup>Partnership to End Addiction, New York, NY, USA

## Corresponding Author:

Megan A. O’Grady, Department of Public Health Sciences, School of Medicine, University of Connecticut, 263 Farmington Avenue, Farmington, CT 06030-6325, USA.  
Email: ogrady@uchc.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

asked about alcohol use and advised to cut down by a health professional<sup>6</sup> and less than 10% of individuals with AUD receive treatment.<sup>3</sup> Even when patients are being routinely screened for UAU, few who screen positive receive recommended care as part of standard clinical practice.<sup>4</sup>

Primary care practices (PCPs) play an increasingly important role in providing care for UAU.<sup>7-11</sup> The U.S. Preventive Service Task Force recently renewed their recommendation that adults be screened and provided a brief intervention for UAU in PCPs.<sup>5</sup> Such screening and brief intervention (SBI) models in PCPs have been shown to reduce alcohol use among people who are not dependent on alcohol.<sup>12,13</sup> In addition, models of care that also address those with more severe alcohol use are being developed and tested within PCPs.<sup>14-18</sup> These models tend to utilize care management or collaborative care approaches that provide patients with ongoing care coordination, pharmacotherapy for AUD, and evidence-based therapy using multidisciplinary teams. Results have been mixed in randomized trials testing these models. Notably, even in studies with positive outcomes, initial engagement and sustaining patients in ongoing care (eg, therapy sessions and clinic visits) was challenging.<sup>9,15,17</sup>

Providing AUD care in PCPs differs from specialty treatment settings because patients may not be seeking treatment for AUD, may not be aware that such care is available within their PCP, or may be surprised when they screen positive on a universal screen conducted during a routine PCP visit. Recent studies have found that patient-level characteristics (eg, demographics, stigma) are related to engagement in substance use disorder (SUD) care in PCPs.<sup>9,19</sup> These findings suggest that reasons for engagement and acceptability need to be better understood to improve interventions and their implementation. As integration of AUD treatment within primary care continues to be developed, tested, and expanded, additional studies are needed to learn how to best engage patients and understand how acceptable they find services to ensure that these services can be implemented and sustained effectively.<sup>20</sup>

Several systematic reviews have shown that patient views are rarely evaluated in integrated care programs.<sup>21-23</sup> However, patient evaluations of treatment programs play a substantial role in predicting AUD treatment outcomes and should be more frequently considered.<sup>24</sup> Patient perspectives are needed to identify strategies to improve outcomes and inform implementation of treatment for UAU in PCPs.<sup>25,26</sup> While staff and provider views of implementing SUD screening and care in PCPs have been explored fairly extensively;<sup>27-30</sup> there are limited qualitative studies that examine patient perspectives of UAU programs in PCPs, especially those that examine patients' real-world experiences in receiving this care. Of the few qualitative studies that have focused on PCP patients, they have mostly focused on SBI rather than expanded care models.<sup>8,28,31-35</sup>

This study will address these gaps in knowledge by examining patient perspectives and experiences in a care

program for UAU in a PCP using qualitative methods. Specifically, we assess reasons for program participation and program acceptability (ie, perception that the program is agreeable, palatable or satisfactory). Understanding reasons for participation and program acceptability can contribute to future research and implementation efforts of programs for UAU in PCPs.<sup>36,37</sup> The program under examination in this study is Project ReDUCE (**R**educing **D**rinking **U**sing **C**ollaborative **E**fforts), a collaborative care model to treat UAU in a PCP. Three research questions guided this study: (1) What were the reasons patients participated in ReDUCE? (2) What were patients' experiences in ReDUCE? and (3) How acceptable did patients find ReDUCE?

## Method

### Design, Setting, and Study Procedures

The current study was conducted as part of a larger project to develop and test Project ReDUCE, a collaborative care model to treat UAU in a PCP. The data examined in this study was collected as part of an initial feasibility phase in which the collaborative care model was adapted and implemented in a PCP.

The study was conducted in a Level 3 Patient-Centered Medical Home (PCMH) in the Northeast that has approximately 32,000 visits per year. The practice has an ongoing screening, brief intervention and referral to treatment (SBIRT)<sup>38</sup> program in which every patient is screened for risky alcohol use using the AUDIT-C screening tool<sup>39</sup> at each visit and connected with a PCP-based health coach for follow-up if they screen positive. Study recruitment was conducted predominantly via the SBIRT program such that during SBIRT procedures, PCMH health coaches identified potential participants and offered them the opportunity to participate. In addition, patients identified by practice physicians during routine visits could be referred. If interested, participants were consented using a written consent procedure and then completed additional measures to further determine eligibility and alcohol use severity (eg, Timeline Follow Back for the past 60 days, the Composite International Diagnostic Interview-Alcohol Module). Patients were eligible to participate if they regularly received care at that PCMH. They also needed to (1) have estimated weekly alcohol consumption of greater than 14 drinks (men) or 7 drinks (women) or at least 1 binge drinking day per week; (2) demonstrate observable fluency in English; and (3) have no recognizable cognitive impairment. Twenty-six patients were enrolled; 2 were lost to follow-up, leaving 24 participants in this study.

### Core Program Elements

Eligible participants were initially classified into 1 of 4 groups that determined the treatment options they would be offered (see Table 1): (1) At-Risk Drinkers (AR); (2)

**Table 1.** Drinking Classification and Treatment Offering.

Drinking classification	Criteria	Core elements of treatment <sup>a</sup>
At-risk (AR)	> 14 SD men or 7 drinks/women per week; or binge drinking defined as > 5 SD/men or > 4 SD/women per day	SBIRT plus ongoing monitoring by PCP
Problem (PD)	≥ 24 SD/men or ≥ 14 SD/women per week; regular binge drinking defined as > 5 SD/men or > 4 SD/women per day at least > 3 times per month	Choice of either 4 sessions of Motivational Enhancement Therapy or 12 weeks of daily naltrexone with medication management
AUD with physiological withdrawal (AD-W)	Met DSM-5 AUD criteria with physiological withdrawal	Offered outpatient detoxification as part of standard PCP care, unless contraindicated. Upon completion of detoxification, offered 12 weeks of daily naltrexone plus medication management or 12 weekly sessions of Modified Behavioral Self-Control Therapy
AUD with chronic or complex presentation (AD-CMPLX)	Met DSM-5 criteria AUD, reported inpatient treatment for alcohol use and/or other mental health issues within last 5 years, experience social problems (ie, unstable housing) suggesting need for more intensive treatment, or deemed not appropriate for outpatient detoxification and in need higher level of care	Referred to specialty SUD treatment based on routine SBIRT protocol and ongoing monitoring by PCP

Abbreviations: SD, standard drinks.

<sup>a</sup>All received daily text message drinking tracking surveys.

Problem Drinkers (PD); (3) Alcohol Use Disorder with Physiological Withdrawal Drinkers (AD-W); or (4) Alcohol Use Disorder with Chronic or Complex Presentation (AD-CMPLX). Classifications were based on guidance from the National Institute on Alcohol Abuse and Alcoholism (NIAAA), Substance Abuse and Mental Health Services Administration (SAMHSA), the American Society of Addiction Medicine (ASAM) Levels of Care, and our team's previous research.<sup>40-42</sup> A core element of the program was that services were integrated within the PCP and PCP providers monitored patients' progress during regular PCP visits. Other core elements included: daily text message drinking tracking surveys that took about 1-minute to complete, medication management using Naltrexone, and therapy sessions (Motivational Enhancement Therapy, Modified Behavioral Self-Control Therapy) as outlined in Table 1. Treatment options were based on risk stratification with increasing services as risk level increased. Patient preference regarding therapy versus medication management with Naltrexone, as appropriate, was the main driver of initial intervention for PD and AD-W patients. All participants were offered stepped-up care after discussion with the clinician about best options if they were deemed to be a treatment non-responder at their week 4 treatment appointment; participants also had a great deal of input and could choose a different treatment (eg, therapy instead of medication) in consultation with their treatment team at any point. Participants who chose behavioral interventions were able

to select whether they wanted to attend these appointments in the PCP or via televideo. Participants were also able to choose whether abstinence or reducing/moderating drinking was their goal.

Specifically relevant to the data being examined in this paper, during the final assessment at study week 12, a research assistant administered a patient feedback interview, which consisted of semi-structured questions that took 15 to 20 minutes. These interviews were conducted mainly in person in the PCMH in a private room and patients received a \$20 incentive for the entire final assessment. The research assistant (MR), a trained interviewer and master's-level mental health clinician took detailed notes during these interviews, and these notes made up the data for this study. IRB approval for this study was given by the Feinstein Institutes for Medical Research at Northwell Health Institutional Review Board.

### Participants and Measures

Participants included 24 primary care patients (see Table 2). To address research question 1, we adapted questions from a previous qualitative study on barriers to SUD treatment.<sup>43</sup> These questions included: What is the main reason you agreed to be part of the program? Why did you feel that now was a good time to receive treatment? For research question 2 and 3, we adapted questions from a study on integrated medical and SUD treatment:<sup>44</sup> How would you describe

**Table 2.** Participant Demographics and Drinking Patterns.

	Mean (SD) or %	Mean (SD) or %	Mean (SD) or %
	Men (n = 13)	Women (n = 11)	Total (n = 24)
Age	52.53 (14.01)	52.45 (19.24)	52.50 (16.23)
Male gender	100	0	54
Hispanic/Latino	31	27	29
Race			
Asian/Pacific Islander	23	0	13
Black	8	0	4
White	38	73	54
Other/multi-racial	31	27	29
Education			
Less than high school	8	9	8
High school or GED	38	9	26
Some college	23	46	33
Bachelor or graduate degree	31	36	33
Drinking classification			
At-risk	15	27	21
Problem drinking	8	64	33
Alcohol dependence-withdrawal	69	0	38
Alcohol dependence-complex	8	9	8
Baseline AUDIT <sup>a</sup> score	17.17 (6.32)	13.00 (5.39)	15.17 (6.14)
Baseline drinks per week	27.67 (19.64)	13.36 (4.65)	20.83 (16.00)

<sup>a</sup>AUDIT, alcohol use disorders identification test (range = 0-40; 8-14 = hazardous or harmful alcohol use; 15+ indicates the likelihood of alcohol use disorder).

your experience in the program? Did the program help reduce your drinking? What aspects of the program did you like the most? Least? The interview guide included probes to gain information about specific aspects of the program (eg, receiving alcohol treatment in their regular doctor's office).

### Analysis

The coding team was made up of the first author (MO), a health services research scientist and PhD social psychologist with training in qualitative methods, and a research associate with a Master's in Public Health with qualitative research experience (CM). Both coders were not affiliated with the PCMH or health system in which the study took place. Atlas.ti was used to organize, manage, and examine the data. Conventional qualitative content analysis<sup>45</sup> was used to analyze the data following the process outlined in Erlingsson and Brysiewicz.<sup>46</sup> First, the coders read and re-read each interview to get a sense of the whole dataset and to document initial impressions. Second, the first author broke text down into meaning units, the smallest text unit that contains insights the researcher is interested.<sup>47</sup> Third, codes and definitions were developed into a codebook. Codes were developed inductively based on the team's initial read of the notes from all the interviews and

open-coding of 3 interviews. Fourth, each interview was coded by both coders. The codebook was continually refined throughout the coding process through discussion and consensus; the coding team met weekly to discuss coding impressions, compare codes, and refine the codebook. Reliability between the 2 coders averaged 89%. Any coding disagreements were resolved through discussion. In the fifth and final step, once coding was completed, codes were sorted into categories.


### Results

We identified 331 meaning units in the data. The meaning units were coded using 28 different codes that were organized into 7 categories (see Figure 1). In the results below, drinking classification of the participant is included with representative quotes.

#### *What Were the Reasons Patients Participated in ReDUCE?*

**External factors.** Patients cited a number of external factors that influenced their interest in engaging in ReDUCE. First, participants cited concerns from important others, such as family. For example, one participant engaged in the program to "ease concern from mother about drinking" (AR).





**Figure 1.** Qualitative categories.

Another participant said “family-daughter, wife, mother all worried” (AD-W).

Second, participants indicated that healthcare providers within the PCP weighed into their decision to engage. Patients suggested that their primary care physicians often connected their alcohol use to their health when suggesting the program. A patient said, I “came out of the hospital—doctor connected health issues to alcohol and ReDUCE presented itself, so wanted to go for it” (AR). And another patient said, “Dr. [x] presented it and suggested at recent visit because I was using drinking to relax and associated it with some health issues” (PD). Health coaches were also an influential provider within the PCP. One patient said, the “opportunity presented itself when the health coach came in during doctor visit” (AD-W). Another said, the “health coach approached me during visit in a caring, energetic way. The health coach did not sell the program, instead presented in a way that felt right place-right time” (AD-W). The program was often offered opportunistically during patients’ regular doctor visit, which participants seemed to welcome. As a participant stated, “when this was offered during my visit I felt very happy” (PD). Another said that during their regular doctor visit the program “came my way, was honest with health coach about drinking” (PD).

Other external factors included affordability and the desire to help others. One participant said, I am participating “for my addiction, it’s hard to find affordable rehab” (AD-W). Another said, I “can see it costing less to treat in primary care versus rehab” (PD). The desire to help others seemed to stem from participating in a research study that may eventually benefit others. One participated “to help self and to help others—we all benefit” (AD-CMPLX).

*Internal factors.* Internal factors also played into participants’ decision to participate. First, participants’ concerns about their own drinking as well as being in a stage in which they were already contemplating quitting or cutting down were key factors. For example, one participant said they were “drinking too much and worried it was going to get really out of control” (PD). Another said, I “needed the help. I have been saying ‘How can I stop!’ but felt hopeless—did not know how to stop” (PD).

A second internal factor was health concerns participants had connected to their drinking. Some had specific concerns due to a co-morbid chronic disease, while others had more general desires to become healthier or felt they needed to make changes because their future health was at risk. One participant said, “I felt I was reaching a point with my health. . .starting to feel really concerned about where I was going to go if I continued on that path” (AD-W). Another said they participated to “understand how drinking is bad for my health, especially diabetes—did not realize alcohol was bad for diabetes” (AR). An additional participant indicated, “I know it affects my health, it was time to make changes for my health, wanted to work on both drinking and eating” (AD-W).

### *What Were Patients’ Experiences in ReDUCE?*

*Affective responses.* Affective responses (ie, emotions, moods and feelings) to the program seemed to shape patients’ experiences. For example, when asked about their experience the majority of patients had generally positive feelings, such as “enlightening!” (AD-CMPLX), “pleasant and fun” (AR), “glad I did this, really good experience” (PD), and “positive, no judgment, eye opening and empowering” (PD). Participants’ feelings were also more specifically related to the flexibility of the program. For example, one patient said the program was “not a burden, [it was] accessible to fit into life” (PD). Another said, the program “felt structured but flexible” (PD). When participants did not have a positive experience, it was typically framed as a problem with specific patient fit. For example, a patient said the program was “supportive, pleasant, but not right for me” (AD-W). Another said their experience was, “not great, not for me” (PD).

*Therapeutic benefits.* Participants experienced therapeutic benefits from ReDUCE. First, participants reported gaining new insights and perspectives about their drinking. For example, one patient said “by sitting down and talking it out, I realized how much alcohol was permeating all areas of life” (PD). Another noted that it was “harder to reduce than anticipated, [the program] made me more aware and want to continue to try harder” (AR).

Second, participants also experienced feelings of support from program staff. For example, a patient said the

“health coach and therapist both used a compassionate, realistic approach and were exemplary; they delivered messages in a great way” (AD-W). Another patient said that the program was “easy going, no pressure, and I appreciated the support” (PD).

### *How Acceptable was ReDUCE to Patients?*

**Core program elements.** Acceptability was driven by core program elements, such as daily text message tracking surveys, integration within the PCP, medication and therapy sessions. For example, the majority of participants found the daily text tracking survey helpful. One participant said, “text messages in the morning remind me not to drink” (AD-W). When participants did not drink, these seemed especially influential, as one participant noted they were a “huge help—I loved entering zeros” (AD-W). However, a few participants found these cumbersome or non-therapeutic. For example a participant said “daily was a little excessive and it was not user friendly” (AR). Another said the text surveys were “not helpful because they triggered thoughts to drink” (AD-W).

Other core program elements, such as its integration within the PCP and AUD medication, were related to acceptability. Many participants liked that the program was integrated into the PCP due to the convenience and affordability. One patient said that receiving alcohol care in their regular doctor’s office made them able to “make appointments for diabetes and come in for alcohol treatment at the same time” (AD-W). Another said they liked receiving care in their regular doctor’s office because they “didn’t want to go to meetings or regular treatment” (AD-W). Negative comments related to integration in the PCP were typically related to transportation or scheduling difficulties due to patient personal circumstances.

Patients also discussed AUD medications and therapy, 2 other core program components. One patient said that medications “helped cut cravings and reduce [drinking]” (AD-W). Another said they “definitely liked being offered meds” (PD). Only 1 participant had a negative comment about the medications due to side effects, “medication made me feel weird, hazy, cloudy, including feeling down” (AD-W). Therapy was an important factor in patients’ views on acceptability with the program and reactions to the therapy component were somewhat mixed. For example, 1 patient said “I was able to open up about my struggles” (PD). Another said, “talking about it and being open about it was 90% helpful” (AD-W). However, other patients were not as positive, perhaps because the therapy portion of the program was not what they expected in a setting like a PCP. For example, a patient said therapy “felt like buying time, treading water, sessions felt like nothing” and that there were “too few sessions in a sterile therapy environment—did not connect with the therapy techniques” (PD).

**Impact on drinking cognition and behavior.** Acceptability was also related to impacts on drinking cognition, such as gains in information or awareness. For example, a participant said, “I became more conscious of drinking in general. I began thinking about drinking differently after sessions and tracking more. I became more aware and this changed how I typically drink” (PD). Another participant said that the program was helpful because of it “making me self-aware of drinking over the course of the week—being conscious about drinking. Before this, I just drank, didn’t think anything of it” (PD). Another participant said that they “realized they had strength within themselves to change” (PD). Several participants mentioned that receiving information about healthy drinking guidelines was useful. The majority of participants responded affirmatively when asked whether the program helped them reduce their drinking.

**Therapeutic approach.** Finally, acceptability with the program was influenced by the program’s therapeutic approaches. One approach used in ReDUCE was being patient-centered by tailoring the program to patient needs and offering some patients a choice between medication and therapy. One patient said, I “liked having the choice, I was not locked into one versus the other, glad I chose therapy first and now considering medications” (AD-W). While another said, I “would not have done it if meds were mandatory” (PD). Flexible modalities (eg, telemedicine vs in-person) also contributed to the view of being patient-centered.

Another key therapeutic approach was offering the opportunity to cut down or moderate drinking rather than mandating abstinence-only goals. One participant said they, “definitely would not have joined if told to quit” (PD). Another agreed that it was “helpful to have the option to reduce, less rigid and easier to take on” (AD-W). Another said of this approach: “loved this—still working toward reducing and now want to quit” (AR). A few participants found this approach not structured enough, as 1 patient mentioned moderation was “not helpful, it enabled drinking when shouldn’t have been” (AD-W).

Therapeutic alliances with providers were important to patient views of program acceptability. One patient noted that “meeting with the therapist and health coach kept me going; connection with staff chemistry—this is most important” (AD-CMPLX). Another participant noted that “meeting with [health coach and therapist] was very helpful; was not talking with anyone and finally opened up about drinking” (PD). Another patient stated that interacting with the staff was “very supportive, comfortable—appreciated staff and what was trying to be done” (AD-W).

## **Discussion**

In this study, we examined perspectives and experiences of patients who participated in an integrated care program for

UAU within a PCP by focusing on reasons for participation as well as perceived acceptability of the program. Generally, we found that participants found the program acceptable and that they felt it helped them decrease their drinking due to the core elements of the program, perceived cognitive and behavioral improvements, and therapeutic approaches utilized. Patients experienced affective and therapeutic benefits and we found that patients initially engaged in the program for both internal and external reasons.

Internal factors such as concern for health and external influences such as family or healthcare practitioner concern contributed to patients' interest in engaging in the program. Findings related to health concerns and pressure from family members are supported by previous qualitative research on older adults' perspectives on alcohol treatment in PCPs.<sup>48</sup> However, what our study highlights that other studies have not is the importance of the PCP provider in encouraging participants to engage in alcohol-related care, especially during routine PCP visits. By framing the conversation with patients as one about health, PCPs implementing AUD care programs could successfully engage patients into integrated programs. PCP providers should receive training on the linkages between alcohol and health and understand the importance of their role as well as the importance of family members in encouraging patients to engage in care. Offering a program opportunistically during routine medical visits can be both an important aspect of care and a challenge given that patients with UAU may not be aware of the problem or ready to address it. Multiple touchpoints with PCP providers may be required to fully engage patients; however, this repeated engagement is especially well suited to the ongoing relationship between PCP physicians and their patients. Future research should examine the touchpoints needed to fully engage patients as well as provider experiences in this ongoing engagement.

In SUD treatment broadly, patient satisfaction and acceptability has been linked to improved patient retention and outcomes.<sup>49-51</sup> Findings from this study suggested acceptability was generally high and that it was affected by 3 main factors: elements of the multi-faceted program, impact on drinking behavior and cognition, and therapeutic approaches employed in this program.

The ReDUCE program offered a variety of elements integrated into the PCP (daily drinking tracking surveys, therapy, medication) and patients seemed to weigh the importance of each of these depending on their individual needs, goals, and current drinking pattern. Related specifically to AUD medication, previous research suggests that patients generally appreciate having the option to discuss medication with PCPs,<sup>52</sup> but that they may need multiple therapy sessions prior to initiating.<sup>9</sup> This was reflected by some patients in this study who said they would not have participated if medication was the only option or that they were considering medications only after receiving therapy.

Further, a few patients did not find the daily text message drinking tracking surveys helpful because they felt they potentially triggered thoughts to drink or were excessive. Others responded positively because the messages served as a reminder to not drink or were positively reinforcing when patients were able to enter zeros. Given the large interest in technology-based care for alcohol and other substance use<sup>53,54</sup> and in particular text-messaging interventions,<sup>55</sup> further studies should examine patient experience and implementation of such technology in primary care and other healthcare settings. Similarly, not all patients felt that the therapy component was helpful for them or what they were expecting. Ensuring that patients fully understand therapy goals and what to expect may improve acceptability. Additional research should clarify patient expectations about behavioral health therapy within PCPs. Overall, our findings suggest that a rigid, one-size fits all program may not be attractive to patients. Therefore, it may be important in PCP-based integrated programs to offer a flexible menu of options to patients and recognize that patient preference could change over the course of their treatment.

The program's therapeutic approach also influenced acceptability. First, being patient-centered was valued, including allowing patients to explore the menu of options available to them given their individual preferences and needs, providing flexibility in terms of interaction with program staff (eg, telehealth), and choice of moderation vs. abstinence goal. Patient-centered care is a key element of SUD treatment and has been found to improve patient outcomes,<sup>56,57</sup> and previous research has found that patients value being involved in decisions about their treatment and having autonomy.<sup>25</sup> Therefore, programs should include training and support to program staff on providing patient-centered care. Second, therapeutic alliance between patient and provider was mentioned frequently by patients when discussing acceptability as well as engagement and experience in the program. Therapeutic alliance has long been an important predictor of outcomes in AUD care.<sup>58-60</sup> Previous quantitative research has found that patients have more positive attitudes toward alcohol-related interventions in PCPs if they are generally satisfied with their PCP provider.<sup>61</sup> Qualitative research also highlights the importance of the patient-provider relationship in helping patients address their alcohol use in the PCP setting.<sup>62</sup> The importance of supportive providers who are comfortable providing AUD care in PCPs cannot be underestimated. Future research on AUD programs in PCPs would benefit from a mixed-methods examination of patient experience, therapeutic alliance, and drinking outcomes to further understand these dynamics. Patients also discussed acceptability in terms of receiving important information and tools that increased their awareness about the impacts of their drinking. This finding is supported by previous research in AUD specialty care that patients felt being offered advice, tools, and resources



improved their motivation to change their drinking and extends this finding to PCP settings.<sup>25</sup> Therefore, PCP providers must be equipped with information and tools to assist patients in achieving their drinking goals.

We did not find any high-level differences between men and women or between drinking classification groups. However, it should be noted that there was not an even distribution of men and women within the drinking classification groups. Further, men reported higher weekly drinking levels and AUDIT scores at baseline than women. Given the small sample size, we were not able to fully explore gender and drinking classification differences in experience and engagement in treatment services for UAU in primary care; future research should examine this.

While there were important findings related to patient engagement, experience, and acceptability, there are limitations. At the PCP's request, we did not audio record interviews due to their concerns that recording was not part of standard practice in the PCP and that it may cause too much discomfort for patients given that this was the patient's regular PCP. As a result, important context or information may not have been captured in our findings. To balance the ethical commitment to participants and the PCP and methodological rigor, the interviewer took detailed notes mainly by writing down direct quotes from participants. It is possible the interviewer missed words from some of the quotes, reflecting notes rather than full direct quotes. While audio recording and transcribing is a suggested practice to support reliability in qualitative research, some studies have justified that audio recording can negatively impact the quality of interviews and have shown that note-taking can be rigorous as well.<sup>63-66</sup> In addition, the interviewer served as one of several part-time health coaches in the PCMH where the study was conducted. Therefore, she may have known some of the participants prior to the study in this role, leading to the possibility that this introduced bias into the interviews. However, she was a trained interviewer and research assistant who was aware of and made efforts to minimize this potential bias. This study was conducted among patients who agreed to participate in the program and consented to study data collection, it does not provide insight into why people chose not to participate or barriers to engagement outside of a research study.

There were also strengths in that we took several measures to demonstrate study trustworthiness.<sup>47</sup> For example, 2 people coded each transcript. A detailed codebook was maintained and all decisions and activities throughout the analysis were carefully documented using an audit trail. While we did not return the interview notes or findings to participants to review, the interviewer, who did not take part in the coding and analysis, was presented with the results during analysis to ensure that we captured patient's thoughts accurately. Finally, we included all patients who participated in the ReDUCE feasibility study in the qualitative interview portion of the study, rather than a sub-sample.

The findings of this study illuminate factors important in reaching patients in AUD care programs integrated in PCPs as well as factors that may influence patient experience and acceptability. One recurring theme was the importance of PCP providers in both engaging patients as well as providing support and building a therapeutic alliance. In addition, offering flexible and comprehensive programs with multiple elements and both abstinence and moderation goals could improve patient acceptability in AUD care programs in PCPs. PCPs will need to be thoughtful about the resources needed to implement programs in terms of staffing, training, and support.

### Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by the National Institute on Alcohol Abuse and Alcoholism [Grant #: R34AA024589]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health/National Institute on Alcohol Abuse and Alcoholism.

### ORCID iD

Megan A. O'Grady  <https://orcid.org/0000-0002-1121-734X>

### References

1. White AM, Castle I-JP, Hingson RW, Powell PA. Using death certificates to explore changes in alcohol-related mortality in the United States, 1999 to 2017. *Alcohol: Clin Exp Res.* 2020;44:178-187.
2. Substance Abuse and Mental Health Services Administration. *Treatment Episode Data Set (TEDS): 2005-2015. State Admissions to Substance Abuse Treatment Services.* SAMHSA; 2017.
3. 2018 National Survey on Drug Use and Health (NSDUH). Table 5.4A—Alcohol Use Disorder in Past Year among Persons Aged 12 or Older, by Age Group and Demographic Characteristics: Numbers in Thousands, 2017 and 2018; 2018.
4. Hepner KA, Paddock SM, Watkins KE, et al. Rates and impact of adherence to recommended care for unhealthy alcohol use. *J Gen Intern Med.* 2019;34:256-263.
5. United States Preventative Service Task Force. Screening and behavioral counseling interventions to reduce unhealthy alcohol use in adolescents and adults: US preventive services task force recommendation Statement. *JAMA.* 2018;320:1899-1909.
6. McKnight-Eily LR, Okoro CA, Mejia R, et al. Screening for excessive alcohol use and brief counseling of adults - 17 states and the District of Columbia, 2014. *MMWR Morb Mortal Wkly Rep.* 2017;66:313-319.
7. Saitz R, ed. *Addressing Unhealthy Alcohol Use in Primary Care.* Springer-Verlag; 2013.



8. Lock CA. Alcohol and brief intervention in primary health care: what do patients think? *Prim Health Care Res Dev*. 2004;5:162-178.
9. Ober AJ, Watkins KE, McCullough CM, Setodji CM, Osilla K, Hunter SB. Patient predictors of substance use disorder treatment initiation in primary care. *J Subst Abuse Treat*. 2018;90:64-72.
10. Wallhed Finn S, Hammarberg A, Andreasson S. Treatment for alcohol dependence in primary care compared to outpatient specialist treatment—a randomized controlled trial. *Alcohol Alcohol*. 2018;53:376-385.
11. Kamstrup-Larsen N, Broholm-Jørgensen M, Dalton SO, Larsen LB, Thomsen JL, Tolstrup JS. Why do general practitioners not refer patients to behaviour-change programmes after preventive health checks? A mixed-method study. *BMC Fam Pract*. 2019;20:135.
12. Álvarez-Bueno C, Rodríguez-Martín B, García-Ortiz L, Gómez-Marcos MÁ, Martínez-Vizcaíno V. Effectiveness of brief interventions in primary health care settings to decrease alcohol consumption by adult non-dependent drinkers: a systematic review of systematic reviews. *Prev Med*. 2015;76:S33-S38.
13. Kaner EF, Beyer FR, Muirhead C, et al. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev*. 2018;2:Cd004148.
14. Bradley KA, Bobb JF, Ludman EJ, et al. Alcohol-related nurse care management in primary care: a randomized clinical trial. *JAMA Intern Med*. 2018;178:613-621.
15. Watkins KE, Ober AJ, Lamp K, et al. Collaborative care for opioid and alcohol use disorders in primary care: the SUMMIT randomized clinical trial. *JAMA Intern Med*. 2017;177:1480-1488.
16. Oslin DW, Lynch KG, Maisto SA, et al. A randomized clinical trial of alcohol care management delivered in Department of Veterans Affairs primary care clinics versus specialty addiction treatment. *J Gen Intern Med*. 2014;29:162-168.
17. Saitz R, Cheng DM, Winter M, et al. Chronic care management for dependence on alcohol and other drugs: the AHEAD randomized trial. *JAMA*. 2013;310:1156-1167.
18. Glass JE, Bobb JF, Lee AK, et al. Study protocol: a cluster-randomized trial implementing Sustained Patient-centered Alcohol-related Care (SPARC trial). *Implement Sci*. 2018;13:108.
19. Kline-Simon AH, Stumbo SP, Campbell CI, et al. Patient characteristics associated with treatment initiation and engagement among individuals diagnosed with alcohol and other drug use disorders in emergency department and primary care settings. *Subst Abuse*. 2019;40:278-284.
20. Buck JA. The looming expansion and transformation of public substance abuse treatment under the Affordable Care Act. *Health Aff (Millwood)*. 2011;30:1402-1410.
21. Briggs CJ, Capdegelle P, Garner P. Strategies for integrating primary health services in middle- and low-income countries: effects on performance, costs and patient outcomes. *Cochrane Database Syst Rev*. 2001;4:Cd003318.
22. Rawaf S, Amati F, McDonald A, Majeed A, Dubois E. Implementation and evaluation of patient-centred care in experimental studies from 2000-2010: systematic review. *Int J Pers Cent Med*. 2011;1:10.
23. Ouwens M, Wollersheim H, Hermens R, Hulscher M, Grol R. Integrated care programmes for chronically ill patients: a review of systematic reviews. *Int J Qual Health Care*. 2005;17:141-146.
24. Kirouac M, Witkiewitz K, Donovan DM. Client evaluation of treatment for alcohol use disorder in COMBINE. *J Subst Abuse Treat*. 2016;67:38-43.
25. McCallum SL, Mikocka-Walus AA, Gaughwin MD, Andrews JM, Turnbull DA. 'I'm a sick person, not a bad person': patient experiences of treatments for alcohol use disorders. *Health Expect*. 2016;19:828-841.
26. Williams EC, Matson TE, Harris AHS. Strategies to increase implementation of pharmacotherapy for alcohol use disorders: a structured review of care delivery and implementation interventions. *Addict Sci Clin Pract*. 2019;14:6.
27. Storholm ED, Ober AJ, Hunter SB, et al. Barriers to integrating the continuum of care for opioid and alcohol use disorders in primary care: a qualitative longitudinal study. *J Subst Abuse Treat*. 2017;83:45-54.
28. Rahm AK, Boggs JM, Martin C, et al. Facilitators and barriers to implementing Screening, Brief Intervention, and Referral to Treatment (SBIRT) in primary care in integrated health care settings. *Subst Abuse*. 2015;36:281-288.
29. Nygaard P, Aasland OG. Barriers to implementing screening and brief interventions in general practice: findings from a qualitative study in Norway†. *Alcohol Alcohol*. 2010;46:52-60.
30. Williams EC, Achtmeyer CE, Young JP, et al. Barriers to and facilitators of alcohol use disorder pharmacotherapy in primary care: a qualitative study in five VA clinics. *J Gen Intern Med*. 2018;33:258-267.
31. Hutchings D, Cassidy P, Dallolio E, Pearson P, Heather N, Kaner E. Implementing screening and brief alcohol interventions in primary care: views from both sides of the consultation. *Prim Health Care Res Dev*. 2006;7:221-229.
32. Tam CW, Leong LH, Zwar N. Let's listen to patients' and GPs' perspectives on alcohol-screening research. *Aust Fam Physician*. 2015;44:427-428.
33. O'Donnell A, Hanratty B, Schulte B, Kaner E. Patients' experiences of alcohol screening and advice in primary care: a qualitative study. *BMC Fam Pract*. 2020;21:68.
34. Khadjesari Z, Stevenson F, Toner P, Linke S, Milward J, Murray E. 'I'm not a real boozier': a qualitative study of primary care patients' views on drinking and its consequences. *J Public Health (Oxf)*. 2019;41:e185-e191.
35. Cucciare MA, Lewis ET, Hoggatt KJ, et al. Factors affecting women's disclosure of alcohol misuse in primary care: a qualitative study with U.S. military veterans. *Womens Health Issues*. 2016;26:232-239.
36. Holtrop JS, Rabin BA, Glasgow RE. Qualitative approaches to use of the RE-AIM framework: rationale and methods. *BMC Health Serv Res*. 2018;18:177.
37. Proctor E, Silmere H, Raghavan R, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health*. 2011;38:65-76.
38. O'Grady MA, Kapoor S, Kwon N, et al. Substance use screening and brief intervention: evaluation of patient and implementation differences between primary care and emergency department settings. *J Eval Clin Pract*. 2019;25:441-447.

39. Bush K, Kivlahan DR, McDonnell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol use disorders identification test. *Arch Intern Med.* 1998;158:1789-1795.
40. Morgenstern J, Irwin TW, Wainberg ML, et al. A randomized controlled trial of goal choice interventions for alcohol use disorders among men who have sex with men. *J Consult Clin Psychol.* 2007;75:72-84.
41. Morgenstern J, Kuerbis AN, Chen AC, Kahler CW, Bux DA Jr, Kranzler HR. A randomized clinical trial of naltrexone and behavioral therapy for problem drinking men who have sex with men. *J Consult Clin Psychol.* 2012;80:863-875.
42. Morgenstern J, Kuerbis A, Houser J, et al. Dismantling motivational interviewing: effects on initiation of behavior change among problem drinkers seeking treatment. *Psychol Addict Behav.* 2017;31:751-762.
43. Jackson A, Shannon L. Barriers to receiving substance abuse treatment among rural pregnant women in Kentucky. *Matern Child Health J.* 2012;16:1762-1770.
44. Drainoni M-L, Farrell C, Sorensen-Alawad A, Palmisano JN, Chaisson C, Walley AY. Patient perspectives of an integrated program of medical care and substance use treatment. *AIDS Patient Care STDs.* 2014;28:71-81.
45. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res.* 2005;15:1277-1288.
46. Erlingsson C, Brysiewicz P. A hands-on guide to doing content analysis. *Afr J Emergency Med.* 2017;7:93-99.
47. Bengtsson M. How to plan and perform a qualitative study using content analysis. *NursingPlus Open.* 2016;2:8-14.
48. Haighton C, Wilson G, Ling J, McCabe K, Crosland A, Kaner E. A qualitative study of service provision for alcohol related health issues in mid to later life. *PLoS One.* 2016;8-14.
49. Kelly SM, O'Grady KE, Brown BS, Mitchell SG, Schwartz RP. The role of patient satisfaction in methadone treatment. *Am J Drug Alcohol Abuse.* 2010;36:150-154.
50. Kendra MS, Weingardt KR, Cucciare MA, Timko C. Satisfaction with substance use treatment and 12-step groups predicts outcomes. *Addict Behav.* 2015;40:27-32.
51. Zhiwei Z, Gerstein DR, Friedmann PD. Patient satisfaction and sustained outcomes of drug abuse treatment. *J Health Psychol.* 2008;13:388-400.
52. Haley SJ, Pinsker EA, Gerould H, Wisdom JP, Hagedorn HJ. Patient perspectives on alcohol use disorder pharmacotherapy and integration of treatment into primary care settings. *Subst Abuse.* 2019;40:501-509.
53. Kiluk BD, Ray LA, Walthers J, Bernstein M, Tonigan JS, Magill M. Technology-delivered cognitive-behavioral interventions for alcohol use: a meta-analysis. *Alcohol Clin Exp Res.* 2019;43:2285-2295.
54. Carreiro S, Newcomb M, Leach R, Ostrowski S, Boudreaux ED, Amante D. Current reporting of usability and impact of mHealth interventions for substance use disorder: a systematic review. *Drug Alcohol Depend.* 2020;215:108201.
55. Bendtsen M. Text messaging interventions for reducing alcohol consumption among harmful and hazardous drinkers: protocol for a systematic review and meta-analysis. *JMIR Res Protoc.* 2019;8:e12898.
56. Davis EL, Kelly PJ, Deane FP, et al. The relationship between patient-centered care and outcomes in specialist drug and alcohol treatment: a systematic literature review. *Subst Abuse.* 2020;41:216-231.
57. Barrio P, Gual A. Patient-centered care interventions for the management of alcohol use disorders: a systematic review of randomized controlled trials. *Patient Prefer Adherence.* 2016;10:1823-1845.
58. Barber JP, Luborsky L, Gallop R, et al. Therapeutic alliance as a predictor of outcome and retention in the National Institute on Drug Abuse Collaborative Cocaine Treatment study. *J Consult Clin Psychol.* 2001;69:119-124.
59. Connors GJ, Carroll KM, DiClemente CC, Longabaugh R, Donovan DM. The therapeutic alliance and its relationship to alcoholism treatment participation and outcome. *J Consult Clin Psychol.* 1997;65:588.
60. Meier PS, Barrowclough C, Donmall MC. The role of the therapeutic alliance in the treatment of substance misuse: a critical review of the literature. *Addiction.* 2005;100:304-316.
61. Fankhaenel T, Samos F, Luck-Sikorski C, Thiel C, Klement A, Frese T. Patient satisfaction as a moderator of risky alcohol consumers' attitude towards screening and brief intervention: a cross sectional survey. *Alcohol Alcohol.* 2018;53:403-407.
62. Field CA, Klimas J, Barry J, et al. Problem alcohol use among problem drug users in primary care: a qualitative study of what patients think about screening and treatment. *BMC Fam Pract.* 2013;14:98.
63. Hamza DM, Greenshaw AJ, Hamza SM, Silverstone PH. Qualitative findings from administrators of the EMPATHY (Empowering a multimodal pathway toward healthy youth) programme using the SBIRT framework. *Br J Guid Counsell.* Published online November 3, 2019. doi:10.1080/03069885.2019.1686121
64. Hayes T, Mattimoe R. Chapter 21 - To tape or not to tape: reflections on methods of data collection. In: Humphrey C, Lee B, eds. *The Real Life Guide to Accounting Research.* Elsevier; 2004: 359-372.
65. Clausen AS. The individually focused interview: methodological quality without transcription of audio recordings. *Qual Rep.* 2012;17:1-17.
66. Al-Yateem N. The effect of interview recording on quality of data obtained: a methodological reflection. *Nurse Res.* 2012;19:31-35.