

## Review Article

# Lipid Rafts: Linking Alzheimer's Amyloid- $\beta$ Production, Aggregation, and Toxicity at Neuronal Membranes

Jo V. Rushworth<sup>1</sup> and Nigel M. Hooper<sup>1,2</sup>

<sup>1</sup>*Institute of Molecular and Cellular Biology, Astbury Centre for Structural Molecular Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK*

<sup>2</sup>*Institute of Molecular and Cellular Biology, Faculty of Biological Sciences, LIGHT Laboratories, Clarendon Way, University of Leeds, Leeds LS2 9JT, UK*

Correspondence should be addressed to Nigel M. Hooper, n.m.hooper@leeds.ac.uk

Received 14 October 2010; Accepted 3 November 2010

Academic Editor: Katsuhiko Yanagisawa

Copyright © 2011 J. V. Rushworth and N. M. Hooper. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lipid rafts are membrane microdomains, enriched in cholesterol and sphingolipids, into which specific subsets of proteins and lipids partition, creating cell-signalling platforms that are vital for neuronal functions. Lipid rafts play at least three crucial roles in Alzheimer's Disease (AD), namely, in promoting the generation of the amyloid- $\beta$  ( $A\beta$ ) peptide, facilitating its aggregation upon neuronal membranes to form toxic oligomers and hosting specific neuronal receptors through which the AD-related neurotoxicity and memory impairments of the  $A\beta$  oligomers are transduced. Recent evidence suggests that  $A\beta$  oligomers may exert their deleterious effects through binding to, and causing the aberrant clustering of, lipid raft proteins including the cellular prion protein and glutamate receptors. The formation of these pathogenic lipid raft-based platforms may be critical for the toxic signalling mechanisms that underlie synaptic dysfunction and neuropathology in AD.

## 1. Introduction

Alzheimer's Disease (AD) is a progressive, neurodegenerative brain disorder which affects over 37 million people worldwide with an estimated global cost of over \$600 billion in 2010 [1, 2]. AD is a growing socioeconomic and financial burden due to its strong correlation with ageing; around 1 in 3 people aged over 80 years have AD, which means that a rapid rise in AD cases is anticipated as life expectancy continues to increase. Although several therapeutics are currently available to slow disease progression, there is currently no way to halt or prevent AD [3].

AD is characterized by the presence of extracellular senile plaques and intracellular neurofibrillary tangles in the brain. The major constituents of senile plaques are the amyloid- $\beta$  ( $A\beta$ ) peptides, which are derived from the proteolytic processing of the amyloid precursor protein (APP) within lipid rafts [4]. The  $A\beta$  peptide, notably  $A\beta_{1-42}$ , is highly aggregation prone and self-assembles to form a heterogeneous mixture of oligomers and protofibrils, ultimately depositing

as fibrils in senile plaques. An accumulating body of evidence indicates that soluble  $A\beta$  oligomers, which correlate strongly with disease onset and severity, are the major neurotoxic species in AD [5–8]. Although  $A\beta$  oligomers are neurotoxic at nanomolar concentrations and cause AD-related memory deficits, the cellular mechanisms of toxicity are poorly characterised. Recently, several neuronal receptors which bind  $A\beta$  oligomers have been identified, including the cellular prion protein (PrP<sup>C</sup>) [9] and glutamate receptors [10, 11] among others. Interestingly, these receptors reside primarily within, or partition into, cholesterol-rich microdomains within the plasma membrane known as lipid rafts.

The three steps which underlie  $A\beta$  oligomer-mediated neuropathology in AD, are (1)  $A\beta$  production, (2)  $A\beta$  assembly into oligomers and (3)  $A\beta$  oligomers interacting with neuronal receptors. These steps therefore represent potential sites of therapeutic intervention in AD. Crucially, all three of these processes occur in lipid raft domains of the plasma membrane which are considered to play a key role in the development of AD [12]. In this paper, we will

outline the pivotal role that lipid rafts play in linking together the generation, self-assembly and toxicity of  $A\beta$  oligomers, which underlie the development of the neuropathology in AD. A major focus will be upon the interaction between  $A\beta$  oligomers and their putative cellular receptors.

## 2. Lipid Rafts

*2.1. Lipid Rafts as Essential Neuronal Signalling Platforms.* The multitude of different lipids and proteins within the plasma membrane were once thought to be distributed homogeneously across the entire lipid bilayer, as proposed by the fluid mosaic model in 1972 [13]. However, the plasma membrane is now known to be more akin to a sea of disordered phospholipids, in which float microdomains with distinct lipid compositions, known as lipid rafts. Lipid rafts are small (10–200 nm), heterogeneous and highly dynamic assemblies that are enriched in specific components, namely cholesterol and sphingolipids (Figure 1) [14, 15]. Biochemically, lipid rafts are defined by their relative insolubility in nonionic detergents at low temperature, conferring upon them the alternative name, detergent-resistant membranes (DRMs). Lipid rafts are also known as liquid-ordered domains because the highly saturated sphingolipid acyl chains enable closer lipid packing, and therefore more restricted lateral movement, than the mainly unsaturated acyl chains of the phospholipids in the surrounding nonraft regions of the membrane.

Functionally, lipid rafts serve to compartmentalise cellular processes by concentrating certain proteins and lipids within the same microenvironment. Lipid rafts are particularly enriched in glycosyl-phosphatidylinositol (GPI)-anchored and acylated proteins due to the preferential intercalation of the saturated acyl chains into the liquid-ordered environment [16]. Other proteins can also associate with lipid rafts either directly or through binding to other cofactors or ligands [17]. The dynamic clustering and pinching off of lipid rafts regulates the spatial and temporal assembly of signalling and trafficking molecules, forming short-lived but vital signalling platforms [17]. Lipid rafts are implicated in various essential cellular functions, including signal transduction, cell adhesion and protein/lipid sorting [18]. Of particular relevance here are cell signalling, sorting and axon guidance, as these processes are essential for neural development and synaptic plasticity [19, 20]. Crucially, neuronal lipid rafts are also required for the maintenance of dendritic spines and healthy synapses, which are vital for neural communication including learning and memory; processes which fail in AD [21]. The observation that lipid rafts are much more abundant in mature hippocampal neurons than in other cell types emphasises their physiological importance within the memory centre of the healthy brain, and may explain why hippocampal neurons are a primary target for  $A\beta$  oligomer toxicity and destruction in AD [22].

*2.2.  $A\beta$  Production Is Lipid Raft Dependent.* Lipid rafts are involved in the regulation of APP processing and the generation of the  $A\beta$  peptide which is the driving force

in AD pathology [23, 24]. For comprehensive reviews detailing the involvement of membrane rafts in AD and  $A\beta$  production, see [25–27]. The  $A\beta$  peptide is produced by the lipid raft dependent amyloidogenic processing of its precursor protein, APP (Figure 1) [4]. The amyloidogenic cleavage of full-length APP is initiated by the  $\beta$ -site APP cleaving enzyme-1 (BACE1), a transmembrane aspartic metalloprotease. A large, soluble ectodomain (sAPP $\beta$ ) is released to leave behind a membrane-anchored C-terminal fragment (C99) which retains the intact  $A\beta$  sequence. The second amyloidogenic cleavage of APP involves a  $\gamma$ -secretase complex which contains presenilin-1 or presenilin-2 (the catalytic component), presenilin enhancer-2 (PEN2), nicastrin and anterior pharynx defective-1 (APH1). The  $\gamma$ -secretase complex cleaves the remaining C99 stub to release  $A\beta$  peptides of between 39–42 residues in length, depending upon the precise cleavage site, along with the APP intracellular domain (AICD).

Although the majority of full-length APP is localised to nonraft regions of the plasma membrane, where non-amyloidogenic cleavage by the  $\alpha$ -secretases ADAM 9, 10, and 17 [28] precludes  $A\beta$  formation, a subset of both APP and BACE1 partitions into lipid rafts along with  $\gamma$ -secretase components. Both BACE1 and the  $\gamma$ -secretase subunits undergo posttranslational S-palmitoylation which aids their targeting to lipid raft domains [25]. In the case of APP, a direct interaction with cholesterol—the major component of lipid rafts—was recently identified [29]. High cholesterol increases the partitioning of APP, along with BACE1 and  $\gamma$ -secretase components, into lipid rafts [30]. A large body of evidence points towards lipid rafts being the physiological site of amyloidogenic  $A\beta$  production by BACE1 and the  $\gamma$ -secretase complex. For example, both the copatching of APP and BACE1 by cross-linking antibodies [31] and the exclusive targeting of BACE1 to lipid rafts by the addition of a GPI-anchor [32] significantly increased APP cleavage at the  $\beta$ -secretase site. Furthermore, enrichments in lipid raft components, namely cholesterol and ganglioside GM1, promote the generation of  $A\beta$  [31, 33]. All four of the  $\gamma$ -secretase subunits are also enriched and active within lipid raft fractions derived from human brain [34, 35] and lipid raft-type membranes *in vitro* [36, 37]. In the brain, the majority of  $A\beta$  is found within detergent-resistant, glycolipid-enriched rafts, along with  $\gamma$ -secretase components [38].

*2.3. Depleting Lipid Raft Components Modulates  $A\beta$  Production.* The composition of lipid rafts purified from AD brains has been shown to be abnormal, with the rafts being more ordered and more viscous [39], which implies that the modulation of lipid raft composition may present a therapeutic avenue for modulating AD-related neuropathology. This has led to a number of researchers investigating whether depleting lipid raft components could lower  $A\beta$  production and therefore prevent AD. Cholesterol, being a major component of lipid rafts and a risk factor for AD, was the obvious choice to target [40]. For a recent review of the involvement of cholesterol in AD, see [41].




FIGURE 1: Lipid rafts facilitate the production, aggregation, neuronal binding and toxicity of A $\beta$  oligomers. (a) The A $\beta$  peptide is produced by the lipid raft dependent sequential cleavage of APP, first by BACE1 and then by the  $\gamma$ -secretase complex; (b) Lipid raft components including cholesterol and sialic acid-containing gangliosides promote the aggregation of A $\beta$  to form soluble oligomers; (c) A $\beta$  oligomers bind to specific neuronal receptors within pathogenic lipid rafts, including PrP<sup>C</sup> and the NMDA and mGluR5 receptors. The resulting perturbations in neuronal function and survival underlie the memory impairments and cognitive decline which characterise Alzheimer's disease.

Cholesterol depletion has indeed been shown to reduce APP partitioning into lipid rafts which precludes its interaction with BACE1 and  $\gamma$ -secretase components, thus lowering A $\beta$  production [42]. Hypercholesterolaemia is linked to increased A $\beta$  production and deposition in the brain, both in humans [43–45] and in rodents [46–48] and is linked to an increased risk of developing AD. Cholesterol depletion also lowers A $\beta$  production in cultured cells [31] and one study showed that a 70% reduction in cholesterol in living hippocampal neurons was sufficient to completely abolish A $\beta$  production [49].

Taking this into account, cholesterol-lowering drugs known as statins have been evaluated as potential anti-AD drugs, with conflicting results [50]. Some retrospective epidemiological studies have shown that the administration of statins, which lower cholesterol levels, can reduce the incidence of dementia, including AD [51–53]. Cholesterol inhibitors can also lower A $\beta$  levels in cultured neuroblastoma cells [54]. However, other studies have shown no correlation between statin usage and dementia [55] and the effect of statins upon disease progression and cognitive decline in AD patients has been challenged [56]. Intriguingly, it was revealed recently that A $\beta$  production actually reduces cholesterol in cultured cells of neuronal origin by increasing efflux, possibly acting as a chaperone to remove excess cholesterol from the brain to the circulation [57].

Although a reduction in cholesterol may go some way towards reducing A $\beta$  levels in the brain, much longer-term epidemiological studies and clinical trials initiated before significant neuronal loss and cognitive function are apparent

are required in order to further elucidate the effects of lowering cholesterol levels upon AD onset and neuropathology. Lipid rafts contain many essential components other than cholesterol, such as sphingolipids, and it is likely that the modulation of just one factor will not completely abolish A $\beta$  production *in vivo*. It is important to remember that cholesterol metabolism in the brain is largely isolated from the rest of the body by the blood-brain barrier. As nearly all of the cholesterol in the brain is synthesised *in situ*, the modulation of cholesterol levels within neurons represents a more difficult pharmaceutical challenge and the blood-brain barrier permeability of the drugs used needs to be considered [29]. Furthermore, even if cholesterol depletion mediates a reduction in A $\beta$  levels, A $\beta$  oligomers effect neurotoxicity and memory impairments at low nanomolar concentrations [58]. Therefore, residual levels of A $\beta$  production may be sufficient for continued A $\beta$  oligomer-mediated toxicity.

### 3. Lipid Raft Components Promote A $\beta$ Oligomerisation

**3.1. A $\beta$  Oligomers Are the Major Neurotoxic Species in AD.** The A $\beta$  peptide is natively unfolded and, under certain conditions, it aggregates to form a heterogeneous mixture of soluble oligomers, protofibrils and fibrils. It was accepted for a long time that the A $\beta$  fibrils that deposit in neuritic plaques, which are observed *post mortem* in diseased brains, were responsible for neurotoxicity in AD [59]. A $\beta$  fibrils have been reported to induce neuronal dysfunction and

cell death, although fibrils are less potent neurotoxins than soluble forms of A $\beta$  [60, 61]. Interestingly, fibrils have been found to become more neurotoxic upon fragmentation [62], raising the possibility that soluble species released from fibril ends may underlie their neurotoxicity. A plethora of studies have now demonstrated that levels of soluble A $\beta$  oligomers in the brain correlate much better than plaques or fibrils with AD onset, progression and severity [5, 6, 8, 63, 64]. Within the last fifteen years, a large number of studies from research groups worldwide have reported the existence of many different oligomeric assemblies from various sources, including AD brain and cerebrospinal fluid (CSF) samples, secreted into the conditioned medium of cultured cells or prepared artificially from recombinant or synthetic A $\beta$  peptides [65]. A heterogeneous range of sizes and peptide conformations have been observed among these natural and artificial A $\beta$  oligomers, including dimers and trimers [66, 67], tetramers, hexamers and the dodecameric A $\beta$ \*56 [64], globulomers [68], ring-shaped annular protofibrils [69] and higher molecular weight A $\beta$ -derived diffusible ligands (ADDLs) which can comprise hundreds of monomeric subunits [9, 70] (Figure 2). However, despite the disparity in size and source, A $\beta$  oligomers appear to share important functional properties. Notably, both natural and synthetic A $\beta$  oligomer preparations bind to hippocampal neurons and cells of neuronal lineage, causing a loss of dendritic spines, neurotoxicity, the inhibition of long-term synaptic potentiation (LTP: an electrophysiological correlate of learning and memory) and impairments in working memory at nanomolar concentrations [64, 67, 68, 70–73]. The preferential binding and toxicity of A $\beta$  oligomers towards neurons in the hippocampus may explain why A $\beta$  oligomers correlate with AD severity and disease progression [9, 68, 70]. However, the cellular mechanisms by which these effects are modulated remain poorly understood.

*3.2. A $\beta$  Oligomerisation Is Modulated by Lipid Raft Components.* A $\beta$  is a physiological peptide which is present in the brain tissue and CSF of healthy subjects throughout life, without necessarily causing neurodegeneration [74–76]. Many studies have shown that monomeric, nonaggregated A $\beta$  does not cause the neurotoxic effects that are mediated by A $\beta$  oligomers. In fact, monomeric A $\beta$  has recently been reported to have neuroprotective roles in the brain [77, 78]. The aggregation of A $\beta$  is necessary for its toxicity [79] and the emerging picture is that soluble A $\beta$  oligomers are the proximate neurotoxins in AD [8, 80]. The aggregation of A $\beta$  is therefore a critical step in the development of AD pathogenesis, and one in which lipid rafts appear to play a fundamental role.

Neuronal sensitivity to A $\beta$ -induced toxicity has been found to be dependent upon A $\beta$  binding to the cell membrane [81] and A $\beta$  has been identified in lipid rafts from cultured cells and from human and rodent brains. Soluble A $\beta$  dimers accumulate rapidly, and have been found at elevated levels, in lipid raft fractions isolated from human and transgenic mouse model AD brains [82]. Importantly,

A $\beta$  has been shown to accumulate in presynaptic terminals in AD cortex where it colocalises with the lipid raft markers cholesterol and ganglioside GM1 [83]. Taken together, these data suggest that A $\beta$  accumulation and aggregation within lipid rafts may underlie AD neuropathology.

As cholesterol is a major component of lipid rafts, it was postulated to facilitate A $\beta$  oligomerisation on neuronal membranes. The brain is particularly enriched in cholesterol, harbouring over 23% of the body's total complement but comprising only around 2% of total body mass [84]. However, the role of cholesterol in promoting the assembly of A $\beta$  is controversial and conflicting evidence has been presented in recent years. The main difficulty is being able to distinguish between the key role of cholesterol in building the lipid raft domains necessary for A $\beta$  production and the suggested role of cholesterol in promoting A $\beta$  oligomerisation. As discussed previously, raised cholesterol has been linked to AD; is this solely due to an increase in total lipid raft composition of the plasma membrane which increases amyloidogenic processing of APP to yield more A $\beta$  peptide or due to a direct effect on A $\beta$  oligomerisation?

A growing body of evidence suggests that certain components of lipid raft domains may play a much more sinister role in catalysing the conversion of the aggregation-prone A $\beta$  peptide to its neurotoxic, oligomeric states. Cholesterol is known to modulate the interaction of the A $\beta$  peptide with lipid bilayers [85]. Further, A $\beta$  oligomers isolated from AD patients associate with DRMs in a cholesterol-dependent manner, and cholesterol depletion reduces the aggregation of A $\beta$  [86]. It is currently unknown, however, whether this latter effect is due to a direct interaction between A $\beta$  and cholesterol, or due to the overall depletion in lipid raft domains and/or the subsequent change in composition and properties brought on by a reduction in cholesterol. Conversely, a recent study revealed that increasing the level of cholesterol in human neuroblastoma cells actually reduced the ability of synthetic A $\beta$  oligomers to bind [87], in spite of the colocalisation of the A $\beta$  oligomers with the lipid raft component ganglioside GM1. These data agree with the authors' previous finding that an increased level of membrane cholesterol exerts a protective effect against A $\beta$  oligomer toxicity [88]. In the more recent study [89] it was proposed that a fluctuation in cholesterol levels may alter the physical properties of lipid rafts thereby modulating oligomer binding.

Cholesterol can also facilitate A $\beta$  aggregation through the structural modification of other lipid raft components. A recent study using reconstituted membranes revealed a structural role for cholesterol in modulating the conformation of glycosphingolipids. Depending on the type of glycosphingolipid, cholesterol can either facilitate (such as for ganglioside GM1) or inhibit the interaction of A $\beta$  peptides with lipid rafts through fine-tuning of the glycosphingolipid conformation [90]. This reinforces the notion that A $\beta$  binding to, and aggregation upon, neuronal lipid raft domains cannot be ascribed to a single component, but rather that multiple players are likely to be involved.

In fact, mounting evidence suggests that gangliosides within lipid rafts appear to be the main driving force




FIGURE 2:  $A\beta$  oligomers are the key neurotoxic assemblies in Alzheimer's Disease. The  $A\beta$  peptide is natively unfolded yet conformationally plastic and prone to aggregation. In response to various stimuli, including elevated concentration,  $A\beta$  undergoes complex conformational rearrangements to form oligomer-competent or fibril-competent intermediates. A variety of  $A\beta$  oligomers can form which include low- $n$  oligomers (dimer and trimers), globulomers, hexameric and dodecameric ( $A\beta^*56$ ) states, higher molecular weight species such as  $A\beta$ -derived diffusible ligands (ADDLs) and ring-shaped annular protofibrils (APFs). Some oligomers are stable, off-pathway intermediates whereas others undergo further conformational changes and aggregation to form larger protofibrils and fibrils. Fibrils of  $A\beta$  are insoluble and deposit within extracellular senile plaques.  $A\beta$  oligomers are soluble and represent the active neurotoxic species in AD. The specific binding of  $A\beta$  oligomers to neurons, particularly in the hippocampus, triggers the memory impairments, loss of synaptic functionality and neuronal death which characterise AD.

behind the oligomerisation of  $A\beta$  on neuronal membranes. The development of AD within certain brain regions has been found to correlate with increased ganglioside levels [91]. Gangliosides are glycosphingolipids with one or more sialic acid moieties attached to the sugar chain. Gangliosides are found predominantly in the central nervous system, where they are enriched in lipid rafts due to the preferential packing of their saturated acyl chains within the liquid-ordered phase. A study in 1995 revealed that a population of membrane-bound  $A\beta$  tightly bound to gangliosides exists in AD brains [92]. More recently, exogenously-applied  $A\beta$  was shown to bind to neuronal membranes and to redistribute into lipid rafts where it colocalised with ganglioside GM1 in a time-dependent manner [93]. GM1 facilitated the binding and accumulation of  $A\beta$  oligomers at lipid raft domains and appeared to be required for the  $A\beta$  oligomer-mediated lipid peroxidation of DRMs [94]. Ganglioside GM1 contains just one sialic acid moiety and plays important physiological roles in neuronal function.  $A\beta$  appears to interact with the sialic acid moiety of gangliosides such as GM1 and these bound aggregates can go on to seed further  $A\beta$  aggregation [95]. The interaction between sialic acid and  $A\beta$  induces a conformational rearrangement of the  $A\beta$  peptide chain [96] which may potentiate  $A\beta$  oligomerisation. DRMs derived from ganglioside-rich rat brain, but not from liver, were found to promote the oligomerisation of  $A\beta$  [97]. Further, this study revealed that the removal of cholesterol or protein from these raft fractions did not prevent  $A\beta$  aggregation, providing evidence that neither cholesterol nor protein is essential for this process. However, lipid raft fractions containing very low levels of gangliosides still retained

some  $A\beta$  oligomerisation ability, and therefore ganglioside-independent aggregation mechanisms cannot be ruled out.

#### 4. $A\beta$ Oligomers Bind to Neuronal Receptors within Lipid Rafts

**4.1.  $A\beta$  Oligomers Bind to High Affinity Protein Receptors.** When the first synthetic  $A\beta$  oligomers were prepared from  $A\beta_{1-42}$  peptide by the Klein laboratory in 1998, it was observed that their binding to hippocampal neurons and cultured nerve cells was abolished by treating the cells with trypsin [70]. This, coupled with the low oligomer concentration (5 nM) required for neurotoxicity, implied that specific protein receptors were responsible for the binding of  $A\beta$  oligomers and for the subsequent transduction and amplification of neurotoxicity. Indeed, a recent study found that  $A\beta$  oligomer binding to neurons was saturable with an estimated apparent  $K_d$  of  $\sim 0.4$  nM [9]. This finding implied that one or more high-affinity receptors are responsible for  $A\beta$  oligomer binding and subsequent neurotoxicity. Immunofluorescence microscopy has revealed that  $A\beta$  oligomers bind to dendritic spines of hippocampal neurons where they colocalise with postsynaptic markers [9, 98, 99]. Interestingly,  $A\beta$  oligomer binding to neurons has a punctate appearance [100], which is reminiscent of the appearance of lipid raft localised proteins [101]. Several putative neuronal receptors for  $A\beta$  have been identified in recent years, namely proteins that are related to mechanisms of memory and neuroprotection in the brain. Noteworthy, all of these receptors either reside primarily within, or can partition into, lipid raft domains at the surface of neurons. Lipid rafts may therefore hold the key to

understanding how the deleterious effects of A $\beta$  oligomers are transduced through binding to specific receptors within these microdomains.

**4.2. The Cellular Prion Protein (PrP<sup>Sc</sup>).** In 2009, Laurén and colleagues reported that the cellular prion protein (PrP<sup>C</sup>) is a specific, high-affinity neuronal receptor for A $\beta$ <sub>1–42</sub> oligomers [9]. PrP<sup>C</sup> is a GPI-anchored protein that is expressed at high levels in the brain, particularly at synapses and axons, where it resides in lipid rafts. The misfolded form of the prion protein (PrP<sup>Sc</sup>) is infamous for being the causative agent in Mad Cow Disease (Bovine Spongiform Encephalopathy, BSE) and its human equivalent, Creutzfeldt-Jakob Disease (CJD). Although the correctly-folded PrP<sup>C</sup> is critical for prion disease pathogenesis, its physiological function remains enigmatic, with potential neuroprotective roles in oxidative stress defence, metal ion homeostasis and anti-apoptosis [102]. In a search to identify neuronal receptors for A $\beta$  oligomers, Laurén et al. [9] screened a mouse brain expression library of 225,000 cDNA constructs from which only two positive clones, both encoding full-length PrP<sup>C</sup>, were isolated that were able to bind A $\beta$  oligomers with high affinity and specificity. Interestingly, the PrP<sup>C</sup> homologues Shadoo and Doppel were found not to bind A $\beta$  oligomers to any significant degree. A further, more focussed screen of 352 clones encoding transmembrane proteins identified amyloid- $\beta$  precursor-like protein 1 (APLP1) and transmembrane protein 30B (TMEM30B) as weak A $\beta$  receptors, although their specificity for oligomeric A $\beta$  was poor. The  $\alpha$ 7 nicotinic acetylcholine receptor (nAChR $\alpha$ 7) and the receptor for advanced glycation end products (RAGE) were also assayed due to their previously reported affinities for A $\beta$  peptides [103, 104], although neither displayed high-affinity A $\beta$  oligomer binding. Therefore, PrP<sup>C</sup> was the only identified receptor to display both high affinity and high specificity for A $\beta$  oligomers.

A direct interaction between PrP<sup>C</sup> and A $\beta$  oligomers was confirmed and the core oligomer binding region of PrP<sup>C</sup> was narrowed down to amino acids 95–110, a positively charged cluster rich in lysine residues [9]. PrP<sup>C</sup> was also shown to mediate the inhibition of LTP that is induced when hippocampal slices were incubated with A $\beta$  oligomers at nanomolar concentrations [9]. A follow-up *in vivo* study revealed that the presence of PrP<sup>C</sup> is required for the A $\beta$  oligomer-mediated memory impairments in an AD model mouse [105]. Taken together, these data indicate a strong association between A $\beta$  oligomers binding to PrP<sup>C</sup> within lipid rafts of hippocampal neurons and the induction of memory deficits that are characteristic of AD.

Nevertheless, there has been some dispute over the role of PrP<sup>C</sup> in transducing the deleterious effects of A $\beta$  oligomers *in vivo*, as other studies have reported data which oppose this theory. First, Balducci and colleagues reported that although A $\beta$  oligomers bind tightly to PrP<sup>C</sup> they cause impairments in long-term memory in mice independently of PrP<sup>C</sup> [106]. In this study, the effects of synthetic A $\beta$  oligomers upon wild-type mice were observed, whereas Gimbel et al. [72] utilised a mouse model expressing a familial AD mutant APP. Further,

the synthetic depsipeptide and the oligomer preparation method utilised by Balducci et al. [106] differed from those used by Gimbel and coworkers [72], raising the possibility that PrP<sup>C</sup> does not have the same binding affinity for all types of A $\beta$  oligomers. Second, the Aguzzi group crossed an AD mouse model, which suffers from A $\beta$ -dependent memory deficits in the form of LTP impairment, with mice expressing either wild-type PrP<sup>C</sup>, a secreted form of PrP<sup>C</sup> (lacking its GPI anchor) or no PrP<sup>C</sup> [107]. They found that the presence or absence of wild-type PrP<sup>C</sup> had no effect upon the A $\beta$ -mediated inhibition of LTP. However, expression of the secreted form of PrP<sup>C</sup> was found to suppress the impairment in LTP, which the authors proposed may be due to the potential chelation and subsequent degradation of A $\beta$  oligomers by soluble PrP<sup>C</sup> in the extracellular milieu. Third, Kessels and coworkers reported the influence of PrP<sup>C</sup> upon hippocampal neurons expressing a C-terminally truncated form of APP in a viral expression construct [108]. The same loss of dendritic spines and inhibition of LTP were observed in the presence and absence of PrP<sup>C</sup>, suggesting that A $\beta$ -mediated synaptic defects do not require PrP<sup>C</sup>. However, Laurén and colleagues have emphasised the differences in the model system utilised by Kessels and coworkers in their study which may account for the opposing data, namely the viral expression of APP, a higher concentration of A $\beta$  oligomers and a difference in the observed suppression of synaptic plasticity [109].

Further investigation is needed to clarify the role of PrP<sup>C</sup> in modulating the A $\beta$  oligomer-mediated impairments in memory and LTP. Differences in the oligomer preparations, age and genotype of the mouse models, the nature of the promoter elements driving gene expression and the particular memory tests employed by the different authors may account for the discrepancies in the data.

The binding of A $\beta$  oligomers to PrP<sup>C</sup> is not the first time that PrP<sup>C</sup> has been linked to AD. Senile plaques from a subset of AD patients were observed to contain PrP<sup>C</sup> [110] and abundant A $\beta$  deposits have been observed in some CJD cases [111]. Furthermore, the Met/Val 129 polymorphism in the *PRNP* gene that encodes PrP<sup>C</sup> is a risk factor for early-onset AD [112]. In 2007, we demonstrated that PrP<sup>C</sup> negatively modulates A $\beta$  production through inhibition of the APP cleaving enzyme, BACE1 [113]. These data, along with the recent discovery that PrP<sup>C</sup> binds to A $\beta$  oligomers and transduces their deleterious effects, raises the intriguing possibility of a feedback loop [114]. We propose that, physiologically, PrP<sup>C</sup> maintains A $\beta$  production at a low level through BACE1 inhibition, but in AD this interaction may be disrupted by A $\beta$  oligomers binding to PrP<sup>C</sup> and causing its segregation from BACE1. Therefore, A $\beta$  oligomers binding to PrP<sup>C</sup> may also promote their own production through the ablation of BACE1 inhibition by PrP<sup>C</sup>. More recently, levels of PrP<sup>C</sup> have been shown to be reduced in AD brains [115, 116] possibly arguing against PrP<sup>C</sup> being involved in mediating the neurotoxic effects of A $\beta$  oligomers, at least in the terminal stages of the disease.

It is important to note that Laurén and colleagues reported that the removal of PrP<sup>C</sup> from hippocampal neurons only reduced A $\beta$  oligomer binding by approximately

50% [9]. This suggests that other receptors not identified in the expression library screen due to nonpreferential binding conditions or a low affinity for the particular type of A $\beta$  oligomers that were used, and/or nonprotein lipid raft components, may play equally crucial roles in A $\beta$  oligomer binding and neurotoxicity. Glutamate receptors, which possibly exist in a complex with PrP<sup>C</sup> [117], represent a candidate interacting partner for A $\beta$  oligomers which could explain the deleterious effects upon hippocampal synaptic plasticity.

**4.3. Glutamate Receptors.** Synaptic failure and impairments in synaptic plasticity are hallmarks of early AD neuropathology [100, 118, 119]. LTP and long-term depression (LTD) are mechanistic dimmer switches which facilitate synaptic plasticity by strengthening or weakening communication across a synapse, respectively, with LTP being essential for hippocampal-dependent learning and memory [120, 121]. Numerous lines of study have confirmed that soluble A $\beta$  oligomers from various sources, including those isolated from AD brains, disrupt hippocampal LTP *in vitro* and *in vivo* and cause impairments in learning and memory [9, 67, 70, 107, 122, 123]. Although not all studies agree, it has also been demonstrated that A $\beta$  oligomers can provoke LTD which opposes LTP [67, 124, 125]. Neuronal receptors which modulate LTP and/or LTD are therefore likely candidates for the specific binding of A $\beta$  oligomers. Glutamate receptors are central to the modulation of LTP and LTD. Additionally, glutamate receptor dysfunction has been implicated in AD which is characterised by memory deficits caused by impaired synaptic plasticity [126]. Glutamate receptors consist of two classes; ionotropic (cation-specific ion channels) and metabotropic (G-protein-coupled). Members of both classes have been implicated as neuronal receptors for A $\beta$  oligomers.

**4.3.1. NMDA Ionotropic Glutamate Receptors.** N-methyl-D-aspartate receptors (NMDARs) constitute a major class of glutamate receptors in the mammalian brain which localise to the postsynaptic membrane of excitatory synapses [127]. These ion channels play key roles in excitatory synaptic transmission and synaptic plasticity [128]. The membrane channel is usually blocked by Mg<sup>2+</sup> ions which are displaced when synaptic transmission results in depolarisation and glutamate release and binding. NMDAR channel opening leads to the rapid influx of Ca<sup>2+</sup> which triggers LTP induction [129]. Longer-term effects which maintain the reinforced synapse include the activation of  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPA), altered gene expression and kinase activity and the growth of new dendritic spines. Interestingly, NMDAR activation can also stimulate LTD, having the opposing effect of synapse weakening, and this appears to depend upon the nature of the stimulus and the subtype of NMDAR involved [130].

NMDARs localise to lipid raft domains where they interact with flotillins [131, 132] although they can move laterally between raft and nonraft domains in response to cues including phosphorylation [133]. Statins, which deplete

cellular cholesterol thus reducing lipid raft formation, have been shown to reduce the localisation of NMDARs to lipid raft domains, which has a neuroprotective effect [134].

Mounting evidence points towards a central role for NMDARs in the modulation of A $\beta$  oligomer toxicity. Soluble A $\beta$  oligomers inhibit NMDAR-dependent LTP [70, 135] and exhibit postsynaptic binding to hippocampal neurons which express NMDAR subunits GluN1 and GluN2B [100]. A reduction in NMDAR subunits GluN1 and GluN2B has previously been observed in the hippocampus of AD brains [136]. Crucially, a recent study has confirmed that A $\beta$  oligomer-mediated early synaptic dysfunction depends upon the activation of GluN2B-containing NMDARs [10]. A $\beta$  oligomers were found to decrease the NMDAR-dependent influx of Ca<sup>2+</sup> into dendritic spines [137], and to reduce dendritic spine and synapse density [10] in a mechanism which involve the subsequent phosphorylation of tau [138]. NMDAR antagonists, including one which is specific for GluN2B subunits, were able to reverse the A $\beta$ -induced loss of dendritic spine density [100, 137, 139]. These effects are consistent with A $\beta$  oligomers blocking the NMDAR-mediated stimulation of LTP whilst promoting NMDAR-mediated LTD. In addition, A $\beta$  oligomers have been shown to stimulate the excessive generation of reactive oxygen species (ROS) through an NMDAR-dependent mechanism [140], suggesting a link between aberrant ROS regulation and A $\beta$ -induced cognitive impairment.

Furthermore, evidence to confirm a direct interaction between A $\beta$  oligomers and NMDAR subunits has recently been presented. Partial colocalisation was observed between NMDAR GluN2B and A $\beta$  oligomers in hippocampal slices, which increased upon the addition of glutamate, although the maximum colocalisation was less than 50% [141]. Further, A $\beta$  oligomers were recently found to coimmunoprecipitate with NMDAR subunits [117]. However, an indirect model proposed by Venkitaramani and colleagues suggests that the A $\beta$  oligomer-mediated decrease in GluN2B-containing NMDARs results from the former binding to  $\alpha$ -7 nicotinic acetylcholine receptors ( $\alpha$ 7nAChR), which activates striatal-enriched tyrosine phosphatase (STEP), in turn stimulating NMDAR internalisation [142]. More recent data has revealed elevated levels of STEP in a mouse model of AD and in human AD brains, and that the removal of STEP abrogates the A $\beta$ -mediated reduction in NMDARs at the cell surface [143]. Whether or not A $\beta$  oligomers interact with NMDARs directly, growing evidence suggests that NMDARs play an important role in transducing the deleterious effects of A $\beta$  oligomers upon synaptic functionality.

**4.3.2. mGluR5 Metabotropic Glutamate Receptor.** The mGluR5 metabotropic glutamate receptor plays important regulatory roles in neuronal calcium mobilisation and the modulation of LTP and excitatory postsynaptic potentials in hippocampal neurons [144, 145]. Recently, mGluR5 was identified as a novel A $\beta$  oligomer receptor in a study of the behaviour of fluorescently-labelled A $\beta$  oligomers on hippocampal neurons and their interaction with neuronal receptors [117]. The A $\beta$  oligomers bound to excitatory

synapses where their mobility decreased as they aggregated to form larger clusters over time. Consistent with previous data, A $\beta$  oligomers caused a removal in NMDARs from synapses and were found to coimmunoprecipitate with NMDAR subunits. Interestingly, the A $\beta$  oligomers also formed complexes with mGluR5 receptors, which caused their lateral redistribution into dendritic spines followed by Ca<sup>2+</sup> dysregulation. Renner and colleagues also observed a time-dependent increase in lipid raft-localised mGluR5s which suggests that A $\beta$  oligomers reduce the mobility of mGluRs, causing their aberrant aggregation within pathological signalling platforms [117]. When mGluR5 was removed from mouse hippocampal neurons, A $\beta$  oligomer binding was reduced by approximately 80% and the loss of NMDARs from the cell surface was prevented.

Metabotropic glutamate receptors have been implicated previously in the pathogenesis of AD and other neurodegenerative disorders [126]. Impaired mGluR signalling in the cortex of AD patients has been shown to correlate with AD-related neuropathological changes [146]. Interestingly, the stimulation of mGluRs can modulate APP processing [147]. A recent study revealed that the A $\beta$  peptide upregulates the expression of mGluR5s in astrocytes, protective nonneuronal cells which are implicated in AD pathogenesis and inflammation [148]. Increased levels of mGluR5s were observed in the brains of Down's syndrome patients [149]; a disease in which elevated levels of A $\beta$  result from the triplication of the APP gene [150].

**4.3.3. Other Putative Receptors.** Various other lipid raft-associated proteins have been reported to effect A $\beta$ -mediated synaptic dysfunction. For instance, the removal of nerve growth factor receptors (NGFRs), including TrkA and p75 neurotrophin receptor, from cells treated with GM1-induced A $\beta$  oligomers caused a significant reduction in oligomer-mediated cytotoxicity [151]. NGFR dysfunction and aberrant NGF signalling is associated with AD and increased A $\beta$  production [152, 153]. Although no direct interaction has been shown to our knowledge, it is possible that interplay between A $\beta$  oligomers and NGFRs may form part of a positive feedback loop which serves to reinforce A $\beta$  oligomer production, whilst blocking NGF signalling with deleterious effects upon neuronal survival. Physiologically, NGF binds to TrkA causing the translocation and clustering of receptors within lipid rafts [154]. The binding of A $\beta$  oligomers to TrkA and other NGFRs may therefore cause aberrant lipid raft clustering which prevents or disrupts the formation of the normal signalling platforms.

Recent research proposes that impaired insulin signalling may be involved in AD, even leading to the hypothesis that AD represents a third type of diabetes [155]. Insulin receptors, which are robustly expressed in hippocampal neurons, were found to bind A $\beta$  oligomers and to undergo internalisation from dendritic spines [156]. Perturbations in insulin signalling in the brain caused by A $\beta$  oligomers may impair memory and LTP [157]. Interestingly, insulin receptor subunits are also enriched in lipid raft domains in hippocampal neurons [158].

**4.3.4. Multireceptor, Pathogenic Signalling Platforms Are Induced by A $\beta$  Oligomers.** The emerging picture is that lipid rafts accommodate multiple receptors for A $\beta$  oligomers, namely PrP<sup>C</sup> along with NMDAR, mGluR5 and possibly other, lower affinity receptors. Interestingly, there is evidence to suggest that these three lipid raft-associated receptors interact together. Metabotropic glutamate receptors have been found to cocluster with NMDARs [159]. It has also been reported that PrP<sup>C</sup> inhibited NMDAR function in hippocampal neurons and coimmunoprecipitated with NMDAR subunits [160]. The functional and physical links between these A $\beta$  oligomer receptors suggest the existence of a multi-component, A $\beta$  oligomer binding raft complex, comprising of PrP<sup>C</sup>, mGluR5 and NMDAR (Figure 3) [117]. Whether the formation of this complex is required for oligomer binding, or whether the interaction of A $\beta$  oligomers with the individual proteins induces its assembly, is a "chicken and egg" situation. One possible hypothesis is that A $\beta$  oligomers promote the clustering of PrP<sup>C</sup> and glutamate receptors into pathological mega-scaffolds which induce both toxic loss- and gain-of-function downstream effects. For instance, the aberrant localisation of glutamate receptors may impede neuronal signalling mechanisms including LTP, while the clustering or internalisation of NMDARs may promote their LTD-inducing functionality. The combined effects of oligomer binding upon more than one glutamate receptor is likely to be a large disturbance in Ca<sup>2+</sup> homeostasis which results in pathological signalling cascades. Interestingly, the PrP<sup>C</sup>-mediated response to oxidative stress is thought to induce signalling cascades which can modulate Ca<sup>2+</sup> flux and synaptic plasticity [161]. Furthermore, A $\beta$  oligomers may cause the internalisation or loss of function of components such as PrP<sup>C</sup> thus reducing neuroprotection against oxidative stress at the cell surface. The clustering of A $\beta$  oligomers at lipid raft domains may also cause damage to physiologically important signalling rafts, thus impairing neuronal function. Furthermore, the A $\beta$  oligomer-induced redistribution of neuronal proteins into lipid rafts may influence their nonraft interacting partners, with additional deleterious effects upon neuronal function and integrity.

## 5. Conclusions

Neuronal lipid rafts are crucial modulators of A $\beta$  production and aggregation, leading to the accumulation of neurotoxic A $\beta$  oligomers in the brain which drive AD pathology. Recent evidence now incriminates lipid rafts as pathological signalling platforms in which A $\beta$  oligomer receptors, such as PrP<sup>C</sup> and glutamate receptors, cluster. A $\beta$  oligomer binding appears to induce the aberrant localisation of these proteins with deleterious effects upon their physiological functions including hippocampal LTP, which underlies memory, and defence against oxidative stress. In this way, lipid rafts appear to be directly responsible for the transduction of A $\beta$  oligomer-mediated memory impairments and neurotoxicity which characterise AD. Lipid rafts are not only implicated in AD but may also be the key to a range of neurodegenerative






FIGURE 3:  $A\beta$  oligomer binding stimulates the clustering of specific neuronal receptors into aberrant pathogenic signalling platforms at the synapse. (a) Synaptic function and neural communication is maintained by the activity of postsynaptic receptors including the neuroprotective PrP<sup>C</sup> and the NMDA and mGluR5 glutamate receptors, which modulate synaptic plasticity. In the healthy brain, the dynamic translocation of such receptors between lipid raft and nonraft domains of the plasma membrane modulates their activities; (b) in AD, the binding of  $A\beta$  oligomers at postsynaptic membranes causes the redistribution and clustering of receptors including PrP<sup>C</sup>, NMDAR and mGluR5 into pathological signalling platforms [117]. The resulting loss of transient lateral movement and subsequently interaction with other components is proposed to cause a loss of normal functionality combined with aberrant signalling by these receptors. The dysregulation of  $Ca^{2+}$  and inhibition of synaptic long-term potentiation likely underlie the memory deficits which characterise AD. Further, the loss of PrP<sup>C</sup> depletes neuronal protection against oxidative stress which may partially account for the neuronal death that is observed in AD brains.

proteinopathies, including Parkinson's Disease, Huntington's Disease, amyotrophic lateral sclerosis and prion diseases (reviewed in [12]). Indeed, lipid raft disruption protects neurons against the toxicity of other oligomers besides  $A\beta$  [22] and lipid rafts may therefore represent generic platforms for oligomer-mediated neurotoxicity. Understanding the cell biology of the downstream effects of amyloid oligomers binding to neuronal lipid raft proteins may uncover potential therapeutic targets for the prevention of AD and other neurodegenerative diseases.

## Abbreviations

$\alpha 7nAChR$ :  $\alpha$ -7 nicotinic acetylcholine receptor  
 $A\beta$ : Amyloid-beta  
 AD: Alzheimer's Disease  
 ADDL:  $A\beta$ -derived diffusible ligand  
 AICD: APP intracellular domain  
 AMPAR:  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor

APF: Annular protofibril  
 APH1: Anterior pharynx defective-1  
 APLP1: Amyloid- $\beta$  precursor-like protein 1  
 APP: Amyloid precursor protein  
 BACE1: Beta-site APP cleaving enzyme-1  
 BSE: Bovine Spongiform Encephalopathy  
 CJD: Creutzfeldt-Jakob Disease  
 CSF: Cerebrospinal fluid  
 DRM: Detergent-resistant membrane  
 GPI: Glycosyl phosphatidylinositol  
 $K_d$ : Apparent dissociation constant  
 LTD: Long-term synaptic depression  
 LTP: Long-term synaptic potentiation  
 mGluR: Metabotropic glutamate receptor  
 NGFR: Nerve growth factor receptor  
 NMDAR: N-methyl-D-aspartate receptor  
 PEN2: Presenilin enhancer-2  
 PrP<sup>C</sup>: Cellular isoform of the prion protein  
 PrP<sup>Sc</sup>: Scrapie isoform of the prion protein

RAGE: Receptor for advanced glycation end products  
 STEP: Striatal-enriched tyrosine phosphatase  
 TMEM30B: Transmembrane protein 30B.

## Acknowledgments

The authors thank the Wellcome Trust, the Alzheimer's Research Trust and the Medical Research Council of Great Britain for funding. The authors thank Dr I. J. Whitehouse, E. B. C. Glennon and H. O. King for comments.

## References

- [1] C. Mount and C. Downton, "Alzheimer disease: progress or profit?" *Nature Medicine*, vol. 12, no. 7, pp. 780–784, 2006.
- [2] A. Wimo and M. Prince, *World Alzheimer Report 2010—The Global Economic Impact of Dementia*, Alzheimer's Disease International, 2010.
- [3] M. Citron, "Alzheimer's disease: strategies for disease modification," *Nature Reviews Drug Discovery*, vol. 9, no. 5, pp. 387–398, 2010.
- [4] E. R. L. C. Vardy, A. J. Catto, and N. M. Hooper, "Proteolytic mechanisms in amyloid- $\beta$  metabolism: therapeutic implications for Alzheimer's disease," *Trends in Molecular Medicine*, vol. 11, no. 10, pp. 464–472, 2005.
- [5] L. F. Lue, Y. M. Kuo, A. E. Roher et al., "Soluble amyloid  $\beta$  peptide concentration as a predictor of synaptic change in Alzheimer's disease," *American Journal of Pathology*, vol. 155, no. 3, pp. 853–862, 1999.
- [6] C. A. McLean, R. A. Cherny, F. W. Fraser et al., "Soluble pool of A $\beta$  amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease," *Annals of Neurology*, vol. 46, no. 6, pp. 860–866, 1999.
- [7] S. Lesné, L. Kotilinek, and K. H. Ashe, "Plaque-bearing mice with reduced levels of oligomeric amyloid- $\beta$  assemblies have intact memory function," *Neuroscience*, vol. 151, no. 3, pp. 745–749, 2008.
- [8] R. Roychaudhuri, M. Yang, M. M. Hoshi, and D. B. Teplow, "Amyloid  $\beta$ -protein assembly and Alzheimer disease," *Journal of Biological Chemistry*, vol. 284, no. 8, pp. 4749–4753, 2009.
- [9] J. Laurén, D. A. Gimbel, H. B. Nygaard, J. W. Gilbert, and S. M. Strittmatter, "Cellular prion protein mediates impairment of synaptic plasticity by amyloid- $\beta$  oligomers," *Nature*, vol. 457, no. 7233, pp. 1128–1132, 2009.
- [10] R. Ronicke, M. Mikhaylova, S. Ronicke et al., "Early neuronal dysfunction by amyloid beta oligomers depends on activation of NR2B-containing NMDA receptors," *Neurobiology of Aging*. In press.
- [11] E. Alberdi, M. V. Sanchez-Gomez, F. Cavaliere et al., "Amyloid  $\beta$  oligomers induce Ca<sup>2+</sup> dysregulation and neuronal death through activation of ionotropic glutamate receptors," *Cell Calcium*, vol. 47, no. 3, pp. 264–272, 2010.
- [12] C. L. Schengrund, "Lipid rafts: keys to neurodegeneration," *Brain Research Bulletin*, vol. 82, no. 1-2, pp. 7–17, 2010.
- [13] S. J. Singer and G. L. Nicolson, "The fluid mosaic model of the structure of cell membranes," *Science*, vol. 175, no. 4023, pp. 720–731, 1972.
- [14] L. J. Pike, "Rafts defined: a report on the keystone symposium on lipid rafts and cell function," *Journal of Lipid Research*, vol. 47, no. 7, pp. 1597–1598, 2006.
- [15] K. Simons and M. J. Gerl, "Revitalizing membrane rafts: new tools and insights," *Nature Reviews Molecular Cell Biology*, vol. 11, no. 10, pp. 688–699, 2010.
- [16] K. Simons and E. Ikonen, "Functional rafts in cell membranes," *Nature*, vol. 387, no. 6633, pp. 569–572, 1997.
- [17] J. A. Allen, R. A. Halverson-Tamboli, and M. M. Rasenick, "Lipid raft microdomains and neurotransmitter signalling," *Nature Reviews Neuroscience*, vol. 8, no. 2, pp. 128–140, 2007.
- [18] V. Lewis and N. M. Hooper, "The role of lipid rafts in prion protein biology," *Frontiers in Bioscience*, vol. 16, pp. 151–168, 2011.
- [19] H. Kamiguchi, "The region-specific activities of lipid rafts during axon growth and guidance," *Journal of Neurochemistry*, vol. 98, no. 2, pp. 330–335, 2006.
- [20] C. Guirland and J. Q. Zheng, "Membrane lipid rafts and their role in axon guidance," *Advances in Experimental Medicine and Biology*, vol. 621, pp. 144–155, 2007.
- [21] H. Hering, C. C. Lin, and M. Sheng, "Lipid rafts in the maintenance of synapses, dendritic spines, and surface AMPA receptor stability," *Journal of Neuroscience*, vol. 23, no. 8, pp. 3262–3271, 2003.
- [22] F. Malchiodi-Albedi, V. Contruscieri, C. Raggi et al., "Lipid raft disruption protects mature neurons against amyloid oligomer toxicity," *Biochimica et Biophysica Acta*, vol. 1802, no. 4, pp. 406–415, 2010.
- [23] J. A. Hardy and G. A. Higgins, "Alzheimer's disease: the amyloid cascade hypothesis," *Science*, vol. 256, no. 5054, pp. 184–185, 1992.
- [24] J. Hardy, "Alzheimer's disease: the amyloid cascade hypothesis: an update and reappraisal," *Journal of Alzheimer's Disease*, vol. 9, no. 3, pp. 151–153, 2006.
- [25] K. S. Vetrivel and G. Thinakaran, "Membrane rafts in Alzheimer's disease beta-amyloid production," *Biochimica et Biophysica Acta*, vol. 1801, no. 8, pp. 860–867, 2010.
- [26] D. R. Taylor and N. M. Hooper, "Role of lipid rafts in the processing of the pathogenic prion and Alzheimer's amyloid- $\beta$  proteins," *Seminars in Cell and Developmental Biology*, vol. 18, no. 5, pp. 638–648, 2007.
- [27] J. M. Cordy, N. M. Hooper, and A. J. Turner, "The involvement of lipid rafts in Alzheimer's disease," *Molecular Membrane Biology*, vol. 23, no. 1, pp. 111–122, 2006.
- [28] T. M. J. Allinson, E. T. Parkin, A. J. Turner, and N. M. Hooper, "ADAMs family members as amyloid precursor protein  $\alpha$ -secretases," *Journal of Neuroscience Research*, vol. 74, no. 3, pp. 342–352, 2003.
- [29] A. J. Beel, M. Sakakura, P. J. Barrett, and C. R. Sanders, "Direct binding of cholesterol to the amyloid precursor protein: an important interaction in lipid-Alzheimer's disease relationships?" *Biochimica et Biophysica Acta*, vol. 1801, no. 8, pp. 975–982, 2010.
- [30] M. Kosicek, M. Malnar, A. Goate, and S. Hecimovic, "Cholesterol accumulation in Niemann Pick type C (NPC) model cells causes a shift in APP localization to lipid rafts," *Biochemical and Biophysical Research Communications*, vol. 393, no. 3, pp. 404–409, 2010.
- [31] R. Ehehalt, P. Keller, C. Haass, C. Thiele, and K. Simons, "Amyloidogenic processing of the Alzheimer  $\beta$ -amyloid precursor protein depends on lipid rafts," *Journal of Cell Biology*, vol. 160, no. 1, pp. 113–123, 2003.
- [32] J. M. Cordy, I. Hussain, C. Dingwall, N. M. Hooper, and A. J. Turner, "Exclusively targeting  $\beta$ -secretase to lipid rafts by GPI-anchor addition up-regulates  $\beta$ -site processing of the amyloid precursor protein," *Proceedings of the National*

- Academy of Sciences of the United States of America*, vol. 100, no. 20, pp. 11735–11740, 2003.
- [33] Q. Zha, Y. Ruan, T. Hartmann, K. Beyreuther, and D. Zhang, "GM1 ganglioside regulates the proteolysis of amyloid precursor protein," *Molecular Psychiatry*, vol. 9, no. 10, pp. 946–952, 2004.
- [34] K. S. Vetrivel, H. Cheng, W. Lin et al., "Association of  $\gamma$ -secretase with lipid rafts in post-golgi and endosome membranes," *Journal of Biological Chemistry*, vol. 279, no. 43, pp. 44945–44954, 2004.
- [35] J.-Y. Hur, H. Welander, H. Behbahani et al., "Active  $\gamma$ -secretase is localized to detergent-resistant membranes in human brain," *FEBS Journal*, vol. 275, no. 6, pp. 1174–1187, 2008.
- [36] P. C. Fraering, W. Ye, J. M. Strub et al., "Purification and characterization of the human  $\gamma$ -secretase complex," *Biochemistry*, vol. 43, no. 30, pp. 9774–9789, 2004.
- [37] P. Osenkowski, W. Ye, R. Wang, M. S. Wolfe, and D. J. Selkoe, "Direct and potent regulation of  $\gamma$ -secretase by its lipid microenvironment," *Journal of Biological Chemistry*, vol. 283, no. 33, pp. 22529–22540, 2008.
- [38] S. J. Lee, U. Liyanage, P. E. Bickel, W. Xia, P. T. Lansbury, and K. S. Kosik, "A detergent-insoluble membrane compartment contains A $\beta$  in vivo," *Nature Medicine*, vol. 4, no. 6, pp. 730–734, 1998.
- [39] V. Martín, N. Fabelo, G. Santpere et al., "Lipid alterations in lipid rafts from Alzheimer's disease human brain cortex," *Journal of Alzheimer's Disease*, vol. 19, no. 2, pp. 489–502, 2010.
- [40] L. Canevari and J. B. Clark, "Alzheimer's disease and cholesterol: the fat connection," *Neurochemical Research*, vol. 32, no. 4-5, pp. 739–750, 2007.
- [41] J. R. Harris and N. G. Milton, "Cholesterol in Alzheimer's disease and other amyloidogenic disorders," *Sub-Cellular Biochemistry*, vol. 51, pp. 47–75, 2010.
- [42] C. Guardia-Laguarta, M. Coma, M. Pera et al., "Mild cholesterol depletion reduces amyloid- $\beta$  production by impairing APP trafficking to the cell surface," *Journal of Neurochemistry*, vol. 110, no. 1, pp. 220–230, 2009.
- [43] M. A. Pappolla, T. K. Bryant-Thomas, D. Herbert et al., "Mild hypercholesterolemia is an early risk factor for the development of Alzheimer amyloid pathology," *Neurology*, vol. 61, no. 2, pp. 199–205, 2003.
- [44] Y. M. Kuo, M. R. Emmerling, C. L. Bisgaier et al., "Elevated low-density lipoprotein in Alzheimer's disease correlates with brain A $\beta$  1–42 levels," *Biochemical and Biophysical Research Communications*, vol. 252, no. 3, pp. 711–715, 1998.
- [45] M. Kivipelto and A. Solomon, "Cholesterol as a risk factor for Alzheimer's disease—epidemiological evidence," *Acta Neurologica Scandinavica*, vol. 114, no. 185, pp. 50–57, 2006.
- [46] L. M. Refolo, M. A. Pappolla, B. Malester et al., "Hypercholesterolemia accelerates the Alzheimer's amyloid pathology in a transgenic mouse model," *Neurobiology of Disease*, vol. 7, no. 4, pp. 321–331, 2000.
- [47] L. Thirumangalakudi, A. Prakasam, R. Zhang et al., "High cholesterol-induced neuroinflammation and amyloid precursor protein processing correlate with loss of working memory in mice," *Journal of Neurochemistry*, vol. 106, no. 1, pp. 475–485, 2008.
- [48] O. Ghribi, B. Larsen, M. Schrag, and M. M. Herman, "High cholesterol content in neurons increases BACE,  $\beta$ -amyloid, and phosphorylated tau levels in rabbit hippocampus," *Experimental Neurology*, vol. 200, no. 2, pp. 460–467, 2006.
- [49] M. Simons, P. Keller, B. De Strooper, K. Beyreuther, C. G. Dotti, and K. Simons, "Cholesterol depletion inhibits the generation of  $\beta$ -amyloid in hippocampal neurons," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 11, pp. 6460–6464, 1998.
- [50] M. Kivipelto, A. Solomon, and B. Winblad, "Statin therapy in Alzheimer's disease," *Lancet Neurology*, vol. 4, no. 9, pp. 521–522, 2005.
- [51] H. Jick, G. L. Zornberg, S. S. Jick, S. Seshadri, and D. A. Drachman, "Statins and the risk of dementia," *Lancet*, vol. 356, no. 9242, pp. 1627–1631, 2000.
- [52] D. L. Sparks, M. N. Sabbagh, D. J. Connor et al., "Atorvastatin for the treatment of mild to moderate Alzheimer disease: preliminary results," *Archives of Neurology*, vol. 62, no. 5, pp. 753–757, 2005.
- [53] B. Wolozin, W. Kellman, P. Ruosseau, G. G. Cesesia, and G. Siegel, "Decreased prevalence of Alzheimer disease associated with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors," *Archives of Neurology*, vol. 57, no. 10, pp. 1439–1443, 2000.
- [54] G. P. Hooff, I. Peters, W. G. Wood, W. E. Müller, and G. P. Eckert, "Modulation of cholesterol, farnesylpyrophosphate, and geranylgeranylpyrophosphate in neuroblastoma SH-SY5Y-APP695 cells: impact on amyloid beta-protein production," *Molecular Neurobiology*, vol. 41, no. 2-3, pp. 341–350, 2010.
- [55] P. P. Zandi, D. L. Sparks, A. S. Khachaturian et al., "Do statins reduce risk of incident dementia and Alzheimer disease? The Cache county study," *Archives of General Psychiatry*, vol. 62, no. 2, pp. 217–224, 2005.
- [56] S. Hoyer and P. Riederer, "Alzheimer disease—no target for statin treatment. A mini review," *Neurochemical Research*, vol. 32, no. 4-5, pp. 695–706, 2007.
- [57] T. Umeda, H. Mori, H. Zheng, and T. Tomiyama, "Regulation of cholesterol efflux by amyloid  $\beta$  secretion," *Journal of Neuroscience Research*, vol. 88, no. 9, pp. 1985–1994, 2010.
- [58] D. M. Walsh, I. Klyubin, J. V. Fadeeva et al., "Naturally secreted oligomers of amyloid  $\beta$  protein potently inhibit hippocampal long-term potentiation in vivo," *Nature*, vol. 416, no. 6880, pp. 535–539, 2002.
- [59] D. H. Small, S. S. Mok, and J. C. Bornstein, "Alzheimer's disease and A $\beta$  toxicity: from top to bottom," *Nature Reviews Neuroscience*, vol. 2, no. 8, pp. 595–598, 2001.
- [60] E. Tamagno, P. Bardini, M. Guglielmotto, O. Danni, and M. Tabaton, "The various aggregation states of  $\beta$ -amyloid 1–42 mediate different effects on oxidative stress, neurodegeneration, and BACE-1 expression," *Free Radical Biology and Medicine*, vol. 41, no. 2, pp. 202–212, 2006.
- [61] A. Deshpande, E. Mina, C. Glabe, and J. Busciglio, "Different conformations of amyloid  $\beta$  induce neurotoxicity by distinct mechanisms in human cortical neurons," *Journal of Neuroscience*, vol. 26, no. 22, pp. 6011–6018, 2006.
- [62] W. F. Xue, A. L. Hellewell, W. S. Gosal, S. W. Homans, E. W. Hewitt, and S. E. Radford, "Fibril fragmentation enhances amyloid cytotoxicity," *Journal of Biological Chemistry*, vol. 284, no. 49, pp. 34272–34282, 2009.
- [63] J. L. Tomic, A. Pensalfini, E. Head, and C. G. Glabe, "Soluble fibrillar oligomer levels are elevated in Alzheimer's disease brain and correlate with cognitive dysfunction," *Neurobiology of Disease*, vol. 35, no. 3, pp. 352–358, 2009.
- [64] S. Lesne, T. K. Ming, L. Kotilinek et al., "A specific amyloid- $\beta$  protein assembly in the brain impairs memory," *Nature*, vol. 440, no. 7082, pp. 352–357, 2006.

- [65] D. M. Walsh and D. J. Selkoe, "A $\beta$  oligomers—a decade of discovery," *Journal of Neurochemistry*, vol. 101, no. 5, pp. 1172–1184, 2007.
- [66] M. B. Podlisny, B. L. Ostaszewski, S. L. Squazzo et al., "Aggregation of secreted amyloid  $\beta$ -protein into sodium dodecyl sulfate- stable oligomers in cell culture," *Journal of Biological Chemistry*, vol. 270, no. 16, pp. 9564–9570, 1995.
- [67] G. M. Shankar, S. Li, T. H. Mehta et al., "Amyloid- $\beta$  protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory," *Nature Medicine*, vol. 14, no. 8, pp. 837–842, 2008.
- [68] S. Barghorn, V. Nimmrich, A. Striebing et al., "Globular amyloid  $\beta$ -peptide oligomer—a homogenous and stable neuropathological protein in Alzheimer's disease," *Journal of Neurochemistry*, vol. 95, no. 3, pp. 834–847, 2005.
- [69] R. Kaye, A. Pensalfini, L. Margol et al., "Annular protofibrils area structurally and functionally distinct type of amyloid oligomer," *Journal of Biological Chemistry*, vol. 284, no. 7, pp. 4230–4237, 2009.
- [70] M. P. Lambert, A. K. Barlow, B. A. Chromy et al., "Diffusible, nonfibrillar ligands derived from A $\beta$  are potent central nervous system neurotoxins," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 11, pp. 6448–6453, 1998.
- [71] D. B. Freir, R. Fedriani, D. Scully et al., "Abeta oligomers inhibit synapse remodelling necessary for memory consolidation," *Neurobiology of Aging*. In press.
- [72] D. A. Gimbel, H. B. Nygaard, E. E. Coffey et al., "Memory impairment in transgenic alzheimer mice requires cellular prion protein," *Journal of Neuroscience*, vol. 30, no. 18, pp. 6367–6374, 2010.
- [73] T. Tomiyama, S. Matsuyama, H. Iso et al., "A mouse model of amyloid  $\beta$  oligomers: their contribution to synaptic alteration, abnormal tau phosphorylation, glial activation, and neuronal loss in vivo," *Journal of Neuroscience*, vol. 30, no. 14, pp. 4845–4856, 2010.
- [74] C. Haass, M. G. Schlossmacher, A. Y. Hung et al., "Amyloid  $\beta$ -peptide is produced by cultured cells during normal metabolism," *Nature*, vol. 359, no. 6393, pp. 322–325, 1992.
- [75] M. Shoji, "Cerebrospinal fluid Abeta40 and Abeta42: natural course and clinical usefulness," *Frontiers in Bioscience*, vol. 7, pp. 997–1006, 2002.
- [76] C. Vigo-Pelfrey, D. Lee, P. Keim, I. Lieberburg, and D. B. Schenk, "Characterization of  $\beta$ -amyloid peptide from human cerebrospinal fluid," *Journal of Neurochemistry*, vol. 61, no. 5, pp. 1965–1968, 1993.
- [77] D. L. Brody, S. Magnoni, K. E. Schwetye et al., "Amyloid- $\beta$  dynamics correlate with neurological status in the injured human brain," *Science*, vol. 321, no. 5893, pp. 1221–1224, 2008.
- [78] M. L. Giuffrida, F. Caraci, B. Pignataro et al., " $\beta$ -amyloid monomers are neuroprotective," *Journal of Neuroscience*, vol. 29, no. 34, pp. 10582–10587, 2009.
- [79] C. J. Pike, A. J. Walencewicz, C. G. Glabe, and C. W. Cotman, "Aggregation-related toxicity of synthetic  $\beta$ -amyloid protein in hippocampal cultures," *European Journal of Pharmacology*, vol. 207, no. 4, pp. 367–368, 1991.
- [80] G. M. Shankar and D. M. Walsh, "Alzheimer's disease: synaptic dysfunction and A $\beta$ ," *Molecular Neurodegeneration*, vol. 4, no. 1, article no. 48, 2009.
- [81] O. Simakova and N. J. Arispe, "The cell-selective neurotoxicity of the Alzheimer's A $\beta$  peptide is determined by surface phosphatidylserine and cytosolic ATP levels. Membrane binding is required for A $\beta$  toxicity," *Journal of Neuroscience*, vol. 27, no. 50, pp. 13719–13729, 2007.
- [82] T. Kawarabayashi, M. Shoji, L. H. Younkin et al., "Dimeric amyloid beta protein rapidly accumulates in lipid rafts followed by apolipoprotein E and phosphorylated tau accumulation in the Tg2576 mouse model of Alzheimer's disease," *Journal of Neuroscience*, vol. 24, no. 15, pp. 3801–3809, 2004.
- [83] K. H. Gylys, J. A. Fein, F. Yang, C. A. Miller, and G. M. Cole, "Increased cholesterol in A $\beta$ -positive nerve terminals from Alzheimer's disease cortex," *Neurobiology of Aging*, vol. 28, no. 1, pp. 8–17, 2007.
- [84] J. M. Dietschy and S. D. Turley, "Cholesterol metabolism in the central nervous system during early development and in the mature animal," *Journal of Lipid Research*, vol. 45, no. 8, pp. 1375–1397, 2004.
- [85] L. Qiu, A. Lewis, J. Como et al., "Cholesterol modulates the interaction of  $\beta$ -amyloid peptide with lipid bilayers," *Biophysical Journal*, vol. 96, no. 10, pp. 4299–4307, 2009.
- [86] A. Schneider, W. Schulz-Schaeffer, T. Hartmann, J. B. Schulz, and M. Simons, "Cholesterol depletion reduces aggregation of amyloid-beta peptide in hippocampal neurons," *Neurobiology of Disease*, vol. 23, no. 3, pp. 573–577, 2006.
- [87] C. Cecchi, D. Nichino, M. Zampagni et al., "A protective role for lipid raft cholesterol against amyloid-induced membrane damage in human neuroblastoma cells," *Biochimica et Biophysica Acta*, vol. 1788, no. 10, pp. 2204–2216, 2009.
- [88] C. Cecchi, F. Rosati, A. Pensalfini et al., "Seladin-1/DHCR24 protects neuroblastoma cells against A $\beta$  toxicity by increasing membrane cholesterol content," *Journal of Cellular and Molecular Medicine*, vol. 12, no. 5B, pp. 1990–2002, 2008.
- [89] C. Cecchi, D. Nichino, M. Zampagni et al., "A protective role for lipid raft cholesterol against amyloid-induced membrane damage in human neuroblastoma cells," *Biochimica et Biophysica Acta*, vol. 1788, no. 10, pp. 2204–2216, 2009.
- [90] N. Yahi, A. Aulas, and J. Fantini, "How cholesterol constrains glycolipid conformation for optimal recognition of Alzheimer's  $\beta$  amyloid peptide (A $\beta$ <sub>1–40</sub>)," *PLoS ONE*, vol. 5, no. 2, article no. e9079, 2010.
- [91] M. Molander-Melin, K. Blennow, N. Bogdanovic, B. Dellheden, J. E. Månsson, and P. Fredman, "Structural membrane alterations in Alzheimer brains found to be associated with regional disease development; increased density of gangliosides GM1 and GM2 and loss of cholesterol in detergent-resistant membrane domains," *Journal of Neurochemistry*, vol. 92, no. 1, pp. 171–182, 2005.
- [92] K. Yanagisawa, A. Odaka, N. Suzuki, and Y. Ihara, "GM1 ganglioside-bound amyloid  $\beta$ -protein (AB): a possible form of preamyloid in Alzheimer's disease," *Nature Medicine*, vol. 1, no. 10, pp. 1062–1066, 1995.
- [93] K. Matsuzaki, K. Kato, and K. Yanagisawa, "A $\beta$  polymerization through interaction with membrane gangliosides," *Biochimica et Biophysica Acta*, vol. 1801, no. 8, pp. 868–877, 2010.
- [94] M. Zampagni, E. Evangelisti, R. Cascella et al., "Lipid rafts are primary mediators of amyloid oxidative attack on plasma membrane," *Journal of Molecular Medicine*, vol. 88, no. 6, pp. 597–608, 2010.
- [95] A. Kakio, S. I. Nishimoto, K. Yanagisawa, Y. Kozutsumi, and K. Matsuzaki, "Interactions of amyloid  $\beta$ -protein with various gangliosides in raft-like membranes: importance of GM1 ganglioside-bound form as an endogenous seed for Alzheimer amyloid," *Biochemistry*, vol. 41, no. 23, pp. 7385–7390, 2002.

- [96] J. McLaurin, T. Franklin, P. E. Fraser, and A. Chakrabarty, "Structural transitions associated with the interaction of Alzheimer  $\beta$ -amyloid peptides with gangliosides," *Journal of Biological Chemistry*, vol. 273, no. 8, pp. 4506–4515, 1998.
- [97] S. I. Kim, J. S. Yi, and Y. G. Ko, "Amyloid  $\beta$  oligomerization is induced by brain lipid rafts," *Journal of Cellular Biochemistry*, vol. 99, no. 3, pp. 878–889, 2006.
- [98] P. N. Lacor, M. C. Buniel, L. Chang et al., "Synaptic targeting by Alzheimer's-related amyloid  $\beta$  oligomers," *Journal of Neuroscience*, vol. 24, no. 45, pp. 10191–10200, 2004.
- [99] G. A. Krafft and W. L. Klein, "ADDLs and the signaling web that leads to Alzheimer's disease," *Neuropharmacology*, vol. 59, no. 4–5, pp. 230–242, 2010.
- [100] P. N. Lacor, M. C. Buniel, P. W. Furlow et al., "A $\beta$  oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease," *Journal of Neuroscience*, vol. 27, no. 4, pp. 796–807, 2007.
- [101] D. R. Taylor, N. T. Watt, W. S. S. Perera, and N. M. Hooper, "Assigning functions to distinct regions of the N-terminus of the prion protein that are involved in its copper-stimulated, clathrin-dependent endocytosis," *Journal of Cell Science*, vol. 118, no. 21, pp. 5141–5153, 2005.
- [102] D. R. Taylor and N. M. Hooper, "The prion protein and lipid rafts (Review)," *Molecular Membrane Biology*, vol. 23, no. 1, pp. 89–99, 2006.
- [103] S. D. Yan, X. Chen, J. Fu et al., "RAGE and amyloid- $\beta$  peptide neurotoxicity in Alzheimer's disease," *Nature*, vol. 382, no. 6593, pp. 685–691, 1996.
- [104] H.-Y. Wang, D. H. S. Lee, M. R. D'Andrea, P. A. Peterson, R. P. Shank, and A. B. Reitz, " $\beta$  - Amyloid<sub>1–42</sub> binds to  $\alpha 7$  nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology," *Journal of Biological Chemistry*, vol. 275, no. 8, pp. 5626–5632, 2000.
- [105] D. A. Gimbel, H. B. Nygaard, E. E. Coffey et al., "Memory impairment in transgenic alzheimer mice requires cellular prion protein," *Journal of Neuroscience*, vol. 30, no. 18, pp. 6367–6374, 2010.
- [106] C. Balducci, M. Beeg, M. Stravalaci et al., "Synthetic amyloid- $\beta$  oligomers impair long-term memory independently of cellular prion protein," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 5, pp. 2295–2300, 2010.
- [107] A. M. Calella, M. Farinelli, M. Nuvolone et al., "Prion protein and Ab-related synaptic toxicity impairment," *EMBO Molecular Medicine*, vol. 2, no. 8, pp. 306–314, 2010.
- [108] H. W. Kessels, L. N. Nguyen, S. Nabavi, and R. Malinow, "The prion protein as a receptor for amyloid- $\beta$ ," *Nature*, vol. 466, no. 7308, pp. E3–E4, 2010.
- [109] J. Laurén, D. A. Gimbel, H. B. Nygaard, J. W. Gilbert, and S. M. Strittmatter, "Lauren et al. reply," *Nature*, vol. 466, no. 7308, pp. E4–E5, 2010.
- [110] T. Voigtlander, S. Kloppel, P. Birner et al., "Marked increase of neuronal prion protein immunoreactivity in Alzheimer's disease and human prion diseases," *Acta Neuropathologica*, vol. 101, no. 5, pp. 417–423, 2001.
- [111] L. Debatin, J. Streffer, M. Geissen, J. Matschke, A. Aguzzi, and M. Glatzel, "Association between deposition of beta-amyloid and pathological prion protein in sporadic Creutzfeldt-Jakob disease," *Neurodegenerative Diseases*, vol. 5, no. 6, pp. 347–354, 2008.
- [112] M. Riemenschneider, N. Klopp, W. Xiang et al., "Prion protein codon 129 polymorphism and risk of Alzheimer disease," *Neurology*, vol. 63, no. 2, pp. 364–366, 2004.
- [113] E. T. Parkin, N. T. Watt, I. Hussain et al., "Cellular prion protein regulates  $\beta$ -secretase cleavage of the Alzheimer's amyloid precursor protein," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 26, pp. 11062–11067, 2007.
- [114] K. A. B. Kellett and N. M. Hooper, "Prion protein and Alzheimer disease," *Prion*, vol. 3, no. 4, pp. 190–194, 2009.
- [115] J. L. Velayos, A. Irujo, M. Cuadrado-Tejedor, B. Paternain, F. J. Molerés, and V. Ferrer, "The cellular prion protein and its role in Alzheimer disease," *Prion*, vol. 3, no. 2, pp. 110–117, 2009.
- [116] I. J. Whitehouse, C. D. Jackson, A. J. Turner, and N. M. Hooper, "Prion protein is reduced in aging and in sporadic but not in familial Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 22, pp. 1023–1031, 2010.
- [117] M. Renner, P. N. Lacor, P. T. Velasco et al., "Deleterious effects of amyloid beta oligomers acting as an extracellular scaffold for mGluR5," *Neuron*, vol. 66, no. 5, pp. 739–754, 2010.
- [118] D. J. Selkoe, "Alzheimer's disease is a synaptic failure," *Science*, vol. 298, no. 5594, pp. 789–791, 2002.
- [119] M. J. Rowan, I. Klyubin, W. K. Cullen, and R. Anwyl, "Synaptic plasticity in animal models of early Alzheimer's disease," *Philosophical Transactions of the Royal Society B*, vol. 358, no. 1432, pp. 821–828, 2003.
- [120] M. A. Lynch, "Long-term potentiation and memory," *Physiological Reviews*, vol. 84, no. 1, pp. 87–136, 2004.
- [121] S. J. Martin and R. G. M. Morris, "New life in an old idea: the synaptic plasticity and memory hypothesis revisited," *Hippocampus*, vol. 12, no. 5, pp. 609–636, 2002.
- [122] M. Townsend, G. M. Shankar, T. Mehta, D. M. Walsh, and D. J. Selkoe, "Effects of secreted oligomers of amyloid  $\beta$ -protein on hippocampal synaptic plasticity: a potent role for trimers," *Journal of Physiology*, vol. 572, no. 2, pp. 477–492, 2006.
- [123] D. M. Walsh, I. Klyubin, J. V. Fadeeva, M. J. Rowan, and D. J. Selkoe, "Amyloid- $\beta$  oligomers: their production, toxicity and therapeutic inhibition," *Biochemical Society Transactions*, vol. 30, no. 4, pp. 552–557, 2002.
- [124] J. H. Kim, R. Anwyl, Y. H. Suh, M. B. A. Djamgoz, and M. J. Rowan, "Use-dependent effects of amyloidogenic fragments of  $\beta$ -amyloid precursor protein on synaptic plasticity in rat hippocampus in vivo," *Journal of Neuroscience*, vol. 21, no. 4, pp. 1327–1333, 2001.
- [125] H. Hsieh, J. Boehm, C. Sato et al., "AMPA removal underlies Abeta-induced synaptic depression and dendritic spine loss," *Neuron*, vol. 52, no. 5, pp. 831–843, 2006.
- [126] H. G. Lee, X. Zhu, M. J. O'Neill et al., "The role of metabotropic glutamate receptors in Alzheimer's disease," *Acta Neurobiologiae Experimentalis*, vol. 64, no. 1, pp. 89–98, 2004.
- [127] M. R. Hynd, H. L. Scott, and P. R. Dodd, "Glutamate-mediated excitotoxicity and neurodegeneration in Alzheimer's disease," *Neurochemistry International*, vol. 45, no. 5, pp. 583–595, 2004.
- [128] P. Paoletti and J. Neyton, "NMDA receptor subunits: function and pharmacology," *Current Opinion in Pharmacology*, vol. 7, no. 1, pp. 39–47, 2007.
- [129] P. Ascher and L. Nowak, "The role of divalent cations in the N-methyl-D-aspartate responses of mouse central neurones in culture," *Journal of Physiology*, vol. 399, pp. 247–266, 1988.
- [130] R. C. Malenka and M. F. Bear, "LTP and LTD: an embarrassment of riches," *Neuron*, vol. 44, no. 1, pp. 5–21, 2004.
- [131] C. C. Swanwick, M. E. Shapiro, Z. Yi, K. Chang, and R. J. Wenthold, "NMDA receptors interact with flotillin-1 and -2,

- lipid raft-associated proteins," *FEBS Letters*, vol. 583, no. 8, pp. 1226–1230, 2009.
- [132] H. Hering, C. C. Lin, and M. Sheng, "Lipid rafts in the maintenance of synapses, dendritic spines, and surface AMPA receptor stability," *Journal of Neuroscience*, vol. 23, no. 8, pp. 3262–3271, 2003.
- [133] S. Besshoh, S. Chen, I. R. Brown, and J. W. Gurd, "Developmental changes in the association of NMDA receptors with lipid rafts," *Journal of Neuroscience Research*, vol. 85, no. 9, pp. 1876–1883, 2007.
- [134] J. Ponce, N. P. De La Ossa, O. Hurtado et al., "Simvastatin reduces the association of NMDA receptors to lipid rafts: a cholesterol-mediated effect in neuroprotection," *Stroke*, vol. 39, no. 4, pp. 1269–1275, 2008.
- [135] H. W. Wang, J. F. Pasternak, H. Kuo et al., "Soluble oligomers of  $\beta$  amyloid (1–42) inhibit long-term potentiation but not long-term depression in rat dentate gyrus," *Brain Research*, vol. 924, no. 2, pp. 133–140, 2002.
- [136] A. J. Mishizen-Eberz, R. A. Rissman, T. L. Carter, M. D. Ikonovic, B. B. Wolfe, and D. M. Armstrong, "Biochemical and molecular studies of NMDA receptor subunits NR1/2A/2B in hippocampal subregions throughout progression of Alzheimer's disease pathology," *Neurobiology of Disease*, vol. 15, no. 1, pp. 80–92, 2004.
- [137] G. M. Shankar, B. L. Bloodgood, M. Townsend, D. M. Walsh, D. J. Selkoe, and B. L. Sabatini, "Natural oligomers of the Alzheimer amyloid- $\beta$  protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway," *Journal of Neuroscience*, vol. 27, no. 11, pp. 2866–2875, 2007.
- [138] C. Tackenberg and R. Brandt, "Divergent pathways mediate spine alterations and cell death induced by amyloid- $\beta$ , wild-type tau, and R406W tau," *Journal of Neuroscience*, vol. 29, no. 46, pp. 14439–14450, 2009.
- [139] R. Ronicke, M. Mikhaylova, and S. Ronicke, "Early neuronal dysfunction by amyloid beta oligomers depends on activation of NR2B-containing NMDA receptors," *Neurobiology of Aging*. In press.
- [140] F. G. De Felice, P. T. Velasco, M. P. Lambert et al., "A $\beta$  oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine," *Journal of Biological Chemistry*, vol. 282, no. 15, pp. 11590–11601, 2007.
- [141] A. Deshpande, H. Kawai, R. Metherate, C. G. Glabe, and J. Busciglio, "A role for synaptic zinc in activity-dependent A $\beta$  oligomer formation and accumulation at excitatory synapses," *Journal of Neuroscience*, vol. 29, no. 13, pp. 4004–4015, 2009.
- [142] D. V. Venkitaramani, J. Chin, W. J. Netzer et al., " $\beta$ -amyloid modulation of synaptic transmission and plasticity," *Journal of Neuroscience*, vol. 27, no. 44, pp. 11832–11837, 2007.
- [143] P. Kurup, Y. Zhang, J. Xu et al., "A $\beta$ -mediated NMDA receptor endocytosis in Alzheimer's disease involves ubiquitination of the tyrosine phosphatase STEP<sub>61</sub>," *Journal of Neuroscience*, vol. 30, no. 17, pp. 5948–5957, 2010.
- [144] A. Lau and M. Tymianski, "Glutamate receptors, neurotoxicity and neurodegeneration," *Pflugers Archiv European Journal of Physiology*, vol. 460, no. 2, pp. 525–542, 2010.
- [145] J. Wu, S. Harney, M. J. Rowan, and R. Anwyl, "Involvement of group I mGluRs in LTP induced by strong high frequency stimulation in the dentate gyrus in vitro," *Neuroscience Letters*, vol. 436, no. 2, pp. 235–238, 2008.
- [146] J. L. Albasanz, E. Dalfo, I. Ferrer, and M. Martín, "Impaired metabotropic glutamate receptor/phospholipase C signaling pathway in the cerebral cortex in Alzheimer's disease and dementia with Lewy bodies correlates with stage of Alzheimer's-disease-related changes," *Neurobiology of Disease*, vol. 20, no. 3, pp. 685–693, 2005.
- [147] R. K. K. Lee, R. J. Wurtman, A. J. Cox, and R. M. Nitsch, "Amyloid precursor protein processing is stimulated by metabotropic glutamate receptors," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 92, no. 17, pp. 8083–8087, 1995.
- [148] C. S. Casley, V. Lakics, H. G. Lee et al., "Up-regulation of astrocyte metabotropic glutamate receptor 5 by amyloid- $\beta$  peptide," *Brain Research*, vol. 1260, pp. 65–75, 2009.
- [149] A. Oka and S. Takashima, "The up-regulation of metabotropic glutamate receptor 5 (mGluR5) in Down's syndrome brains," *Acta Neuropathologica*, vol. 97, no. 3, pp. 275–278, 1999.
- [150] G. G. Glenner and C. W. Wong, "Alzheimer's disease and Down's syndrome: sharing of a unique cerebrovascular amyloid fibril protein," *Biochemical and Biophysical Research Communications*, vol. 122, no. 3, pp. 1131–1135, 1984.
- [151] N. Yamamoto, Y. Fukata, M. Fukata, and K. Yanagisawa, "GM1-ganglioside-induced A $\beta$  assembly on synaptic membranes of cultured neurons," *Biochimica et Biophysica Acta*, vol. 1768, no. 5, pp. 1128–1137, 2007.
- [152] S. E. Counts and E. J. Mufson, "The role of nerve growth factor receptors in cholinergic basal forebrain degeneration in prodromal Alzheimer disease," *Journal of Neuropathology and Experimental Neurology*, vol. 64, no. 4, pp. 263–272, 2005.
- [153] P. Calissano, C. Matrone, and G. Amadoro, "Nerve growth factor as a paradigm of neurotrophins related to Alzheimer's disease," *Developmental Neurobiology*, vol. 70, no. 5, pp. 372–383, 2010.
- [154] A. S. Limpert, J. C. Karlo, and G. E. Landreth, "Nerve growth factor stimulates the concentration of TrkA within lipid rafts and extracellular signal-regulated kinase activation through c-Cbl-associated protein," *Molecular and Cellular Biology*, vol. 27, no. 16, pp. 5686–5698, 2007.
- [155] Z. Kroner, "The relationship between Alzheimer's disease and diabetes: type 3 diabetes?" *Alternative Medicine Review*, vol. 14, no. 4, pp. 373–379, 2009.
- [156] W.-Q. Zhao, F. G. De Felice, S. Fernandez et al., "Amyloid beta oligomers induce impairment of neuronal insulin receptors," *FASEB Journal*, vol. 22, no. 1, pp. 246–260, 2008.
- [157] W. Q. Zhao and D. L. Alkon, "Role of insulin and insulin receptor in learning and memory," *Molecular and Cellular Endocrinology*, vol. 177, no. 1-2, pp. 125–134, 2001.
- [158] C. Wu, S. Butz, Y. S. Yingt, and R. G. W. Anderson, "Tyrosine kinase receptors concentrated in caveolae-like domains from neuronal plasma membrane," *Journal of Biological Chemistry*, vol. 272, no. 6, pp. 3554–3559, 1997.
- [159] J. Perroy, F. Raynaud, V. Homburger et al., "Direct interaction enables cross-talk between ionotropic and group I metabotropic glutamate receptors," *Journal of Biological Chemistry*, vol. 283, no. 11, pp. 6799–6805, 2008.
- [160] H. Khosravani, Y. Zhang, S. Tsutsui et al., "Prion protein attenuates excitotoxicity by inhibiting NMDA receptors," *Journal of General Physiology*, vol. 131, no. 6, p. i5, 2008.
- [161] N. Vassallo and J. W. Herms, "Cellular prion protein function in copper homeostasis and redox signalling at the synapse," *Journal of Neurochemistry*, vol. 86, no. 3, pp. 538–544, 2003.