



Asian Focus

# Survival after radical cystectomy for bladder cancer: Multicenter comparison between minimally invasive and open approaches



Weibin Xie <sup>a,b,1</sup>, Junming Bi <sup>a,b,1</sup>, Qiang Wei <sup>c</sup>, Ping Han <sup>c</sup>, Dongkui Song <sup>d</sup>, Lei Shi <sup>d</sup>, Dingwei Ye <sup>e,f</sup>, Yijun Shen <sup>e,f</sup>, Xin Gou <sup>g</sup>, Weiyang He <sup>g</sup>, Shaogang Wang <sup>h</sup>, Zheng Liu <sup>h</sup>, Jinhai Fan <sup>i</sup>, Kaijie Wu <sup>i</sup>, Zhiwen Chen <sup>j</sup>, Xiaozhou Zhou <sup>j</sup>, Chuize Kong <sup>k</sup>, Yang Liu <sup>k</sup>, Chunxiao Liu <sup>l</sup>, Abai Xu <sup>l</sup>, Baiye Jin <sup>m</sup>, Guanghou Fu <sup>m</sup>, Wei Xue <sup>n</sup>, Haige Chen <sup>n</sup>, Tiejun Pan <sup>o</sup>, Zhong Tu <sup>o</sup>, Tianxin Lin <sup>a,b,\*</sup>, Jian Huang <sup>a,b,\*</sup>

<sup>a</sup> Department of Urology, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, China

<sup>b</sup> Guangdong Provincial Key Laboratory of Malignant Tumor Epigenetics and Gene Regulation, Sun Yat-Sen Memorial Hospital, Sun Yat-Sen University, Guangzhou, China

<sup>c</sup> Department of Urology, West China Hospital, Sichuan University, Chengdu, China

<sup>d</sup> Department of Urology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China

<sup>e</sup> Department of Urology, Fudan University Shanghai Cancer Center, Shanghai, China

<sup>f</sup> Department of Oncology, Shanghai Medical College, Fudan University, Shanghai, China

<sup>g</sup> Department of Urology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China

<sup>h</sup> Department of Urology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

<sup>i</sup> Department of Urology, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China

<sup>j</sup> Department of Urology, First Affiliated Hospital of Army Medical University, Urology Institute of People Liberation Army, Chongqing, China

<sup>k</sup> Department of Urology, The First Hospital of China Medical University, Shenyang, China

<sup>l</sup> Department of Urology, Zhujiang Hospital of Southern Medical University, Guangzhou, China

<sup>m</sup> Department of Urology, The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou, China

<sup>n</sup> Department of Urology, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

<sup>o</sup> Department of Urology, General Hospital of Central Theater Command of the People's Liberation Army, Wuhan, China

\* Corresponding authors. Department of Urology, Sun Yat-Sen Memorial Hospital, Sun Yat-Sen University, Guangzhou, China.

E-mail addresses: [lintx@mail.sysu.edu.cn](mailto:lintx@mail.sysu.edu.cn) (T. Lin), [urohjh@sina.com](mailto:urohjh@sina.com) (J. Huang).

Peer review under responsibility of Second Military Medical University.

<sup>1</sup> These authors contributed equally to this study.

Received 12 November 2019; received in revised form 1 March 2020; accepted 26 April 2020  
Available online 14 June 2020

## KEYWORDS

Bladder cancer;  
Radical cystectomy;  
Minimally invasive  
surgery;  
Robotic surgery;  
Laparoscopy

**Abstract** *Objective:* To investigate oncological outcomes in patients with bladder cancer who underwent minimally invasive radical cystectomy (MIRC) or open radical cystectomy (ORC).

*Methods:* We identified patients with bladder cancer who underwent radical cystectomy (RC) in 13 centers of the Chinese Bladder Cancer Consortium (CBCC). Perioperative outcomes were compared between MIRC and ORC. The influence of surgical approaches on overall survival (OS) and cancer-specific survival (CSS) in the entire study group and subgroups classified according to pathologic stage or lymph node (LN) status was assessed with the log-rank test. Multivariable Cox proportional hazard models were used to evaluate the association among OS, CSS and risk factors of interest.

*Results:* Of 2 098 patients who underwent RC, 1 243 patients underwent MIRC (1 087 laparoscopic RC and 156 robotic-assisted RC, respectively), while 855 patients underwent ORC. No significant differences were noted in positive surgical margin rate and 90-day postoperative mortality rate. MIRC was associated with less estimated blood loss, more LN yield, higher rate of neobladder diversion, longer operative time, and longer length of hospital stay. There was no significant difference in OS and CSS according to surgical approaches ( $p=0.653$ , and  $0.816$ , respectively). Subgroup analysis revealed that OS and CSS were not significantly different regardless of the status of extravesical involvement or LN involvement. Multivariable Cox regression analyses showed that the surgical approach was not a significant predictor of OS and CSS.

*Conclusions:* Our study showed that MIRC was comparable to conventional ORC in terms of OS and CSS.

© 2020 Editorial Office of Asian Journal of Urology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Radical cystectomy (RC) with pelvic lymph node dissection (PLND) is the standard treatment for non-metastatic high-risk bladder cancer [1]. Although open radical cystectomy (ORC) has been the gold standard procedure, this complicated procedure is associated with significant perioperative morbidity and mortality [2]. Minimally invasive approaches such as laparoscopic radical cystectomy (LRC) and robotic-assisted radical cystectomy (RARC) have been increasingly performed in order to minimize surgical morbidity and improve recovery. Many studies have documented the surgical feasibility and potential benefits of LRC and RARC for bladder cancer. Minimally invasive surgery appeared to offer advantages to patients in terms of overall lower perioperative complications, decreased pain, reduced scarring, and faster recovery [3–5]. However, oncologic outcomes following minimally invasive radical cystectomy (MIRC) are still controversial because of the lack of adequate comparative data and sufficient follow-up. In recent years, several small sample RCTs reported no major difference in short-term oncologic outcomes [6–8], while a retrospective study based on the Surveillance, Epidemiology, and End Results (SEER) database reported that RARC was associated with a benefit in 2-year overall survival (OS) compared with ORC [9]. Recently, two studies published in

the *New England Journal of Medicine* found that minimally invasive radical hysterectomy was associated with shorter OS than open surgery among women with stage IA2 or IB1 cervical carcinoma [10,11]. Though the surgical procedures of cervical carcinoma were different from bladder cancer, these surprising results motivated us to reassess the value of minimally invasive surgery on cancer control.

Here, 13 centers of the Chinese Bladder Cancer Consortium (CBCC) [12] conducted a retrospective study to compare oncological outcomes in Asian patients with urothelial carcinoma of the bladder underwent MIRC or ORC.

## 2. Patients and methods

### 2.1. Patients selection

This study was approved by our institutional review board. Data of patients who were treated between 2007 and 2016 by experienced surgeons in 13 high-level centers of the CBCC were acquired from the CBCC database. We included patients with pathologic stage pT1-pT4a urothelial bladder cancer who had undergone RC and standard or extended PLND as primary treatment. The choice of surgical approach was mainly determined by the surgeon's preference and the patient's motivation. We excluded patients

for whom the surgical approach was unknown, those who had evidence of distant metastasis at the time of surgery, those who had a preexisting cancer diagnosis, and those who had a prior history of pelvic radiotherapy or surgery. Following exclusions, 2 098 patients remained for further analyses (Fig. 1).

## 2.2. Measures

Cancer registrars documented the primary surgical approaches as open, laparoscopic, or robotic-assisted. Patients who had undergone LRC or RARC with PLND were categorized as having undergone MIRC, even when the urinary diversion was performed as open procedures or when conversion to open surgery occurred. Demographic, perioperative, pathologic, and follow-up data were recorded by cancer registrars and ascertained through the end of 2018. Perioperative outcomes included the type of PLND, type of urinary diversion, operative time, length of hospital stays (LOS), positive surgical margins (PSM), adjuvant chemotherapy, number of lymph node (LN) yield, estimated blood loss (EBL), 90-day postoperative mortality. Oncologic outcomes included OS and CSS. OS and CSS were defined as the time from RC to death due to any cause and bladder cancer, respectively.

## 2.3. Statistical analysis


Mean and standard deviation (SD) or median and interquartile range (IQR) were used for the descriptions of

quantitative variables, and frequency and percentage were used for qualitative variables. Continuous variables were compared using the Student *t*-test or non-parametric tests, whereas categorical variables were compared using the Chi-square test. The influence of surgical approaches on OS and CSS in the entire study group and pathologic stage or LN status subgroups was analyzed using the Kaplan-Meier method, and differences were assessed with the log-rank test. Multivariable Cox proportional hazard models were used to evaluate the association among OS, CSS and risk factors of interest. A *p*-value <0.05 was considered statistically significant. Statistical analyses were conducted in Stata/SE 15.0 (Stata Corp LP, TX, USA).

## 3. Results

### 3.1. Patient and disease characteristics

A total of 2 098 patients were included in this study. Among these patients, 1 243 (59.2%) underwent minimally invasive surgery (1 087 LRC and 156 RARC, respectively), while 855 (40.8%) patients underwent open surgery. Patients and tumor characteristic were shown in Table 1. Quantity of surgery rapidly increased for both MIRC and ORC. Baseline characteristics including age, gender, body mass index (BMI), and ASA score were similar between MIRC and ORC ( $p=0.865, 0.917, 0.821, 0.147$ , respectively). There were no significant differences in pathologic stage, pathologic grade, and pathologic LN involvement rate between two treatment groups. The proportion of patients with



**Figure 1** Flowchart diagram depicting patient selection. PLND, pelvic lymph node dissection; CBCC, Chinese Bladder Cancer Consortium; MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.

**Table 1** Patient and tumor characteristics.

Characteristics	Minimally invasive RC <i>n</i> =1 243	Open RC <i>n</i> =855	<i>p</i> -Value
Year of surgery, <i>n</i> (%)			
2007–2010	118 (9.5)	137 (16.0)	<0.001
2011–2013	290 (23.3)	306 (35.8)	
2014–2016	835 (67.2)	412 (48.2)	
Age, year	63.4±10.1	63.5±10.2	0.865
Male, <i>n</i> (%)	1087 (87.5)	749 (87.6)	0.917
BMI, kg/m <sup>2</sup>	22.8±2.4	22.9±2.2	0.821
ASA score, <i>n</i> (%)			0.147
I+II	1079 (86.8)	713 (83.4)	
III+IV	164 (13.2)	142 (16.6)	
Pathologic stage, <i>n</i> (%)			0.587
<T2	346 (27.8)	244 (28.5)	
T2	566 (45.5)	390 (45.6)	
T3	239 (19.2)	148 (17.3)	
T4	92 (7.4)	73 (8.5)	
Pathologic grade, <i>n</i> (%)			
High grade	1065 (85.7)	745 (87.1)	0.341
Low grade	178 (14.3)	110 (12.9)	
Pathological nodal involvement, <i>n</i> (%)	193 (15.5)	120 (14.0)	0.346

RC, radical cystectomy; BMI, body mass index; ASA, the American Society of Anesthesiologists.

extravesical involvement ( $\geq pT3$ ) in MIRC and ORC groups was 26.6% and 25.95%, respectively, while LN-positive rate was 15.5% and 14.0%, respectively.

### 3.2. Perioperative outcome

Perioperative variables were shown in Table 2. The vast majority of patients received standard PLND. There was no significant difference in the proportion of patients who

underwent standard PLND (93.2% vs. 94.9%,  $p=0.130$ ). The types of urinary diversion were different between MIRC and ORC. A higher proportion of patients underwent MIRC received a diversion of neobladder (45.9% vs. 23.6%,  $p<0.001$ ). Conversion from minimally invasive surgery to open surgery was rare overall (2.2% of the cases). Minimally invasive surgery was associated with longer operative time and longer hospital stays. PSM occurred in 61 (4.9%) patients underwent MIRC and 43 (5.0%) patients underwent

**Table 2** Perioperative outcomes.

	Minimally invasive RC <i>n</i> =1 243	Open RC <i>n</i> =855	<i>p</i> -Value
PLND, <i>n</i> (%)			
Standard PLND	1 159 (93.2)	811 (94.9)	0.130
Extended PLND	84 (6.8)	44 (5.2)	
Urinary diversion, <i>n</i> (%)			<0.001
Ileal conduit	461 (37.1)	339 (39.7)	
Neobladder	571 (45.9)	202 (23.6)	
Ureterocutaneostomy	208 (16.7)	311 (36.4)	
Other	3 (0.2)	3 (0.4)	
Open conversion, <i>n</i> (%)	27 (2.2)	/	/
Operative time, min	360.6±120.3	288.6±90.3	<0.001
Hospital stays, day	28.0±9.9	25.0±10.0	<0.001
Positive surgical margin, <i>n</i> (%)	61 (4.9)	43 (5.0)	0.900
Adjuvant chemotherapy, <i>n</i> (%)	251 (20.2)	192 (22.5)	0.212
Lymph nodes yield	16.2±7.9	13.2±5.0	<0.001
Estimate blood loss, mL	402.7±517.8	623.1±692.9	<0.001
90-day postoperative mortality, <i>n</i> (%)	34 (2.7%)	24 (2.8%)	0.922

PLND, pelvic lymph node dissection; RC, radical cystectomy.

/, not required to provide.


ORC ( $p=0.900$ ). No difference was found in the percentage of patients who received adjuvant chemotherapy ( $p=0.212$ ). MIRC yielded more LN than ORC with a mean of 16.2 versus 13.2 ( $p<0.001$ ). ORC was associated with more estimate blood loss than MIRC. However, 90-day post-operative mortality was equivalent (1.1% vs. 1.2%,  $p=0.927$ ).

### 3.3. Oncologic outcome

The median follow-up duration was 43.2 (IQR: 28.3–54.7) months and 60.3 (IQR: 38.2–76.9) months for MIRC and ORC, respectively ( $p<0.001$ ). Patients underwent ORC had relatively longer follow-up duration. Overall, 476 (38.3%) patients in the MIRC group and 352 (41.2%) in the ORC group had died at the time of analysis, among whom 398 (32.0%), and 285 (33.3%), respectively, died from bladder cancer. There was no significant difference in OS and CSS according to surgical approaches ( $p=0.850$ , and 0.976, respectively, Fig. 2). Subgroup analysis by pathologic stage and LN status revealed that OS and CSS were similar for patients with organ-confined cancer ( $\leq pT2$ ) ( $p=0.257$ , and 0.277, respectively, Fig. 3), patients with extravesical involvement ( $\geq pT3$ ) ( $p=0.329$ , and 0.327, respectively Fig. 4), patients without LN involvement (LN negative) ( $p=0.361$  and 0.363, respectively, Fig. 5), and patients with LN involvement (LN-positive) ( $p=0.175$  and 0.119, respectively, Fig. 6). Multivariable Cox regression analyses revealed that pathologic tumor stage was independently associated with a significantly increased risk of death from all causes (Hazard ratio [HR]=1.84,  $p<0.001$ , Table 3) and death from bladder cancer (HR=2.41,  $p<0.001$ ). Other predictive factors for all-cause death and cancer-cause death included LN involvement (HR=2.46,  $p<0.001$ , and HR=2.62,  $p<0.001$ , respectively), PSM (HR=1.71,  $p=0.004$ , and HR=1.78,  $p=0.005$ , respectively), and age (HR=1.03,  $p<0.001$ , and HR=1.03,  $p<0.001$ , respectively). ASA score  $\geq 3$  was a risk factor for all-cause death (HR=1.28,  $p=0.028$ ) but not for cancer-cause death (HR=1.22,  $p=0.115$ ). The surgical approach was not an independent predictive factor for neither all-cause death nor cancer-cause death.


## 4. Discussion

Due to data limitations, LRC and RARC were still considered as investigational procedures for which no advantages in oncologic outcome could be shown as compared to ORC. Therefore, we conducted this multicenter study of minimally invasive surgery vs. open surgery, with a maximum follow-up of more than 10 years. A total of 2 098 patients who met the criterion were included. To our knowledge, this is the largest series studied oncologic outcome between MIRC and ORC in the Asian population. The baseline characteristics of the patients and tumor were well matched. All patients underwent RC and PLND by experienced surgeons from high-volume centers of the CBCC. Therefore, we minimized the bias introduced by patient selection and surgeon's experience and provided sound evidence comparing MIRC and ORC.



**Figure 2** Survival of the whole patient group underwent radical cystectomy, stratified by surgical approach (MIRC or ORC). (A) Overall survival (OS); (B) Cancer-specific survival (CSS). MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.


Consistent with other studies [13,14], we found that MIRC was associated with less EBL. The type of urinary diversion was determined by the preference of surgeons. Neobladder, which was a very complex procedure, was performed more often in MIRC than ORC (45.9% vs. 23.6%,  $p<0.001$ ). Thus, it was reasonable that the operative time was significantly longer in MIRC compared with ORC. The LOS was determined by several factors, including recovery protocol and postoperative complications. Because the CBCC database does not contain enough information about postoperative complications, we cannot determine the main reason of longer LOS for patients who underwent MIRC. Previous study comparing cancer control in patients with neobladder and ileal conduit showed no difference in CSS between the two groups when adjusting for pathological stage [15]. Neobladder in male and female patients does not compromise the oncological outcome of cystectomy.



**Figure 3** Survival of patients with organ-confined cancer ( $\leq pT2$ ), stratified by surgical approach (MIRC or ORC). (A) Overall survival (OS); (B) Cancer-specific survival (CSS). MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.


There were no significant differences in PSM rate between MIRC and ORC. PSM is an independent predictor of metastatic progression in patients undergoing radical cystectomy, which increased the risk of metastatic progression at 5 years from 32% to 74% [16]. The incidence of PSM ranged from 1% to 12.7% of the ORC cases, from 2% to 6% of the LRC cases, and from 0%–10.6% of the RARC cases [13,17]. In our trial, the PSM rate was 4.9% (61/1 263) in MIRC, and 5.0% (43/855) in ORC, respectively, which were similar to reported studies. Tumor infiltration may be the main limitation in achieving a negative margin [7,13]. PSM was associated with extravesical involvement tumor but not surgical approaches.

PLND is essential for both accurate staging and adequate local and regional control [18]. Several studies have taken LN yield as an indicator of the surgical quality of RC [4,19]. Although extensive PLND may yield more LN, it meanwhile associated with higher morbidity. It was still controversial for the optimal range of PLND [20,21]. It was widely




**Figure 4** Survival of patients with extravesical involvement ( $\geq pT3$ ), stratified by surgical approach (MIRC or ORC). (A) Overall survival (OS); (B) Cancer-specific survival (CSS). MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.

accepted that qualified RC should include at least standard PLND. We conducted standard PLND as routine and extensive PLND if necessary, for example, when patients were diagnosed with non-organ-confined or LN-positive cancer preoperatively or intraoperatively. The proportion of extensive PLND was similar between MIRC and ORC. Our result showed that MIRC yielded more LN than ORC, with a mean of 16.2 versus 13.2. Previous study [9] reported that RARC was associated with higher LN count than ORC, with a median of 17 versus 12, which was consistent with our study. Herr et al. [22] proposed that a PSM rate of less than 10% with a median of 10–14 LNs examined is a reasonable overall standard for RC. PSM rates for MIRC and ORC in this study were 4.9% and 5.0%, respectively ( $p=0.633$ ). Median LN yields were significantly lower in ORC than in MIRC ( $p<0.001$ ), but all gave adequate yields of  $>10$  LNs. We consider that on the basis of qualified PLND, LN yielded was not associated with better survival.



**Figure 5** Survival of patients without lymph node involvement (LN-positive), stratified by surgical approach (MIRC or ORC). (A) Overall survival (OS); (B) Cancer-specific survival (CSS). MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.

Long-term survival is the final touchstone for the oncologic efficiency of minimally invasive RC. A European LRC multicenter study [23] reported that actuarial recurrence-free survival (RFS), CSS and OS rates were 66%, 75% and 62% at 5 years and 62%, 55%, 38% at 10 years with a median follow-up of 50 months, which was comparable to our previously reported Asian LRC single-surgeon study [19]. Ten-year oncologic outcomes of RARC based on the International Robotic Cystectomy Consortium (IRCC) database seemed comparable to previously reported open series [24]. However, few studies have reported a comparison of survival between MIRC and ORC. In the present study, we noted that CSS and OS were not affected by surgical technique with a mean follow-up of 50.2 months, which was consistent with previous studies. Lin et al. [7] found no significant difference in RFS and OS between LRC and ORC with a median follow-up of 26 months. Recently, two RCTs also reported the oncologic outcome of RARC vs. ORC [6,8].



**Figure 6** Survival of patients with lymph node involvement (LN+), stratified by surgical approach (MIRC or ORC). (A) Overall survival (OS); (B) Cancer-specific survival (CSS). MIRC, minimally invasive radical cystectomy; ORC, open radical cystectomy.

The RAZOR trial [8] revealed that RARC was non-inferior to ORC for 2-year progression-free survival (PFS). Its update still showed no difference in recurrence, 3-year PFS and 3-year OS between RARC and ORC [25]. Bochner et al. [6] demonstrated no difference in RFS, OS, or CSS between RARC and ORC with a median follow-up of 4.9 years. Retrospective comparative studies also found that MIRC and ORC provide similar oncologic outcomes [26–28]. Most of the previous studies showed that MIRC was equivalent to ORC on cancer control. However, Hanna et al. [9] reported that RARC was associated with a 7.7% higher OS rate at 2 years based on the SEER database. In contrast to other studies, we additionally performed subgroup analysis according to pathologic stage and LN status, confirmed that there was no significant difference between minimally invasive and open approach with regard to OS and CSS.

The present study showed that pathologic tumor stage, LN status, and presences of PSM predicted OS and CSS by

**Table 3** Multivariable Cox proportional hazard regression analyses.

characteristics	OS			CSS		
	HR	95% CI	p-Value	HR	95% CI	p-Value
Year of surgery						
2007–2010	Reference					
2011–2013	1.03	0.88–1.21	0.703	1.01	0.85–1.16	0.889
2014–2016	0.93	0.76–1.08	0.232	0.96	0.81–1.15	0.686
Age, year	1.03	1.02–1.04	<0.001	1.03	1.01–1.04	<0.001
Gender						
Male	Reference					
Female	1.15	0.85–1.55	0.498	1.18	0.84–1.66	0.342
BMI kg/m <sup>2</sup>	0.96	0.93–1.00	0.071	0.99	0.95–1.04	0.802
ASA group						
I+II	Reference					
III+IV	1.28	1.03–1.60	0.028	1.22	0.95–1.57	0.115
Surgical approach						
Open	Reference					
Minimally invasive	0.95	0.77–1.17	0.653	0.97	0.77–1.23	0.816
Pathologic stage						
≤T2	Reference					
≥T3	1.84	1.46–2.31	<0.001	2.41	1.86–3.13	<0.001
Pathologic grade						
LG	Reference					
HG	1.26	0.88–1.80	0.209	1.07	0.72–1.59	0.749
Pathological nodal involvement						
No	Reference					
Yes	2.46	1.93–3.13	<0.001	2.62	2.00–3.42	<0.001
Positive surgical margin						
No	Reference					
Yes	1.71	1.19–2.46	0.004	1.78	1.19–2.66	0.005
PLND						
Standard PLND	Reference					
Extensive PLND	1.26	0.93–1.72	0.135	1.30	0.92–1.83	0.139

BMI, body mass index; OS, overall survival; CSS, cancer-specific survival; HR, hazard ratio; CI, confidence interval; ASA, the American Society of Anesthesiologists; PLND, pelvic lymph node dissection.

multivariable cox regression. Age was also a predictive factor associated with OS and CSS because elderly patients were more likely to have high-risk tumor and inactive follow-up. Consistent with our study, several previous studies [16,19,29] also demonstrated that these prognostic factors were associated with oncologic outcomes in patients who underwent RC. However, the type of surgical approach for RC was not a significant predictor of OS or CSS.

This study has several limitations. Foremost, the retrospective nature of this study should be considered. There might have been a selection bias and surgeon bias. The type of surgical approach is mainly determined by surgeon's preference and patient's characteristic. The uneven distribution among ORC and MIRC between centers could have affected the observed differences. The study period is long, and it remains unclear whether this factor would influence the outcomes. Moreover, the database does not provide enough information about postoperative complications, recovery protocol, extracorporeal or intracorporeal diversion, so we cannot provide a complete depiction of some perioperative outcomes. Additionally, although most of these patients received regular follow-up and physical

check, we still cannot accurately document the date of recurrence and metastasis of some patients, so RFS and cancer-free survival were not examined in our study.

It is so far the study with the longest follow-up period and the largest sample-size comparing minimally invasive and open radical cystectomy in the Asian population. All the surgeons are from Chinese high-level urological centers and have extensive experience in minimally invasive and open surgery. Although our results showed no difference in the survival of bladder cancer between these two groups, it does not mean that minimally surgery is generally equivalent to open surgery. It requires more skill, more training, and a longer learning curve. Therefore, the interpretation of the results must be more careful.

## 5. Conclusion

Minimally invasive cystectomy, such as RARC or LRC, was comparable to ORC with regard to OS and CSS. MIRC can be performed as an alternative to ORC. Future large sample size RCTs with longer follow-up are needed to provide more convincing survival outcomes.

## Author contributions

*Study concept and design:* Tianxin Lin, Jian Huang.

*Data acquisition:* Weibin Xie, Junming Bi, Qiang Wei, Ping Han, Dongkui Song, Lei Shi, Dingwei Ye, Yijun Shen, Xin Gou, Weiyang He, Shaogang Wang, Zheng Liu, Jinhai Fan, Kaijie Wu, Zhiwen Chen, Xiaozhou Zhou, Chuize Kong, Yang Liu, Chunxiao Liu, Abai Xu, Baiye Jin, Guanghou Fu, Wei Xue, Haige Chen, Tiejun Pan, Zhong Tu.

*Data analysis:* Weibin Xie, Junming Bi.

*Drafting of manuscript:* Weibin Xie.

*Critical revision of the manuscript:* Tianxin Lin, Jian Huang.

## Conflicts of interest

The authors declare no conflict of interest.

## Acknowledgements

This study was supported by the National Natural Science Foundation of China (Grant No.81825016, 81772719, 81772728, 81572514), the Key Areas Research and Development Program of Guangdong (Grant No. 2018B010109006), and Medical Scientific Research Foundation of Guangdong Province (Grant No. A2018388).

## References

- [1] Milowsky MI, Rumble RB, Booth CM, Gilligan T, Eapen LJ, Hauke RJ, et al. Guideline on muscle-invasive and metastatic bladder cancer (European association of urology guideline): American society of clinical oncology clinical practice guideline endorsement. *J Clin Oncol* 2016;34:1945–52.
- [2] Lowrance WT, Rumohr JA, Chang SS, Clark PE, Smith Jr JA, Cookson MS. Contemporary open radical cystectomy: analysis of perioperative outcomes. *J Urol* 2008;179:1313–8.
- [3] Modi PK, Hollenbeck BK, Oerline M, Weizer AZ, Montgomery JS, Kaffenberger SD, et al. Real-world impact of minimally invasive *versus* open radical cystectomy on perioperative outcomes and spending. *Urology* 2019;125:86–91.
- [4] Nix J, Smith A, Kurpad R, Nielsen ME, Wallen EM, Pruthi RS. Prospective randomized controlled trial of robotic versus open radical cystectomy for bladder cancer: perioperative and pathologic results. *Eur Urol* 2010;57:196–201.
- [5] Rai BP, Bondad J, Vasdev N, Adshead J, Lane T, Ahmed K, et al. Robotic versus open radical cystectomy for bladder cancer in adults. *Cochrane DB Syst Rev* 2019;4:CD011903. <https://doi.org/10.1002/14651858.CD011903.pub2>.
- [6] Bochner BH, Dalbagni G, Marzouk KH, Sjoberg DD, Lee J, Donat SM, et al. Randomized trial comparing open radical cystectomy and robot-assisted laparoscopic radical cystectomy: oncologic outcomes. *Eur Urol* 2018;74:465–71.
- [7] Lin T, Fan X, Zhang C, Xu K, Liu H, Zhang J, et al. A prospective randomised controlled trial of laparoscopic vs open radical cystectomy for bladder cancer: perioperative and oncologic outcomes with 5-year follow-up. *Br J Canc* 2014;110:842–9.
- [8] Parekh DJ, Reis IM, Castle EP, Gonzalgo ML, Woods ME, Svatek RS, et al. Robot-assisted radical cystectomy versus open radical cystectomy in patients with bladder cancer (RAZOR): an open-label, randomised, phase 3, non-inferiority trial. *Lancet* 2018;391:2525–36.
- [9] Hanna N, Leow JJ, Sun M, Friedlander DF, Seisen T, Abdollah F, et al. Comparative effectiveness of robot-assisted vs. open radical cystectomy. *Urol Oncol* 2018;36:88.e81–9. <https://doi.org/10.1016/j.urolonc.2017.09.018>.
- [10] Melamed A, Margul DJ, Chen L, Keating NL, Del Carmen MG, Yang J, et al. Survival after minimally invasive radical hysterectomy for early-stage cervical cancer. *N Engl J Med* 2018;379:1905–14.
- [11] Ramirez PT, Frumovitz M, Pareja R, Lopez A, Vieira M, Ribeiro R, et al. Minimally Invasive *versus* abdominal radical hysterectomy for cervical cancer. *N Engl J Med* 2018;379:1895–904.
- [12] Li K, Lin T, Xue W, Mu X, Xu E, Yang X, et al. Current status of diagnosis and treatment of bladder cancer in China -analyses of Chinese Bladder Cancer Consortium database. *Asian J Urol* 2015;2:63–9.
- [13] Iwata T, Kimura S, Foerster B, Fossati N, Briganti A, Karakiewicz PI, et al. Oncologic outcomes after robot-assisted versus open radical cystectomy: a systematic review and meta-analysis. *World J Urol* 2019;37:1557–70.
- [14] Tang K, Li H, Xia D, Hu Z, Zhuang Q, Liu J, et al. Laparoscopic versus open radical cystectomy in bladder cancer: a systematic review and meta-analysis of comparative studies. *PLoS One* 2014;9:e95667. <https://doi.org/10.1371/journal.pone.0095667>.
- [15] Yossepowitch O, Dalbagni G, Golijanin D, Donat SM, Bochner BH, Herr HW, et al. Orthotopic urinary diversion after cystectomy for bladder cancer: implications for cancer control and patterns of disease recurrence. *J Urol* 2003;169:177–81.
- [16] Dotan ZA, Kavanagh K, Yossepowitch O, Kaag M, Olgac S, Donat M, et al. Positive surgical margins in soft tissue following radical cystectomy for bladder cancer and cancer specific survival. *J Urol* 2007;178:2308–13.
- [17] Chade DC, Laudone VP, Bochner BH, Parra RO. Oncological outcomes after radical cystectomy for bladder cancer: open versus minimally invasive approaches. *J Urol* 2010;183:862–9.
- [18] Abol-Enein H, El-Baz M, Abd El-Hameed MA, Abdel-Latif M, Ghoneim MA. Lymph node involvement in patients with bladder cancer treated with radical cystectomy: a patho-anatomical study-a single center experience. *J Urol* 2004;172:1818–21.
- [19] Huang J, Lin T, Liu H, Xu K, Zhang C, Jiang C, et al. Laparoscopic radical cystectomy with orthotopic ileal neobladder for bladder cancer: oncologic results of 171 cases with a median 3-year follow-up. *Eur Urol* 2010;58:442–9.
- [20] Bruins HM, Veskimae E, Hernandez V, Imamura M, Neuberger MM, Dahm P, et al. The impact of the extent of lymphadenectomy on oncologic outcomes in patients undergoing radical cystectomy for bladder cancer: a systematic review. *Eur Urol* 2014;66:1065–77.
- [21] Gschwend JE, Heck MM, Lehmann J, Rubben H, Albers P, Wolff JM, et al. Extended *versus* limited lymph node dissection in bladder cancer patients undergoing radical cystectomy: survival results from a prospective, randomized trial. *Eur Urol* 2018;75:604–11.
- [22] Herr H, Lee C, Chang S, Lerner S. Standardization of radical cystectomy and pelvic lymph node dissection for bladder cancer: a collaborative group report. *J Urol* 2004;171:1823–8.
- [23] Albinssi S, Rassweiler J, Abbou CC, Cathelineau X, Chlosta P, Fossion L, et al. Long-term analysis of oncological outcomes after laparoscopic radical cystectomy in Europe: results from a multicentre study by the European Association of Urology (EAU) section of Uro-technology. *BJU Int* 2015;115:937–45.
- [24] Hussein AA, Elsayed AS, Aldhaam NA, Jing Z, Osei J, Kaouk J, et al. Comparison of long-term oncologic outcomes among historical open and minimally invasive retrospective studies. *J Urol* 2019;202:927–35.
- [25] Venkatramani V, Reis IM, Castle EP, Gonzalgo ML, Woods ME, Svatek RS, et al. Predictors of recurrence, progression-free

- and overall survival following open *versus* robotic radical cystectomy: analysis from the RAZOR Trial with a 3-year follow-up. *J Urol* 2020;203:522–9.
- [26] Gandaglia G, Karl A, Novara G, de Groot R, Buchner A, D'Hondt F, et al. Perioperative and oncologic outcomes of robot-assisted vs. open radical cystectomy in bladder cancer patients: a comparison of two high-volume referral centers. *Eur J Surg Oncol* 2016;42:1736–43.
- [27] Kim TH, Sung HH, Jeon HG, Seo SI, Jeon SS, Lee HM, et al. Oncological outcomes in patients treated with radical cystectomy for bladder cancer: comparison between open, laparoscopic, and robot-assisted approaches. *J Endourol* 2016;30:783–91.
- [28] Faraj KS, Abdul-Muhsin HM, Rose KM, Navaratnam AK, Patton MW, Eversman S, et al. Robot assisted radical cystectomy vs open radical cystectomy: over 10 years of the Mayo clinic experience. *Urol Oncol* 2019;37:862–9.
- [29] Zhang L, Wu B, Zha Z, Qu W, Zhao H, Yuan J, et al. Clinicopathological factors in bladder cancer for cancer-specific survival outcomes following radical cystectomy: a systematic review and meta-analysis. *BMC Cancer* 2019;19:716. <https://doi.org/10.1186/s12885-019-5924-6>.