

A new species of *Labidocera* (Copepoda, Calanoida, Pontellidae) collected from Okinawa, southwestern Japan, with establishment of five Indo-West Pacific species groups in the *L. detruncata* species complex

Takeshi Hirabayashi¹, Susumu Ohtsuka¹

¹ Takehara Marine Science Station, Setoichi Field Center, Graduate School of Biosphere Science, Hiroshima University, 5-8-1 Minato-machi, Takehara, Hiroshima 725-0024, Japan

Corresponding author: Takeshi Hirabayashi (1810take@gmail.com)

Academic editor: Danielle Defaye | Received 25 June 2014 | Accepted 12 September 2014 | Published 16 October 2014

<http://zoobank.org/933AE05C-5587-40C8-BFA6-56A2080573F2>

Citation: Hirabayashi T, Ohtsuka S (2014) A new species of *Labidocera* (Copepoda, Calanoida, Pontellidae) collected from Okinawa, southwestern Japan, with establishment of five Indo-West Pacific species groups in the *L. detruncata* species complex. ZooKeys 447: 21–34. doi: 10.3897/zookeys.447.8171

Abstract

Labidocera churaumi sp. n. is described from Okinawa, southwestern Japan. The female of the new species differs from other congeners in genital compound somite with right postero-lateral and left antero-lateral processes. The male is distinguished from other congeners by the structure of the fifth leg. This new species is assigned to a newly proposed species group, the *L. madurae* species group, within the *L. detruncata* species complex. In this species complex five Indo-West Pacific species groups are recognized (*cervi*, *detruncata*, *gangetica*, *madurae*, and *pavo*) and defined on the basis of difference in sexual dimorphism.

Keywords

Indo-West Pacific, *Labidocera*, Okinawa, Pontellidae, species group, taxonomy

Introduction

We have been carrying out intensive faunistic surveys of marine invertebrates around the Nansei Islands, in the subtropical region of southwestern Japan, since 1988 and have discovered many new crustacean taxa, especially copepods. For example the copepod order Platycopioidea was first reported from the Indo-Pacific in this region

in 1994 (Ohtsuka and Boxshall 1994). The systematics of shallow- and deep-water copepods from these waters, have been a major focus of research and some of discoveries have been important in elucidating aspects of the evolutionary history and zoogeography of copepods (Barr and Ohtsuka 1989, Ohtsuka et al. 1991, 1996, 1998, 2002, 2003, 2004, 2005, Barthélémy et al. 1998).

In May 2011 we found an undescribed species of the calanoid genus *Labidocera* Lubbock, 1853 (Family Pontellidae) from Okinawa Island and neighboring islands. It clearly belongs to the *L. detruncata* species complex (Fleminger 1967, Greenwood and Othman 1979, Mulyadi 2002), due to its characteristic sexual dimorphism. This species complex mainly consists of coastal species from tropical and subtropical Indo-West Pacific waters and also two Atlantic species (Fleminger 1967, Greenwood and Othman 1979, Mulyadi 2002). The present paper provides a description of the new species, and remarks on species groups within the *detruncata* species complex.

Materials and methods

A fish collection light (KU-5MB, MW50S-G, KOTO electric Co., Ltd.) was deployed at Naha Port (May 2011) and Tokashiki Port (May 2011) after sunset. Conical plankton nets (diameter 30 cm, mesh size 0.1 mm) were towed around the light several times. Copepod specimens were fixed with 10% neutralized formalin/seawater or 70% ethanol immediately after collection. Copepods were dissected under a binocular microscope and examined and illustrated using a compound microscope fitted with differential interference contrast lighting (Optiphot, Nikon Co., Ltd.) and a drawing tube.

In describing the features of the new species, we have followed the terminology of Huys and Boxshall (1991). We followed Fleminger (1967) and Fleminger et al. (1982) about species complexes (=superspecies) and groups in the genus *Labidocera*.

Type specimens are deposited at the Kitakyushu Natural History and Human History Museum (KMNH IvR 500,734 – KMNH IvR 500,783).

Systematics

Order Calanoida Sars, 1903

Family Pontellidae Dana, 1853

Genus *Labidocera* Lubbock, 1853

Labidocera churaumi sp. n.

<http://zoobank.org/E4C58DC0-DDE3-4808-B7D1-5925D444280B>

Figs 1–4

Material examined. Tokashiki Port, Tokashiki Island, Okinawa Prefecture, Japan, (26°12'0.98"N; 127°22'8.77"E), 21 May 2011 (8 ♀♀, 3 ♂♂); (26°12'1.21"N;




Figure 1. *Labidocera churaumi* sp. n., female (Paratype). **A** habitus, dorsal view **B** habitus, lateral view **C** anterior part of cephalosome, lateral view **D** urosome with outline of attached spermatophore and coupler, dorsal view **E** left antennule **F** antenna.

127°22'10.20"E), 27 May 2012 (21♀♀, 11♂♂). Naha New Port, Okinawa Prefecture, Japan, (26°14'8.22"N; 127°40'47.56"E), 20 May 2011, (1 ♀, 6 ♂♂).

Types. Holotype : 1♀, Tokashiki Port, 27 May 2012, whole specimen (KMNH IvR 500,759). Allotype: 1♂Tokashiki Port, 27 May 2012, whole specimen (KMNH

IvR 500,783). Paratypes: 1♀, 6♂♂, Naha New Port, 20 May 2011, whole specimen (♀ KMNH IvR 500,734; ♂♂ KMNH IvR 500,764-KMNH IvR 500,769); 8♀♀, 3♂♂, Tokashiki Port, 21 May 2011 partly dissected and mounted on 11 glass slides (♀♀KMNH IvR 500,735-KMNH IvR 500,742; ♂♂KMNH IvR 500,770-KMNH IvR 500,772); 20♀♀, 10♂♂, Tokashiki Port, 27 May 2012, whole specimen (♀♀KMNH IvR 500,743-KMNH IvR 500,758 and KMNH IvR 500,760-KMNH IvR 500,763; ♂♂KMNH IvR 500,773-KMNH IvR 500,782).

Type locality. Tokashiki Port, Tokashiki Island, Okinawa Prefecture, Japan (26°12'1.21"N; 127°22'10.20"E).

Female. Body (Fig. 1A, B) length of females ranging between: 2225 and 2790 μm (average 2475 μm, n=29), measured from frontal margin of cephalosome to end of caudal rami excluding caudal setae. Ratio of prosome to urosome lengths 4:1, prosome length to width ratio 2.85:1. Cephalic profile rounded in dorsal view, without lateral cephalic hooks. Paired dorsal eyes with cuticular lenses; protuberant ventral eye extending anteroventrally between rostral processes (Fig. 1C). Rostrum bifid, directed posteroventrally. Posterior margins of prosome almost symmetrical in dorsal view, tapering to simple abbreviated, pointed process at each lateral corner. Urosome 2-segmented of highly characteristic shape. Genital compound somite strongly asymmetrical; anterior left surface with posteriorly-directed rod-like process and posterior right smoothly rounded. Spermatophore (Fig. 1D) attached dorsally to genital compound somite.

Antennules (Fig. 1E) symmetrical, 23-segmented: segments armed as follows (Arabic numbers=setae; sp=spines, aes=aesthetasc): (I) 2+aes, (II-IV) 4+aes, (V) 2+aes, (VI) 2, (VII) 2+aes, (VIII-IX) 4+aes, (X) 2, (XI) 2+aes, (XII) 1+sp, (XIII) 1+sp+aes, (XIV) 1+sp+aes, (XV) 2+aes, (XVI) 2+aes, (XVII) 1+sp+aes, (XVIII) 2+aes, (XIX) 1+sp+aes, (XX) 2+aes, (XXI) 2+aes, (XXII) 1, (XXIII) 1, (XXIV) 1+sp, (XXV) 1+sp+aes, (XXVI-XXVIII) 6. Larger and longer setae on segments 3-6. Antenna (Fig. 1F) biramous: coxa with short plumose distal seta, basis and first endopodal segment fused to form elongate allobasis, setation formula 2, 2. Compound distal endopodal segment with 9 and 7 setae on proximal and distal lobes, respectively; exopod 5-segmented, setation formula 0, 0, 2, 2, 3. Mandible (Fig. 2A) with wide, heavily chitinized gnathobase; mandibular palp biramous, basis robust, armed with 4 inner setae. Endopod 2-segmented, first segment armed with 1 short and 3 long setae; second segment with 7 terminal setae. Exopod 2-segmented, first segment unarmed, second segment with 6 terminal setae. Mandibular gnathobase distal edge bearing 8 teeth comprising: from ventral margin 1 apical, 1 subapical, 3 compound medial, and 3 basal (see Fig. 2A); medial teeth with bifurcated cusps; dorsal end of gnathobase with 1 seta. Maxillule (Fig. 2B) praecoxal arthrite with 15 setal elements, 4 on posterior surface; coxal endite with 2 long and 1 short elements on endite and 9 setae on epipodite; basis with 3 setae on proximal and distal endites; and 1 large seta on basal exite; proximal endopod segment and endopod segment 2 incorporated into basis, proximal endopod segments with 2 setae, endopod segment 2 with 2 setae and distal endopod segment with 5 apical setae; exopod with 10 setae. Maxilla (Fig. 2C) with first praecoxal endite bearing 6 setae, second with 3 seta; coxa with 3 setae each



Figure 2. *Labidocera churaumi* sp. n., female (Paratype). **A** mandible **B** maxillule **C** maxilla **D** maxilliped.

on proximal and distal endites. Basis with 3 setae; endopod 3-segmented, setal formula of endopod: 1, 1, 4. Maxilliped (Fig. 2D) with praecoxa and coxa fused, three syncoxal endites well developed, with setal formula 2, 2, 4; endite setae strong, spinulose. Basis

fringed with medial row of spiniform processes and 2 distal setae. Endopod 4-segmented, setal formula of endopod as: 2, 1, 1, 2.

Legs 1–4 (Fig. 3A–D) with 2-segmented endopods and 3-segmented exopods. Coxae with plumose inner seta. Seta and spine formula (Arabic numbers=setae, Roman numerals=spines) of legs 1–4 as follows:

	Coxa	Basis	Exopod	Endopod
Leg 1	0-1	0-0	I-1; I-1; II, I, 4	0-3; 1, 2, 3
Leg 2	0-1	0-0	I-1; I-1; III, I, 5	0-3; 2, 2, 4
Leg 3	0-1	0-0	I-1; I-1; III, 1, 5	0-3; 2, 2, 4
Leg 4	0-1	0-0	I-1; I-1; III, 1, 5	0-3; 2, 2, 3

Leg 5 (Fig. 3E) biramous, slightly asymmetrical; coxa and intercoxal sclerite fused. Basis subrectangular, with posterior seta. Endopod rounded distally, about 0.3 times as long as exopod. Exopods of both legs 1-segmented, bifurcated tip and with 2 outer spines; outer process on left slightly larger than right and with small spine-like process on proximal part.

Male. Body (Fig. 4A, B) slightly smaller than female (1819–2531 μm , average: 2219 μm , $n=20$). Prosome about 4 times as long as urosome, Urosome (Fig. 4A) symmetrical with 5 somites; anal somite and caudal rami asymmetrical.

Right antennule (Fig. 4C) with 15 segments geniculate between segments 11 and 12, reaching middle of third pedigerous somite. Antennular segments armed as follows (Arabic numbers=setae; sp=spines, aes=aesthetasc): (I) 2+aes, (II–IV) 4+aes, (V) 2+aes, (VI) 2, (VII) 2+aes, (VIII–XIV) 8+6sp+4aes, (XV–XVI) 4+2aes, (XVII) 2+aes, (XVIII) 2+aes, (XIX) 1+aes, (XX) 1+aes, (XXI–XXIII) 2+aes, (XXIV) 2, (XXV) 2+aes, (XXVI–XXVIII) 6; Segments 11 and 12 with row of teeth.

Left antennule, antenna, mouthparts and swimming legs as in female.

Leg 5 (Fig. 4D) asymmetrical. Left leg 5 short; intercoxal sclerite and left coxa fused. Basis cylindrical with seta near base. Exopod 2-segmented: first segment cylindrical; second segment triangular short with protruding hairy medial surface and 3 distal and 1 lateral spines, one of them long. Right leg 5 basis with seta. Exopod 2-segmented, forming chela; thumb of chela large, triangular, arising near base of first exopodal segment. First exopodal segment with 2 small setae. Second exopodal segment elongate and curved, with 3 slender marginal setae.

Remarks. The present new species is similar to *Labidocera madurae* Scott, 1909 and *L. tasmanica* Taw, 1974 in having the following features: (1) the posterolateral margins of the prosome are symmetrical, each triangular with a sharply pointed tip; (2) the female urosome is moderately or markedly asymmetrical; (3) the caudal rami are symmetrical and not highly modified; (4) the endopods of female leg 5 are nearly symmetrical, short, conical, and not bifid at the tip; (5) the thumb of the right leg 5 of the male is triangular with a broad base, and is slightly recurved; (6) the distal part of terminal segment of the left leg 5 of the male bears 3 spines, the outermost of which is the longest. These 3 species constitute a species group within the *Labidocera detruncata* species complex (see Discussion). *Labidocera churaumi* sp. n. can be dis-




Figure 3. *Labidocera churaumi* sp. n., female (Paratype). **A** leg 1 **B** leg 2 **C** leg 3 **D** leg 4 **E** leg 5.

tinguished from *L. madurae* and *L. tasmanica* by: (1) the presence of right postero-lateral and left antero-lateral processes on the female genital compound somite; (2) the exopod of the female leg 5 is very short, only as long as the basis, and has a bi-




Figure 4. *Labidocera churaumi* sp. n., male (Paratype). **A** habitus, dorsal view **B** habitus, lateral view **C** right antennule **D** leg 5.

furcated tip on both sides; (3) the inner margin of the terminal segment of the male left leg 5 has a protrusion at mid-length.

Etymology. The new specific name “*churaumi*” is from an Okinawan dialect, meaning the beautiful seas around the type locality Okinawa.

Discussion

Fleminger (1967) classified species of the genus *Labidocera* into four superspecies (=species complex), the *L. wilsoni* Fleminger & Tan, 1966, *L. detruncata* (Dana, 1849), *L. darwini* Lubbock, 1853, and *L. krøyeri* (Brady, 1883) species complexes, but left some species unassigned. Within the *L. detruncata* species complex he recognized nine species. Subsequently Greenwood and Othman (1979) added further three species to this species complex, and compared nine Indo-West Pacific members of the species complex, but did not consider three species: *L. orsinii* Giesbrecht, 1889, *L. gangetica* Sewell, 1934 and *L. nerii* (Krøyer, 1849) that Fleminger (1967) originally assigned to this species complex. Subsequently Greenwood and Othman (1979) and Othman (1986) added new species: *L. farrani* Greenwood & Othman, 1979 and *L. jaafari* Othman, 1986 to the species complex. Mulyadi (2002) was the first to define the Indo-West Pacific species group within the *L. detruncata* species complex which he referred to as *L. detruncata* species group, in which were accommodated the following ten species: *L. detruncata*, *L. pavo* Giesbrecht, 1889, *L. bataviae* Scott, 1909, *L. madurae* Scott, 1909, *L. cervi* Krämer, 1895, *L. caudata* Nicholls, 1944, *L. sinilobata* Shen & Lee, 1963, *L. tasmanica* Taw, 1974, *L. farrani* Greenwood & Othman, 1979, and *L. jaafari*. One remaining issue is that two Atlantic species (*L. orsinii* and *L. nerii*) and the Indian species (*L. gangetica*) were not assigned to this species complex by Mulyadi (2002). Here we reinstate Fleminger's (1967) inclusion of the two Atlantic species (*L. orsinii* and *L. nerii*) and the Indian species (*L. gangetica*).

Therefore Mulyadi's (2002) definition of the Indo-West Pacific *L. detruncata* species group needs some emendations as follows: (1) the posterolateral prosomal corners of both sexes protrude posteriorly into a pointed tip; (2) the female urosome is slightly or distinctly asymmetrical, about 1/6 to 1/4 as long as prosome, and 2- or 3-segmented; (3) the rostrum of both sexes is widely divided; (4) the caudal rami of the female are slightly or remarkably asymmetrical, broadened, with or without one or more thickened setae; (5) the exopods of female fifth legs are asymmetrical, with each bearing 3 lateral and 2 terminal processes; (6) the endopods of female fifth legs are either simply conical (rarely bifid) distally or are totally reduced; (7) the thumb of the right leg 5 of the male is conical or spatulate; (8) the finger of the right leg 5 of the male is slender; (9) the terminal segment of the left leg 5 of the male bears 1 outer and 3 slender terminal (rarely thick in *L. caudata*) elements. Since the two Atlantic and *L. gangetica* comply with this emended diagnosis, Fleminger's (1967) original *L. detruncata* species complex (Mulyadi's (2002) Indo-West Pacific *L. detruncata* species group plus *L. orsinii*, *L. nerii* and *L. gangetica*) is well defined by this amended diagnosis.

Although Mulyadi (2002) defined the Indo-West Pacific *L. detruncata* species group (the ten species listed above), it can be further subdivided into the following five newly proposed species groups on the basis of variation in the differing expressions of sexual dimorphism. *Labidocera sinilobata*, *L. jaafari* and *L. gangetica* share synapomorphies in variation in the differing expressions of sexual dimorphism, viz., (1) the absence of endopods from leg 5 of the female, (2) the thumb and finger of the right leg 5 of the male slender, (3) there is a rounded process present basal to the thumb of the right male




Figure 5. Distribution of *Labidocera detruncata* species complex based on previous records and present study.

leg 5, and (4) there is a protrusion on the inner surface of the terminal segment of the left male leg 5. These three species have a restricted distribution in the subtropical and tropical, coastal waters of the Indo-West Pacific (Sewell 1912, Shen and Lee 1963, Silas and Pillai 1973, Othman 1986, Mulyadi 2002, Razouls et al. 2014), and are referred to here as *L. gangetica* species group (Fig. 5).

Two species, *L. cervi* and *L. caudata* from Oceania are unusual in both having a triangular process distally on the terminal segment of the male left leg 5. The first exopodal segment of male leg 5 bears an outer subterminal process in both species, although it is not certain whether these are homologous. In the female the exopod of leg 5 bears 3 distinct lateral and 2 terminal prominences, while the endopod is simply spiniform. The posterolateral prosomal corners of *L. cervi* are remarkably large compared to those of *L. caudata*. These two species are considered here as the *L. cervi* species group. McKinnon and Kimmerer (1984) pointed out the similarity in the fifth legs of both sexes of *L. caudata* and *L. madurae*, but we regard the unique structure of the terminal exopodal segment of the male left leg 5 as a robust synapomorphy to define the species group.

Labidocera detruncata is most closely related to *L. farrani* in sharing the following synapomorphies: (1) the complex, dorsal swelling on the genital compound somite of the female; (2) the anal somite of the female protruded mid-posteriorly; (3) the caudal rami of the female are widely separated, and the right ramus is larger than the left; (4) the basis of the female leg 5 is swollen; (5) the male right leg 5 has a spatulate thumb; (6) the terminal segment of the male left leg 5 has 4 elements, the second outermost of which is the longest. These two species belong to *L. detruncata* species group *sensu stricto*. *Labidocera detruncata* is widely distributed in oceanic waters of the Indo-Pacific

and West Atlantic regions, while *L. farrani* has a distribution in coastal waters of Indo-West Pacific (Silas and Pillai 1973, Greenwood and Othman 1979, Mulyadi 2002, Jeong et al. 2009, Razouls et al. 2014) (Fig. 5).

Labidocera pavo and *L. bataviae* share the following features in the female: (1) the female caudal rami are broadly separated and posterolaterally expanded; (2) the exopod of the female leg 5 is slender, with 3 lateral and 2 terminal distinct prominences; (3) the endopod of the female leg 5 is short, at most 1/3 to 1/4 as long as the exopod; (4) the thumb of the first exopodal segment of the male right leg 5 is bifid; (5) the terminal exopodal segment of the male left leg 5 is slender, and carries 3 fine elements. These two species belong to the *L. pavo* species group and they are broadly distributed in coastal waters of the temperate to tropical Indo-Pacific regions (Silas and Pillai 1973, Razouls et al. 2014) (Fig. 5).

As already mentioned in “Remarks”, *L. madurae*, *L. tasmanica* and *L. churaumi* sp. n. together belong to the *L. madurae* species group. This species group has a restricted distribution in tropical to temperate waters of the Indo-Pacific (Scott 1909, Silas and Pillai 1973, Taw 1974, Razouls et al. 2014, present study) (Fig. 5). *Labidocera detruncata* is widely distributed in the Indo-West Pacific, while *L. pavo* has a narrow coastal distribution in the region. In addition *L. sinilobata* is restricted to the West Pacific, whereas *L. gangetica* occurs in the Indian Ocean. Such restricted distributions within this species complex suggest us the possibility of parallel speciation due to the isolation mechanism by the existence of Sundaland during the glacial periods (cf. Fleminger 1986).

Key to Indo-West Pacific species groups in *Labidocera detruncata* species complex

- 1 Endopods of female leg 5 absent; thumb and finger of male leg 5 slender..... ***L. gangetica* species group**
- Endopods of female leg 5 present; thumb and finger of male leg 5 not slender..... **2**
- 2 Female genital compound somite with complex swelling dorsally; apical segment of male left leg 5 with 4 terminal elements, second outermost the longest.....***L. detruncata* species group**
- Female genital compound somite lacking dorsal swelling; apical segment of male left leg 5 with 4 or fewer elements terminally **3**
- 3 Female caudal rami asymmetrical, widely separated, perpendicular to anal somite; thumb of first exopodal segment of male right leg 5 bifid ***L. pavo* species group**
- Female caudal rami symmetrical, neither widely separated nor perpendicular to anal somite; thumb of first exopodal segment of male right leg 5 not bifid **4**
- 4 Posterolateral prosomal corners of female asymmetrical with right longer than left; terminal exopodal segment of male left leg 5 with distal triangular process.....***L. cervi* species group**
- Posterolateral prosomal corners of female symmetrical; terminal exopodal segment of male leg left 5 lacking of distal triangular process..... ***L. madurae* species group**

Key to species of *Labidocera madurae* species group

Female

- 1 Genital compound somite with right postero-lateral and left antero-lateral processes *L. churaumi* sp. n.
- Genital compound somite without right postero-lateral and left antero-lateral processes 2
- 2 Genital compound somite more than twice as long as wide, furnished with anterior triangular small process on each side *L. tasmanica*
- Genital compound somite as long as wide, and produced mid-laterally *L. madurae*

Male

- 1 Inner margin of terminal segment of left leg 5 with protrusion at mid-length *L. churaumi* sp. n.
- Inner margin of terminal segment of left leg 5 without protrusion 2
- 2 Terminal segment of right leg 5 with expanded basal region with serrated margin *L. tasmanica*
- Terminal segment of right leg 5 without expanded basal region *L. madurae*

Acknowledgements

We would like to express our sincere thanks to Prof. G. A. Boxshall for his critical comments on an earlier draft. Thanks are due to the captain and crew of TRV Toyoshio-maru, Hiroshima University, for their cooperation at sea. This study was partially supported by a grant-in-aid from the Japan Society of the Promotion of Science, awarded to the second author (No. 25304031).

References

- Barr DJ, Ohtsuka S (1989) *Pseudocyclops lepidotus*, a new species of demersal copepod (Calanoida: Pseudocyclopidae) from the Northwestern Pacific. *Proceedings of the Biological Society of Washington* 102(2): 331–338.
- Barthélémy R, Ohtsuka S, Cuoc C (1998) Description and female genital structures of a new species of the demersal calanoid copepod *Ridgewayia* from southern Japan. *Journal of Natural History* 32: 1303–1318. doi: 10.1080/00222939800770651
- Fleminger A (1967) Taxonomy, distribution and polymorphism in the *Labidocera jollae* group with remarks on evolution within the group (Copepoda: Calanoida). *Proceedings of the US National Museum* 120: 1–61. doi: 10.5479/si.00963801.120-3567.1

- Fleminger A (1986) The Pleistocene equatorial barrier between the Indian and Pacific Oceans and a likely cause for Wallace's Line. UNESCO Technical Papers on Marine Science 49: 84–97.
- Greenwood JG, Othman BHR (1979) Description of *Labidocera farrani* sp. nov., a pontellid copepod known from eastern and northern Australian waters, (Crustacea, Copepoda). Journal of Plankton Research 1(3): 231–239. doi: 10.1093/plankt/1.3.231
- Huys R, Boxshall GA (1991) Copepod evolution. The Ray Society, London, 468 pp.
- Jeong HG, Suh HL, Jeong SB, Yoon TH, Soh HY (2009) *Labidocera* Species (Copepoda: Pontellidae) in Waters of the Tsushima Warm Current with Notes on Their Genital Structure and Zoogeography. Zoological Studies 48(4): 508–523.
- McKinnon AD, Kimmerer WJ (1984) Description of the male of *Labidocera caudata* Nicholls (Copepoda: Pontellidae) with remarks on the female. Proceedings of the Royal Society of Victoria 96(3): 169–172.
- Mulyadi (2002) The calanoid copepods family Pontellidae from Indonesian waters, with notes on its species-groups. Treubia 32: 1–167.
- Nicholls AG (1944) Littoral Copepoda from South Australia (II) Calanoida, Cyclopoida, Notodelphyoida, Monstrilloida and Caligoida. Records of the South Australian Museum 8: 1–62.
- Ohtsuka S, Boxshall GA (1994) *Platycopia orientalis*, new species (Copepoda: Platycopioida), from the North Pacific, with descriptions of copepodid stages. Journal of crustacean biology 14(1): 151–167. doi: 10.2307/1549062
- Ohtsuka S, Boxshall GA (2004) A new species of the deep-sea copepod genus *Scutogerulus* (Calanoida: Arietellidae) from the hyperbenthic waters of Okinawa, Japan. Systematics and Biodiversity 2: 49–55. doi: 10.1017/S1477200004001331
- Ohtsuka S, Boxshall GA, Fosshagen A (2003) A new species of *Neoscolecithrix* (Crustacea; Copepoda; Calanoida) from off Okinawa, southwestern Japan, with comments on the generic position in the superfamily Clausocalanoidea. Bulletin of National Science Museum Tokyo Series A 29(2): 53–63.
- Ohtsuka S, Boxshall GA, Shimomura M (2005) Three new species of deep-sea hyperbenthic Aetideid copepods (Crustacea) collected from Nansei Islands, southwestern Japan. Deep-sea fauna and pollutants in Nansei Islands by K. Hasegawa, G. Shinohara and M. Takeda National Science Museum Monograph 29: 224–247.
- Ohtsuka S, Fosshagen A, Go A (1991) The hyperbenthic calanoid copepod *Paramisophria* from Okinawa, south Japan. Zoological science 8: 793–804.
- Ohtsuka S, Fosshagen A, Soh HY (1996) Three new species of the demersal calanoid copepod *Placocalanus* (Ridgewayiidae) from Okinawa, southern Japan. Sarsia 81: 247–263.
- Ohtsuka S, Nishida S, Nakaguchi K (2002) Three new species of the genus *Macandrewella* (Copepoda: Calanoida: Scolecitrichidae) from the Pacific Ocean, with notes on distribution and feeding habits. Journal of Natural History 36(5): 531–564. doi: 10.1080/00222930010015861
- Ohtsuka S, Soh HY, Ueda H (1998) *Platycopia compacta* sp. n., the second species of Platycopioida (Crustacea: Copepoda) in the Indo-Pacific Region, with remarks on development, feeding, swimming, and zoogeography. Zoological Science 15: 415–424. doi: 10.2108/zsj.15.415

- Othman BHR (1986) A new species of *Labidocera* (Copepoda, Calanoida) from peninsular Malaysia. *Malayan Nature Journal* 39: 193–201.
- Razouls C, de Bovée F, Kouwenberg J, Desreumaux N (2012) Diversity and geographic distribution of marine planktonic copepods. <http://copepodes.obs-banyuls.fr/en> [accessed on March 21, 2014]
- Scott A (1909) The Copepoda of the Siboga Expedition. 1. Free-swimming, littoral and semi-parasitic Copepoda. *Siboga Expedition Monographs* 29: 1–323.
- Sewell RBS (1912) Notes on the Surface-living Copepoda of the Bay of Bengal I and II. *Records of the Indian Museum* 7: 313–382.
- Shen CJ, Lee FS (1963) The estuarine Copepoda of Chiekong and Zaikong Rivers, Kwangtung Province, China. *Acta Zoologica Sinica* 15(4): 571–596.
- Silas EG, Pillai PP (1973) The calanoid copepod family Pontellidae from the Indian Ocean. *Journal of the Marine Biology Association of India* 15: 771–858.
- Taw N (1974) A new species of *Labidocera* (Copepoda: Calanoida) from Tasmania and its postnaupliar developmental stages. *Australian Journal of Marine and Freshwater Research* 25(2): 261–272. doi: 10.1071/MF9740261