



HHS Public Access

Author manuscript

Newborn (Clarksville). Author manuscript; available in PMC 2022 December 19.

Published in final edited form as:

Newborn (Clarksville). 2022 ; 1(2): 215–218. doi:10.5005/jp-journals-11002-0034.

Risk Prediction for Stillbirth and Neonatal Mortality in Low-resource Settings

Vivek V Shukla¹, Waldemar A Carlo²

¹Division of Neonatology, University of Alabama at Birmingham, Birmingham, Alabama, United States of America

²Division of Neonatology, University of Alabama at Birmingham, Birmingham, Alabama, United States of America

Abstract

High stillbirth and neonatal mortality are major public health problems, particularly in low-resource settings in low- and middle-income countries (LMIC). Despite sustained efforts by national and international organizations over the last several decades, quality intrapartum and neonatal care is not universally available, especially in these low-resource settings. A few studies identify risk factors for adverse perinatal outcomes in low-resource settings in LMICs. This review highlights the evidence of risk prediction for stillbirth and neonatal death. Evidence using advanced machine-learning statistical models built on data from low-resource settings in LMICs suggests that the predictive accuracy for intrapartum stillbirth and neonatal mortality using prenatal and pre-delivery data is low. Models with delivery and post-delivery data have good predictive accuracy of the risk for neonatal mortality. Birth weight is the most important predictor of neonatal mortality. Further validation and testing of the models in other low-resource settings and subsequent development and testing of possible interventions could advance the field.

Keywords

Low- and middle-income countries; Mortality fetal; Mortality neonatal; Neonatal; Newborn infant; Preterm infants; Perinatal mortality; Resuscitation; Stillbirth

Introduction

Advances in perinatal and neonatal care, including the implementation of programs directed at improving perinatal care, have decreased stillbirth and neonatal mortality globally.¹ However, global stillbirth and neonatal mortality rates remain high with an

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and non-commercial reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Corresponding Author: Vivek V Shukla, Division of Neonatology, University of Alabama at Birmingham, Birmingham, Alabama, United States of America, Phone: +1 2059344680, vshukla@uabmc.edu.

Conflict of interest: None

increasing proportion of the under-5 child mortality.^{2,3} Stillbirth and neonatal mortality are concentrated in low-resource settings of LMICs.^{4,5} Approximately 98% of all stillbirths and neonatal deaths occur in these low-resource settings,⁶ despite the sustained focus of many local or global public health organizations over the last few decades.⁷

A large number of deliveries occur at home in the absence of trained birth attendants that can provide an appropriate level of care.^{7,8} Inadequate access to appropriate health care at delivery^{9–11} is one of the leading factors responsible for preventable stillbirth and neonatal mortality.¹² Lack of optimal resources combined with varying social issues, cultural practices, and health literacy lead to wide healthcare disparities. The barriers to healthcare at different stages¹³ and available resources vary between sites^{14–16} making universal access to quality perinatal health care at delivery a difficult aim to achieve.

Timely identification of at-risk pregnancies and neonates is critical for reducing stillbirth and neonatal mortality in low-resource settings. It has been estimated that 0.8 million stillbirths and 1.9 million neonatal deaths per year could be prevented by optimizing the coverage and quality of prenatal, intrapartum, and neonatal care.¹⁰ Unfortunately, studies focused on the quantification of stillbirth risk have been hampered by small event rates, a limited range of predictors that typically exclude obstetric history, lack of validation, and restriction to a single classifier (logistic regression). Consequently, predictive performance remains low, and risk quantification has not been adopted into antenatal practice.¹⁷ Improved risk stratification and triaging of pregnancies and neonates at higher risk of mortality could lead to optimal resource utilization, ensuring an appropriate level of care for at-risk pregnancies and neonates. Analyses of data of pregnancies and neonates with a higher risk of stillbirth or neonatal mortality can inform the field and lead to improved resource allocation to improve care. A recently published study suggests that models that include delivery and post-delivery variables had good predictive accuracy for the risk of neonatal mortality and that birth weight was the most important predictor for neonatal mortality.¹⁸ The current review highlights the available evidence for stillbirth and neonatal mortality risk prediction and identifies possible future research and implementation directions.

Risk Prediction for Stillbirth and Neonatal Mortality

Many studies report associations of pregnancy, delivery, and neonatal variables with stillbirth or neonatal mortality.^{19,20} Specific variable associations with mortality do not help in identifying individual-level risk assessment. Multivariable risk prediction models provide individual-level risk assessment, which could be helpful for clinical interpretation and intervention.^{21,22} Only a few investigators have published studies designed to develop models to predict the risk for stillbirth or neonatal mortality. The majority of the modeling studies for neonatal mortality risk prediction used data from extremely low birth weight neonates (ELBWs) who received intensive care in high-income countries (HIC).^{23–27} The prediction models are specific for the type of population and settings from which they are developed, so the models from HICs cannot be extrapolated to low-resource settings of LMICs.

There is a paucity of outcome prediction models for intrapartum stillbirth and neonatal mortality in either HICs or LMICs. A few outcome prediction modeling studies from

HICs^{23,24} used machine learning-based modeling techniques as they are thought to perform better than conventional models when applied to relatively large data sets because these models may improve the ability to delineate complex relationships and identify novel interactions between variables. However, machine-learning models had not been performed with data from low resource settings of LMICs.²⁸

Stillbirth and Neonatal Mortality Risk Prediction in Low-resource Settings in LMICs

Prediction models for intrapartum stillbirth and neonatal mortality in low-resource settings in LMICs have been reported recently.¹⁸ This study had the largest sample size stillbirth or neonatal mortality prediction study to date with 502,648 prospectively enrolled pregnancies over 9 years from six countries in South Asia (India and Pakistan), Africa (Democratic Republic of Congo, Zambia, and Kenya), and Latin America (Guatemala) using a high-quality population-based vital registry (Global Network Maternal Neonatal Health Registry, GN Registry).¹⁹ The investigators reported a comparison of predictive accuracies of conventional and advanced machine learning predictive modeling techniques to identify the best predictive model for intrapartum stillbirth and neonatal mortality in low-resource settings. The availability of a large sample size allowed building models using rigorous modeling techniques with independent training, test, and validation data sets to ensure their generalizability. Using the most important predictive variables and variable interactions from the best-identified model, which was a machine learning model, mortality risk prediction scores were developed. The study also identified important predictors for intrapartum stillbirth and neonatal mortality (Tables 1 and 2). This study determined that models based on prenatal or pre-delivery data have low predictive accuracy for intrapartum stillbirths, whereas neonatal mortality models that include delivery and post-delivery data had a good predictive accuracy of the risk for neonatal mortality. A similar finding of improved predictive accuracy with the addition of delivery and post-delivery variables has been reported in relatively small studies from both HICs²³ and LMICs.²² Birth weight was identified as the most important predictor of neonatal mortality (Table 2). Birth weight has also been reported in small studies to be associated with the risk for neonatal mortality in studies from HIC,^{23,26,29,30} although these studies did not identify that birth weight was the best predictor. An innovative easy to use mortality risk-prediction tool using birth weight as a continuous measure was developed for use by healthcare workers for point of care risk assessment (Fig. 1).

Future Directions for Retrospective or Prospective Assessment of the Risk Score and Tool

The study from the GN Registry¹⁸ was done using a prospectively collected high-quality population-level research database, so the study results are likely to be reproducible in communities in resource-limited regions similar to those of the study settings. However, retrospective and prospective studies could be done to evaluate the predictive accuracy of the risk score and risk assessment tool in communities outside of the GN Registry and adapt them as needed. The score was developed using readily available clinical data; data on stratification of individual risk factors as per illness severity, details of treatments received, laboratory variables, and clinical response were not included for ease of use and because of unavailability of such complex variables. It may be possible to further improve the score predictive accuracy by including complex variables, but such an attempt should be made

carefully as that may need model reconfiguration with changes in the variable coefficients.³¹ The incremental benefit of such an exercise may not be reasonable as the risk prediction score and tool had good predictive accuracy using readily available clinical variables. After ensuring the applicability of the risk score and risk tool, implementation research assessing mortality risk-based triage, referral, and management for reducing the number of stillbirths and neonatal deaths in low-resource settings of LMICs can be tested. Pre-delivery estimates of birth weight could also be tested as a strategy for pre-delivery triage and referral.

Conclusion

Timely identification of at-risk pregnancies and neonates is critical for the reduction of stillbirth and neonatal mortality in low-resource settings of LMICs. Risk stratification and triage of pregnancies and neonates at higher risk of mortality could help to reduce the burden of stillbirth and neonatal mortality in low-resource settings LMICs. The available risk assessment score and tool can be tested further using retrospectively or prospectively collected data. Implementation research should also be used to assess and improve the utility of these models. A population-based, multi-country, and high-quality database would assist and promote future outcome prediction research in low-resource settings.

References

1. Hug L, Alexander M, You D, et al. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: a systematic analysis. *Lancet Glob Health* 2019;7(6):e710–e720. DOI: 10.1016/S2214-109X(19)30163-9. [PubMed: 31097275]
2. Burstein R, Henry NJ, Collison ML, et al. Mapping 123 million neonatal, infant and child deaths between 2000 and 2017. *Nature* 2019;574(7778):353–358. DOI: 10.1038/s41586-019-1545-0. [PubMed: 31619795]
3. Levels and trends in child mortality. UN Inter-agency Group for Child Mortality Estimation. <https://data.unicef.org/wp-content/uploads/2018/10/Child-Mortality-Report-2018.pdf>.
4. Blencowe H, Cousens S, Jassir FB, et al. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health* 2016;4(2):e98–e108. DOI: 10.1016/S2214-109X(15)00275-2. [PubMed: 26795602]
5. Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet* 2015;385(9966): 430–440. DOI: 10.1016/S0140-6736(14)61698-6. [PubMed: 25280870]
6. Lawn JE, Cousens S, Zupan J, et al. 4 Million neonatal deaths: When? Where? Why? *Lancet* 2005;365(9462):891–900. DOI: 10.1016/S0140-6736(05)71048-5. [PubMed: 15752534]
7. Wall SN, Lee AC, Niermeyer S, et al. Neonatal resuscitation in low-resource settings: what, who, and how to overcome challenges to scale up? *Int J Gynaecol Obstet* 2009;107(Suppl 1):S47–S62, S63–64. DOI: 10.1016/j.ijgo.2009.07.013. [PubMed: 19815203]
8. Montagu D, Yamey G, Visconti A, et al. Where do poor women in developing countries give birth? A multi-country analysis of demographic and health survey data. *PLoS One* 2011;6(2):e17155. DOI: 10.1371/journal.pone.0017155. [PubMed: 21386886]
9. Darmstadt GL, Lee AC, Cousens S, et al. 60 Million non-facility births: who can deliver in community settings to reduce intrapartum-related deaths? *Int J Gynaecol Obstet* 2009;107(Suppl 1):S89–S112. DOI: 10.1016/j.ijgo.2009.07.010. [PubMed: 19815200]
10. Bhutta ZA, Das JK, Bahl R, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *Lancet* 2014;384(9940):347–370. DOI: 10.1016/S0140-6736(14)60792-3. [PubMed: 24853604]

11. Lawn JE, Blencowe H, Pattinson R, et al. Stillbirths: Where? When? Why? How to make the data count? *Lancet* 2011;377(9775):1448–1463. DOI: 10.1016/S0140-6736(10)62187-3. [PubMed: 21496911]
12. Kruk ME, Gage AD, Joseph NT, et al. Mortality due to lowquality health systems in the universal health coverage era: a systematic analysis of amenable deaths in 137 countries. *Lancet* 2018;392(10160):2203–2212. DOI: 10.1016/S0140-6736(18)31668-4. [PubMed: 30195398]
13. Thaddeus S, Maine D. Too far to walk: maternal mortality in context. *Soc Sci Med* 1994;38(8):1091–1110. DOI: 10.1016/0277-9536(94)90226-7. [PubMed: 8042057]
14. Geleto A, Chojenta C, Musa A, et al. Barriers to access and utilization of emergency obstetric care at health facilities in sub-Saharan Africa: a systematic review of literature. *Syst Rev* 2018;7(1):183. DOI: 10.1186/s13643-018-0842-2. [PubMed: 30424808]
15. Nimbalkar AS, Shukla VV, Phatak AG, et al. Newborn care practices and health seeking behavior in urban slums and villages of Anand, Gujarat. *Indian Pediatr* 2013;50(4):408–410. DOI: 10.1007/s13312-013-0116-y. [PubMed: 23024104]
16. Enweronu-Laryea C, Dickson KE, Moxon SG, et al. Basic newborn care and neonatal resuscitation: a multi-country analysis of health system bottlenecks and potential solutions. *BMC Pregnancy Childbirth* 2015;15(Suppl 2):S4. DOI: 10.1186/1471-2393-15-S2-S4. [PubMed: 26391000]
17. Malacova E, Tippaya S, Bailey HD, et al. Stillbirth risk prediction using machine learning for a large cohort of births from Western Australia, 1980–2015. *Sci Rep* 2020;10(1):5354. DOI: 10.1038/s41598-020-62210-9. [PubMed: 32210300]
18. Shukla VV, Eggleston B, Ambalavanan N, et al. Predictive modeling for perinatal mortality in resource-limited settings. *JAMA Netw Open* 2020;3(11):e2026750. DOI: 10.1001/jamanetworkopen.2020.26750. [PubMed: 33206194]
19. Goudar SS, Carlo WA, McClure EM, et al. The maternal and newborn health registry study of the Global Network for Women’s and Children’s Health Research. *Int J Gynaecol Obstet* 2012;118(3):190–193. DOI: 10.1016/j.ijgo.2012.04.022. [PubMed: 22738806]
20. Joyce R, Webb R, Peacock JL. Associations between perinatal interventions and hospital stillbirth rates and neonatal mortality. *Arch Dis Child Fetal Neonatal Ed* 2004;89(1):F51–F56. DOI: 10.1136/fn.89.1.f51. [PubMed: 14711857]
21. Ratnasiri AWG, Lakshminrusimha S, Dieckmann RA, et al. Maternal and infant predictors of infant mortality in California, 2007–2015. *PLoS One* 2020;15(8):e0236877. DOI: 10.1371/journal.pone.0236877. [PubMed: 32760136]
22. Houweling TAJ, van Klaveren D, Das S, et al. A prediction model for neonatal mortality in low- and middle-income countries: an analysis of data from population surveillance sites in India, Nepal and Bangladesh. *Int J Epidemiol* 2019;48(1):186–198. DOI: 10.1093/ije/dyy194. [PubMed: 30325465]
23. Ambalavanan N, Carlo WA, Bobashev G, et al. Prediction of death for extremely low birth weight neonates. *Pediatrics* 2005;116(6): 1367–1373. DOI: 10.1542/peds.2004-2099. [PubMed: 16322160]
24. Ambalavanan N, Baibergenova A, Carlo WA, et al. Early prediction of poor outcome in extremely low birth weight infants by classification tree analysis. *J Pediatr* 2006;148(4):438–444. DOI: 10.1016/j.jpeds.2005.11.042. [PubMed: 16647401]
25. Ambalavanan N, Carlo WA, Tyson JE, et al. Outcome trajectories in extremely preterm infants. *Pediatrics* 2012;130(1):e115–e125. DOI: 10.1542/peds.2011-3693. [PubMed: 22689874]
26. Tyson JE, Parikh NA, Langer J, et al. Intensive care for extreme prematurity—moving beyond gestational age. *N Engl J Med* 2008;358(16):1672–1681. DOI: 10.1056/NEJMoa073059. [PubMed: 18420500]
27. Rysavy MA, Horbar JD, Bell EF, et al. Assessment of an updated neonatal research network extremely preterm birth outcome model in the Vermont Oxford Network. *JAMA Pediatr* 2020;174(5):e196294. DOI: 10.1001/jamapediatrics.2019.6294. [PubMed: 32119065]
28. Paul AK, Schaefer M. Safeguards for the use of artificial intelligence and machine learning in global health. *Bull World Health Organ* 2020;98(4):282–284. DOI: 10.2471/BLT.19.237099. [PubMed: 32284653]

29. Medvedev MM, Brotherton H, Gai A, et al. Development and validation of a simplified score to predict neonatal mortality risk among neonates weighing 2000 g or less (NMR-2000): an analysis using data from the UK and The Gambia. *Lancet Child Adolesc Health* 2020;4(4):299–311. DOI: 10.1016/S2352-4642(20)30021-3. [PubMed: 32119841]
30. Salas AA, Carlo WA, Ambalavanan N, et al. Gestational age and birthweight for risk assessment of neurodevelopmental impairment or death in extremely preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2016;101(6):F494–F501. DOI: 10.1136/archdischild-2015-309670. [PubMed: 26895876]
31. Saravanou A, Noelke C, Huntington N. Predictive modeling of infant mortality. *Data Min Knowl Disc* 2021;35:1785–1807. 10.1007/s10618-020-00728-2.




Fig. 1: Probability of mortality as a function of birth weight. The risk for neonatal mortality increased with decreasing birth weight. Birth weight was the most important predictor of neonatal mortality in both the delivery/day 1 and post-delivery/day 2 scenarios and explained a large percentage of the variance of mortality. The probability of neonatal mortality with decreasing in birth weight occurred in both the delivery/day 1 and post-delivery/day 2 scenarios. Reproduced with permission¹⁸

Table 1:

Top predictors for intrapartum stillbirth and neonatal mortality

Rank	Predictor	AUC	AUC increase
Prenatal			
1	Cluster perinatal mortality	0.6	X
2	Gestational age at enrollment	0.61	0.008
3	Maternal age	0.61	0.003
4	Birth order	0.62	0.008
5	Parity	0.63	0.009
Pre-delivery			
1	Antepartum hemorrhage	0.56	X
2	Cluster perinatal mortality	0.64	0.086
3	Gestational age at enrollment	0.64	0.002
4	Hypertension/pre-eclampsia/eclampsia	0.66	0.018
5	Maternal age	0.66	0.002

Predictors are added consecutively using the gradient boosted ensemble model; then AUC is calculated. Intrapartum stillbirth was defined as non-macerated stillbirth occurring presumably during labor, and neonatal mortality was defined as mortality from birth to 28 days. Cluster perinatal mortality is the perinatal mortality rate within each distinct geographical area (cluster) of the sites as defined by the GN Registry. Hypertensive disease/severe pre-eclampsia/eclampsia is defined as blood pressure >140/90 mm Hg, proteinuria, and seizures¹⁸

Table 2:

Top predictors for neonatal mortality

<i>Delivery/day 1</i>			
1	Birth weight	0.78	X
2	Bag and mask resuscitation	0.81	0.039
3	Gestational age	0.81	0.003
4	Cluster perinatal mortality	0.82	0.001
5	Maternal age	0.82	0.004
<i>Post-delivery/day 2</i>			
1	Birth weight	0.76	X
2	Neonatal hospitalization	0.81	0.05
3	Neonatal antibiotics	0.84	0.032
4	Gestational age	0.84	0.003
5	Bag and mask resuscitation	0.85	0.01

Predictors are added consecutively using the gradient boosted ensemble model; then AUC is calculated. Neonatal mortality was defined as mortality from birth to 28 days. Cluster perinatal mortality is the perinatal mortality rate within each distinct geographical area (cluster) of the sites as defined by the GN Registry¹⁸