



# Synthetic and biological surfactant effects on freshwater biofilm community composition and metabolic activity

Stephanie P. Gill<sup>1</sup> · William R. Hunter<sup>2</sup> · Laura E. Coulson<sup>3,4</sup> · Ibrahim M. Banat<sup>5</sup> · Jakob Schelker<sup>3,6</sup>

Received: 27 July 2022 / Revised: 2 September 2022 / Accepted: 8 September 2022 / Published online: 19 September 2022  
© The Author(s) 2022

## Abstract

Surfactants are used to control microbial biofilms in industrial and medical settings. Their known toxicity on aquatic biota, and their longevity in the environment, has encouraged research on biodegradable alternatives such as rhamnolipids. While previous research has investigated the effects of biological surfactants on single species biofilms, there remains a lack of information regarding the effects of synthetic and biological surfactants in freshwater ecosystems. We conducted a mesocosm experiment to test how the surfactant sodium dodecyl sulfate (SDS) and the biological surfactant rhamnolipid altered community composition and metabolic activity of freshwater biofilms. Biofilms were cultured in the flumes using lake water from Lake Lunz in Austria, under high (300 ppm) and low (150 ppm) concentrations of either surfactant over a four-week period. Our results show that both surfactants significantly affected microbial diversity. Up to 36% of microbial operational taxonomic units were lost after surfactant exposure. Rhamnolipid exposure also increased the production of the extracellular enzymes, leucine aminopeptidase, and glucosidase, while SDS exposure reduced leucine aminopeptidase and glucosidase. This study demonstrates that exposure of freshwater biofilms to chemical and biological surfactants caused a reduction of microbial diversity and changes in biofilm metabolism, exemplified by shifts in extracellular enzyme activities.

## Key points

- *Microbial biofilm diversity decreased significantly after surfactant exposure.*
- *Exposure to either surfactant altered extracellular enzyme activity.*
- *Overall metabolic activity was not altered, suggesting functional redundancy.*

**Keywords** Biofilm · Surfactants · Freshwater · Metabolic · Sequencing · Metagenomics

## Introduction

In aquatic ecosystems, microorganisms live preferentially in microbial biofilms and act as major drivers of the biogeochemical cycling of carbon, nitrogen, and other nutrients (Paul et al. 1991; Decho 2000; Dang and Lovell 2016; Flemming et al. 2016; Battin et al. 2016). Microbial biofilms play an important role in the carbon cycle in aquatic environments such as streams and rivers (Lyon and Ziegler 2009; Qu et al. 2017). Microbial biofilms resident in fluvial sediments can also substantially reduce inorganic nitrogen (N) loads in N-rich aquatic ecosystems through denitrification (Zhang et al. 2016; Cui et al. 2017). In N-limited aquatic systems, they may help to facilitate the transfer of N to higher trophic levels (Dodds et al. 2000).

While microbial biofilms play an important ecological role, they are of concern in industrial and medical applications. For example, biofilms can facilitate the growth of

✉ Stephanie P. Gill  
gill-S5@ulster.ac.uk

<sup>1</sup> Department of Geography and Environmental Studies, Ulster University, Coleraine BT52 1SA, N. Ireland, UK

<sup>2</sup> Fisheries and Aquatic Ecosystems Branch, Agri-Food and Biosciences Institute, Belfast, N. Ireland, UK

<sup>3</sup> WasserCluster Lunz, Lunz am See, Austria

<sup>4</sup> Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Life Sciences, Vienna, Austria

<sup>5</sup> School of Biomedical Sciences, Ulster University, Coleraine, N. Ireland, UK

<sup>6</sup> Department of Functional and Evolutionary Ecology, University of Vienna, Vienna, Austria

antibiotic resistant bacteria (Lindsay and von Holy 2006; Srey et al. 2013; Bowler et al. 2020). Synthetic surfactants, such as sodium dodecyl sulfate (SDS), are widely used in personal care and cleaning products. They reduce biofilm growth in a wide range of industrial, medical, and environmental contexts (Waalder et al. 1993; Landa et al. 1999; Rigotti et al. 2017; Sloup et al. 2016; Ueda et al. 2019). Surfactants are amphiphilic chemicals that reduce surface tension within a liquid, sometimes through the formation of micelles. Micelles occur when molecules form an aggregate with hydrophobic tails on the inside (Schmitz 2018) and hydrophilic heads on the outside of an amphiphilic compound (Milovanovic et al. 2017).

Research on the effects of surfactants on microbial biofilms has primarily focused on single-species biofilms or those commonly found in medical and industrial environments. Direct applications of the anionic synthetic surfactant SDS have shown significant disruption in the growth and development of *Pseudomonas aeruginosa* biofilms (Díaz De Rienzo et al. 2016; Nguyen et al. 2020). Effects on other single species or medical biofilms have also been investigated including methicillin-resistant *Staphylococcus aureus* (Ueda et al. 2019), *Pseudomonas fluorescens* (Simões et al. 2008), and *Salmonella enterica* serovar *typhimurium* (Mireles II et al. 2001). Inhibited biofilm formation was seen to occur within all investigations.

The use of synthetic surfactants has been questioned due to the potential environmental impact, specifically the longevity and toxicity to aquatic biota. For example, previous research has shown that synthetic surfactants, such as Triton X-100 (Octylphenol Ethoxylate), are not easily biodegradable under either aerobic or anaerobic conditions (Mohan et al. 2006). Other synthetic surfactants such as SDS have also been noted as toxic to a variety of microorganisms such as *Vibrio fischeri*, fish and shellfish such as *Clarias batrachus* and *Mytilus galloprovincialis*, yeast, and also small invertebrates, such as *Daphnia magna*, and *Tigriopus fulvius* (Mariani et al. 2006; Lima et al. 2011; Franzetti et al. 2012; Messina et al. 2014; Cao et al. 2020; Kumar et al. 2020). Research has therefore shifted towards identifying a natural alternative that may be less harmful to the aquatic environment.

Biofilm-dwelling microorganisms naturally produce a number of surfactants to control the surrounding extracellular polymeric substance (EPS) matrix. The most common substances of this group are rhamnolipids and sophorolipids which are glycolipids formed by the combination of either a long-chain aliphatic acid or hydroxyaliphatic acid with a carbohydrate (Desai and Banat 1997; Marchant and Banat 2012). When exposing single-cell biofilms of *P. aeruginosa* to rhamnolipids, growth and development were disrupted, and cell viability was reduced (De Rienzo et al. 2015). Disruptions by rhamnolipids have also been shown for single

species biofilms of *Staphylococcus aureus* (Quinn et al. 2013; Aleksic et al. 2017; Ceresa et al. 2021a; Tambone et al. 2021), *Bacillus pumilus* (Dusane et al. 2010), and *Rhodococcus erythropolis* (Schreiberova et al. 2012). Yet, rhamnolipids are biodegradable in a variety of environments, including under aerobic and anaerobic conditions (Mohan et al. 2006; Naughton et al. 2019; Ramos da Silva et al. 2019). Biosurfactants in general have been reported to act as antibiofilm agents (Banat et al. 2014). Given that previous research primarily addressed the effects of biological surfactants on single species biofilms of bacteria relevant to medical settings, there remains a knowledge gap on how the exposure of multi-species freshwater biofilms to surfactants affects biofilm microbial community composition and function.

There is a growing body of evidence that anthropogenic trace contaminants, and contaminants of emerging concern (CECs), such as pharmaceuticals, herbicides, and antimicrobial agents can have major impacts upon the composition and functioning of aquatic microbial communities (Rosi-Marshall et al. 2013; Argudo et al. 2020; Harjung et al. 2020; McClean and Hunter 2020; Ke et al. 2020; Kusi et al. 2020). Exposure to contaminants such as the antibacterial agent Triclosan (Hay et al. 2001; Drury et al. 2013) and the herbicide Diuron (Sumpono et al. 2003; Seghers et al. 2003) has been observed to place microbial communities under a selective pressure that favors microorganisms that can use these contaminants as an additional carbon source. Given that levels of surfactants and their metabolites are consistently among the top contaminants identified within wastewater influent and groundwater (Lara-Martín et al. 2014; Schaidler et al. 2014), it is reasonable to infer that they could represent a major control upon the microbial communities within human affected inland waters.

This study investigated the effects of the synthetic surfactant SDS and the biological surfactant rhamnolipid on microbial biofilm community composition, microbial diversity, and metabolic activity. Using recirculating flume mesocosms, we aimed to answer the following overarching scientific question: What are the effects of a synthetic and biological surfactant on a natural aquatic microbial biofilm, and how do these effects compare? We hypothesized that biofilm exposure to a surfactant will decrease bacterial biodiversity, as the surfactants impact microbial community composition and cause the loss of sensitive bacterial taxa as previous research has shown that chemical exposure can place microbial communities under selective pressure. Secondly, we hypothesize that the resultant changes in microbial community composition will alter biofilm metabolic activity. We planned to test these hypotheses by quantifying changes in community composition through 16 s rRNA analysis and changes in biofilm structure and metabolic activity through different analytical methods, such as changes in gross

primary productivity (GPP), community respiration (CR), ash free dry mass (AFDM), Carbon: Nitrogen ratios (C:N), and extracellular enzyme activity (EEA).

## Materials and methods

### Experimental setup

We set up 15 recirculating experimental flumes near Lunz Am See, Austria, following the method described by Roche et al. (2017). Each flume was 2.5 m long and had an approximate slope of 0.8%. In each flume, ~5 L of water recirculated, which was renewed with unfiltered lake water twice per week. We added rhamnolipid (RL) (R90-100G, Sigma-Aldrich, Darmstadt, Germany) and SDS (sodium dodecyl sulfate, Sigma-Aldrich, Darmstadt, Germany) as treatments of 150 and 300 ppm to three flumes each (hereafter referred to as either RL 150, RL 300, SDS 150, or SDS 300), while three flumes acted as controls. After the first week,  $5 \mu\text{g L}^{-1}$  of phosphorus in the form of  $\text{Na}_2\text{HPO}_4$  was also added to help growth due to low phosphorus in the utilized lake water. Surfactant concentrations were chosen to represent the effective dose of rhamnolipid, as well as double that concentration. Due to malfunction of one flume,  $n$  for the rhamnolipid treatments was reduced to 2. Treatments were randomly placed among flumes. Artificial light was spread equally over all flumes and light density averaged  $9.5 \mu\text{mol m}^{-2} \text{s}^{-1}$  and was the maximum even light density achievable with the provided light sources. We used a light cycle of 12:12 and 18:6 (Cheah and Chan 2021) hours of light versus dark early and later in the experiment to mimic natural light cycles at first and then to encourage biofilm growth later in the experiment. Each flume was equipped with microscope slides (6) and clay tiles (8) to quantify biofilm growth.

We measured pH, dissolved oxygen (DO) (mg/L), water temperature ( $^{\circ}\text{C}$ ), and flow (by bucket measurement) every other day to ensure stable conditions. Twice per week we analyzed DOC, ammonium ( $\text{NH}_4$ ),  $\text{NO}_x$ , and  $\text{PO}_4$  of flume water to ensure they remained stable. In addition, water temperature was monitored by automatic loggers in drainage tanks. After 32 days, the experiment was stopped due to adequate biofilm growth observed, and biofilm samples were retrieved from the flumes for further analysis.

### Sample collection

Two microscope slides were removed from each flume and placed in 120-mL Schott bottles with oxygen sensors. Bottles were filled with lake water run through glass fiber filters to remove any microorganisms that may alter the results of the analyses and sealed to remove all air bubbles. Three control bottles with no slides were also set up to test for

potential activity within the filtered lake water with control activity subtracted from treatments during analyses. All bottles were placed in an Imago 500 Controller Environmental Chamber (Snijders Scientific B.V., Tilburg, Netherlands) using a 12:12 light cycle for 72 h. Oxygen measurements were taken at the start and end of each light and dark cycle to obtain data on GPP and CR.

We determined AFDM by weighing 2 clay tiles per flume after drying (9 h at  $70^{\circ}\text{C}$ ), and combustion (4 h at  $450^{\circ}\text{C}$ ). AFDM was obtained by subtracting the remaining ash weight from the dry weight. Two tiles from SDS 150 flumes were damaged during drying, and AFDM could not be determined. Biofilms covering two tiles per flume were then scraped into sterilized and pre-weighed 2-mL Eppendorf tubes using a sterile razor blade, lyophilised ( $-0.001$  mbar;  $-76^{\circ}\text{C}$ ) (Martin Christ, Gefriertrocknungsanlagen GmbH, Osterode am Harz, Germany) and then analyzed on a Thermo Scientific FLASH 2000 Elemental Analyser (Thermo Fisher Scientific GmbH, Bremen, Germany) to quantify carbon (% C) and nitrogen (% N) content. Quantities of C and N by mass were then examined in ratios to identify changes in ecological stoichiometry (the relationship between elements, energy, and organisms and their interactions in an ecological system (Sterner and Elser 2002; Van de Waal et al. 2018)).

EEA of  $\beta$ -D-1,4-glucosidase (glucosidase), phosphatase, and leucine-aminopeptidase were measured in the biofilms following the method described in Romani and Sabater (2001). Four mL of sterile water was added to approximately 0.5 g of biofilm per flume that was scraped from the tiles. The mixture was homogenized by shaking. EEA was measured via artificial substrates: Glucosidase and phosphatase using 4-methylumbelliferyl (MUF)- substrates (MUF- $\beta$ -D-glucopyranoside and MUF-phosphate, respectively (Sigma-Aldrich, Darmstadt, Germany)) and leucine-aminopeptidase using a 7-amino-4-methylcoumarin (AMC)- substrate (L-leucine-AMC, Sigma-Aldrich, Darmstadt, Germany). Artificial substrates were added to the homogenate at a final concentration of MUF-glucosidase;  $135 \text{ mg L}^{-1}$ , MUF-phosphatase;  $102.46 \text{ mg L}^{-1}$ , and AMC-leucine-aminopeptidase;  $129.94 \text{ mg L}^{-1}$  and incubated for 1 h in the dark. Fluorescence was measured at the beginning and end of the incubation with a plate reader (Varioskan Flash, Thermo Fischer Scientific, Vaanta, Finland) at 365/455 nm excitation/emission wavelengths for the MUF-substrates and 380/440 excitation/emission wavelengths for the AMC-substrate.

Two microscope slides per flume were scraped using sterile procedures into a sterile 1.5-mL microcentrifuge tube for 16 s rRNA analysis. DNA was immediately extracted from collected samples using the Dneasy Power Soil Kit (Qiagen GmbH, Hilden, Germany) following the manufacturer's methodology. We included internal replicates per flume ( $n=2$ ), and experimental controls were used to

control for any potential unaccounted sample contamination. Extracted products were sent to LGC Biosearch Technologies (LGC Genomics GmbH, Berlin, Germany). DNA was quantified using gel electrophoresis before amplifying the V3-V4 region of the 16 s rRNA gene with polymerase chain reactions using 341F and 785R primers. Resultant DNA were sequenced using the Illumina Mi-Seq platform (300 bp paired-end reads) for amplicon creation.

We used Mothur version 1.44.3 (Schloss et al. 2009), and the Mothur MiSeq SOP (Kozich et al. 2013) with some modification, to analyze returned reads. A maximum homopolymer of 8 was chosen due to the limitation of the reference database (SILVA version 132) with a 97% similarity and above required for identification of operational taxonomic units (OTUs). UCHIME (Edgar et al. 2011) was also used to identify and remove chimeric fragments. Sequences that could not be directly identified through SILVA were manually run through NCBI's BLAST (Madden 2002). We were able to identify all but one OTU with 97% similarity; OTU005 is thus reported as 'unnamed' in this study.

## Data analysis

All data, except the 16 s rRNA and C:N results, met assumptions of normality and homogeneity of variance. We considered  $p < 0.05$  as significant and analyzed data using R version 4.0.3 (R Core Team 2020) and the R packages car (Fox and Weisberg 2019), vegan (Oksanen et al. 2020), ggplot2 (Wickham 2016), and entropart (Marcon and Herault 2015). Nonmetric multidimensional scaling analyses (NMDS) were run using the PAST 3 software (Hammer et al. 2001). Due to replication error early in the experiment, only the controls and the 300 ppm concentration variables were used in statistical difference testing. All GPP and CR data were normalized to mg O<sub>2</sub> in 120 mL per mg of biofilm in 12 h before being analyzed. C:N data was normalized to mass (mg C/N per mg biofilm). A one way analysis of variance (ANOVA) analysis was used to test for significant differences among treatments and the controls for the GPP, CR, and AFDM. A Kruskal–Wallis test was used to test for significant differences among the C:N ratios. Coefficients of variation were also calculated for individual flumes for each analysis.

Changes in EEA were analyzed through both a distance-based redundancy analysis (db-RDA) and three one-way ANOVAs, one per extracellular enzyme. All the data was used within the db-RDA and only the maximum concentrations of each chemical and control were used for analysis in the ANOVA. Where appropriate, we applied a Tukey HSD post hoc test to the ANOVA output.

Bacterial OTUs were analyzed for alpha and beta diversity as well as differences in the overall bacterial community composition. We used the Chao1 (Chao 1984) and the Shannon–Wiener Diversity Indices to quantify bacterial

Alpha diversity (Jost 2007). Chao 1 and Shannon–Wiener results were then analyzed using an ANOVA followed by a Tukey HSD post-hoc test to identify differences in diversity among treatments. Following this, all singletons and doubletons were removed, and the data was rarefied based on the smallest number of sequences identified within a sample. Beta diversity was then analyzed using a non-metric multidimensional scaling analysis (NMDS) on all data present to identify differences in the biofilm bacterial community among treatments and controls. An analysis of similarities (ANOSIM) was performed with 999 permutations to identify if these differences were significant. Families and genera were compared graphically, focusing on the top 80% of OTUs identified within the biofilms.

## Results


### Primary productivity and community respiration

Lake control bottles had an average primary productivity of  $0.0304 \pm 0.0256$  mg O<sub>2</sub> in 120 mL and average community respiration of  $-0.016 \pm 0.0177$  mg O<sub>2</sub> in 120 mL. We found no significant differences in primary productivity (ANOVA,  $F = 0.022$ ,  $p = 0.978$ ) and community respiration (ANOVA,  $F = 0.148$ ,  $p = 0.865$ ) of the treatments versus the controls (Fig. 1A, B). All treatments showed substantial variation among replicates (Fig. 1), with the most pronounced variability being present in respiration. Coefficients of variation were highest for respiration in the controls (1.56, see Table 1) and lowest (0.021) in the rhamnolipid 150 treatment. Rhamnolipid treated systems overall had the lowest variation for CR, while the controls had the highest.

### Ash free dry mass and C:N ratios

We found no significant difference among treatments and controls (ANOVA  $F = 0.045$ ,  $p = 0.957$ ) for AFDM. The coefficients of variation, however, showed differences in variation among the treatments (Table 1), with higher surfactant concentrations decreasing variation (e.g., RL 150 has greater variation than RL 300) (Fig. 1C). Control variation was comparable to both low dose surfactant variations. However, C (ANOVA  $F = 18.36$ ,  $p = 0.0028$ ) and N (ANOVA  $F = 5.471$ ,  $p = 0.044$ ) content was significantly different by mass among treatments (Fig. 2A and Fig. 2B). Differences lie between the RL 300 and the control for both C (Tukey HSD  $p = 0.0063$ ) and N (Tukey HSD  $p = 0.041$ ) content, and between SDS 300 and the control for C (Tukey HSD  $p = 0.0036$ ) content.

There were also no significant differences among treatments and controls for C:N ratios by mass (Kruskal–Wallis  $p = 0.193$ ) present in the biofilms (Fig. 2C). There were



**Fig. 1** GPP (A), CR (B), and AFDM (C) of microbial biofilms exposed to SDS (sodium dodecyl sulfate) and RL (rhamnolipid) for 32 days at 150 and 300 ppm

**Table 1** Coefficients of variation for AFDM, GPP, and CR data. Data collected from samples of biofilm grown in either the surfactant SDS (sodium dodecyl sulfate) or RL (rhamnolipid) for 32 days

Treatment	Ash free dry mass	Gross primary productivity	CR	C:N
SDS 150	0.70	0.67	0.92	0.373
SDS 300	0.11	0.626	1.09	0.643
RL 150	0.60	0.34	0.021	0.0875
RL 300	0.32	0.52	0.59	0.0698
Control	0.59	0.61	1.56	0.569


differences in variation among treatments and controls (Table 1). Rhamnolipid treated flumes had lower variation than either SDS treated flumes or the controls; however, within each treatment, a higher surfactant concentration increased variation.

**Bacterial community composition**

Shannon–Wiener Diversity was significantly different among treatments (ANOVA  $F = 71.702$ ,  $p < 0.0001$ , Tukey

HSD rhamnolipid and control;  $p = 0.043$ , SDS and control;  $p < 0.0001$ , SDS and rhamnolipid;  $p < 0.001$ ) with a higher diversity in the control flumes, second highest in the rhamnolipid flumes, and lowest in the SDS flumes. Significance was driven by surfactant presence and type, but not concentration. The control had the greatest bacterial diversity, while both rhamnolipid treatments had the next highest, and the lowest belonged to both SDS treatments (Fig. 3A). Chao1 had no significant differences (ANOVA  $F = 3.7748$ ,  $p = 0.087$ ).

By testing beta diversity and community differences on the rarefied data, we identified distinct groupings between all treatments and controls, as evident by the NMDS (Fig. 3B). The ANOSIM results confirmed the differences among both surfactants and the controls ( $R = 0.9112$ ,  $p = 0.001$ ). Community differences were further identified by looking at only the taxa in the top 80% of all taxa present across all treatments, which was 18 taxa. We identified several genera that were heavily dominant in rhamnolipid-treated flumes. These were *Brevundimonas spp.*, OTU005, *Pseudomonas*, and *Arthrobacter*. In SDS flumes, we observed *Brevundimonas terrae*, *Flavobacterium spp.*, *A*, and in the control flumes *Exiguobacterium*,



**Fig. 2** Carbon (**A**), nitrogen (**B**), and C:N ratios (**C**) of microbial biofilms exposed to SDS (sodium dodecyl sulfate) and RL (rhamnolipid) for 32 days at 150 and 300 ppm

*Arenimonas*, and *Stenotrophomona*. We also identified 32 genera only present in the controls, 5 there were exclusive to RL 150, 6 in RL 300, 19 in SDS 150, and 7 in SDS 300 (Fig. 3C).

Total OTUs within each treatment and OTUs within the top 80% of all taxa per treatment were also calculated (Table 2). The control flumes had the greatest number of OTUs present, and the greatest number of OTUs within their top 80% of taxa. The SDS and rhamnolipid flumes had similar numbers of total OTUs present, but the SDS flumes had 80% of their total OTUs within half the number of taxa as the rhamnolipids. This indicated a large difference in evenness between treatments, while the total number of OTUs was fairly similar.


### Extracellular enzyme activity

According to the db-RDA, EEA differed among all treatments and controls, with the exception of the rhamnolipid treatments. The differences in db-RDA-distances were mainly caused by the differences in EEA of leucine aminopeptidase and phosphatase, although the results were not significant (ANOSIM  $F=2.9907$ ,  $p=0.058$ ). While phosphatase did not have any

significant differences among treatments (ANOVA  $F=2.934$   $p=0.129$ ), with the exception of SDS 150 addition of any surfactant decreased the production of phosphatase when compared to the control (Fig. 4A). Rhamnolipid additions yielded a significantly higher production of glucosidase, while SDS additions reduced production (ANOVA  $F=5.741$   $p=0.0404$ , Tukey HSD  $p=0.034$ ) (Fig. 4B). The effects of RL 300 and SDS 300 on leucine aminopeptidase were also significantly different (ANOVA  $F=5.914$   $p=0.038$ , Tukey HSD  $p=0.032$ ). The addition of rhamnolipid yielded a higher amount of the extracellular enzymes when compared to the control and the addition of SDS reduced production (Fig. 4C).

### Discussion

Our experiment showed that exposure to biological and synthetic surfactants alters the metabolic activity and the community composition of a mixed-species freshwater biofilm. The effects of SDS were more pronounced than the effects of rhamnolipid. Our experiment is the first study to our knowledge that examines the effects of these two surfactants on a multi-species near-natural freshwater biofilm, which contrasts with



**Fig. 3** Biofilm bacterial diversity from 16 s rRNA sequencing of biofilm exposed to SDS (sodium dodecyl sulfate) or RL (rhamnolipid) after 32 days at 150 and 300 ppm. **A** shows Shannon Diversity, **B**

shows an NMS2D depicting the community differences in treatments when examining all OTUs present within each sample after rarefaction and **C** depicts relative abundance of OTUs

the commonly performed single-species studies. This study was also novel as it directly compared the effects of a synthetic and a biological surfactant and explicitly studied for the first time the C:N ratios and C and N contents of biofilms that are exposed to surfactants. These results differ from previous research examining the effectiveness of biosurfactants on biofilm removal in medical (Ceresa et al. 2019, 2021b) and industrial (Campos et al. 2013) environments.

### Primary productivity, respiration, ash free dry mass, and C:N ratios

Although our results showed biofilms had no differences in GPP or CR in any treatments or controls, chemical exposure decreased biofilm respiratory activity in previous experiments, including surfactant exposure (Simões et al. 2008) and pharmaceutical exposure (Rosi-Marshall et al. 2013). Biofilm volume has also been reduced after a single surfactant exposure event, with decreases of 26.5–98.1% occurring between 22 and 48 h after exposure (Mireles II et al. 2001; Quinn et al. 2013). Typically, biosurfactants, which are amphiphilic polymers, can affect biofilm surface attachment through altering surface wettability (Rodrigues et al. 2006). Biosurfactants such as rhamnolipids

are able to make cell surfaces more hydrophobic through loss of fatty acids such as lipopolysaccharides (Al-Tahhan et al. 2000). Amphiphilic anionic surfactants work similarly through altering surface characteristics and changing solubility of certain compounds (Cserháti et al. 2002). Surface changes then can alter a biofilm's ability to adhere to a surface, reducing growth and development. However, previous work on surfactant effects has also shown that biofilms demonstrate resistance and can recover full respiratory activity 12 h after the initial exposure (Simões et al. 2008) and biomass 5 h after initial exposure (Nguyen et al. 2020). As our experiment lasted 32 days, we believe that GPP, CR, and AFDM could have stabilized and recovered full respiratory activity as seen in previous studies (Simões et al. 2008; Nguyen et al. 2020).

Altered C:N ratios are indicative of changes in microbial productivity (where reduced C:N ratios may yield greater productivity) (Li et al. 2016) and biofilm formation (where lower C:N ratios increase biofilm volume) (Thompson et al. 2006). As we did not have any alterations in C:N ratios, and no changes in productivity (GPP and CR) or growth (AFDM) among any treatments or controls, our results support these studies. The addition of surfactants provided an added source of C; however, as there were consistent C:N ratios across all flumes, it

**Table 2** Biofilm OTUs per surfactant treatment and their dominant genera. Biofilms were grown in either SDS (sodium dodecyl sulfate) or rhamnolipid for 32 days

Surfactant	Conc. (ppm)	Total OTUs present	OTUs in top 80% per treatment	OTUs lost compared to control	OTUs gained compared to control	Dominant genera in top 80% of all taxa present
Rhamnolipid	150	145	10	58	25	<i>Brevundimonas</i> , <i>Otu005</i> , <i>Pseudomonas</i> , <i>Arthrobacter</i> , <i>Flavobacterium B</i> , <i>Prostheco-bacter</i> , <i>Nocardioides</i>
Rhamnolipid	300	159	14	44	36	<i>Brevundimonas</i> , <i>Otu005</i> , <i>Pseudomonas</i> , <i>Arthrobacter</i> , <i>Flavobacterium B</i> , <i>Prostheco-bacter</i>
SDS	150	176	7	27	37	<i>Brevundimonas terrae</i> , <i>Acinetobacter</i> , <i>Flavobacterium A</i> , <i>Paucibacter</i> , <i>Flavobacterium B</i> , <i>Prostheco-bacter</i>
SDS	300	130	6	73	23	<i>Brevundimonas terrae</i> , <i>Acinetobacter</i> , <i>Flavobacterium A</i> , <i>Prostheco-bacter</i> , <i>Flavobacterium B</i> , <i>Prostheco-bacter</i> , <i>Sphingomonas</i>
Control	0	203	15	–	–	<i>Exiguobacterium</i> , <i>Acinetobacter</i> , <i>Arenimonas</i> , <i>Stenotrophomonas</i> , <i>Nocardioides</i>

can be assumed that our system was C limited. It is also likely that a thickened EPS matrix, which may be used as a defense mechanism in biofilms in response to surfactant exposure, could be another source of extra carbon (Jefferson 2004; Zakaria and Dhar 2020). In contrast, in systems where C is not limiting, an increase in C inputs through surfactants may increase C:N ratios, and therefore negatively affect biofilm productivity and growth.


## Diversity and bacterial community composition

Our controls had taxa present that were minimally identified within the surfactant flumes. These taxa included species (spp.) of *Exiguobacterium*, *Acinetobacter*, *Arenimonas*, and *Stenotrophomonas*. *Exiguobacterium* is currently researched as an emerging human pathogen and while susceptible to antibiotics appears to easily develop antibiotic resistance (Chen et al. 2017). *Acinetobacter* contains species such as *A. baumannii* that are very well known as human pathogens and that demonstrate high antibiotic resistance (Rice 2008; Tewari et al. 2018). However, it appears that some species may not have resistance to biological surfactants as its presence in rhamnolipid treated flumes was extremely low, although certain species of *Acinetobacter* had an increase in abundance within the SDS treated flumes. Similar results appear with *Stenotrophomonas* as it has human pathogenic potential, and species that have developed multi drug resistance, but appears to be sensitive to both biological and synthetic surfactants (Chang et al. 2015). In our study, the addition of either surfactant created an environment unsuitable for these bacteria, demonstrating how surfactant exposure can alter biofilm community composition and potentially help to control certain pathogenic species that are demonstrating resistance to antibiotics. The control flumes had the highest amount of OTUs present ( $n = 203$ ) and the highest diversity, which indicates that both the diversity and abundance of bacteria present in the biofilms decreases after surfactant exposure.

Rhamnolipids are studied for their ability to reduce pre-existing biofilms and prevent future biofilm growth with some success (Quinn et al. 2013; Dusane et al. 2010; De Rienzo et al. 2015; Satpute et al. 2018). Current research has shown that rhamnolipid exposure on soil bacteria can shift community composition, although many studies do not examine differences to the genus level (Lu et al. 2019; Wei et al. 2020; Akbari et al. 2021). Research has also shown that aquatic microbial communities will shift in response to an outside contaminant with an increased abundance of bacteria that are not sensitive to that contaminant or can use it as a C source (Hay et al. 2001; Sumpono et al. 2003; Seghers et al. 2003; Drury et al. 2013). Our study showed that, much like soil communities and aquatic communities exposed to other contaminants, aquatic biofilms experience significant community changes after exposure to rhamnolipids.

Further examination into the bacterial genera indicates large increases in a *Brevundimonas* spp., *Pseudomonas* spp.,





**Fig. 4** EEA of phosphatase (A), glucosidase (B), and leucine aminopeptidase (C) of biofilms after exposure to either SDS (sodium dodecyl sulfate) or RL (rhamnolipid) for 32 days at 150 and 300 ppm (\*indicates significant differences)

*Arthrobacter* spp., and OTU005 when compared to all other treatments and to the controls. One OTU of *Arthrobacter* spp. was only present within the rhamnolipid flumes, with the exception of one control flume that had low levels of that particular OTU. *Brevundimonas* spp., who are opportunistic pathogens (Ryan and Pembroke 2018), have experienced accelerated growth in the presence of rhamnolipids before when testing the effects of rhamnolipids on increasing bioremediation in soils (Lu et al. 2019). It is likely that these genera were able to use rhamnolipid as a C source due to the large C quantities within rhamnolipids (Mnif et al. 2018). Their adaptability may explain why they were able to increase in abundance over time when compared to the other genera present.

SDS caused the largest reduction in microbial diversity when compared with the other treatments and controls. SDS has been researched and used for its ability to degrade biofilms in applications where biofilms are undesirable (Izano et al. 2007; Ueda et al. 2019; Nguyen et al. 2020). This is particularly due to their ability to bind to hydrophobic protein components, introducing a negative charge to the structure which unfolds the bound protein (Gudiksen et al. 2006; Fong and Yildiz 2015). While rhamnolipids are also able to denature proteins, they are considered weak binders, and typically have a reduced denaturing ability when compared to many other surfactants (Otzen 2017). This may explain why biofilms exposed to SDS had lower diversity than those in the rhamnolipid treatments or the controls. SDS also had the lowest amount of OTUs within the top 80% of taxa present in the treatments. We believe that the low diversity, and community domination by few taxa, demonstrates that SDS exposure creates an inhospitable environment to many potential specialist taxa. The few remaining and insensitive taxa can then thrive as more resources are available. Following our findings, these taxa would include *Brevundimonas terrae*, and an *Acinetobacter* and *Flavobacterium* spp. While the rhamnolipid treated flumes also had less diversity than the controls, the rhamnolipid

treatments still appeared to favor more taxa and a higher diversity than the SDS treated flumes.

### Extracellular enzyme activity

Our experiment showed that substantial changes in biofilm EEA can occur after surfactant exposure. Although phosphatase activity, or the separation of phosphate groups from a larger compound such as an acid or protein (Margalef et al. 2017), generally decreased after surfactant exposure, we did have similar levels of activity between the control and SDS 150 flumes. Examination of the bacterial communities within SDS 150 flumes and control flumes shows several parallel genera present that are not as common in the other systems, including *Nocardioidea*. Treatments were laid out randomly across our experimental flumes within the experimental setup, and water was allocated to all systems from the same source, which minimized potential experimental alterations by relative location. We thus assume that all changes in community composition and EEA refer to the actual biofilm response to treatments. The presence of a low concentration SDS allowed for the development of bacterial genera like *Nocardioidea* which typically have both acid and alkaline phosphatase (Xie et al. 2017). However, the addition of high concentration SDS, or rhamnolipid at either a high or low concentration, reduced phosphatase activity.

Beta-glucosidase (rate-limiting enzyme that converts cellulose to glucose (Singh et al. 2016)) and leucine aminopeptidase (enzyme that aids in the hydrolysis of leucine and other N-terminal residues at the end of peptides and proteins (Matsui et al. 2006)) activities were also affected by surfactant exposure. Previous research has shown that rhamnolipids can act as an enhancer for the enzyme beta-glucosidase (Zhang et al. 2009; Yao et al. 2020). Rhamnolipids are able to protect proteins and limit the degradation of cellulase (Zhang et al. 2009; Wang et al.

2011). Yin et al. (2019) showed that the presence of rhamnolipid can increase genes related to leucine and amino acid synthesis. The addition of rhamnolipid to our flumes likely protected cellulase and leucine in the system which led to enhanced beta-glucosidase and leucine aminopeptidase activity. Beta-glucosidase is also known for its ability to degrade *Pseudomonas aeruginosa*, a rhamnolipid producing bacteria (Banar et al. 2019; Pirlar et al. 2020). The flumes treated with rhamnolipid had a higher abundance of *P. aeruginosa* when compared with the other flumes.

We assume therefore, that the higher beta-glucosidase levels within the rhamnolipid flumes were likely caused by a combination of the presence of rhamnolipid to act as an enhancer, and more abundant *P. aeruginosa*. However, we observed somewhat contrasting results in the SDS treated flumes, where a decrease in enzyme activity occurred. SDS is an anionic surfactant that can be used in denaturation of polypeptides (Bhuyan 2010; Schlager et al. 2012). SDS is commonly used in an analysis called SDS–polyacrylamide gel electrophoresis (SDS–PAGE) where an SDS infused gel is used to separate proteins and enzymes based on their molecular weight. This analytical method has been used with experiments examining beta-glucosidase previously (Laemmli 1970; Bai et al. 2013), which demonstrates that SDS is able to denature beta-glucosidase. It is possible that the denaturing ability of SDS has degraded beta-glucosidase and leucine aminopeptidase, leading to reduced enzyme activity, and likely also reduced biofilm metabolic activity.

Our experiment showed that both the biologic surfactant rhamnolipid and the synthetic surfactant, SDS, can significantly affect a mixed freshwater microbial biofilm in terms of community composition and EEA. Our original hypotheses predicted changes to bacterial biodiversity, and biofilm metabolic activity after exposure to either surfactant. Rhamnolipids acted as an enhancer for beta-glucosidase and leucine aminopeptidase, increasing EEA within the biofilm, while SDS likely denatured the enzymes and reduced EEA within the biofilm. While there were no differences in GPP and CR, EEA changes can be used as a proxy for community-scale metabolic activity within the biofilms. There were alterations in biofilm metabolic activity post surfactant exposure, supporting one of our original hypotheses. Both surfactants also decreased the diversity of the biofilm bacterial community, which supports our other original hypothesis. There were unique genera in the biofilms of each treatment, including the controls which demonstrates that the presence of either surfactant creates an environment that is inhospitable to sensitive taxa. Although we saw differences in EEA and diversity, there were no differences in GPP and CR, potentially demonstrating functional redundancy within mixed microbial biofilms. These results highlight several important areas of concern. Firstly, rhamnolipids may not be a suitable alternative to SDS for use environmental biofilm

control because of its lower effectiveness in open environmental settings. Secondly, in natural ecosystems where biofilms are necessary for key ecosystem processes, such as degradation of organic matter, we have shown that the presence of these surfactants (for example through the release from WWTPs) may lead to negative effects on microbial diversity and metabolic activity within biofilms. Finally, our results demonstrate that functional redundancy may exist within diverse biofilms. Surfactant exposure altered community composition and EEA, but the overall biofilm did not experience any negative respiratory effects. Therefore, while the function of the biofilm may be altered, enough bacteria are still taking part in respiratory cycles that the overall metabolic activity of the biofilm is not disrupted.

**Acknowledgements** We thank Gertraud Steniczka and Hermann Hofreiter for help in the laboratory and with setting up the experiment.

**Author contribution** SG, WH, LC, IB, and JS were all equally involved in the design, execution, and data analysis within the study. SG wrote the first draft of the manuscript with the exception of the extracellular enzyme analysis methodology which was written by LC. All authors contributed to commenting on and editing the manuscript and have read and approved of the final manuscript.

**Funding** SG is funded by an Ulster University Vice Chancellors Doctoral Research Fellowship, and received additional support through an Ulster University Broadening Horizons Travel Bursary. Analytical costs were partly supported by the HYDRO-DIVERSITY project funded by the Environmental Systems Sciences Program of the Austrian Academy of Sciences (ÖAW) to JS, and core funding of the AFBI Aquatic Chemistry Laboratory (WH).

**Data availability** Data can be accessed at <https://zenodo.org/record/5153485#.YQgBzEAo9PY>. Access is restricted and must be requested from the page to prevent work from being copied before publishing. After publishing work, will be publicly accessible.

## Declarations

**Ethics approval** Not applicable.

**Consent for publication** Not applicable.

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Akbari A, Kasprzyk A, Galvez R, Ghoshal S (2021) A rhamnolipid biosurfactant increased bacterial population size but hindered hydrocarbon biodegradation in weathered contaminated soils. *Sci Total Environ*. <https://doi.org/10.1016/j.scitotenv.2021.145441>
- Aleksic I, Petkovic M, Jovanovic M, Miliwojevic D, Vasiljevic B, Nikodinovic-Runic J, Senerovic L (2017) Anti-biofilm properties of bacterial di-Rhamnolipids and their semi-synthetic amide derivatives. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2017.02454>
- Al-Tahhan R, Sandrin TR, Bodour AA, Maier RM (2000) Rhamnolipid-induced removal of lipopolysaccharide from *Pseudomonas aeruginosa*: effect on cell surface properties and interaction with hydrophobic substrates. *Appl Environ Microbiol*. <https://doi.org/10.1128/AEM.66.8.3262-3268.2000>
- Argudo M, Gich F, Bonet B, Espinosa C, Gutiérrez M, Guasch H (2020) Responses of resident (DNA) and active (RNA) microbial communities in fluvial biofilms under different polluted scenarios. *Chemosphere*. <https://doi.org/10.1016/j.chemosphere.2019.125108>
- Bai H, Wang H, Sun J, Irfan M, Han M, Huang Y, Han X, Yang Q (2013) Production, purification and characterization of novel beta glucosidase from newly isolated *Penicillium simplicissimum H-11* in submerged fermentation. *EXCLI journal*. <https://doi.org/10.17877/DE290R-7346>
- Banar M, Emaneini M, Beigverdi R, Fanaei Pirlar R, Node Farahani N, van Leeuwen WB, Jabalameli F (2019) The efficacy of lyticase and  $\beta$ -glucosidase enzymes on biofilm degradation of *Pseudomonas aeruginosa* strains with different gene profiles. *BMC Microbiol* <https://doi.org/10.1186/s12866-019-1662-9>
- Banat IM, Díaz De Rienzo MA, Quinn GA (2014). Microbial biofilms: biosurfactants as antibiofilm agents. *Appl Microbiol Biotechnol*. <https://doi.org/10.1007/s00253-014-6169-6>
- Battin TJ, Besemer K, Bengtsson MM, Romani AM, Packmann AI (2016) The ecology and biogeochemistry of stream biofilms. *Nat Rev Microbiol*. <https://doi.org/10.1038/nrmicro.2016.15>
- Bhuyan AK (2010) On the mechanism of SDS-induced protein denaturation. *Biopolymers*. <https://doi.org/10.1002/bip.21318>
- Bowler P, Murphy C, Wolcott R (2020) Biofilm exacerbates antibiotic resistance: is this a current oversight in antimicrobial stewardship? *Antimicrob Resist Infect Control*. <https://doi.org/10.1186/s13756-020-00830-6>
- Cao C, Cao Z, Yu P, Zhao Y (2020) Genome-wide identification for genes involved in sodium dodecyl sulfate toxicity in *Saccharomyces cerevisiae*. *BMC Microbiol*. <https://doi.org/10.1186/s12866-020-1721-2>
- Campos JM, Montenegro Stamford TL, Sarubbo LA, de Luna JM, Rufino RD, Banat IM (2013) Microbial biosurfactants as additives for food industries. *Biotechnol Progress* 29:1097–1108. <https://doi.org/10.1002/btpr.1796>
- Ceresa C, Tessarolo F, Maniglio D, Tambone E, Carmagnola I, Fedeli E, Caola I, Nollo G, Chiono V, Allegrone G, Rinaldi M, Fracchia L (2019) Medical-grade silicone coated with Rhamnolipid R89 is effective against *Staphylococcus* spp biofilms. *Molecules* 24:3843. <https://doi.org/10.3390/molecules24213843>
- Ceresa C, Rinaldi M, Tessarolo F, Maniglio D, Fedeli E, Tambone E, Caciagli P, Banat IM, Diaz De Rienzo MA, Fracchia L (2021) Inhibitory effects of lipopeptides and glycolipids on *C albicans* - *Staphylococcus* spp dual-species biofilms. *Front Microbiol*. 11:545654. <https://doi.org/10.3389/fmicb.2020.545654>
- Ceresa C, Fracchia L, Fedeli E, Porta C, Banat IM (2021b) Recent advances in biomedical, therapeutic and pharmaceutical applications of microbial surfactants. *Pharmaceutics* 13(4):466. <https://doi.org/10.3390/pharmaceutics13040466>
- Chao A (1984) Nonparametric estimation of the number of classes in a population. *Scand J Stat* 11:265–270
- Chang YT, Lin CY, Chen YH, Hsueh P (2015) Update on infections caused by *Stenotrophomonas maltophilia* with particular attention to resistance mechanisms and therapeutic options. *Front Microbiol*. <https://doi.org/10.3389/fmicb.2015.00893>
- Cheah YT, Chan DJC (2021) Physiology of microalgal biofilm: a review on prediction of adhesion on substrates <https://doi.org/10.1080/21655979.2021.1980671>
- Chen X, Wang L, Zhou J, Wu H, Li D, Cui Y, Lu B (2017) *Exiguobacterium* sp. *A1b/GX59* isolated from a patient with community-acquired pneumonia and bacteremia: genomic characterization and literature review. *BMC Infect Dis*. <https://doi.org/10.1186/s12879-017-2616-1>
- Cserháti T, Forgács E, Oros G (2002) Biological activity and environmental impact of anionic surfactants. *Environ Int* [https://doi.org/10.1016/S0160-4120\(02\)00032-6](https://doi.org/10.1016/S0160-4120(02)00032-6)
- Cui B, Liu X, Yang Q, Li J, Zhou X, Peng Y (2017) Achieving partial denitrification through control of biofilm structure during biofilm growth in denitrifying biofilter. *Bioresour Technol* <https://doi.org/10.1016/j.biortech.2017.04.034>
- Dang H, Lovell CR (2016) Microbial surface colonization and biofilm development in marine environments. *Microbiol Mol Biol Rev* <https://doi.org/10.1128/MMBR.00037-15>
- Decho AW (2000) Microbial biofilms in intertidal systems: an overview. *Cont Shelf Res* [https://doi.org/10.1016/S0278-4343\(00\)00022-4](https://doi.org/10.1016/S0278-4343(00)00022-4)
- Desai JD, Banat IM (1997) Microbial production of surfactants and their commercial potential. *Microbiol Mol Biol Rev* <https://doi.org/10.1128/61.1.47-64.1997>
- Díaz De Rienzo MA, Stevenson PS, Marchant R, Banat IM (2016) *Pseudomonas aeruginosa* biofilm disruption using microbial surfactants. *J Appl Microbiol* <https://doi.org/10.1111/jam.13049>
- Dodds WK, Evans-White M, Gerlanc NM, Gray L, Gudder DA, Kemp MJ, Lopez AL, Stagliano D, Strauss EA, Tank JL, Whiles MR, Wollheim WM (2000) Quantification of the nitrogen cycle in a prairie stream. *Ecosystems* 3:574–589
- Drury B, Scott J, Rosi-Marshall EJ, Kelly JJ (2013) Triclosan exposure increases triclosan resistance and influences taxonomic composition of benthic bacterial communities. *Environ Sci Technol* <https://doi.org/10.1021/es401919k>
- Dusane DH, Zinjjarde SS, Venugopalan VP, Mclean RJ, Weber MM, Rahman PKSM (2010) Quorum sensing: implications on Rhamnolipid biosurfactant production. *Biotechnol Genet Eng Rev* <https://doi.org/10.1080/02648725.2010.10648149>
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* <https://doi.org/10.1093/bioinformatics/btr381>
- Flemming H, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S (2016) Biofilms: an emergent form of bacterial life. *Nat Rev Microbiol* <https://doi.org/10.1038/nrmicro.2016.94>
- Fong JNC, Yildiz FH (2015) Biofilm Matrix Proteins. *Microbiol Spectr* 3:10. <https://doi.org/10.1128/microbiolspec.MB-0004-2014>
- Fox J, Weisberg S (2019) *An R Companion to applied regression*, Third Edition. Thousand Oaks California, USA: Sage. Accessed: February 2022. <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>
- Franzetti A, Gandolfi I, Raimondi C, Bestetti G, Banat IM, Smyth TJ, Papacchini M, Cavallo M, Fracchia L (2012) Environmental fate, toxicity, characteristics and potential applications of novel bioemulsifiers produced by *Variovorax paradoxus 7bCT5*. *Bioresour Technol* <https://doi.org/10.1016/j.biortech.2012.01.005>
- Gudiksen KL, Gitlin I, Whitesides GM (2006) Differentiation of proteins based on characteristic patterns of association and denaturation in solutions of SDS. *Proc Natl Acad Sci USA* <https://doi.org/10.1073/pnas.0602816103>
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: Paleontological statistics software package for education and data analysis. *Palaeontol Electron* <https://doi.org/10.1016/j.bcp.2008.05.025>

- Harjung A, Attermeyer K, Aigner V, Krlovic N, Steniczka G, Švecová H, Schagerl M, Schelker J (2020) High anthropogenic organic matter inputs during a festival increase river heterotrophy and refractory carbon load. *Environ Sci Technol* <https://doi.org/10.1021/acs.est.0c02259>
- Hay AG, Dees PM, Saylor GS (2001) Growth of a bacterial consortium on triclosan. *FEMS Microbiol Ecol* [https://doi.org/10.1016/S0168-6496\(01\)00127-1](https://doi.org/10.1016/S0168-6496(01)00127-1)
- Izano EA, Wang H, Ragunath C, Ramasubbu N, Kaplan JB (2007) Detachment and killing of *Aggregatibacter actinomycetemcomitans* biofilms by Dispersin B and SDS. *J Dent Res* <https://doi.org/10.1177/154405910708600707>
- Jefferson KK (2004) What drives bacteria to produce a biofilm? *FEMS Microbiol Lett* <https://doi.org/10.1111/j.1574-6968.2004.tb09643.x>
- Jost L (2007) Partitioning diversity into independent alpha and beta components. *Ecol* <https://doi.org/10.1890/06-1736.1>
- Ke Y, Chen J, Hu X, Tong T, Huang J, Xie S (2020) Emerging perfluoroalkyl substance impacts soil microbial community and ammonia oxidation. *Environ Pollut* <https://doi.org/10.1016/j.envpol.2019.113615>
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD (2013) Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ Microbiol* <https://doi.org/10.1128/AEM.01043-13>
- Kumar A, Yadav CB, Bajpeyee AK (2020) Glucose, protein, cholesterol and immunohaematological alterations with acute (SDS) Sodium Dodecyl Sulfate toxicity in *Clarias batrachus*. *J Exp Zoo (india)* 23:401–404
- Kusi J, Scheuerman PR, Maier KJ (2020) Emerging environmental contaminants (silver nanoparticles) altered the catabolic capability and metabolic fingerprinting of microbial communities. *Aquat Toxicol* <https://doi.org/10.1016/j.aquatox.2020.105633>
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680–685
- Landa AS, Van de Belt-Gritter B, Van der Mei HC, Busscher HJ (1999) Recalcitrance of *Streptococcus mutans* biofilms towards detergent-stimulated detachment. *Eur J Oral Sci* <https://doi.org/10.1046/j.0909-8836.1999.eos107402.x>
- Lara-Martín PA, González-Mazo E, Petrovic M, Barceló D, Brownawell BJ (2014) Occurrence, distribution and partitioning of nonionic surfactants and pharmaceuticals in the urbanized Long Island Sound Estuary (NY). *Mar Pollut Bull* <https://doi.org/10.1016/j.marpolbul.2014.01.022>
- Li Y, Wu J, Shen J, Liu S, Wang C, Chen D, Huang T, Zhang J (2016) Soil microbial C:N ratio is a robust indicator of soil productivity for paddy fields. *Sci Rep* <https://doi.org/10.1038/srep35266>
- Lima TMS, Procópio LC, Brandão FD, Leão BA, Tótoia MR, Borges AC (2011) Evaluation of bacterial surfactant toxicity towards petroleum degrading microorganisms. *Bioresour* <https://doi.org/10.1016/j.biortech.2010.09.109>
- Lindsay D, von Holy A (2006) Bacterial biofilms within the clinical setting: what healthcare professionals should know. *J Hosp Infect* <https://doi.org/10.1016/j.jhin.2006.06.028>
- Lu L, Zhang J, Peng C (2019) Shift of soil polycyclic aromatic hydrocarbons (PAHs) dissipation pattern and microbial community composition due to Rhamnolipid supplementation. *Water Air Soil Pollut* <https://doi.org/10.1007/s11270-019-4118-9>
- Lyon DR, Ziegler SE (2009) Carbon cycling within epilithic biofilm communities across a nutrient gradient of headwater streams. *Limnol Oceanogr* <https://doi.org/10.4319/lo.2009.54.2.0439>
- Madden T (2002) Chapter 16: The BLAST sequence analysis tool. In: *The NCBI Handbook* [Internet]. National Center for Biotechnology Information, Bethesda (MD). Available from: <https://www.ncbi.nlm.nih.gov/books/NBK21097/>
- Marchant R, Banat IM (2012) Biosurfactants: a sustainable replacement for chemical surfactants? *Biotechnol Lett* 34:1597–1605
- Margalef O, Sardans J, Fernández-Martínez M, Molowny-Horas R, Janssens IA, Ciais P, Goll D, Richter A, Obersteiner M, Asensio D, Peñuelas J (2017) Global patterns of phosphatase activity in natural soils. *Sci Rep* <https://doi.org/10.1038/s41598-017-01418-8>
- Mariani L, De Pascale D, Faraponova O, Tornambè A, Sarni A, Giuliani S, Ruggiero G, Onorati F, Magaletti E (2006) The use of a test battery in marine ecotoxicology: the acute toxicity of Sodium Dodecyl Sulfate. *Enviro Toxicol* <https://doi.org/10.1002/tox.20204>
- Matsui M, Fowler JH, Walling LL (2006) Leucine aminopeptidases: diversity in structure and function. *Biol Chem* <https://doi.org/10.1515/BC.2006.191>
- McClellan P, Hunter WR (2020) 17 $\alpha$ -ethynylestradiol (EE2) limits the impact of ibuprofen upon respiration by streambed biofilms in a sub-urban stream. *Environ Sci Pollut Res* 27:37149–37154
- Messina CM, Faggio C, Laudicella VA, Sanfilippo M, Trischitta F, Santulli A (2014) Effect of sodium dodecyl sulfate (SDS) on stress response in the Mediterranean mussel (*Mytilus Galloprovincialis*): regulatory volume decrease (Rvd) and modulation of biochemical markers related to oxidative stress. *Aquat Toxicol* <https://doi.org/10.1016/j.aquatox.2014.10.001>
- Milovanovic M, Arsenijevic A, Milovanovic J, Kanjevac T, Arsenijevic N (2017) Nanoparticles in antiviral therapy. *Antimicrob Nanoarchitectonics*. <https://doi.org/10.1016/B978-0-323-52733-0.00014-8>
- Mireles II JR, Toguchi A, Harshey RM (2001) *Salmonella enterica* Serovar Typhimurium swarming mutants with altered biofilm-forming abilities: Surfactin inhibits biofilm formation. *J Bacteriol* <https://doi.org/10.1128/JB.183.20.5848-5854.2001>
- Mnif I, Ellouze-Chaabouni S, Ghribi D (2018) Glycolipid biosurfactants, main classes, functional properties and related potential applications in environmental biotechnology. *J Polym Environ* <https://doi.org/10.1007/s10924-017-1076-4>
- Mohan PK, Nakhla G, Yanful EK (2006) Biokinetics of biodegradation of surfactants under aerobic, anoxic and anaerobic conditions. *Water Res (Oxford)* <https://doi.org/10.1016/j.watres.2005.11.030>
- Naughton PJ, Marchant R, Naughton V, Banat IM (2019). Microbial biosurfactants: current trends and applications in Agricultural and Biomedical industries. *J Appl Microbiol*. <https://doi.org/10.1111/jam.14243>
- Nguyen BVG, Nagakubo T, Toyofuku M, Nomura N, Utada AS (2020) Synergy between Sophorolipid biosurfactant and SDS increases the efficiency of *P. aeruginosa* biofilm disruption. *Langmuir*. <https://doi.org/10.1021/acs.langmuir.0c00643>
- Otzen DE (2017) Biosurfactants and surfactants interacting with membranes and proteins: same but different? *Biochim Biophys Acta Biomembr* <https://doi.org/10.1016/j.bbmem.2016.09.024>
- Paul BJ, Duthie HC, Taylor WD (1991) Nutrient cycling by biofilms in running waters of differing nutrient status. *J North Am Benthol Soc* <https://doi.org/10.2307/1467762>
- Qu X, Ren Z, Zhang H, Zhang M, Zhang Y, Liu X, Peng W (2017) Influences of anthropogenic land use on microbial community structure and functional potentials of stream benthic biofilms. *Sci Rep*. <https://doi.org/10.1038/s41598-017-15624-x>
- Quinn GA, Maloy AP, Banat MM, Banat IM (2013) A comparison of effects of broad-spectrum antibiotics and biosurfactants on established bacterial biofilms *Curr Microbiol*. <https://doi.org/10.1007/s00284-013-0412-8>
- Ramos da Silva A, Manresa MÁ, Pinazo A, García MT, Pérez L (2019) Rhamnolipids functionalized with basic amino acids: synthesis, aggregation behavior, antibacterial activity and biodegradation studies *Colloids Surf B*. <https://doi.org/10.1016/j.colsurfb.2019.05.037>
- Rigotti RT, Corrêa JAF, Maia NJL, Cesaro G, Rosa EAR, Macedo, Renata Ernlund Freitas de, Luciano FB (2017) Combination of natural antimicrobials and sodium dodecyl sulfate for disruption of biofilms formed by contaminant bacteria isolated from sugarcane mills. *Innov Food Sci Emerg Technol* <https://doi.org/10.1016/j.ifset.2017.01.007>

- Rice LB (2008) Federal funding for the study of antimicrobial resistance in nosocomial pathogens: No ESKAPE J Infect Dis. <https://doi.org/10.1086/533452>
- Roche KR, Drummond JD, Boano F, Packman AI, Battin TJ, Hunter WR (2017) Benthic biofilm controls on fine particle dynamics in streams Water Resour Res. <https://doi.org/10.1002/2016WR019041>
- Rodrigues L, Banat IM, Teixeira J, Oliveira R (2006) Biosurfactants: potential applications in medicine J Antimicrob Chemother. <https://doi.org/10.1093/jac/dk1024>
- Romaní AM, Sabater S (2001) Structure and activity of rock and sand biofilms in a Mediterranean stream Ecol (durham). [https://doi.org/10.1890/0012-9658\(2001\)082\[3232:SAORA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[3232:SAORA]2.0.CO;2)
- Rosi-Marshall E, Kincaid DW, Bechtold HA, Royer TV, Rojas M, Kelly JJ (2013) Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial communities in stream biofilms Ecol Appl. <https://doi.org/10.1890/12-0491.1>
- Ryan MP, Pembroke JT (2018) Brevundimonas spp: emerging global opportunistic pathogens Virulence. <https://doi.org/10.1080/21505594.2017.1419116>
- Satpute SK, Mone NS, Das P, Banpurkar AG, Banat IM (2018) Lactobacillus acidophilus derived biosurfactant as a biofilm inhibitor: a promising investigation using microfluidic approach Appl Sci. <https://doi.org/10.3390/app8091555>
- Schaidler LA, Rudel RA, Ackerman JM, Dunagan SC, Brody JG (2014) Pharmaceuticals, perfluorosurfactants, and other organic wastewater compounds in public drinking water wells in a shallow sand and gravel aquifer Sci Total Environ. <https://doi.org/10.1016/j.scitotenv.2013.08.067>
- Schlager B, Straessle A, Hafen E (2012) Use of anionic denaturing detergents to purify insoluble proteins after overexpression BMC Biotechnol. <https://doi.org/10.1186/1472-6750-12-95>
- Schloss PD, Westcott SL, Thomas R, Hall JR, Martin H, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Blaz S, Thallinger GG, Horn Van DJ, Weber CF (2009) Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities Appl Environ Microbiol. <https://doi.org/10.1128/AEM.01541-09>
- Schmitz KS (2018) Chapter 4: Life Science. Physical Chemistry. Elsevier, Boston, pp 755–832
- Schreiberova O, Hedbavna P, Aejkova A, Jirka[macron] V, Masak J (2012) Effect of surfactants on the biofilm of Rhodococcus erythropolis, a potent degrader of aromatic pollutants New Biotechnol. <https://doi.org/10.1016/j.nbt.2012.04.005>
- Seghers D, Verthé K, Reheul D, Bulcke R, Siciliano SD, Verstraete W, Top EM (2003) Effect of long-term herbicide applications on the bacterial community structure and function in an agricultural soil FEMS Microbiol Ecol. [https://doi.org/10.1016/S0168-6496\(03\)00205-8](https://doi.org/10.1016/S0168-6496(03)00205-8)
- Simões M, Simões LC, Pereira MO, Vieira MJ (2008) Sodium dodecyl sulfate allows the persistence and recovery of biofilms of Pseudomonas fluorescens formed under different hydrodynamic conditions Biofouling. <https://doi.org/10.1080/08927010701730311>
- Singh G, Verma AK, Kumar V (2016) Catalytic properties, functional attributes and industrial applications of  $\beta$ -glucosidases. 3 Biotech. <https://doi.org/10.1007/s13205-015-0328-z>
- Sloup RE, Cieza RJ, Needle DB, Abramovitch RB, Torres AG, Waters CM (2016) Polysorbates prevent biofilm formation and pathogenesis of Escherichia coli O104:H4 Biofouling. <https://doi.org/10.1080/08927014.2016.1230849>
- Srey S, Jahid IK, Ha S (2013) Biofilm formation in food industries: a food safety concern Food Control. <https://doi.org/10.1016/j.foodcont.2012.12.001>
- Sturner RW, Elser JJ (2002) Ecological stoichiometry. Princeton University Press
- Sumpono PP, Belan A, Forestier C, Lavedrine B, Bohatier J (2003) Effect of Diuron on aquatic bacteria in laboratory-scale wastewater treatment ponds with special reference to Aeromonas species studied by colony hybridization Chemosphere. [https://doi.org/10.1016/S0045-6535\(02\)00621-5](https://doi.org/10.1016/S0045-6535(02)00621-5)
- Tambone E, Bonomi E, Ghensi P, Maniglio D, Ceresa C, Agostinaccio F, Caciagli P, Nollo G, Piccoli F, Caola I, Fracchia L, Tessarolo F (2021) Rhamnolipid coating reduces microbial biofilm formation on titanium implants: an in vitro study BMC Oral Health. <https://doi.org/10.1186/s12903-021-01412-7>
- Tewari R, Chopra D, Wazahat R, Dhingra S, Dudeja M (2018) Antimicrobial susceptibility patterns of an emerging multidrug resistant nosocomial pathogen: Acinetobacter baumannii. Malays J Med Sci. <https://doi.org/10.21315/mjms2018.25.3.13>
- Thompson LJ, Gray V, Lindsay D, Holy Von A (2006) Carbon: nitrogen: phosphorus ratios influence biofilm formation by Enterobacter cloacae and Citrobacter freundii J Appl Microbiol. <https://doi.org/10.1111/j.1365-2672.2006.03003.x>
- Ueda Y, Mashima K, Miyazaki M, Hara S, Takata T, Kamimura H, Takagi S, Jimi S (2019) Inhibitory effects of polysorbate 80 on MRSA biofilm formed on different substrates including dermal tissue Sci Rep. <https://doi.org/10.1038/s41598-019-39997-3>
- Waal Van de DB, Elser JJ, Martiny AC, Sturner RW, Cotner JB (2018) Editorial: Progress in ecological stoichiometry Front Microbiol. <https://doi.org/10.3389/fmicb.2018.01957>
- Waal SM, Rölla G, Skjörland KK, Ogaard B (1993) Effects of oral rinsing with triclosan and sodium lauryl sulfate on dental plaque formation: a pilot study. Scand J Dent Res 101:192–195
- Wang H, Fan B, Li C, Liu S, Li M (2011) Effects of Rhamnolipid on the cellulase and xylanase in hydrolysis of wheat straw. Bioresour Technol 102:6515–6521
- Wei Z, Wang JJ, Gaston LA, Li J, Fultz LM, DeLaune RD, Dodla SK (2020) Remediation of crude oil-contaminated coastal marsh soil: integrated effect of biochar, Rhamnolipid biosurfactant and nitrogen application. J Hazard Mater. <https://doi.org/10.1016/j.jhazmat.2020.122595>
- Wickham H (2016) Ggplot2: Elegant graphics for data analysis. Springer-Verlag, New York, USA
- Xie F, Yang Y, Ma H, Quan S, Yue D, Sun Y, Feng F, Zhen J, Liu D (2017) Nocardioides phosphati sp. nov., an actinomycete isolated from a phosphate mine. Int J Syst Evol. <https://doi.org/10.1099/ijsem.0.001754>
- Yao X, Xiao L, Liu H, Zhou M, Zhou Z, Ju X, Li L (2020) Rhamnolipid enhanced beta-glucosidase from Paenibacillus sp. LLZ1 for in situ lignocellulose saccharification in ionic liquids. Biomass Convers Biorefin. <https://doi.org/10.1007/s13399-020-01002-7>
- Yin Y, Gu J, Wang X, Zhang Y, Zheng W, Chen R, Wang X (2019) Effects of rhamnolipid and Tween-80 on cellulase activities and metabolic functions of the bacterial community during chicken manure composting Bioresour Technol. <https://doi.org/10.1016/j.biortech.2019.121507>
- Zakaria BS, Dhar BR (2020) Changes in syntrophic microbial communities, EPS matrix, and gene-expression patterns in biofilm anode in response to silver nanoparticles exposure Sci Total Environ. <https://doi.org/10.1016/j.scitotenv.2020.139395>
- Zhang Q, He G, Wang J, Cai W, Xu Y (2009) Mechanisms of the stimulatory effects of Rhamnolipid biosurfactant on rice straw hydrolysis Appl Energy. <https://doi.org/10.1016/j.apenergy.2009.04.030>
- Zhang S, Pang S, Wang P, Wang C, Guo C, Addo FG, Li Y (2016) Responses of bacterial community structure and denitrifying bacteria in biofilm to submerged macrophytes and nitrate Sci Rep. <https://doi.org/10.1038/srep36178>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.