Enhancing the effects of nusinersen with cybernic treatment using Hybrid

Assistive Limb (HAL) in spinal muscular atrophy: a real-world case series and

exploratory cohort analysis

**Authors:** Takashi Nakajima<sup>1</sup>, Toshio Saito<sup>2</sup>, Akihiro Hashiguchi<sup>3</sup>, Taiki Nakabayashi<sup>4</sup>, Kazuki Kodera<sup>4</sup>,

Kota Utsumi<sup>1</sup>, Takeshi Kanayama<sup>1</sup>, Haruka Urabe<sup>1</sup> and Satoru Kinoshita<sup>4</sup>

Affiliations: <sup>1</sup>Department of Neurology, NHO Niigata National Hospital, Kashiwazaki, Japan; <sup>2</sup>Division of

Child Neurology, Department of Neurology, NHO Osaka Toneyama Medical Centre, Toneyama, Japan;

<sup>3</sup>Department of Neurology and Geriatrics, Kagoshima University Graduate School of Medical and Dental

Sciences, Kagoshima, Japan; <sup>4</sup>Department of Paediatrics, NHO Niigata National Hospital, Kashiwazaki,

Japan

\* Correspondence: Takashi Nakajima: nakajima-md@mqb.biglobe.ne.jp

Supplementary material

1

## STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No.	Recommendation	Page No.
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	1-2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	2-6
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	(a) Cohort study—Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up  Case-control study—Give the eligibility criteria, and the sources and methods of case ascertainment and control selection.  Give the rationale for the choice of cases and controls  Cross-sectional study—Give the eligibility criteria, and the sources and methods of selection of participants	7
		(b) Cohort study—For matched studies, give matching criteria and number of exposed and unexposed Case-control study—For matched studies, give matching criteria and the number of controls per case	7
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	9
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	8-9
Bias	9	Describe any efforts to address potential sources of bias	N/A
Study size	10	Explain how the study size was arrived at	9-10

Continued on next page

Quantitative	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and	10
variables		why	
Statistical	12	(a) Describe all statistical methods, including those used to control for confounding	10
methods		(b) Describe any methods used to examine subgroups and interactions	10
		(c) Explain how missing data were addressed	10
		(d) Cohort study—If applicable, explain how loss to follow-up was addressed	N/A
		Case-control study—If applicable, explain how matching of cases and controls was addressed	
		Cross-sectional study—If applicable, describe analytical methods taking account of sampling strategy	
		$(\underline{e})$ Describe any sensitivity analyses	N/A
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed	11
		eligible, included in the study, completing follow-up, and analysed	
		(b) Give reasons for non-participation at each stage	N/A
		(c) Consider use of a flow diagram	N/A
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	11-12, 29
		(b) Indicate number of participants with missing data for each variable of interest	29
		(c) Cohort study—Summarise follow-up time (eg, average and total amount)	11-12, 29
Outcome data	15*	Cohort study—Report numbers of outcome events or summary measures over time	12-13, 30, 31
		Case-control study—Report numbers in each exposure category, or summary measures of exposure	N/A
		Cross-sectional study—Report numbers of outcome events or summary measures	N/A
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence	12-13, 31
		interval). Make clear which confounders were adjusted for and why they were included	
		(b) Report category boundaries when continuous variables were categorized	N/A
		© If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A

Continued on next page

Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	
Key results	18	Summarise key results with reference to study objectives	14-17
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	17-18
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	18-19
Generalisability	21	Discuss the generalisability (external validity) of the study results	14-17
Other informati	on		
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	20

<sup>\*</sup>Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at http://www.plosmedicine.org/, Annals of Internal Medicine at http://www.annals.org/, and Epidemiology at http://www.epidem.com/). Information on the STROBE Initiative is available at www.strobe-statement.org.