



ORIGINAL ARTICLE

Association of polymorphism in heat shock protein 70 genes with type 2 diabetes in Bangladeshi population

Md. Moniruzzaman¹  | Irfan Ahmed¹ | Saaimatul Huq¹ | Md. Shakur All Mahmud² | Sonya Begum³ | U.S. Mahzabin Amin¹  | Md. Hadisur Rahman¹ | Palash Kumar Sarker⁴ | Mohammad Uzzal Hossain⁵ | Keshob Chandra Das¹ | Md. Salimullah¹

¹Molecular Biotechnology Division, National Institute of Biotechnology, Dhaka, Bangladesh

²Department of Biochemistry and Molecular Biology, Tejgaon College, Dhaka, Bangladesh

³Department of Biotechnology and Genetic engineering, Mawlana Bhashani Science and Technology University, Tangail, Bangladesh

⁴Microbial Biotechnology Division, National Institute of Biotechnology, Dhaka, Bangladesh

⁵Bioinformatics Division, National Institute of Biotechnology, Dhaka, Bangladesh

Correspondence

Md. Salimullah, Director General (Additional Charge) & Chief Scientific Officer, Molecular Biotechnology Division, National Institute of Biotechnology, Ganakbari, Ashulia, Savar, Dhaka-1349, Bangladesh.
Email: salim2969@gmail.com

Funding information

The work has been supported by the fund of Special allocation of Ministry of Science and technology, Government of the people's Republic of Bangladesh in the fiscal year of 2015- 16. The project ID was MEDI; S- 250.

Abstract

Background: Type 2 diabetes mellitus (T2DM) is a chronic disorder of which stress is a major contributor. Under stressful condition, body synthesizes a family of molecular chaperone called Heat-shock proteins (HSPs). Current study assessed the frequency and association of *HSP70-hom + 2,437 T/C* polymorphism with T2DM risk among Bangladeshis.

Methods: This polymorphism was selected through bioinformatics analyses and identified by PCR-RFLP method.

Results: Bioinformatics analysis identified this SNP as missense mutation which could destabilize the final HSP product. Heterozygous mutant (CT) genotype was significantly associated with T2DM incidence among the studied populations ($p = .015$). Further analysis revealed a strong association with female patients ($p = .002$), while the male group showed no association ($p = .958$). Moreover, the C allele was significantly associated among all diabetic patients ($p = .016$) and particularly in the female patient group ($p = .001$). However, under stressful condition, males with CT genotype were at high risk for T2DM incidence whereas, females with CT genotype showed no significant association.

Conclusions: *HSP70-hom + 2,437 T/C* polymorphism was found to be significantly associated with T2DM incidence in the Bangladeshi population in both stress-dependent and independent manners.

KEYWORDS

bioinformatics, diabetes mellitus, heat-shock proteins, polymorphism, stress

1 | INTRODUCTION

Type 2 diabetes mellitus (T2DM) is an ever-increasing chronic metabolic disorder with a strong genetic predisposition.

Globally, 1 of 11 adults bearing diabetes while it is forecasted to be 1 in 10 by the year 2040 (IDF Atlas, 2015). The high incidence of diabetes creates a major problem in developing countries like Bangladesh where the prevalence of diabetes

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. *Molecular Genetics & Genomic Medicine* published by Wiley Periodicals, Inc.

are estimated to elevate from 9.2% to 13% by 2030 (IDF Atlas, 2015; Rahman et al., 2015).

Stress is one of the major contributors to diabetes. Oxidative stress has been proposed as the root cause underlying the development of insulin resistance, β -cell dysfunction, impaired glucose tolerance, and T2DM through the production of reactive oxygen species. It has also been implicated in the progression of long-term diabetes-associated complications (S. J. Kelly & Ismail, 2015; Pouwer, Kupper, & Adriaanse, 2010). Stresses such as heat shock, ischemia, and other types of cellular stresses induce synthesis of one of the most ubiquitous and highly conserved molecular chaperone, heat shock proteins (HSPs) (Gehrmann et al., 2005; Lindquist, 1986). The 70-kDa HSP (HSP70) family is the most abundant in eukaryotic cells and is essential for cell survival under stressful conditions (Kelly, 2002). In humans, there are three main genes (*HSPA1A*, *HSPA1B*, and *HSPAIL*) of *HSP70* family and the coding proteins are defined as HSP70-1, HSP70-2, and HSP70-hom, respectively. Through their effect on insulin sensitivity, HSPs have also been reported to be involved in diabetes (Macario & De Macario, 2000).

HSP70 gene polymorphism is reported to be associated with an increased risk of developing various disorders including T2DM (Chouchane et al., 2001; Ghanayem et al., 2014; He, Deng, & Luo, 2014; Pociot, Rønningen, & Nerup, 1993; Umapathy, Krishnamoorthy, & Muthukumar, 2012; Umapathy et al., 2012). One of the most studied single-nucleotide polymorphisms (SNPs) of *HSPAIL* is located at position +2437 of HSP70-hom (rs2227956). This is a nonsynonymous missense mutation observed in the coding region of HSP70-hom leading to the change in amino acid from threonine to methionine that affects the stability and activity of this protein (Singh et al., 2006). Moreover, this SNP could affect *HSP70* expression or function and further contribute to disease susceptibility and stress tolerance (Favatier, Bornman, Hightower, Günther, & Polla, 1997). Several epidemiological studies have been done to deduce the relationship of this polymorphism with obesity, DM and Diabetes nephropathy (DN) (Ghanayem et al., 2014). These studies revealed both significant and non-significant association between *HSP70-hom* +2437 T/C polymorphism and T2DM (Ghanayem et al., 2014; Umapathy et al., 2012). The SNP, *HSP70-hom* +2437 T/C was picked up for this study based on bioinformatics analysis which was authenticated by previous association studies. The soundness of this particular SNP study was confirmed through the measurement of its deleterious effect (Mooney, 2005) on the stabilization of the protein using different bioinformatics tools. We have further investigated the frequency of this polymorphism and its association with T2DM in the Bangladeshi mainstream population. To our knowledge, this is the first genetic association study on this SNP ever conducted on Bangladeshi patients with diagnosed T2DM cases. Also, we investigated the cumulative effect of stress and polymorphism on T2DM prevalence in the studied population.

2 | MATERIAL AND METHODS

2.1 | Ethical compliance

The case-control study approved by the ethics committee on National Institute of Biotechnology (NIB). The approval ID is NIBREC2015-02.

2.2 | Selection and in-silico validation of SNP *HSP70-hom* +2437 (T/C)

The information on target SNP (rs2227956) was derived from Online Mendelian Inheritance in Man (OMIM) and NCBI dbSNP (Database of Single-Nucleotide Polymorphism) (Hamosh, Scott, Amberger, Bocchini, & McKusick, 2005; Sherry et al., 2001). Protein sequence and its structure were retrieved by using Uniprot (Universal Protein Resource) and Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCSB PDB), respectively (Consortium, 2016; Rose et al., 2010). A web server called ConSurf was utilized to predict evolutionary conserved regions of the *HSPAIL* protein. A program called MODELLER 9v11 (Šali, Potterton, Yuan, van Vlijmen, & Karplus, 1995) through HHPred (Söding, Biegert, & Lupas, 2005) was used to determine the 3D structure of Wild-type and Mutant Protein encoded by *HSPAIL* gene. To find out the structure of the closest-related proteins, we submitted the protein sequence to HHPred Modeller tool and performed BLAST against the Protein Database (PDB). The best three-dimensional models were selected using ProCheck (Laskowski, MacArthur, Moss, & Thornton, 1993) and Verify3D (Lüthy, Bowie, & Eisenberg, 1992). Another tool named Yet Another Scientific Artificial Reality Application (YASARA) was introduced to calculate the energy minimization of both wild and possible mutant-type protein models (Kuszewski, Gronenborn, & Clore, 1997). This program yields the energy (START vs. END) that ensures accuracy of designed molecules. To confirm the variability in 3D structure of the proteins, an extensive virtual screening was performed with a suitable drug. At the same time, the drug's chemical structure, molecular weight, generic name, and binding capacity were also confirmed by bioinformatics analysis.

2.3 | Study population

The case-control study comprised of 216 clinically confirmed T2DM patients (mean age 50.38 ± 12.35 years) using glucose oxidase-peroxidase (GOD-POD) method and oral glucose tolerance test (OGTT) according to the WHO guideline from Diabetes and health awareness center, Kustia, Bangladesh and Medical Centre of National Institute of Biotechnology, Dhaka, Bangladesh

and 126 healthy controls (mean age 44.58 ± 12.4 years) from the general Bangladeshi population. The control respondent was further confirmed by measuring fasting plasma glucose (via the glucose oxidase-peroxidase [GOD-POD] method) (Meites & Saniel-Banrey, 1973). Informed written consent was taken from all subjects enrolled in the study in accordance with the Helsinki declaration. All participants also completed a questionnaire and participated in an interview. The administered questionnaire included information about age, sex, BMI, food habit, physical exercise, and history of first degree relatives, etc. As all patients included in this study were pre-diagnosed and most of the demographic parameters were in control, the information (except age and sex) were not included in this study.

2.4 | Sample size calculation and power of study

Based on the preliminary result of the pilot study using 40 samples of each group sample size was calculated with an estimated p value of <0.05 , 95% confidence interval (CI), and a power of 80 percent. The formula used for calculating the sample size is as follows (Charan & Biswas, 2013).

$$N = \frac{r+1}{r} \frac{(P^*)(1-P^*)(Z_{\alpha} + Z_{\beta/2})^2}{(P_1 - P_2)^2}$$

2.5 | Categorization of stress

The respondents were administered a detailed questionnaire comprised of information about life event (close relative's death, accident, failure in examination etc.), mental problem/depression, work stress, worries about future planning, financial stress, and daily life stress (Knol et al., 2006; Kumari, Head, & Marmot, 2004; Norberg et al., 2007; Rod, Grønbaek, Schnohr, Prescott, & Kristensen, 2009; Toshihiro, 2008). Numerical score had been given to every stress to assess the level of stress the participants suffered. Following this, the stress score was categorized into nonstress (0–1 point) and stress (2–6 points) groups. Further, the stress group was divided into low (2–3 points) and high (4–6 points) stress according to the modified method of Rod NH *et al* (Rod et al., 2009).

2.6 | Blood sampling and DNA extraction

Three milliliter peripheral blood was drawn in EDTA containing vacutainer (BD, Oxford, UK) Tube. Erythrocytes

of the samples were lysed by osmotic shock using 20 mM Tris-HCl (pH 8.0). DNA was extracted using Genomic DNA isolation kit (FavorPrep, Favorgen, Taiwan). The DNA concentration was determined using a Nanodrop spectrophotometer (Thermo Scientific, USA). Acceptable DNA samples (A260/A280 of 1.8–2.0) were diluted to 50 ng/μl and stored at -80°C until used.

2.7 | Genotyping

Genotype of *HSP70-hom* (+2437 T/C) was determined by the polymerase chain reaction–restriction fragment length polymorphism (PCR-RFLP) method using the following primers 5'GGACAAGTCTGAGAAGGTACAG-3' (forward) and 5'-GTAAGTTAGATTCAGGTCTGG-3' (reverse). Briefly, the PCR mixture contained a 1x reaction buffer (Thermo Fisher Scientific, Waltham, MA USA), 2.0 mM MgCl_2 , 0.2 mM dNTP, 20 picomoles of the two primers, 50–100 ng of genomic DNA and 1.125 unit of *Taq* DNA polymerase (Thermo Fisher Scientific, Waltham, MA USA) in a total volume of 25 μl. Amplification was accomplished by following the thermal cycling condition: 95°C for 5 min (one cycle); 95°C for 1 min, 61°C for 1 min, and 72°C for 1 min (35 cycles); 72°C for 7 min (one cycle), and hold at 4°C . The amplification product (862 bp) was visualized and documented using Gel documentation (proteinsimple, Santa Clara, CA USA) after resolving the product along with DNA size markers (1 Kb + DNA ladder, Invitrogen, USA) in 1% agarose gel and staining the gel with ethidium bromide. The resulting amplicons were subjected to restriction digestion using restriction endonuclease *NcoI* (New England Biolabs, UK) at 37°C for 2 hrs and then resolved in 1.5% gel.

2.8 | Statistical Analysis

Different statistical analyses using SPSS (SPSS v20, IBM) and MedCalc (Medcalc, Mariakerke, Belgium) softwares were performed to validate the results generated from the study. A robust test estimating χ^2 value at the 0.05% level of significance was carried out to check whether the genotypic frequencies of the polymorphism in *HSPA1L* deviated from Hardy–Weinberg equilibrium. The value was considered as rejected if it crossed the standard mark ($\chi^2 > 3.84$). Odds ratio (OR) with 95% confidence interval (CI) was calculated for the determination of the association of genotype and allele type with T2DM. Risk analysis by relative risk (RR) estimation with 95% CI confirmed the risk of disease occurrence for having specific genotypes. The above results having a p value ≤ 0.05 were accepted as statistically significant.

3 | RESULTS

3.1 | In-silico identification and validation of *HSP70-hom* +2437 (T/C) as a potent T2DM marker

Out of 1,192 SNPs found in *HSPA1L* gene, 359 are missense. The target SNP (rs2227956) is one of these missense SNPs. The protein encoded by *HSPA1L* gene consists of 641 amino acids in which amino acid at 493 position changes from threonine (T) to methionine (M) due to this SNP. Amino acid conservation analysis by ConSurf also predicted this T amino acid as a variable one (Figure S1). We built the 3D protein structures for both wild type and possible mutant sequence (T → M) based on the PDB id 5e84_A which shows 100% identity with HSPA1L protein. The tertiary structure having a minimum energy was predicted and the quality of the model was verified through different bioinformatics tools (Figure S2, S3, S4). Our analysis revealed decreased free energy for all mutant-type proteins (−1122.1 KJ/mol) than wild types (−1111.1 KJ/mol) (Figure S5). Drug binding analysis with an experimental drug DB07045 also identified the structural variation between wild type and mutant (T493M) model (Table S1). In wild-type Thr493, Asp154, Asn507, and Leu488 were the interacted amino acid residues with the drug whereas amino acids Ile503, Ile505, Arg511, lys509, and Gln158 interacted with the mutant model (Figure 1). The

interacting amino acid Thr493 is in the active site of the protein (Figure S6).

3.2 | Genotypic distribution of *HSP70-hom* (+2437)

DNA was isolated from peripheral blood of 216 clinically confirmed diabetes patients comprised of 78 males and 138 females and 126 closely age-matched clinically confirmed healthy control subjects, including 56 males and 70 females. The good quality isolated DNA was subjected to PCR-RFLP methods. The digestion generated two fragments of 616 bp and 246 bp size upon the presence of the desired SNP (Figure 2). Genotypic distribution of *HSP70-hom* (+2,437) polymorphism in the studied population followed the principle of Hardy–Weinberg equilibrium (Table S2).

Genotype frequency among the male, female, and all subjects with and without type 2 diabetes was measured among the studied subjects (Table 1). Frequencies of subjects bearing the CT and CC genotypes were higher in the diabetic group than in the control when all subjects are considered. Multivariate logistic regression was done to assess risk level of genotypes and alleles of the studied SNP (Table 2). The frequency difference of CC genotype was not statistically significant ($p = .725$), while the CT genotype increased the risk of type 2 diabetes with a significant level ($p = .015$) of difference. When the frequencies of CT and CC genotypes of male and female subjects were separately compared with




FIGURE 1 Drug–protein interacting residues in (a) wild type and (b) mutated model of *HSPA1L* protein. Active site amino acid, Thr 493 interacts with drug only in the case of wild protein (highlighted)




FIGURE 2 Restriction enzyme digestion pattern of *HSP70-hom* (+2437) polymorphism. For each possible pattern (TT, CT, and CC) PCR amplified and digested products are shown

TABLE 1 Genotype and allele frequency of SNP (rs2227956) in the study subjects

Catagories	Genotype/Allele	Control (%)	Patient (%)
All Subjects	CT	36 (28.6)	86 (39.8)
	TT	85 (67.5)	116 (53.7)
	CC	5 (3.9)	14 (6.5)
	C	46 (18.25)	74 (26.39)
	T	206 (81.75)	318 (73.61)
Male	CT	14 (29.16)	31 (39.74)
	TT	32 (66.66)	45 (57.69)
	CC	2 (4.16)	2 (2.56)
	C	18 (18.75)	35 (22.44)
	T	78 (81.25)	121 (77.56)
Female	CT	22 (28.21)	55 (39.85)
	TT	53 (67.95)	71 (51.44)
	CC	3 (3.84)	12 (8.69)
	C	28 (17.95)	79 (28.62)
	T	128 (82.05)	197 (71.38)

the corresponding control, the same scenario was found as that of all subjects. However, in the case of female subjects increased prevalence of CT genotype in patient was highly significant ($p = .002$). In comparison, the relation is not significant in the case of male respondents ($p = .958$).

The comparative analysis of allele frequency between diabetic group and control subjects showed that the C allele is significantly associated among all the patients (OR = 1.61, CI: 1.1–2.4, $p = .016$) and frequency difference was strongly significant for female patients group ($p = .001$) (Table 2). However, allele frequency of C did not differ by diabetes status among male subjects ($p = .880$).

3.3 | Association of *HSP70-hom* (+2437) polymorphism and stress with T2DM

The risk estimation analysis based on different combination of genotypic ratio is shown in Table S3. Higher p value made the ratio of CC/CT (OR = 1.17, $p = .78$) and CC/TT (OR = 2.05, $p = .18$) among the diabetic patients compared to the nondiabetic control group not significant, although OR and risk ratio (RR) were the highest for the CC/TT (OR = 2.05, RR = 1.28) among the three combination. However, the diabetes patient had higher ratio for CT/TT compare to the control and the result was statistically significant (OR = 1.75, $p = .02$; RR = 1.22, $p = .02$). The ratio of (CC + CT)/TT and CC/(CT + TT) among diabetes patient were substantially higher than that of control respondents, although the difference for CC/(CT + TT) was not quite statistically significant (OR = 1.68, $p = .33$; RR = 1.18, $p = .25$),

but the higher ratio for (CC + CT)/TT was statistically significant (OR = 1.79, $p = .01$; RR = 1.23, $p = .01$).

Whether age and stress variables are risk factors for type 2 diabetes incidence was assessed by multivariate logistic regression (Table S4). Subjects in the age groups of (40–60) and >60 years had 1.78 \times ($p = .005$) and 3.19 \times ($p = .006$) greater risk for type 2 diabetes respectively than group of <40 years. Overall, patients under stressful condition are more likely to develop T2DM than that of nonstressed respondent ($p = .000$). Moreover, when stress is divided into two groups- low stress and high stress, we found that both males ($p = .000$) and females ($p = .000$) with high stress were at high risk of diabetes mellitus, whereas the association between low stress and T2DM incidence was significant only among males (Male: $p = .002$; Female: $p = .115$). The distribution and association of the genotypes, age, and stress with T2DM have been summarized in Table 3 and Figure 3. There was no difference in T2DM incidence between CT ($p = .030$) and TT/CC ($p = .034$) genotype containing people who were in age group of 40–60 years (Table 3). In contrast, people who were more than 60 years old with CT genotype (OR = 4.636, $p = .029$) were more prone to T2DM than that of TT/CC genotype (OR = 3.714, $p = .007$) subjects (Table 3).

When we studied cumulative effect of the polymorphism with stress and gender on the onset of T2DM, it was found that female with CT (OR = 2.741, $p = .099$) or TT/CC (OR = 2.677, $p = .006$) genotype showed almost same level of risk, albeit relation with CT genotype was statistically insignificant. However, male with CT genotype (OR = 10.971, $p = .000$) were at significantly higher risk than TT/CC genotype (OR = 2.861, $p = .027$) bearing male. Moreover, sex independent analysis of all patients with CT genotype (OR = 5.324, $p = .000$) under stress was at significantly more risk to T2DM in comparison to TT/CC genotype (OR = 2.747, $p = .000$).

4 | DISCUSSION

The rapidly increasing number of diabetic patients becomes a global burden especially for health sector in low- and middle-income countries including Bangladesh (Bleich, Koehlmoos, Rashid, Peters, & Anderson, 2011). Many reasons such as obesity, lack of physical activity, food habit, sedentary job nature and genetic makeup are factors accounting for developing diabetes (Lyssenko & Laakso, 2013; Vilchis-Gil, Galván-Portillo, Klünder-Klünder, Cruz, & Flores-Huerta, 2015). Another cause is stress, which plays important role in the etiology of T2DM (S. J. Kelly & Ismail, 2015; Pouwer et al., 2010). Type 2 diabetic patients not only have to cope with this chronic disease, they are also at increased risk for several diseases like coronary heart disease, peripheral vascular

TABLE 2 Multivariate logistic regression for association of SNP (rs2227956) with Type 2 diabetes

Catagories	Genotype/ Allele	Unadjusted OR (95% CI); p-value	P-value	Adjusted OR (95% CI);p-value ^a	P-value
All Subjects	CC	2.08 (0.72–6.00); 0.175	0.031	1.22 (0.40–3.68); 0.725	0.0508
	CT	1.83 (1.13–2.96); 0.015		1.91 (1.14–3.22); 0.015	
	TT	Ref		Ref	
	C allele	1.61(1.1–2.4); 0.016	–	–	–
Male	CC	0.74 (0.10–5.57); 0.774	0.917	0.45 (0.05–3.65); 0.452	0.739
	CT	1.10 (0.54–2.25); 0.798		1.02 (0.46–2.29); 0.958	
	TT	Ref		Ref	
	C allele	1.00 (0.54–1.7); 0.880	–	–	–
Female	CC	2.97 (0.80–11.07); 0.105	0.005	1.96 (0.50–7.60); 0.332	0.009
	CT	2.92 (1.47–5.81); 0.002		3.03 (1.48–6.20); 0.002	
	TT	Ref		Ref	
	C allele	2.4 (1.4–4.1); 0.001	-	–	–


^aAdjusted for confounding factors (Age, Sex, stress)**TABLE 3** Association of HSP70-hom (+2,437) genotypes with age and stress

Status			Coefficient	OR	95% CI	p Value	
Diabetic patients	CT (n = 122)	Age, ≥61	1.534	4.636	1.171– 18.363	0.029	
		Age, 40–60	0.941	2.564	1.094– 6.010	0.030	
		Age, ≤39	Reference				
	TT/CC (n = 220)	Age, ≥61	1.312	3.714	1.439– 9.590	0.007	
		Age, 40–60	0.717	2.048	1.054– 3.978	0.034	
		Age, ≤39	Reference				
*Corresponding control is the reference category							
Diabetic patients	Male	Stress	2.395	10.971	2.959– 40.679	0.000	
		CT	Nonstress	Reference			
	Female	Stress	1.008	2.741	0.826– 9.091	0.099	
		CT	Nonstress	Reference			
	All Patients	Stress	1.672	5.324	2.271– 12.480	0.000	
		CT	Nonstress	Reference			
	Male	Stress	1.051	2.861	1.128– 7.257	0.027	
		TT/CC	Nonstress	Reference			
	Female	Stress	0.985	2.677	1.322– 5.421	0.006	
		TT/CC	Nonstress	Reference			
	All Patients	Stress	1.01	2.747	1.575– 4.972	0.000	
		TT/CC	Nonstress	Reference			
	*Corresponding control is the reference category						

disease, retinopathy, nephropathy, and neuropathy (Pouwer et al., 2010). To fight and control T2DM we have to seek out an alternative way of diagnosis and treatment based on patient's genetic information. This requires a deep insight into the etiology of this disease including associated single nucleotide polymorphism (SNP).

Various stresses trigger secretion of cortisol and norepinephrine which in turn raise blood glucose level by stimulating hepatic gluconeogenesis and insulin resistance (Bleich et al., 2011; Knol et al., 2006; Kumari et al., 2004; Norberg et al., 2007; Rod et al., 2009; Toshihiro et al., 2008; Zhang et al., 2006). It has been shown in the current study that highly

FIGURE 3 The distribution and association of SNP and stress on T2DM incidence



stressful life leading women experienced $1.25 \times$ higher T2DM prognosis than that of men, whereas women were in $1.26 \times$ lower risk to diabetic mellitus incidence than that of men who were in low level of stress (Table S4).

Age also plays a vital role in the onset of diabetes (Cowie & Eberhardt, 1995). In south-east Asia almost 97% diabetic patients are 40 years old or more (IDF Atlas, 2017). In Bangladesh, the reported age of diabetes is ≥ 40 years in 71% urban and 85% rural female, while in the case of male the proportion is 85.5% urban and 86.5% in rural population (IDF Atlas, 2017). The current study also pinpointed an exponential increase in the risk of onset of T2DM with the increase of age when 40 years was chosen as the reference (Table S4).

Among the genetic factors particularly SNPs alleged for diabetics, *HSP70-hom* + 2437T/C was targeted for the current study. However, there are contrasting association studies on this SNP (Buraczynska, Swatowski, Buraczynska, Dragan, & Ksiazek, 2009; Mir, Pugazhendhi, Paul, Nair, & Ramakrishna, 2009; Pociot et al., 1993). Therefore, before the analysis of the blood samples, we confirmed its suitability as a candidate SNP marker for T2DM by building 3D models for wild and mutant type proteins (Figure 2). The de-stability of structural complexes due to free energy change of the built models indicated significant change in the structure of the protein that can hamper its natural

function. Furthermore, active site analysis confirmed the structural variation and drug ineffectiveness against mutant model as active site amino acid Thr-493 does not interact with the drug (Figure 1).

In the purpose to identify diagnostic biomarker and prognostic significance, we analyzed the distribution of CC, CT and TT and the frequency of C and T allele among the study population. All the analysis done in this report was stratified according to gender. The genotypic distributions of three genotypes (TT, CT and CC) showed a good congruence with Hardy–Weinberg law, implying the consistent flow of these genotypes in Bangladeshi population including male and female groups. The χ^2 values calculated for all genotypes of target SNP were satisfactory, even in patient and control subjects of the study population (Table S1).

The presence of CT genotypes increased the relative risk for T2DM by 1.91 fold among all the patients. In contrast, the ratio of risk to get T2DM rose to $3.03 \times$ in the case of females who had the CT genotype. However, the male patients with the same genotype showed no association (Table 2). Interestingly, a study conducted in India had found the highest frequency for TT with significant p value where CT was less significant (Umapathy et al., 2012). Likewise, two studies conducted among the South and North Indian population found the association of diabetic foot ulcer with TT genotype and T allele (Umapathy et al., 2012; Zubair

& Ahmad, 2018). However, another research on diabetic foot ulcer revealed no significant association with the polymorphism among Tamil Nadu populace of India (Mir et al., 2009). Our results are found to be consistent with an earlier report which revealed that T allele conferred a significant protection for Diabetic macroalbuminuria subjects (Dhamodharan et al., 2017). This variation between two adjacent populations indicated the transformation of ancestral T allele to C allele in the Bangladeshi population and this incorporation of C allele might have some effect on disease prognosis. The impact of the C allele over the T allele was also determined by odd ratio estimation and allele frequency analysis. The risk estimation analysis for allele variation yielded a similar result as assumed previously from genotypic-association analysis. The association of the C allele was observed in patients compared to the control group. The most significant association of the C allele was found for female patients indicating that the C allele might be a potential risk factor for T2DM development especially in female. But the OR and p value could not produce the considerable signal in case of male patients (Table 2). When we pursued the cumulative effect of genotypes with stress and age on T2DM patients separately, it revealed that among the female patients, the effect of stress on T2DM incidence was genotype independent (Table 3) despite the fact that the CT genotype increased risk for T2DM independently (Table 2). On the other hand, existence of the CT genotype in male patients raised the probability of getting T2DM by about 4 × more than that of CC/TT genotypes presence (Table 3), though the CT genotype did not have an effect on male T2DM patients individually (Table 2). In the same way, the effect of age on T2DM prevalence was almost independent to genotype (Table 3), despite a gradual rise in T2DM incidence with the increasing age of patients (Table S4).

From this analysis, a strong association of C allele or the CT genotype with T2DM is evident. However, when we compared CC with TT, we found the values of the odds ratio (OR) and relative risk (RR) were maximum though the CC genotype was not significantly present in the Bangladeshi people. At the same time, we found that the OR and RR values for the CC/CT ratio were close to 1, suggesting that only one C allele in genotype is enough for exerting its effect on developing type 2 diabetes mellitus (Table S3).

In conclusion, our study shows that the CT genotype of *HSP70-hom +2,437 C/T* polymorphism represents a stress-independent risk factor for T2DM incidence among women. This study also suggests that under stressful condition, males are susceptible to T2DM in the presence of the CT genotype (Figure 3). To our knowledge, it is the first study that showed the cumulative effect of stress and rs2227956 polymorphism on T2DM onset. To generalize the association as the potential biomarker, the findings should be replicated

through more studies with a large number of samples and in different populations.

ACKNOWLEDGMENT

The Authors are highly grateful to the ministry of science and technology (MOST), Government of the People's Republic of Bangladesh for their extended support to National Institute of Biotechnology (NIB) during this work.

CONFLICT OF INTEREST

The authors declare that they have no competing interest.

AUTHOR CONTRIBUTION

SH contributed to conception, study design, and sample collection of the project. MM contributed to the sample collection, data producing, data analysis, interpreting the findings, manuscript writing, reviewing, and editing of the study. IA contributed to the sample collection, data producing and analysis and draft preparation of manuscript. MUH performed bioinformatics analysis, collected sample and produced data. MSAM and SB collected sample and produced data. MHR, PKS, and KCD contributed to sample collection and reviewing the manuscript. MS contributed to conception and critically reviewed the manuscript.

ORCID

Md. Moniruzzaman  <https://orcid.org/0000-0002-2700-8139>

U.S. Mahzabin Amin  <https://orcid.org/0000-0002-6664-6432>

REFERENCES

- Bleich, S. N., Koehlmoos, T. L., Rashid, M., Peters, D. H., & Anderson, G. (2011). Noncommunicable chronic disease in Bangladesh: Overview of existing programs and priorities going forward. *Health Policy, 100*(2–3), 282–289. <https://doi.org/10.1016/j.healthpol.2010.09.004>
- Buraczynska, M., Swatowski, A., Buraczynska, K., Dragan, M., & Ksiazek, A. (2009). Heat-shock protein gene polymorphisms and the risk of nephropathy in patients with type 2 diabetes. *Clinical Science, 116*(1), 81–86.
- Charan, J., & Biswas, T. (2013). How to calculate sample size for different study designs in medical research? *Indian Journal of Psychological Medicine, 35*(2), 121. <https://doi.org/10.4103/0253-7176.116232>
- Chouchane, L., Danguir, J., Beji, C., Bouassida, K., Camoin, L., Sfar, H., ... Strosberg, A. D. (2001). Genetic variation in the stress protein hsp70-2 gene is highly associated with obesity. *International Journal of Obesity, 25*(4), 462. <https://doi.org/10.1038/sj.ijo.0801545>
- Consortium, U (2016). UniProt: The universal protein knowledgebase. *Nucleic Acids Research, 45*(D1), D158–D169.
- Cowie, C. C., & Eberhardt, M. S. (1995). Sociodemographic characteristics of persons with diabetes. *Diabetes in America, 2*, 85–116.
- Dhamodharan, U., Ezhilarasi, K., Ponjayanthi, B., Sireesh, D., Ramkumar, K. M., & Viswanathan, V. (2017). Association of A1538G and C2437T

- single nucleotide polymorphisms in heat shock protein-70 genes with diabetic nephropathy among South Indian population. *Bioscience Reports*, 37(2), BSR20160605. <https://doi.org/10.1042/BSR20160605>
- Favatiere, F., Bornman, L., Hightower, L. E., Günther, E., & Polla, B. S. (1997). Variation in hsp gene expression and Hsp polymorphism: Do they contribute to differential disease susceptibility and stress tolerance? *Cell Stress & Chaperones*, 2(3), 141. [https://doi.org/10.1379/1466-1268\(1997\)002<0141:VIHGEA>2.3.CO;2](https://doi.org/10.1379/1466-1268(1997)002<0141:VIHGEA>2.3.CO;2)
- Gehrmann, M., Marienhagen, J., Eichholtz-Wirth, H., Fritz, E., Ellwart, J., Jäättelä, M., ... Multhoff, G. (2005). Dual function of membrane-bound heat shock protein 70 (Hsp70), Bag-4, and Hsp40: Protection against radiation-induced effects and target structure for natural killer cells. *Cell Death and Differentiation*, 12(1), 38. <https://doi.org/10.1038/sj.cdd.4401510>
- Ghanayem, N. M., Khamis, S. S. A., El-Shafie, M. K., El-Ghobashi, Y. A. E. S., El-Shazly, R. M. A. M., & Habib, M. S. E. D. M. (2014). The potential association of heat shock protein 70-2+1267 A/G gene polymorphism with nephropathy in type II diabetic patients. *Menoufia Medical Journal*, 27(3), 589. <https://doi.org/10.4103/1110-2098.145521>
- Hamosh, A., Scott, A. F., Amberger, J. S., Bocchini, C. A., & McKusick, V. A. (2005). Online mendelian inheritance in man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Research*, 33(suppl_1), D514–D517.
- He, L., Deng, T., & Luo, H.-S. (2014). Heat shock protein 70 gene polymorphisms and cancer risk: A meta-analysis. *The Scientific World Journal*, 2014, 1–9.
- IDF Diabetes Atlas Eighth Edition. (2017). <https://www.idf.org/e-library/epidemiology-research/diabetesatlas/134-idf-diabetes-atlas-8th-edition.html> [accessed 18 February 2018].
- IDF Diabetes Atlas Seventh Edition. (2015). <https://www.idf.org/e-library/epidemiology-research/diabetes-atlas/13-diabetes-atlas-seventh-edition.html> [accessed 02 February 2018].
- Kelly, K. (2002). Stress response proteins and renal ischemia. *Minerva urologica e nefrologica*. *The Italian Journal of Urology and Nephrology*, 54(2), 81–91.
- Kelly, S. J., & Ismail, M. (2015). Stress and type 2 diabetes: A review of how stress contributes to the development of type 2 diabetes. *Annual Review of Public Health*, 36, 441–462. <https://doi.org/10.1146/annurev-publhealth-031914-122921>
- Knol, M., Twisk, J. W., Beekman, A. T., Heine, R., Snoek, F. J., & Pouwer, F. (2006). Depression as a risk factor for the onset of type 2 diabetes mellitus. *A meta-analysis*. *Diabetologia*, 49(5), 837.
- Kumari, M., Head, J., & Marmot, M. (2004). Prospective study of social and other risk factors for incidence of type 2 diabetes in the Whitehall II study. *Archives of Internal Medicine*, 164(17), 1873–1880. <https://doi.org/10.1001/archinte.164.17.1873>
- Kuszewski, J., Gronenborn, A. M., & Clore, G. M. (1997). Improvements and extensions in the conformational database potential for the refinement of NMR and X-ray structures of proteins and nucleic acids. *Journal of Magnetic Resonance*, 125(1), 171–177. <https://doi.org/10.1006/jmre.1997.1116>
- Laskowski, R. A., MacArthur, M. W., Moss, D. S., & Thornton, J. M. (1993). PROCHECK: A program to check the stereochemical quality of protein structures. *Journal of Applied Crystallography*, 26(2), 283–291. <https://doi.org/10.1107/S0021889892009944>
- Lindquist, S. (1986). The heat-shock response. *Annual Review of Biochemistry*, 55(1), 1151–1191. <https://doi.org/10.1146/annurev.bi.55.070186.005443>
- Lüthy, R., Bowie, J. U., & Eisenberg, D. (1992). Assessment of protein models with three-dimensional profiles. *Nature*, 356(6364), 83. <https://doi.org/10.1038/356083a0>
- Lyssenko, V., & Laakso, M. (2013). Genetic screening for the risk of type 2 diabetes: Worthless or valuable? *Diabetes Care*, 36(Supplement 2), S120–S126. <https://doi.org/10.2337/dcS13-2009>
- Macario, A., & De Macario, E. C. (2000). Stress and molecular chaperones in disease. *International Journal of Clinical and Laboratory Research*, 30(2), 49–66. <https://doi.org/10.1007/s005990070016>
- Meites, S., & Saniel-Banrey, K. (1973). Modified glucose oxidase method for determination of glucose in whole blood. *Clinical Chemistry*, 19(3), 308–311.
- Mir, K., Pugazhendhi, S., Paul, M., Nair, A., & Ramakrishna, B. (2009). Heat-shock protein 70 gene polymorphism is associated with the severity of diabetic foot ulcer and the outcome of surgical treatment. *British Journal of Surgery: Incorporating European Journal of Surgery and Swiss Surgery*, 96(10), 1205–1209. <https://doi.org/10.1002/bjs.6689>
- Mooney, S. (2005). Bioinformatics approaches and resources for single nucleotide polymorphism functional analysis. *Briefings in Bioinformatics*, 6(1), 44–56. <https://doi.org/10.1093/bib/6.1.44>
- Norberg, M., Stenlund, H., Lindahl, B., Andersson, C., Eriksson, J. W., & Weinehall, L. (2007). Work stress and low emotional support is associated with increased risk of future type 2 diabetes in women. *Diabetes Research and Clinical Practice*, 76(3), 368–377. <https://doi.org/10.1016/j.diabres.2006.09.002>
- Pociot, F., Rønningen, K., & Nerup, J. (1993). Polymorphic analysis of the human MHC-linked heat shock protein 70 (HSP70-2) and HSP70-Hom genes in insulin-dependent diabetes mellitus (IDDM). *Scandinavian Journal of Immunology*, 38(5), 491–495. <https://doi.org/10.1111/j.1365-3083.1993.tb02593.x>
- Pouwer, F., Kupper, N., & Adriaanse, M. C. (2010). Does emotional stress cause type 2 diabetes mellitus? A review from the European Depression in Diabetes (EDID) Research Consortium. *Discovery Medicine*, 9(45), 112–118.
- Rahman, M. S., Akter, S., Abe, S. K., Islam, M. R., Mondal, M. N. I., Rahman, J. S., & Rahman, M. M. (2015). Awareness, treatment, and control of diabetes in Bangladesh: A nationwide population-based study. *PLoS ONE*, 10(2), e0118365. <https://doi.org/10.1371/journal.pone.0118365>
- Rod, N. H., Grønbaek, M., Schnohr, P., Prescott, E., & Kristensen, T. (2009). Perceived stress as a risk factor for changes in health behaviour and cardiac risk profile: A longitudinal study. *Journal of Internal Medicine*, 266(5), 467–475. <https://doi.org/10.1111/j.1365-2796.2009.02124.x>
- Rose, P. W., Beran, B., Bi, C., Bluhm, W. F., Dimitropoulos, D., Goodsell, D. S., ... Westbrook, J. D. (2010). The RCSB protein data bank: Redesigned web site and web services. *Nucleic Acids Research*, 39(suppl_1), D392–D401.
- Šali, A., Potterton, L., Yuan, F., van Vlijmen, H., & Karplus, M. (1995). Evaluation of comparative protein modeling by MODELLER. *Proteins: Structure, Function, and Bioinformatics*, 23(3), 318–326. <https://doi.org/10.1002/prot.340230306>
- Sherry, S. T., Ward, M.-H., Kholodov, M., Baker, J., Phan, L., Smigielski, E. M., & Sirotkin, K. (2001). dbSNP: The NCBI database of genetic variation. *Nucleic Acids Research*, 29(1), 308–311. <https://doi.org/10.1093/nar/29.1.308>
- Singh, R., Kølvråa, S., Bross, P., Jensen, U. B., Gregersen, N., Tan, Q., ... Rattan, S. I. S. (2006). Reduced heat shock response in human mononuclear cells during aging and its association with

- polymorphisms in HSP70 genes. *Cell Stress & Chaperones*, 11(3), 208. <https://doi.org/10.1379/CSC-184R.1>
- Söding, J., Biegert, A., & Lupas, A. N. (2005). The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Research*, 33(suppl_2), W244–W248. <https://doi.org/10.1093/nar/gki408>
- Toshihiro, Á., Saito, K., Takikawa, S., Takebe, N., Onoda, T., & Satoh, J. (2008). Psychosocial factors are independent risk factors for the development of Type 2 diabetes in Japanese workers with impaired fasting glucose and/or impaired glucose tolerance 1. *Diabetic Medicine*, 25(10), 1211–1217.
- Umapathy, D., Krishnamoorthy, E., & Muthukumar, P. (2012). Is HSP70-hom (C2437T) single nucleotide polymorphism (SNP) associated with diabetic foot ulcer (DFU) among South Indian population? *Journal of Diabetic. Foot Complications*, 4, 57–62.
- Umapathy, D., Krishnamoorthy, E., Muthukumar, P., Rajaram, R., Padmalayam, I., & Viswanathan, V. (2012). Association of A1538G and C2437T single nucleotide polymorphisms in heat shock protein 70 genes with type 2 diabetes. *Laboratory Medicine*, 43(6), 250–255.
- Vilchis-Gil, J., Galván-Portillo, M., Klünder-Klünder, M., Cruz, M., & Flores-Huerta, S. (2015). Food habits, physical activities and sedentary lifestyles of eutrophic and obese school children: A case-control study. *BMC Public Health*, 15(1), 124. <https://doi.org/10.1186/s12889-015-1491-1>
- Zhang, J., Niaura, R., Dyer, J. R., Shen, B.-J., Todaro, J. F., McCaffery, J. M., ... Ward, K. D. (2006). Hostility and urine norepinephrine interact to predict insulin resistance: The VA normative aging study. *Psychosomatic Medicine*, 68(5), 718–726. <https://doi.org/10.1097/01.psy.0000228343.89466.11>
- Zubair, M., & Ahmad, J. (2018). Heat shock protein 70 gene single nucleotide polymorphism and diabetic foot ulcer. Is there any relationship? *Journal of Clinical Medicine*, 7(8), 187.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Moniruzzaman Md, Ahmed I, Huq S, et al. Association of polymorphism in heat shock protein 70 genes with type 2 diabetes in Bangladeshi population. *Mol Genet Genomic Med.* 2020;8:e1073. <https://doi.org/10.1002/mgg3.1073>