

Research article

Open Access

Cytosolic phospholipase A₂: a member of the signalling pathway of a new G protein α subunit in *Sporothrix schenckii*

Shirley Valentín-Berriós¹, Waleska González-Velázquez², Lizaida Pérez-Sánchez², Ricardo González-Méndez³ and Nuri Rodríguez-del Valle^{*2}

Address: ¹Department of Microbiology, San Juan Bautista School of Medicine, Box 4968, Caguas, PR 00726-4968, USA, ²Department of Microbiology and Medical Zoology, Medical Sciences Campus, University of Puerto Rico, PO Box 365067, San Juan, PR 00936-5067, USA and ³Department of Radiological Sciences, Medical Sciences Campus, University of Puerto Rico, PO Box 365067, San Juan, PR 00936-5067, USA

Email: Shirley Valentín-Berriós - svalentin@sanjuanbautista.edu; Waleska González-Velázquez - waleska.gonzalez@upr.edu; Lizaida Pérez-Sánchez - lizaida.perez@upr.edu; Ricardo González-Méndez - ricardo.gonzalez7@upr.edu; Nuri Rodríguez-del Valle* - nuri.rodriguez@upr.edu

* Corresponding author

Published: 19 May 2009

Received: 16 December 2008

BMC Microbiology 2009, 9:100 doi:10.1186/1471-2180-9-100

Accepted: 19 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2180/9/100>

© 2009 Valentín-Berriós et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Sporothrix schenckii* is a pathogenic dimorphic fungus, the etiological agent of sporotrichosis, a lymphocutaneous disease that can remain localized or can disseminate, involving joints, lungs, and the central nervous system. Pathogenic fungi use signal transduction pathways to rapidly adapt to changing environmental conditions and *S. schenckii* is no exception. *S. schenckii* yeast cells, either proliferate (yeast cell cycle) or engage in a developmental program that includes proliferation accompanied by morphogenesis (yeast to mycelium transition) depending on the environmental conditions. The principal intracellular receptors of environmental signals are the heterotrimeric G proteins, suggesting their involvement in fungal dimorphism and pathogenicity. Identifying these G proteins in fungi and their involvement in protein-protein interactions will help determine their role in signal transduction pathways.

Results: In this work we describe a new G protein α subunit gene in *S. schenckii*, *ssg-2*. The cDNA sequence of *ssg-2* revealed a predicted open reading frame of 1,065 nucleotides encoding a 355 amino acids protein with a molecular weight of 40.9 kDa. When used as bait in a yeast two-hybrid assay, a cytoplasmic phospholipase A₂ catalytic subunit was identified as interacting with SSG-2. The *sspla₂* gene, revealed an open reading frame of 2538 bp and encoded an 846 amino acid protein with a calculated molecular weight of 92.62 kDa. The principal features that characterize cPLA₂ were identified in this enzyme such as a phospholipase catalytic domain and the characteristic invariable arginine and serine residues. A role for SSPLA₂ in the control of dimorphism in *S. schenckii* is suggested by observing the effects of inhibitors of the enzyme on the yeast cell cycle and the yeast to mycelium transition in this fungus. Phospholipase A₂ inhibitors such as AACOCF₃ (an analogue of archidonic acid) and isotetrandrine (an inhibitor of G protein PLA₂ interactions) were found to inhibit budding by yeasts induced to re-enter the yeast cell cycle and to stimulate the yeast to mycelium transition showing that this enzyme is necessary for the yeast cell cycle.

Conclusion: A new G protein α subunit gene was characterized in *S. schenckii* and protein-protein interactions studies revealed this G protein alpha subunit interacts with a cPLA₂ homologue. The PLA₂ homologue reported here is the first phospholipase identified in *S. schenckii* and the first time a PLA₂ homologue is identified as interacting with a G protein α subunit in a pathogenic dimorphic fungus, establishing a relationship between these G proteins and the pathogenic potential of fungi. This cPLA₂ homologue is known to play a role in signal transduction and fungal pathogenesis. Using cPLA₂ inhibitors, this enzyme was found to affect dimorphism in *S. schenckii* and was found to be necessary for the development of the yeast or pathogenic form of the fungus.

Background

Sporothrix schenckii is a dimorphic fungus that produces lymphocutaneous lesions in humans and animals. It is the etiologic agent of sporotrichosis, a subcutaneous lymphatic mycosis with a worldwide distribution [1]. In its saprophytic form it develops hyaline, regularly septated hyphae and pyriform conidia which can be found single or in groups in a characteristic daisy-like arrangement. The yeast or parasitic form shows ovoid cells with single or multiple budding.

In *S. schenckii*, dimorphism is both a proliferative and morphogenetic process. We have reported that in response to different environmental stimuli, *S. schenckii* unbudded synchronized yeast cells, either proliferate (yeast cell cycle) or engage in a developmental program that includes proliferation accompanied by morphogenesis (yeast to mycelium transition). Dimorphism in *S. schenckii*, depends on transmembrane signalling pathways that respond to cell density [2,3], external pH [2,3], cyclic nucleotides [4] and extracellular calcium concentration [5].

Dimorphism is an adaptation response to changing environmental conditions. The morphology displayed by dimorphic fungi is probably the result of the stimulation of membrane receptors by extracellular ligands. Heterotrimeric ($\alpha\beta\gamma$) guanine nucleotide binding proteins have been associated with membrane receptors and with morphogenetic transition signalling in many eukaryotes, and play a crucial role in fungal morphogenesis as well [6]. They constitute a family of GTP hydrolases involved in signal transduction pathways. These proteins are coupled to membrane receptors (GPCR) that recognize different extracellular signals. The α subunits of the heterotrimeric G proteins bind GTP. The interaction of a ligand with the GPCR initiates the exchange of bound GDP for GTP in the α subunit resulting in the dissociation of the heterotrimer into α -GTP and $\beta\gamma$ subunits. The dissociated α -GTP subunit and the $\beta\gamma$ dimer, relay signals to different targets resulting in changes in cytoplasmic ionic composition or in second messenger levels (e.g., cAMP) that ultimately lead to a cellular response [7-10].

Genes encoding proteins that are similar to the $G\alpha$ class of the heterotrimeric G proteins have been described in filamentous fungi such as *Aspergillus nidulans* [11] and *Neurospora crassa* [12-14], as well as in fungal plant pathogens like *Cryphonectria parasitica* [15,16], *Ustilago maydis* [17] and *Magnaporthe grisea* [18], among others. In *S. schenckii*, a 41 kDa $G\alpha$ subunit homologous to the $G\alpha_i$ subunit and sensitive to inhibition by pertussis toxin was described previously by us [19]. This was the first $G\alpha_i$ subunit described in a pathogenic dimorphic fungus.

In higher eukaryotes, members of the $G\alpha$ class are known to regulate adenylate cyclase [20], cGMP phosphodiesterase [21], phosphoinositide-3-kinase [22], calcium and potassium channels [22-24], and the activity of phospholipases [9,25-28]. In fungi, $G\alpha$ subunits have been shown to regulate adenylate cyclase, morphogenesis and pathogenicity [6,14,29,30]. Most of the studies related to determining the role of the heterotrimeric G protein subunits in fungi involved the observation of the morphological effects produced in the fungus when these genes are deleted [6,12,14,18]. Nevertheless, the full scope of the processes that $G\alpha$ subunits regulate in fungi is still not known and interactions between these subunits and cellular proteins have seldom been reported in pathogenic fungi.

A large number of G protein coupled receptors have been observed to induce activation of phospholipase A_2 in higher eukaryotic systems [31]. The PLA_2 superfamily can be classified according to cellular location or biological properties [32]. The phospholipase A superfamily includes the calcium dependent-secretory PLA_2 (s PLA_2), the calcium independent-intracellular PLA_2 (i PLA_2) and the cytosolic PLA_2 (c PLA_2). They differ in terms of calcium requirements, substrate specificity, molecular weight and lipid modification. The s PLA_2 is usually a 13 to 15 kDa protein while the c PLA_2 is a 85 kDa protein in human macrophages. The c PLA_2 possesses characteristics that suggest that it is associated to receptor-activated signal transduction cascades [33]. This PLA_2 is known to translocate to the membrane in response to an increase in intracellular calcium concentration [34]. Cytosolic PLA_2 hydrolyses the *sn*-2 position of phospholipids, resulting in the release of lysophospholipids and free fatty acids. The most commonly released fatty acid is arachidonic acid, which in turn is converted to eicosanoids that regulate multiple processes including calcium channels, mitogenic signals and most important, the inflammatory response of macrophages [31,32,35,36].

The present study was undertaken to identify the presence of and characterize additional $G\alpha$ subunits in *S. schenckii*, to identify any important interacting partners of the new $G\alpha$ subunit, and finally to determine the involvement if any of the interacting protein, in this case c PLA_2 , in the control of dimorphism in this fungus. Here we give details of the identification and sequencing of the *ssg-2* gene, including gene organization, the presence and position of introns, derived amino acid sequence and conserved polypeptide-encoded domains. Using SSG-2 as bait, we identified a c PLA_2 homologue interacting with this G protein α subunit. We give the genomic sequence of this gene and the complete derived amino acid sequence. We also report the effects on the yeast to mycelium transition and

the yeast cell cycle of phospholipase effectors in *S. schenckii*.

This work constitutes the first report of the presence of multiple G protein α subunits in *S. schenckii*, the presence of a cPLA₂ homologue interacting with this G protein α subunit, and the involvement of cPLA₂ in the control of dimorphism in *S. schenckii*. In addition to being a very important determinant of pathogenicity in fungi and other organisms, cPLA₂ is shown to have a direct effect in the control of dimorphism in this fungus. This information will ultimately help us construct the signal transduction pathway leading from the G proteins onward and the role of G proteins and its interacting partners in fungal pathogenesis.

Results

Identification of the *ssg-2* gene

Most fungal G α subunit genes vary only slightly in size within the region encoding the GESGKST and KWIHCF motifs where primers for PCR are usually made because of the conserved nature of these regions. In the region comprised between these primers size variations are usually due to the presence of introns of slightly different sizes. Two PCR products were obtained when using fungal DNA as template and the GESGKST/KWIHCF primer pair one belonging to *ssg-1* and the other to *ssg-2* of approximately 620 and 645 bp, respectively. The *ssg-2* PCR product (645 bp) established the presence of a new gene encoding another G α subunit in *S. schenckii*. Figure 1A shows the sequencing strategy used for the identification of this new G protein α subunit gene. Once the coding sequence was completed, it was confirmed using yeast cDNA as template and the MGACMS/KDSGIL primer pair. A 1,065 bp ORF was obtained, containing the coding region of the *ssg-2* cDNA as shown in Figure 1B. Using the same primer pair and genomic DNA as template a 1,333 bp PCR product was obtained. Sequencing of this PCR product confirmed the sequences obtained previously and showed the presence and position of 4 introns. These introns had the consensus GT/AG junction splice site and interrupted the respective codons after the second nucleotide. The first intron interrupted the codon for G42 and consisted of 82 bp, the second intron interrupted the codon for Y157 and consisted of 60 bp, the third intron interrupted the codon for H200 and consisted of 60 bp, the fourth intron starts interrupted the codon H323 and consisted of 67 bp. With the exception of the regions where introns were present in the genomic sequence of the *ssg-2* gene, the cDNA sequence and genomic sequence were identical. The overlapping of these two sequences confirmed the presence of the introns in the genomic sequence. The cDNA and genomic sequence of *ssg-2* have GenBank accession numbers [AF454862](#) and [AY078408](#), respectively.

Bioinformatic characterization of SSG-2

The derived amino acid sequence (GenBank accession number [AAL57853](#)) revealed a G α subunit of 355 amino acids as shown in Figure 1B. The calculated molecular weight of the *ssg-2* gene product was 40.90 kDa. Blocks analysis of the amino acid sequence of SSG-2 revealed a G-protein alpha subunit signature from amino acids 37 to 276 with an E value of $5.2e^{-67}$ and a fungal G-protein alpha subunit signature from amino acids 61 to 341 with an E value of $3.3e^{-28}$ [37]. SSG-2 has the motifs encoding the GTPase domain [38] with the corresponding consensus sequences involved in GTP binding shaded in gray in Figure 1B. The phosphate binding loop which includes the sequence GXGXXGKS is found in SSG-2 as GSGESGKS. The magnesium binding residues with the consensus sequence DXXG is present as DVGG in SSG-2, while the guanine ring binding sites are those with the consensus sequence NKXD is present as NKVD. The TXAT consensus sequence is present as TQAT in SSG-2. Another region involved in phosphate binding includes the consensus sequence RXXT that in SSG-2 is present as RTKT. In addition to these conserved domains, the protein derived from the *ssg-2* cDNA sequence has the N-terminal glycine that is myristoylated in G α subtypes and is needed for membrane association. The 5 residues that identify the adenylate cyclase interaction site according to BLAST analysis [39] are in red in Figure 1, these include I187, K212, I215, H216, and E 219. The putative receptor binding site includes amino acids L318 to R334 and is shown in blue letters in Figure 1[39].

The derived amino acid sequence alignment of SSG-2 to that of the several fungal homologues is shown in Figure 2. This figure shows more than 85% identity to MAGA of *M. grisea* [18], CPG-2 of *C. parasitica* [16] and GNA-3 of *N. crassa* [14]. Table 1 summarizes the percent identity of SSG-2 to some members of the fungal G α homologues and SSG-1.

Yeast two-hybrid screening

Two independent yeast two-hybrid screenings, using different *S. schenckii* yeast cells cDNA libraries were done with the complete coding sequence of SSG-2 as bait. In both screenings, 3 blue colonies growing in quadruple drop out (QDO) medium (SD/-Ade/-His/-Leu/-Trp/X- α -gal) were identified as containing the same PLA₂ homologue insert. The expression of the Ade⁺, His⁺ phenotypes and α galactosidase activity are considered by the manufacturer as corroborative of true interactions. The inserts from all three colonies were found to contain the carboxy-terminal residues of a protein homologous to PLA₂'s from *A. nidulans*. Our results indicated that the last 162 amino acids of the *S. schenckii* cPLA₂ homologue interacted with SSG-2.




Figure 1
cDNA and derived amino acid sequences of the *S. schenckii* *ssg-2* gene. Figure 1A shows the sequencing strategy used for *ssg-2*. The size and location in the gene of the various fragments obtained from PCR and RACE are shown. The black boxes indicate the size and relative position of the introns. Figure 1B shows the cDNA and derived amino acid sequence of the *ssg-2* gene. Non-coding regions are given in lower case letters, coding regions and amino acids are given in upper case letters. The sequences that make up the GTPase domain are shaded in gray, the five residues that identify the adenylate cyclase interaction site are given in red and the putative receptor binding site is shown in blue.




Figure 2
Amino acid sequence alignments of SSG-2 with other Gα subunit homologues. The predicted amino acid sequence of *S. schenckii* SSG-2 and SSG-1, *C. parasitica* CPG2, *N. crassa* GNA3, *R. necatrix* WGA1, *E. nidulans* GANB, and *M. grisea* MAGA were aligned as described in Methods. In the alignment, black shading with white letters indicates 100% identity, gray shading with white letters indicates 75–99% identity, gray shading with black letters indicates 50–74% identity.

Co-immunoprecipitation (Co-IP)

The SSG-2-SSPLA₂ interaction was corroborated by co-immunoprecipitation. Figure 3 shows the confirmation of the interaction observed in the yeast two-hybrid assay between SSG-2 and SSPLA₂ by co-immunoprecipitation and Western blot analysis. Lane 1 shows the band

obtained using anti-cMyc antibody that recognizes SSG-2. This band is of the expected size (62 kDa) considering that SSG-2 was expressed fused to the GAL-4 binding domain. The two high molecular weight bands present belong to the anti-cMyc antibodies used for precipitation. Lane 2 shows the results obtained in the Western blot when the

Table 1: Comparison of G protein alpha subunit homologues to SSG-2 of *S. schenckii*

UniProt AC	Name	Length	Organism Name	Overlap	%iden	E-value	Score
Q8TF91	SSG2	355	<i>Sporothrix schenckii</i>	355	100	0	729
O13314	MAGA	356	<i>Magnaporthe grisea</i>	355	88	0	642
Q00581	CPG2	355	<i>Cryphonectria parasitica</i>	355	87	0	640
Q9HFW7	GNA3	356	<i>Neurospora crassa</i>	356	85	e-177	623
Q9HFA3	WGA1	356	<i>Rosellinia necatrix</i>	355	84	e-175	619
Q9UVK8	GANB	356	<i>Emericella nidulans</i>	356	77	e-160	567
O74259	SSG1	353	<i>Sporothrix schenckii</i>	353	50	2e-93	346

SSG-1 is included as reference. Analysis was carried out using iProtClass database and the BLAST algorithm. Overlap refers to the number of residues used to determine SSG-2% identity when doing pairwise comparisons.




Figure 3
Western Blots results from SSG-2/SSPLA₂ co-immunoprecipitation. Whole cell free extracts of *S. cerevisiae* cells containing PGBKT7 and PGADT7 plasmids with the complete SSG-2 coding region fused to the GAL4 activation domain and cMyc, and the initial SSPLA₂ coding fragment identified in the yeast two-hybrid assay fused to the GAL4 DNA binding domain and HA, respectively, were co-immunoprecipitated as described in Methods. The co-precipitated proteins were separated using 10% SDS polyacrylamide electrophoresis and transferred to nitrocellulose. The nitrocellulose strips were probed with anti-cMyc antibodies (Lane 1) and anti HA antibodies (Lane 3). Lanes 2 and 4 are negative controls where no primary antibody was added. The antigen-antibody reactions were detected using the Immun-Star™ AP chemiluminescent protein detection system. Pre-stained molecular weight markers were included in outside lanes of the gel and also transferred to nitrocellulose, the position of the molecular weight markers is indicated in the figure.

primary anti-cMyc antibody was not added (negative control). Lane 3 shows the band obtained using anti-HA antibody that recognizes the original SSPLA₂ fragment isolated from the yeast two-hybrid clone. This band is of the expected size (35.9 kDa) considering that only the last 162 amino acids of the protein were present and that this fragment was fused to the GAL-4 activation domain. Lane 4 shows the results obtained in the Western blot when the primary anti-HA antibody was not added (negative control).

Sequencing of the *sspla₂* gene

Figure 4A shows the sequencing strategy used for the *sspla₂* gene. The DNA sequence of *sspla₂* gene was completed using genome walking and PCR. Figure 4B shows the genomic and derived amino acid sequence of the *sspla₂* homologue. The genomic sequence has 2648 bp with an open reading frame of 2538 bp encoding an 846 amino acid protein with a predicted molecular weight of 92.6 kDa. The GenBank numbers for the genomic and derived

amino acid sequence are [FJ357242.1](#) and [ACI04517.1](#), respectively.

Bioinformatic characterization of SSPLA₂

The PANTHER Classification System identified this protein as a member of the cytosolic phospholipase A2 family (PTHR10728) (residues 132–827) with an extremely significant E value of 6.4×10^{-97} [40]. BLAST analysis of the derived amino acid sequence of the *S. schenckii* SSPLA₂, showed a phospholipase domain extending from amino acids 177 to 750 [39]. Pfam analysis shows similar results, and in this domain the PLA₂ signature GXSG [G, S] (Pfam: Family PLA2_B PF 01735) is present as GVSGS in the active site (highlighted green in Figure 4B) [41,42]. The amino acids needed for catalytic activity R235, S263 and D553 are given in red in this same figure [43]. S263 is essential for the formation of arachidonyl serine needed for the transfer of the arachidonyl group to glycerol or to water. The amino acids D511 to L523, D583 to G595 and D738 to A750 (highlighted in yellow) comprise putative EF hand domains of the protein (76% identity, probability, 3.33×10^{-6}). In Figure 4B a putative calmodulin binding domain was identified from amino acids Q806 to L823 using the Calmodulin Target Database [44] and highlighted in gray. A serine protease, subtilase family, aspartic acid active site motif was identified using Scan Prosite with an E value of 5.283×10^{-7} from amino acids 549 to 559 and is shaded in blue green in Figure 4B[45]. This motif is characteristic of both yeast and fungal cPLA₂ homologues [43].

Figure 5 shows the multiple sequence alignment of the derived amino acid sequence of *S. schenckii* PLA₂ homologue to that of other PLA homologues or hypothetical proteins from *N. crassa*, *A. nidulans*, *M. grisea*, *Chaetomium globosum*, *Podospora anserina* and *Gibberella zeae*. This figure shows that the important domains are very similar, although variations occur in the N terminal and C terminal regions. The alignment shown includes only the catalytic domain, the complete alignment is given as additional material (Additional file 1).

Effects of PLA₂ effectors on the yeast to mycelium transition and the yeast cell cycle

S. schenckii is not a genetically manageable organism, therefore, effectors of PLA₂ were tested for their effects on the yeast to mycelium transition and the yeast cell cycle. Arachidonic acid is the primary product of cPLA₂ activity on phospholipids, while AACOCF₃ and isotetrandrine are inhibitors of PLA₂ activity. AACOCF₃ is a known competitive inhibitor of PLA₂ [46]. It is an analogue of arachidonic acid and presumably binds directly to the active site of the enzyme. It is a potent and selective inhibitor of cytosolic phospholipase A [46]. Isotetrandrine on the




Figure 4
Genomic and derived amino acid sequences of the *S. schenckii* *sspla*₂ gene. Figure 4A shows the sequencing strategy used for sequencing the *sspla*₂ gene. The size and location in the gene of the various fragments obtained from PCR and RACE are shown. Figure 4B shows the genomic and derived amino acid sequence of the *sspla*₂ gene. Non-coding regions are given in lower case letters, coding regions and amino acids are given in upper case letters. The invariant amino acids required for phospholipase activity are shown in red. The potential EF hands are shaded in yellow and the putative calmodulin binding domain is shaded in gray. The cPLA₂ signature motif is shaded in green and the serine proteases, subtilase family, aspartic acid active site motif is shaded in blue green.




Figure 5
Amino acid sequence alignments of SSPLA₂ with other PLA₂ homologues. The *S. schenckii* SSPLA₂ was aligned to other PLA₂ fungal homologues as described in Methods. The fungal PLA₂ used for the alignment were: *E. nidulans* (PLA₂), *M. grisea* (hypothetical protein), *N. crassa* (PLA₂), *C. globosum* (hypothetical protein), *P. anserina* (hypothetical protein) and *G. zeae* (PLA₂). The alignment was done using MCOFFEE and visualized using the program GeneDoc. Only the catalytic core of these proteins is shown in this alignment, from amino acids 192 to 611 (in reference to the multiple alignment position). The black shading with white letters indicates 100% identity, gray shading with white letters indicates 75–99% identity, gray shading with black letters indicates 50–74% identity.

other hand is an alkaloid that has been reported to interfere with G protein activation of PLA₂ [47]. Figure 6 shows the percentage of yeast cells forming germ tubes in the presence and absence of arachidonic acid, AACOCF₃ and isotetrandrine. This figure shows that these latter compounds significantly stimulated the yeast to mycelium transition at 6 and 9 h of incubation when the control cells are in the process of DNA synthesis and germ tube emergence [2]. The percent stimulation was approximately 68% and 33% at 6 h and 9 h of incubation in the presence of both AACOCF₃ and isotetrandrine. In the presence of arachidonic acid a slight (25%) non-significant inhibition was observed at 6 h of incubation. The

degree of stimulation caused by the addition of AACOCF₃ and isotetrandrine was similar even though the mechanism of action of these compounds is completely different.

Figure 7 shows the percentage of budding in yeast cells induced to re-enter the cell cycle in the presence and absence of arachidonic acid, AACOCF₃ and isotetrandrine. The percent inhibition observed in the presence of both AACOCF₃ and isotetrandrine was approximately 60% and 40% at 9 h of incubation, respectively. Arachidonic acid on the other hand significantly stimulated budding at 6 h of incubation (percent stimulation was




Figure 6
Effects of SSPLA₂ effectors on the yeast to mycelium transition. Yeast cells grown, harvested, synchronized and selected by filtration as described in Methods were induced to form germ tubes in a basal medium with glucose at pH 4.0 and incubated at 25°C in the presence and absence of arachidonic acid (40 μM), AACOCF3 (100 μM; Nonadeca-4,7,10,13-tetraenyl-trifluoro-methyl ketone) and isotretandrine (50 μM; 6,6',7,12-tetra methoxy-2,2'-dimethyl-berbaman). All values are given as the average percentage ± one SD of at least three independent experiments. The Student's t test was used to determine the statistical significance of the data at a 95% confidence level. Values that differ significantly from those of the control at 95% confidence level are marked with an asterisk.




Figure 7
Effects of SSPLA₂ effectors on the yeast budding cycle. Yeast cells grown, harvested, synchronized and selected by filtration as described in Methods were induced to re-enter the budding cycle in a basal medium with glucose at pH 7.2 and incubated at 25°C in the presence and absence of arachidonic acid (40 μM), AACOCF3 (100 μM; Nonadeca-4,7,10,13-tetraenyl-trifluoro-methyl ketone) and isotretandrine (50 μM; 6,6',7,12-tetra methoxy-2,2'-dimethyl-berbaman). All values are given as the average percentage ± one SD of at least three independent experiments. The Student's t test was used to determine the statistical significance of the data at a 95% confidence level. Values that differ significantly from those of the control at 95% confidence level are marked with an asterisk.

50%). At this time interval, control cells are initiating DNA synthesis [3].

Discussion

The heterotrimeric G protein family ranks among the most important protein families identified as intracellular recipients of external signalling. The present study was conducted in order to describe new Gα subunit encoding genes in *S. schenckii*, identify any important protein interacting with this G alpha subunit and determine the effects on dimorphism in *S. schenckii* of the protein or proteins identified.

The results presented here, together with our previous report [19] corroborate the existence of more than one heterotrimeric G protein α subunit gene in *S. schenckii*. Unpublished results indicate that this protein is one of at least 3 Gα subunits present in *S. schenckii*. In this sense, *S. schenckii* is behaving more like the filamentous fungi and plant pathogens such as *N. crassa* [14], *C. parasitica* [48] and *M. grisea* [18], where genes that encode 3 different Gα subunits similar to the Gα class of animals rather than to the GPA group present in yeasts and plants. Computational sequence and phylogenetic analysis of the Gα sub-

units in filamentous fungi shows the existence of 3 distinct subfamilies of G protein alpha subunits [19]. According to the classification offered by Li and collaborators, SSG-2 belongs to Group III of the fungal G protein alpha subunits [49]. The Group III considered by them to be Gas analogues because they positively influence cAMP levels although they have more sequence similarity to Gαi [49].

The nucleotide and amino acid sequence analysis of this new G protein α subunit gene are different from the previously identified *ssg-1* gene. The nucleotide conservation of the coding region of *ssg-2* is less than 50% when compared to that of the previously reported *ssg-1* gene, confirming that *ssg-1* and *ssg-2* are two different genes (data not shown). The derived amino acid sequence of *ssg-2* is 50% identical to that of SSG-1, but they have differences in the motifs that are characteristic of the G protein alpha subunits, the most important difference being that SSG-2 lacks the cysteine residue in domain 5 that characterizes the pertussis binding domain of SSG-1 (TCADT). For this reason, SSG-2 belongs to the Gα class but cannot be strictly considered a Gαi, even though it is 46% identical

to mammalian G α class members. This shows the high degree of conservation in G α subunits even among phylogenetically distant organisms.

The work done in order to identify the role of G α subunits in the filamentous fungi has been mainly concerned with the phenotypes observed when these genes are knocked-out (as reviewed by [6]). In this paper a different approach was used. We wanted to identify important protein-protein interactions between SSG-2 and the complex signalling system that regulates the flow of information from the environment through the heterotrimeric G proteins into the cell in *S. schenckii*. Using the yeast two-hybrid technique we identified a cPLA₂ homologue as interacting with SSG-2 in two independent experiments, using two different cDNA libraries. This SSG-2-PLA₂ interaction was also confirmed by co-immunoprecipitation. Up to date, protein-protein interactions of these G α subunits have not been reported in the pathogenic fungi, and the exact proteins with which these G α subunits interact have not been identified. This is the first report of a cytosolic PLA₂ homologue interacting with a G protein α subunit in a pathogenic dimorphic fungus, suggesting a functional relationship between these two important proteins. Other proteins interact with SSG-2 (unpublished results), but the SSG-2-PLA₂ interaction is very important as it connects this G protein α subunit with both pathogenicity and lipid signal transduction in fungi [50].

This PLA₂ homologue belongs to the Group IV PLA₂ family that has been highly conserved throughout evolution. BLAST searches of the amino acid sequence of SSPLA₂ against the *Homo sapiens* database shows that it is phylogenetically related to the human Group IVA PLA₂ family. This same analysis using the fungal databases revealed that SSPLA₂ is more closely related to the phospholipases of the filamentous fungi than to PLAB of yeasts. The similarity to both human and fungal phospholipases is found primarily in the catalytic domain with a great deal of variation contained in the first and last 200 amino acids. In the catalytic domain we find an important difference between SSPLA₂ and the human homologues. The former has one continuous catalytic domain, rather than the more typical cPLA₂ structure where two homologous catalytic domains are present, interspaced with unique sequences [43].

SSPLA₂ lacks the C2 motif found in cPLA₂ of higher eukaryotes. This domain is involved in the translocation of the enzyme to the membrane in response to an increase in intracellular calcium concentration [34]. Nevertheless, SSPLA₂ has 3 putative EF hand motifs suggesting that it could also be calcium modulated. EF hand motifs are also present in the PLA₂ homologues of *M. grisea*, *G. zeae*, *N. crassa* and *A. nidulans* in different areas of these proteins.

It is interesting to note that *A. nidulans* PLA₂ has been reported to be responsive to calcium even though it also lacks a C2 domain [51].

Also contributing to the possible modulation by calcium of this protein is the presence of a putative calmodulin binding domain [44]. As in the case of the EF hand motifs, analysis of the PLA₂ homologues of *M. grisea*, *N. crassa*, *G. zeae* and in *A. nidulans* show the presence of possible calmodulin binding domains in different areas of the proteins [44]. In *S. schenckii* the putative calmodulin binding domain is at the C terminal end of the protein, while in *M. grisea*, *N. crassa* and *G. zeae* it is within the first 150 to 250 amino acids.

In addition to the identification of PLA₂ as interacting with SSG-2, we inquired as to the effects of PLA₂ in *S. schenckii* dimorphism. As mentioned previously, PLA₂ hydrolyses the *sn*-2 position of phospholipids, resulting in the release of lysophospholipids and free fatty acids. The most commonly released fatty acid is arachidonic acid. We tested the effects of exogenously added arachidonic acid on the kinetics of germ tube formation or the yeast cell cycle in *S. schenckii*. Our results show that exogenously added arachidonic acid had no significant effect on the kinetics of the yeast to mycelium transition, but a significant stimulation (50%) in the percentage of budding in cells induced to re-enter the yeast cell cycle was observed at 6 h of incubation in the presence of this compound. The observed stimulation of the yeast cell cycle by arachidonic acid is consistent with the inhibitory effects on this same cycle observed in the presence of AACOCF3 and isotetrandrine in *S. schenckii*, inhibitors of PLA₂. These inhibitors have different mechanisms of action as stated previously. AACOCF3 is a competitive inhibitor of PLA₂ [46] and an analogue of arachidonic acid, while isotetrandrine interferes with G protein activation of PLA₂ [47]. Both AACOCF3 and isotetrandrine increased significantly the percentage of cells with germ tubes at 6 and 9 h after inoculation and decreased budding in cells induced to re-enter the yeast cycle. The AACOCF3 results are consistent with our hypothesis that PLA₂ activity is needed for the yeast cell cycle in *S. schenckii*, specifically at the start of DNA synthesis [3]. Furthermore, the isotetrandrine results support the hypothesis that the interaction of SSG-2 with PLA₂ is required for these processes to occur.

It is of interest to note that we recently reported similar results in the presence of calmodulin inhibitor W7 and inhibitors of calcium-calmodulin kinase in *S. schenckii* [52]. Inhibiting calmodulin or calmodulin-dependent kinase also inhibited the re-entry of yeast cells into the cell cycle. We can speculate that by inhibiting the calmodulin dependent kinase we are also inhibiting the migration of cPLA₂ to the membrane and/or its activation.

We cannot fully ascertain the functional consequences of the observed interaction between PLA₂ and SSG-2 at this time. Future work will help us clarify this relationship. Nevertheless, two important processes that have a bearing in cell cycle progression have been identified as subjected to cPLA₂ activity in other systems: 1) the production of biologically active molecules and 2) membrane remodeling [53].

There is very little information regarding the effects of the primary metabolites released from the action of PLA₂ (arachidonic acid and lysophospholipids) in fungi. Arachidonic acid was reported to stimulate adenylate cyclase [54] in *S. cerevisiae*. If this is also true for *S. schenckii*, addition of arachidonic acid to the medium would be expected to stimulate the yeast cell cycle and this was what we observed. We had previously reported that dibutyryl derivatives of cAMP inhibit the yeast to mycelium transition in *S. schenckii* [4].

On the other hand, membrane remodeling is also an important function of enzymes such as phospholipases. This process is needed for cell cycle progression and fungal morphogenesis [53]. It has been reported in other systems that in order for the cell cycle to occur there must be a careful balance between membrane phospholipid synthesis and degradation. PLA₂ has an important role in the maintenance of this balance [35,55]. The lipid composition of the membrane is also essential for the correct receptor-protein interactions and plays an important role in signal transduction. G proteins are usually in molar excess when compared to the GPCR's and a large number of inactive GDP-bound heterotrimeric G protein molecules must be available in receptor-rich domains associated to membrane lipids [56].

G proteins can also affect PLA₂ activity by a number of different mechanisms such as: increasing the intracellular calcium concentration, transcriptional regulation and stimulation of phosphorylation through different protein kinases such as protein kinase C and MAP kinases (for a review see [57]).

The studies presented here constitute the first evidence of the interaction of G protein subunits of fungi with a phospholipase. These results establish for the first time a relationship between G proteins and the pathogenic determinants of fungi. The identification of such an important protein as partners of a G protein alpha subunit in fungi suggests a mechanism by which these G proteins can control pathogenicity in fungi. The existence of the interaction reported here may offer an explanation as to why fungi with decreased G protein alpha subunits such as *C. parasitica*, hypovirus infection [15] and *M. grisea* with disrupted G α subunit gene, *magB*, exhibit reduced

levels of virulence [18]. This information is essential if we are to understand the disease producing process of fungi. It will also help elucidate the signal transduction pathway leading from the G protein onward and will give us a better insight into signal transduction in pathogenesis and dimorphism in *S. schenckii*.

Conclusion

We have shown the presence of a new G protein α subunit in *S. schenckii*, SSG-2. The cDNA sequence of the *ssg-2* gene encoded a 355 amino acid G α subunit of 40.90 kDa containing the 5 consensus domains present in all G α subunits. The genomic sequence has four introns, whose positions are conserved in the other fungal homologues of this gene.

Yeast two-hybrid analysis using the complete amino acid sequence of SSG-2 identified a PLA₂ homologue as an interacting partner of this G protein subunit. This 846 amino acid protein was encoded by an intronless gene. The 92.62 kDa protein encoded by this gene contained all the domains and amino acid residues that characterize cytosolic phospholipase A₂.

PLA₂ and other phospholipases in fungi have very diverse roles not only as virulence factors but also in membrane homeostasis and signal transduction. Inhibitor studies showed that this PLA₂ homologue and its interaction with SSG-2 were necessary for the re-entry of *S. schenckii* yeast cells into the budding cycle suggesting a role for this important virulence factor in the control of dimorphism in this fungus and for the expression of the yeast form. The effects of PLA₂ on the yeast cell cycle could be viewed as resulting from the generation of lipid messenger molecules or from membrane remodelling that affects the G1->S transition and G protein activity.

The relationship reported here between these two proteins, SSG-2 and SSPLA₂, constitutes the first report of the interaction of a fungal phospholipase and a G protein subunit and the possible involvement of G protein in fungal virulence and morphogenesis.

Methods

Strains and culture conditions

S. schenckii (ATCC 58251) was used for all experiments. The yeast form of this fungus was obtained as described [2]. *S. cerevisiae* strains AH109 and Y187 were supplied with the MATCHMAKER Two-Hybrid System 3 (Clontech Laboratories Inc., Palo Alto, CA).

Nucleic acids isolation

DNA and RNA were obtained from *S. schenckii* yeast cells as described previously using the methods of Sherman [58], and Chomczynski & Sacchi [59], respectively. Poly

A+ RNA was obtained from total RNA using the mRNA Purification Kit from Amersham Biosciences (Piscataway, NJ, USA).

Sequencing the *ssg-2* gene

Polymerase chain Reaction and Rapid amplification of cDNA ends (RACE)

S. schenckii DNA (100 ng) was used as template for polymerase chain reaction (PCR) with primers (100–200 ng) targeted to conserved motifs in $G\alpha$ subunits. The primers used were: GESGKST (fw) 5' ggtgc(c/t)ggtga(a/g)tc(a/c)gg(a/t)aa(a/g)tc 3'; KWIHCF (rev) 5' aagcag tgaatccactc 3'; TQATDT (rev) 5'gtatcggtagcttggtc 3'; MGACMS (fw) 5' atggg ggcttgcagatg 3' and KDSGIL (rev) 5' taggatccggaatctttg 3'. The Ready-to-Go™ PCR Beads (Amersham Biosciences) were used for PCR using the amplification parameters described previously [52].

PCR products were isolated and cloned using the TOPO TA Cloning System (Invitrogen Corp., Carlsbad, CA, USA) [19]. Plasmid preparations were obtained using the Fast Plasmid TM Mini technology from Eppendorf (Brinkmann Instruments, Inc. Westbury, NY, USA).

The 5' and 3' ends of the *S. schenckii* $G\alpha$ subunit gene were obtained using SMART RACE (BD Biosciences, Clontech, Palo Alto CA, USA). All RACE reactions were carried out as described previously [19]. Primers for RACE were designed based on the sequence obtained previously. Nested primers were designed to improve the original amplification reactions. Bands from the 5' and 3' nested PCR, respectively, were excised from the gel, cloned and sequenced [19]. The following primers were used for 3' RACE: GSP2A (fw) 5' cttgaggaaagcagtcagaaccgaatgatg 3' and GSP2C (fw) 5' gtagatcgggcacacctcaactatatact 3'. The following primers were used for 5' RACE: GSP1E (rev) 5' catcattcgggttctgactgtcttctcaag 3'; GSP1D (rev) 5' aaagtcg-cagtagcagcagatctcatcgct 3' and SSG-2 5' UTR primer-1 (rev) 5' tagcagtagaactctgattctgcctg 3' and SSG-2 5' UTR primer-2 (rev) 5' tctcttcttctgctccacctctact 3'.

The complete coding sequence of the *ssg-2* gene from cDNA and genomic DNA were obtained using reverse transcriptase polymerase chain reaction (RT-PCR) and end to end PCR, respectively. The cDNA obtained using the RETROscript™ First Strand Synthesis kit (Ambion, Applied Biosystems, Foster City, CA, USA) was used as template. The following primers were used: MGACMS (fw)/KDSGIL (rev) primer pair. The sequence of these primers were the following: 5' atgggggcttgcagatg 3' and 5' aggataccg-gaatctttg 3', respectively. For the genomic sequence PCR, DNA was used as template and the primers used were the same as those used for RT-PCR. The PCR products containing the entire coding sequence, from both the cDNA and genomic templates were cloned and sequenced.

Sequencing the *sspla₂* gene

Polymerase chain Reaction and Genome Walker

The 5' sequence of the PLA₂ homologue was obtained using a combination of PCR and Genome Walker (Clontech Laboratories Inc., Palo Alto, CA, USA). Genomic DNA was used as template for PCR. For genome walking a Pvu II library of *S. schenckii* genomic DNA done as described by the manufacturer was used as template for the primary specific PCR reactions using the gene specific primers (GSP) and AP1 primer. The primary PCR reactions were used as template for nested PCR using nested gene specific primers (NGSP) and AP2 primer. The primers used were: YARRFA (NGSP, rev) 5' ccgagagacgat-gcaaagcgacggcgta 3'; SLLVFS (GSP and NGSP, rev) 5' agagaagcagaggagact 3'; GLSDEIWARE (rev) 5' ctcgcgcaaatctcgtcggacaggatcc 3'; VHPEVQ (GSP, rev) 5' gaactggactcggggg 3'; LAKYLDLPA (NGSP, rev) 5' ggag-ggaggtcgaggtacttggcgag 3'; (fw) 5' ctcgcaagtacctcgacctcct-gccg 3'; DDVPVIA (rev) 5' aacgcaatcacgggcagctcgtcg 3'; GVSGSGC (fw) 5' ggagtgagcgttcatgctgg 3'; LYFSSFT (rev) 5' taacgacgaaagtagac 3'; PVGVGWPPA (GSP, rev) 5' cggccggcggccagccacaccactgg 3'; PVGVG (fw) 5' ccagtggt-gtgggctg 3'; DDKIEQ (fw) 5' gacgacattgaacaggccaagccgac 3'; DKIEQ (rev) 5' ttcaatcttgcgtccgg 3'; ERHKRERL (rev) 5' cagccgctccttctgctcctc 3'; NOVGGR (NGSP, rev) 5' ctccgacttaataaaat 3'; 3' UTR primer (GSP, rev) 5' atgctgt-gtcgacctccgac 3'. The AP1 and AP2 primers supplied by the manufacturer. The touchdown and nested PCR parameters used were those described previously [60].

DNA sequencing and analysis

All sequencing reactions for the *ssg-2* gene were conducted using the ABI PRISM™ 377 automated DNA sequencer (Applied Biosystems) and the Thermo Sequenase II Dye terminator Cycle Sequencing Premix Kit (Amersham Biosciences) as described previously [19]. Sequencing of the *sspla₂* gene products was done commercially using the SeqWright sequencing service (Fisher Scientific, Houston, TX, USA)

Bioinformatics Sequence Analysis

The theoretical molecular weights were calculated using the on-line ExPASy tool <http://www.expasy.ch/tools/>. On-line Prosite Scan (Proscan Search) search was used to identify potential motifs present in SSG-2 and SSPLA₂ http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=npsa_prosite.html[45]. The protein classification was performed using the PANTHER Gene and Protein Classification System <http://www.pantherdb.org>[40] and on-line Blocks Analysis Server http://blocks.fhcrc.org/blocks/blocks_search.html[37]. The calmodulin-binding domain was identified using the on line Calmodulin Target Database <http://calcium.uhnres.utoronto.ca/ctdb/ctdb/sequence.html>[44]. On-line database searches and comparisons for SSG-2 were performed

using the Integrated Protein Classification (iProClass) database [61] and its BLAST algorithm implementation with a cutoff of 10^{-7} , a low complexity filter and the BLOSUM 62 matrix. The iProClass/UniProt accession numbers of the sequences used for the multiple sequence alignment of G protein subunits were: *S. schenckii* (SSG-2), Q8TF91; *M. grisea* (MAGA), O13314; *C. parasitica* (CPG2), Q00581; *N. crassa* (GNA3) Q9HFW7; *R. necatrix* (WGA1/RGA1), Q9HFA3; *E. nidulans* (GANB), Q9UVK8, and *S. schenckii* (SSG-1), O74259.

On-line database searches and comparisons for SSPLA₂ were performed with the BLAST algorithm <http://www.ncbi.nlm.nih.gov/BLAST/> with a cutoff of 10^{-7} , a low complexity filter and the BLOSUM 62 matrix [39]. The Pfam analysis was done on-line using the using the Wellcome Trust Sanger Institute server <http://pfam.sanger.ac.uk/>[42]. The GenBank accession numbers for the multiple sequence alignment of phospholipases were: *A. nidulans* (PLA₂), [XP_663815](#); *S. schenckii* (SSPLA₂), [ACI04517.1](#); *M. grisea* (hypothetical protein), [XP_363597](#); *N. crassa* (PLA₂), [XP_962511](#); *C. globosum* (hypothetical protein) [XP_001223932](#); *P. anserina* (hypothetical protein) [XP_001909265](#), and *G. zeae* (PLA₂), [XP_382145](#).

Multiple sequence alignments were built using MCOFFEE <http://www.igs.cnrs-mrs.fr/Tcoffee/tcoffee.cgi/index.cgi>[62]. The alignments were visualized using the program GeneDoc <http://www.nrbsc.org/downloads/>.

Yeast two-hybrid

MATCHMAKER Two-Hybrid System 3 was used for the yeast two-hybrid assay (Clontech Laboratories Inc., Palo Alto, CA) using all 3 different reporter genes for the confirmation for truly interacting proteins. For the construction of the bait plasmid, *ssg-2* cDNA was obtained from poly A⁺ RNA, transcribed and amplified by RT-PCR using the Ready-to-Go™ Beads (Amersham Biosciences). The RT-PCR product was amplified using primers containing the gene sequence and an additional sequence containing restriction enzyme sites, Xma I and BamH I at the 5' and 3' ends, respectively. The primers used were: Xma I-MGACMS (fw) 5' ccccggggatggggcttgcagtagt 3' and DSGIL-BamH I (rev) 5' cgcggatccgcgctaggataccggaattctt 3'. The *ssg-2* gene PCR product was cloned in frame into the linearized bait plasmid, pGBKT7 (Clontech Laboratories Inc.) using Quick T4 DNA ligase kit (New England Biolabs Inc., Ipswich, MA, USA) and amplified in *E. coli* by transformation. Sequencing corroborated the sequence, correct orientation, and frame of the inserted gene. The bait containing plasmid was isolated using Fast Plasmid™ Mini technology (Brinkmann Instruments, Inc.) and used to transform competent *S. cerevisiae* yeast cells (Y187). Competent *S. cerevisiae* yeast cells were transformed using the

YEASTMAKER™ Yeast Transformation System 2 from Clontech (BD Biosciences, Clontech Laboratories Inc.). Tests for autonomous gene activation and cell toxicity were carried out also as described by the manufacturer.

Double stranded cDNA was synthesized from *S. schenckii* yeast cells Poly A⁺ RNA using SMART™ Technology Kit (Clontech Laboratories Inc.). The cDNA's were amplified using Long Distance PCR and size selected using the BD CHROMA-SPIN™+TE-400 columns (Clontech Laboratories Inc.).

S. cerevisiae yeast cells AH109 were made competent using the lithium-acetate (LiAc) method mentioned above and transformed with SMART ds cDNA (20 µl) previously amplified by LD-PCR and the linearized pGADT7-Rec (Sma I-linearized plasmid). Transformants were selected in SD/-Leu plates, harvested and used for mating with the bait containing *S. cerevisiae* strain Y187. Mating of *S. cerevisiae* yeast cells strains Y187 (Mat-α) and AH109 (Mat-a) was done according to the manufacturer's instructions. The expression of three reporter ADE2, HIS3 and MEL1 genes in the diploids was used as confirmation for true interacting proteins. Diploids expressing interacting proteins were selected in triple drop out medium (TDO), SD/-Ade/-Leu/-Trp. Colonies growing in TDO medium were tested for growth and α-galactosidase production in quadruple drop out medium (QDO), SD/-Ade/-His/-Leu/-Trp/X-α-gal. Re-plating of these positive colonies into QDO medium was done at least 3 times to verify that they maintain the correct phenotype. Colony PCR was also done to corroborate the presence of both plasmids in the diploid cells using the T7/3'BD sequencing primer pair for the pGBKT7/*ssg-2* plasmid and the T7/3'AD primer pair for the pGADT7-Rec library plasmid. The PCR products obtained with the T7 Sequencing Primer/3'AD Sequencing Primer pair were cloned and sequenced as described above.

Co-immunoprecipitation (Co-IP)

S. cerevisiae diploids obtained in the yeast two-hybrid assay were grown in 125 ml flasks containing 25 ml of QDO for 16 h, harvested by centrifugation and resuspended in 4 ml containing phosphate buffer saline (400 µl) with phosphatase inhibitor (400 µl), deacetylase inhibitor (40 µl) (Active Motif North America, Carlsbad, CA, USA) and protease inhibitors cocktail (40 µl) (EDTA-free, Thermo Scientific, Pierce Biotechnology, Rockford, IL, USA). The cells were frozen in a porcelain mortar in liquid nitrogen, glass beads added and the cells broken as described previously [63]. The cell extract was centrifuged and the supernatant used for Co-IP using the Immunoprecipitation Starter Pack (GE Healthcare, Bio-Sciences AB, Bjorkgatan, Sweden) as described by the manufacturer. Briefly, 500 µl of the cell extract (1–2 µg of protein/ml)

were combined with 1–5 μ l of the anti-cMyc antibody (Clontech, Corp.) and incubated at 4 °C for 4 h, followed by the addition of protein G beads and incubated at 4 °C overnight in a rotary shaker. The suspension was centrifuged and the supernatant discarded, 500 μ l of the wash buffer added followed by re-centrifugation. This was repeated 4 times. The pellet was resuspended in Laemmli buffer (20 μ l) and heated for 5 min at 95 °C, centrifuged and the supernatant used for 10% SDS PAGE at 110 V/1 h. Pre-stained molecular weight standards were electrophoresed in outside lanes of the gel (BioRad Corporation, Hercules, CA, USA).

Western Blots

Western blots were done as described by us previously [63]. The electrophoretically separated proteins were transferred to nitrocellulose membranes using the BioRad Trans Blot System^R for 1 h at 20 volts. After transferring, the nitrocellulose strips were blocked with 3% gelatin in TTBS (20 mM Tris, 500 mM NaCl, 0.05% Tween-20, pH 7.5) at room temperature for 30–60 min. The strips were washed for 5–10 min with TTBS. The TTBS was removed and the strips incubated overnight in the antibody solution containing 20 μ g of antibody, anti-cMyc or anti-HA (Clontech, Corp.) was added to each strip. Controls where the primary antibody was not added were included. The antigen-antibody reaction was detected using the Immun-StarTM AP chemiluminescent protein detection system from BioRad Corporation as described by the manufacturer.

Induction of the yeast to mycelium transition

The yeast form of the fungus was obtained from conidia as described previously [2]. Briefly, yeast cell were grown for 5 days from conidia in 125 ml flasks containing 50 ml of medium M with aeration at 35 °C. These cells were filtered through sterile Whatman #1 filters (GE Healthcare Life Sciences). This procedure increases the concentration of unbudded singlets to approximately 90%. The cells were collected by filtration using Millipore filters GSWP04700 (0.2 μ m) (Millipore Corp. Billerica, MA, USA), washed using basal medium with glucose and used for inoculation to give a final concentration of 10⁵ cells/ml. These cells were induced to form germ tubes in the presence and absence of effectors of PLA₂ activity in a basal medium with glucose at pH 4.0 and 25 °C. Parallel cultures were inoculated with unbudded yeast cells and at 6 and 9 h after inoculation the content of a flask was filtered for the determination of the percentage of cells with germ tubes for each of the substances tested. These same yeast cells were inoculated to give a final concentration of 10⁷ cells/ml and induced to re-enter the yeast cell cycle as described previously in the presence and absence of effectors of PLA₂ in a basal medium with glucose at pH 7.2 and 25 °C with aeration. At 6 and 9 h after inoculation sam-

ples were taken and the percentage of budding cells was recorded.

The following substances were tested for their effects on the yeast to mycelium transition and the yeast cell cycle: arachidonic acid (40 μ M; AACOCF₃ (100 μ M; Nonadeca-4,7,10,13-tetraenyl-trifluoro-methyl ketone) [46] and isotetrandrine (50 μ M; 6,6',7,12-tetra methoxy-2,2'-dimethyl-berbaman) [47]. These substances were obtained from Calbiochem, EMD Biosciences Inc. (Darmstadt, Germany). The results are expressed as the average percentage of cells with germ tubes or buds at 6 and 9 h of incubation \pm one standard deviation of at least three independent determinations. The Student t test was used to determine the statistical significance of the data. A 95% confidence level was used to determine statistical significance.

Authors' contributions

SVB carried out all the molecular biology studies concerning gene cloning and identification of *ssg-2* gene, constructed a yeast cDNA library and did the first yeast two-hybrid analysis. SVB also conducted the PLA₂ inhibition studies. WGV and LPS repeated the yeast two-hybrid analysis with a new cDNA library, identified PLA₂ as an interacting protein for the second time and confirmed the results with co-immunoprecipitation. RGM carried out the sequence alignments and domain characterization of SSG-2 and PLA₂. NRV designed the study, drafted the manuscript, completed the sequenced the *sspla₂* gene, participated in sequence identification, alignments and domain characterization. All authors have read and approved the final manuscript.

Additional material

Additional file 1

Complete multiple sequence alignment of S. schenckii SSPLA₂ to selected cPLA₂ fungal homologues. The complete multiple sequence alignment of fungal cPLA₂ homologues to SSPLA₂ as described in the methods is presented here.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-100-S1.pdf>]

Acknowledgements

The authors wish to acknowledge the technical support of Ms. Claribel González in sequencing the *sspla₂* gene and the cooperation of graduate student Mr. Jorge Rodríguez with the cloning of PCR products. This investigation was supported by the National Institute of General Medicine, Minority Biomedical Research Support Grant 3S06-GM-008224 and partially by the RISE Program grant R25GM061838. RGM acknowledges funding through NIH NIGMS grant T36GM008789-05 and acknowledges the use of the Pittsburgh Supercomputing Center National Resource for Biomedical Supercomputing resources funded through NIH NCRR grant 2 P41 RR06009-16A1.

References

- Travassos LR, Lloyd KO: **Sporothrix schenckii and related species of Ceratocystis.** *Microbiol Rev* 1980, **44(4)**:683-721.
- Betancourt S, Torres-Bauza LJ, Rodriguez-Del Valle N: **Molecular and cellular events during the yeast to mycelium transition in *Sporothrix schenckii*.** *Sabouraudia* 1985, **23(3)**:207-218.
- Resto S, Rodriguez-del Valle N: **Yeast cell cycle of *Sporothrix schenckii*.** *J Med Vet Mycol* 1988, **26(1)**:13-24.
- Rodriguez-Del Valle N, Debs-Elias N, Alsina A: **Effects of caffeine, cyclic 3', 5' adenosine monophosphate and cyclic 3', 5' guanosine monophosphate in the development of the mycelial form of *Sporothrix schenckii*.** *Mycopathologia* 1984, **86(1)**:29-33.
- Serrano S, Rodriguez-del Valle N: **Calcium uptake and efflux during the yeast to mycelium transition in *Sporothrix schenckii*.** *Mycopathologia* 1990, **112(1)**:1-9.
- Lengeler KB, Davidson RC, D'Souza C, Harashima T, Shen WC, Wang P, Pan X, Waugh M, Heitman J: **Signal transduction cascades regulating fungal development and virulence.** *Microbiol Mol Biol Rev* 2000, **64(4)**:746-785.
- Cabrera-Vera TM, Vanhauwe J, Thomas TO, Medkova M, Preininger A, Mazzoni MR, Hamm HE: **Insights into G protein structure, function, and regulation.** *Endocr Rev* 2003, **24(6)**:765-781.
- Oldham WM, Hamm HE: **Structural basis of function in heterotrimeric G proteins.** *Q Rev Biophys* 2006, **39(2)**:117-166.
- McCudden CR, Hains MD, Kimple RJ, Siderovski DP, Willard FS: **G-protein signaling: back to the future.** *Cell Mol Life Sci* 2005, **62(5)**:551-577.
- Dupre DJ, Robitaille M, Rebois RV, Hebert TE: **The role of Gbetagamma subunits in the organization, assembly, and function of GPCR signaling complexes.** *Annu Rev Pharmacol Toxicol* 2009, **49**:31-56.
- Hicks JK, Yu JH, Keller NP, Adams TH: **Aspergillus sporulation and mycotoxin production both require inactivation of the FadA G alpha protein-dependent signaling pathway.** *Embo J* 1997, **16(16)**:4916-4923.
- Baasiri RA, Lu X, Rowley PS, Turner GE, Borkovich KA: **Overlapping functions for two G protein alpha subunits in *Neurospora crassa*.** *Genetics* 1997, **147(1)**:137-145.
- Turner GE, Borkovich KA: **Identification of a G protein alpha subunit from *Neurospora crassa* that is a member of the Gi family.** *J Biol Chem* 1993, **268(20)**:14805-14811.
- Kays AM, Rowley PS, Baasiri RA, Borkovich KA: **Regulation of conidiation and adenylyl cyclase levels by the Galpha protein GNA-3 in *Neurospora crassa*.** *Mol Cell Biol* 2000, **20(20)**:7693-7705.
- Choi GH, Chen B, Nuss DL: **Virus-mediated or transgenic suppression of a G-protein alpha subunit and attenuation of fungal virulence.** *Proc Natl Acad Sci USA* 1995, **92(1)**:305-309.
- Gao S, Nuss DL: **Distinct roles for two G protein alpha subunits in fungal virulence, morphology, and reproduction revealed by targeted gene disruption.** *Proc Natl Acad Sci USA* 1996, **93(24)**:14122-14127.
- Regenfelder E, Spellig T, Hartmann A, Lauenstein S, Bolker M, Kahmann R: **G proteins in *Ustilago maydis*: transmission of multiple signals?** *Embo J* 1997, **16(8)**:1934-1942.
- Liu S, Dean RA: **G protein alpha subunit genes control growth, development, and pathogenicity of *Magnaporthe grisea*.** *Mol Plant Microbe Interact* 1997, **10(9)**:1075-1086.
- Delgado N, Rodriguez-del Valle N: **Presence of a pertussis toxin-sensitive G protein alpha subunit in *Sporothrix schenckii*.** *Med Mycol* 2000, **38(2)**:109-121.
- Taussig R, Iniguez-Lluhi JA, Gilman AG: **Inhibition of adenylyl cyclase by Gi alpha.** *Science* 1993, **261(5118)**:218-221.
- Stryer L: **Cyclic GMP cascade of vision.** *Annu Rev Neurosci* 1986, **9**:87-119.
- Stoyanov B, Volinia S, Hanck T, Rubio I, Loubtchenkov M, Malek D, Stoyanova S, Vanhaesebroeck B, Dhand R, Nurnberg B, et al.: **Cloning and characterization of a G protein-activated human phosphoinositide-3 kinase.** *Science* 1995, **269(5224)**:690-693.
- Brown AM, Birnbaumer L: **Ionic channels and their regulation by G protein subunits.** *Annu Rev Physiol* 1990, **52**:197-213.
- Dasal N: **Ion-channel regulation by G proteins.** *Trends Endocrinol Metab* 2001, **12(9)**:391-398.
- Liu B, Wu D: **Analysis of G protein-mediated activation of phospholipase C in cultured cells.** *Methods Mol Biol* 2004, **237**:99-102.
- Exton JH: **Cell signalling through guanine-nucleotide-binding regulatory proteins (G proteins) and phospholipases.** *Eur J Biochem* 1997, **243(1-2)**:10-20.
- Kurrasch-Orbaugh DM, Parrish JC, Watts VJ, Nichols DE: **A complex signaling cascade links the serotonin2A receptor to phospholipase A2 activation: the involvement of MAP kinases.** *J Neurochem* 2003, **86(4)**:980-991.
- Zhao J, Wang X: **Arabidopsis phospholipase Dalphal interacts with the heterotrimeric G-protein alpha-subunit through a motif analogous to the DRY motif in G-protein-coupled receptors.** *J Biol Chem* 2004, **279(3)**:1794-1800.
- Fang EG, Dean RA: **Site-directed mutagenesis of the magB gene affects growth and development in *Magnaporthe grisea*.** *Mol Plant Microbe Interact* 2000, **13(11)**:1214-1227.
- Ivey FD, Yang Q, Borkovich KA: **Positive regulation of adenylyl cyclase activity by a galphai homolog in *Neurospora crassa*.** *Fungal Genet Biol* 1999, **26(1)**:48-61.
- Balsinde J, Balboa MA, Insel PA, Dennis EA: **Regulation and inhibition of phospholipase A2.** *Annu Rev Pharmacol Toxicol* 1999, **39**:175-189.
- Diaz BL, Arm JP: **Phospholipase A(2).** *Prostaglandins Leukot Essent Fatty Acids* 2003, **69(2-3)**:87-97.
- Mattera R, Hayek S, Summers BA, Grove DL: **Agonist-specific alterations in receptor-phospholipase coupling following inactivation of Gi2alpha gene.** *Biochem J* 1998, **332(Pt 1)**:263-271.
- Hirabayashi T, Murayama T, Shimizu T: **Regulatory mechanism and physiological role of cytosolic phospholipase A2.** *Biol Pharm Bull* 2004, **27(8)**:1168-1173.
- Boonstra J, van Rossum GS: **The role of cytosolic phospholipase A2 in cell cycle progression.** *Prog Cell Cycle Res* 2003, **5**:181-190.
- Chakraborti S: **Phospholipase A(2) isoforms: a perspective.** *Cell Signal* 2003, **15(7)**:637-665.
- Henikoff S, Henikoff JG: **Protein family classification based on searching a database of blocks.** *Genomics* 1994, **19(1)**:97-107.
- Sprang SR: **G protein mechanisms: insights from structural analysis.** *Annu Rev Biochem* 1997, **66**:639-678.
- Altschul SF, Gish W, Miller W, Myers EV, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215(3)**:403-410.
- Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A, Narechania A: **PANTHER: a library of protein families and subfamilies indexed by function.** *Genome Res* 2003, **13(9)**:2129-2141.
- Dessen A, Tang J, Schmidt H, Stahl M, Clark JD, Seehra J, Somers WS: **Crystal structure of human cytosolic phospholipase A2 reveals a novel topology and catalytic mechanism.** *Cell* 1999, **97(3)**:349-360.
- Finn RD, Tate J, Mistry J, Coggill PC, Sammut SJ, Hotz HR, Ceric G, Forslund K, Eddy SR, Sonnhammer EL, et al.: **The Pfam protein families database.** *Nucleic Acids Res* 2008:D281-288.
- Pickard RT, Chiou XG, Striffler BA, DeFelippis MR, Hyslop PA, Tebbe AL, Yee YK, Reynolds LJ, Dennis EA, Kramer RM, et al.: **Identification of essential residues for the catalytic function of 85-kDa cytosolic phospholipase A2. Probing the role of histidine, aspartic acid, cysteine, and arginine.** *J Biol Chem* 1996, **271(32)**:19225-19231.
- Yap KL, Kim J, Truong K, Sherman M, Yuan T, Ikura M: **Calmodulin target database.** *J Struct Funct Genomics* 2000, **1(1)**:8-14.
- Bairoch A, Bucher P, Hofmann K: **The PROSITE database, its status in 1997.** *Nucleic Acids Res* 1997, **25(1)**:217-221.
- Bartoli F, Lin HK, Ghomashchi F, Gelb MH, Jain MK, Apitz-Castro R: **Tight binding inhibitors of 85-kDa phospholipase A2 but not 14-kDa phospholipase A2 inhibit release of free arachidonate in thrombin-stimulated human platelets.** *J Biol Chem* 1994, **269(22)**:15625-15630.
- Akiba S, Kato E, Sato T, Fujii T: **Biscoclaurine alkaloids inhibit receptor-mediated phospholipase A2 activation probably through uncoupling of a GTP-binding protein from the enzyme in rat peritoneal mast cells.** *Biochem Pharmacol* 1992, **44(1)**:45-50.
- Parsley TB, Segers GC, Nuss DL, Dawe AL: **Analysis of altered G-protein subunit accumulation in *Cryphonectria parasitica* reveals a third Galpha homologue.** *Curr Genet* 2003, **43(1)**:24-33.

49. Li L, Wright SJ, Krystofova S, Park G, Borkovich KA: **Heterotrimeric G protein signaling in filamentous fungi.** *Annu Rev Microbiol* 2007, **61**:423-452.
50. Ghannoum MA: **Potential role of phospholipases in virulence and fungal pathogenesis.** *Clin Microbiol Rev* 2000, **13(1)**:122-143.
51. Hong S, Horiuchi H, Ohta A: **Identification and molecular cloning of a gene encoding Phospholipase A2 (plaA) from *Aspergillus nidulans*.** *Biochim Biophys Acta* 2005, **1735(3)**:222-229.
52. Valle-Aviles L, Valentin-Berrios S, Gonzalez-Mendez RR, Rodriguez-Del Valle N: **Functional, genetic and bioinformatic characterization of a calcium/calmodulin kinase gene in *Sporothrix schenckii*.** *BMC Microbiol* 2007, **7**:107.
53. Kohler GA, Brenot A, Haas-Stapleton E, Agabian N, Deva R, Nigam S: **Phospholipase A2 and phospholipase B activities in fungi.** *Biochim Biophys Acta* 2006, **1761(11)**:1391-1399.
54. Resnick RJ, Tomaska L: **Stimulation of yeast adenylyl cyclase activity by lysophospholipids and fatty acids. Implications for the regulation of Ras/effector function by lipids.** *J Biol Chem* 1994, **269(51)**:32336-32341.
55. Zhang XH, Zhao C, Seleznev K, Song K, Manfredi JJ, Ma ZA: **Disruption of G1-phase phospholipid turnover by inhibition of Ca2+-independent phospholipase A2 induces a p53-dependent cell-cycle arrest in G1 phase.** *J Cell Sci* 2006, **119(Pt 6)**:1005-1015.
56. Vogler O, Casas J, Capo D, Nagy T, Borchert G, Martorell G, Escriba PV: **The Gbetagamma dimer drives the interaction of heterotrimeric Gi proteins with nonlamellar membrane structures.** *J Biol Chem* 2004, **279(35)**:36540-36545.
57. Drin G, Scarlata S: **Stimulation of phospholipase Cbeta by membrane interactions, interdomain movement, and G protein binding – how many ways can you activate an enzyme?** *Cell Signal* 2007, **19(7)**:1383-1392.
58. Sherman F, Fink GR, Hicks JB: **Methods in Yeast Genetics.** Cold Spring Harbor, NY; 1986.
59. Chomczynski P, Sacchi N: **Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.** *Anal Biochem* 1987, **162(1)**:156-159.
60. Aquino-Pinero E, Rodriguez-del Valle N: **Characterization of a protein kinase C gene in *Sporothrix schenckii* and its expression during the yeast-to-mycelium transition.** *Med Mycol* 2002, **40(2)**:185-199.
61. Wu CH, Huang H, Nikolskaya A, Hu Z, Barker WC: **The iProClass integrated database for protein functional analysis.** *Comput Biol Chem* 2004, **28(1)**:87-96.
62. Wallace IM, O'Sullivan O, Higgins DG, Notredame C: **M-Coffee: combining multiple sequence alignment methods with T-Coffee.** *Nucleic Acids Res* 2006, **34(6)**:1692-1699.
63. Aquino-Pinero EE, Rodriguez del Valle N: **Different protein kinase C isoforms are present in the yeast and mycelium forms of *Sporothrix schenckii*.** *Mycopathologia* 1997, **138(3)**:109-115.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

