

Clinical Outcome of Floating Hip with Irreducible Hip Dislocation and Distal Third Femur Fracture with Intra-Articular Extension: A Rare Case Report

Nabin Kumar Sahu¹, Ananta Narayan Panda¹, Satya Prasanna Nayak¹, Shakti Swaroop¹,
Sourav Mishra¹, Prasanna Biradar¹

Learning Point of the Article:

Floating hip with hip dislocation is difficult to manage but can reduce the hip dislocation with knee spanning external fixator and manage in stages shall reduce the complications.

Abstract

Introduction: Floating hip with hip dislocation is a very high-energy, devastating, and rare injury whose treatment is very challenging, and the outcome is usually poor.

Case Report: A 35-year-old man presented posterior wall fracture acetabulum and dislocation of the hip with ipsilateral distal third shaft femur fracture with intra-articular extension fracture and un-displaced patella fracture. We achieved a reduction of hip dislocation by a knee-spanning external fixator followed by open reduction and internal fixation with anatomical locking plate for distal third femur fracture with intra-articular extension followed by open reduction and internal fixation for posterior wall of acetabulum with recon plate in Kocher-Langenbeck approach in stages. The patient was able to partial weight bear after 12 weeks of the injury and mobilized independently without any support after 5 months.

Conclusion: Floating hip with hip dislocation is difficult to manage but reducing the hip dislocation with knee spanning external fixator and management in stages will reduce the complications and better outcome.

Keywords: Floating hip, hip dislocation, knee spanning external fixator.

Introduction

A floating hip is a very high-energy, devastating, and rare injury for which treatment is very challenging and the outcome is poor. A floating hip with hip dislocation is a more devastating injury than a floating hip. A floating hip is a rare case with an incidence rate of one in ten thousand fracture patients. A floating joint is a skeletal discontinuity proximal and distal to that joint [1]. The floating hip term can be delineated as a fracture of the pelvis or acetabulum with an ipsilateral femur fracture. Floating hips have been classified into three types by Mueller. Type A is a

combination of acetabulum and femur fractures; type B is a combination of pelvic and femur bone fractures; and type C is a combination of pelvic, acetabulum, and femur fractures [2]. Leibergall is classified into two types: groups A and B. Group A included a femur fracture with an ipsilateral pelvic bone fracture, either an open book fracture or an unstable shear fracture of the pelvis, whereas group B included a femur fracture with an ipsilateral acetabulum fracture [3]. Nathan classified type A as a femur fracture with pelvic ring fracture and type B as a femur fracture with acetabulum fracture [4]. Nathan's irreducible posterior hip dislocation with a floating hip was reduced by open

Access this article online

Website:
www.jocr.co.in

DOI:
<https://doi.org/10.13107/jocr.2024.v14.i08.4632>

Author's Photo Gallery



Dr. Nabin Kumar Sahu



Dr. Ananta Narayan Panda



Dr. Satya Prasanna Nayak



Dr. Shakti Swaroop



Dr. Sourav Mishra



Dr. Prasanna Biradar

¹Department of Orthopaedics, IMS and SUM Hospital, Kalinga Nagar, Bhubaneswar, Odisha, India.

Address of Correspondence:

Dr. Nabin Kumar Sahu,
Department of Orthopaedics, IMS and SUM Hospital, Kalinga Nagar, Bhubaneswar, Odisha, India.
E-mail: nabinsahu08@gmail.com

Submitted: 11/04/2024; Review: 25/05/2024; Accepted: June 2024; Published: August 2024

DOI: <https://doi.org/10.13107/jocr.2024.v14.i08.4632>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License <https://creativecommons.org/licenses/by-nc-sa/4.0/>, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms



Figure 1: X-rays of (a) pelvis bilateral hip showing dislocation of hip with fracture of acetabulum, (b) X-ray full length femur AP and a lateral view showing fracture of distal third shaft femur with extension distally, (c) X-ray knee joint AP and a lateral view showing fracture distal third shaft femur with extension to articular surface.

reduction by the Kocher-Langenbeck approach [4]. We are reporting a case of irreducible hip dislocation with floating hip and distal femur fracture, which is a very rare case, and reporting the method, stages of treatment, and outcome of the case.

Case Report

We present a case of a 35-year-old man who came to our

emergency with a history of road traffic accidents after sustaining multiple fractures and irreducible hip dislocation with type A floating hip as per the Mueller classification of floating hip and type C1 distal femur intercondylar fracture with shaft femur fracture in the lower limb. The patient had no head injury with Glasgow Coma Scale 15/15 without any abdominal, pelvis, or chest injury. X-ray and computed tomography (CT) scan imaging of the pelvis and knee joint




Figure 2: Non contrast computed tomography scan of (a) knee joint coronal view and (b) knee joint axial view showing intra-articular fracture (c) coronal view of pelvis with bilateral hip (d) 3d reconstruction. Showing dislocation of right hip joint with acetabulum fracture and distal third shaft of femur fracture with intra-articular extension.




Figure 3: (a) is Non contrast computed tomography scan axial cut bilateral hip joint and (b) is an X-ray of the bilateral hip joint depicting the reduction of right hip dislocation after knee spanning external fixator on pod-1.

which are shown in Fig. 1 and 2 revealed a posterosuperior hip dislocation with a posterior wall fracture of the acetabulum (Thompson and Epstein classification type II) and an ipsilateral shaft femur with a distal third femur fracture with intra-articular




Figure 5: Post-operative day-2 X-ray of pelvis bilateral hip AP view (a) and oblique view (b) showing the fixation of the acetabulum. X-ray of the femur with knee joint AP, lateral view (c) showing distal third femur shaft fracture with intra-articular extension.




Figure 4: C arm images. Intraoperative c-arm pictures of (a) dislocated right hip (b) after reduction of right hip joint.

of the right lower limb. He had an associated injury, a segmental fracture of the clavicle with ipsilateral upper limb dysfunction with hand power 2/5, which was a partial incomplete brachial plexus injury. His D-dimer value was 1045 which is very high. We planned for emergency hip dislocation reduction by the close reduction method with the help of Schanz pins and an external fixator. We applied traction to the proximal femur with 3 Schanz pins and a fixator and did the reduction maneuver by the Allis method with knee flexion and internal rotation but couldn't reduce the hip dislocation picture [5]. After the failure of the Allis reduction maneuver, hip dislocation was reduced by the closed method after completing the knee-spanning external

fixator while maintaining the limb length of the fractured femur which is shown in intraoperative C-arm pictures in Fig. 3 and 4 and X-ray and CT scan of pelvis with both hips in Fig. 3. On the same anesthesia, intercondylar femur fracture fixation was done by close reduction and internal fixation with two 6.5 mm CC screws to congruent the articular surface. After 3 days, fixators were removed, and the distal third femur fracture with intra-articular extension was fixed with a distal femoral locking plate and two interfragmentary screws and traction was applied to the remaining tibial fixator. On the same anesthesia, the segmental clavicle was fixed with an anatomical plate. After 7 days of injury, the acetabulum was fixed in the Kocher-Langenbeck approach in the left lateral position by a 3.5 mm, 9-hole recon plate, and two interfragmentary screws with the help of a trochanter osteotomy, which was fixed with two 6.5 mm CC screws. Immediate post-operative day 2 X-rays were done which is shown in Fig. 5. Immediate post-operative after 15 days 30–45° by continuous passive movement of the knee joint, and hip flexion were



Figure 6: Clinical images showing functional outcome 6 months post-surgery clinical images- (a) sit cross-legged (b) standing (c) with significant knee and hip flexion (d) full extension.

started till 6 weeks. After 6 weeks, knee bending was increased to 90°–120° till 12 weeks by a physiotherapist. Gait training was started after 12 weeks with support from a physiotherapist followed by weight-bearing mobilization with the help of a stick till 5 months. We followed up on this patient for 6 weeks, 12 weeks, 6 months, and 1 year and found a satisfactory X-ray with a satisfactory clinical outcome. The patient started weight bearing after 12 weeks with support and started squatting and sitting cross-leg after 5 months. He achieved full knee flexion




Figure 7: 6 months post follow-up (a) X-ray full length right femur AP and lateral view (b) X-ray pelvis bilateral hip showing the good union of the distal third femur and intra-articular femur as well as posterior wall acetabulum with myositis ossificans.

and extension, HIP flexion 0–110°, abduction and adduction 0–30° with external and internal rotation 45° after 5 months of surgery as shown in Fig. 6 and found satisfactory union in 6 months of post-operative X-ray which is shown in Fig. 7. He was independently walking after 6 months of surgery. In X-ray myositis ossificans developed in the hip abductors after 6 weeks and was started on indomethacin.

Discussion

Floating hip dislocation is a high-velocity trauma that involves multiple injuries and polytrauma. This is a very rare case and a big challenge to manage trauma surgeons, which needs surgical management in the correct sequence at the right time. Radiologic evaluation by multiple views of X-rays and CT scans is very important to avoid missing diagnoses because hip dislocation is missed by regular X-rays by 50% in floating hip cases [6]. Floating hip dislocation is very difficult

to reduce, and it is an orthopedic emergency to avoid avascular necrosis as a later complication. Many articles describe many techniques for close reduction of hip dislocation by using the Steinmann pin and traction method [3, 7-9]. N.C. Tiedeken et al. describe the open reduction of hip dislocation and the fixation of the acetabulum in irreducible floating hip dislocation [4]. We achieved floating hip dislocation reduction by applying a knee-spanning external fixator by alignment of the limb length, which was failed by the Steinmann pin and Allis maneuver by longitudinal traction and internal rotation [5]. Many complications can be faced during and after treatment of this type of injury [10]. It helps to decrease the chance of avascular necrosis of the head of the femur and improves the general condition of the patient.

Conclusion

Hip dislocation with a floating hip is a very rare case and a very high-energy trauma that is difficult to manage hemodynamically as well as fracture and dislocation. We achieved a favorable outcome for this patient outcome as we hemodynamically optimized immediately and hip dislocation was reduced immediately within 12 h of injury by a knee-spanning external fixator, which failed to reduce several attempts of reduction technique, which reduced the chance of avascular necrosis of the head and arthritis of the hip joint. Subsequent femur and acetabulum fracture fixation was done in stages, which reduced the chance of fat embolism syndrome

and other complications. The surgical line of management of floating hips has been up for debate for decades. No definitive guidelines are available for the order of fixation for these fractures. This is our endeavor to present an organized structural planning for the floating hip.

Clinical Message

Floating hip with hip dislocation is a complex trauma that can be simplified the treatment by reducing the hip dislocation with the help of knee spanning external fixator by maintaining the limb length alignment and fixation of fractures can be done by stages will be better options to decrease the chance of complication in such type of polytrauma case.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given the consent for his/ her images and other clinical information to be reported in the journal. The patient understands that his/ her names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Conflict of interest: Nil **Source of support:** None

References

1. Simpson NS, Jupiter JB. Complex fracture patterns of the upper extremity. *Clin Orthop Relat Res* 1995;318:43-53.
2. Muller EJ, Siebenrock K, Ekkernkamp A, Ganz R, Muhr G. Ipsilateral fractures of the pelvis and the femur-floating hip? A retrospective analysis of 42 cases. *Arch Orthop Trauma Surg* 1999;119:179-82.
3. Liebergall M, Mosheiff R, Safran O, Peyser A, Segal D. The floating hip injury: Patterns of injury. *Injury* 2002;33:717-22.
4. Tiedeken NC, Saldanha V, Handal J, Raphael J. The irreducible floating hip: A unique presentation of a rare injury. *J Surg Case Rep* 2013;2013:rjt075.
5. Allis OH. *An Inquiry into the Difficulties Encountered in the Reduction of Dislocations of the Hip*. Philadelphia, PA: Dornan; 1896.
6. Helal B, Skevis X. Unrecognised dislocation of the hip in fractures of the femoral shaft. *J Bone Joint Surgery. Br Vol* 1967;49:293-300.
7. Chi-Chuan W, Chun-Hsuing S, Lih-Huei C. Femoral shaft fractures complicated by fracture-dislocations of the ipsilateral hip. *J Trauma* 1993;34:70-5.
8. Harper M. Traumatic dislocation of the hip with ipsilateral femoral shaft fracture: A method of treatment. *Injury* 1982;18:391-4.
9. Suzuki T, Shindo M, Soma K. The floating hip injury: Which should we fix first? *Eur J Orthop Surg Traumatol* 2006;16:214-8.
10. Cannada LK, Justin MH, Preston JB, Heidi I, Hassan M, Jason H, et al. Treatment and complications of patients with ipsilateral acetabular and femur fractures. *J Orthop Trauma* 2017;31:650-6.

Conflict of Interest: Nil
Source of Support: Nil

Consent: The authors confirm that informed consent was obtained from the patient for publication of this case report

How to Cite this Article

Sahu NK, Panda AN, Nayak SP, Swaroop S, Mishra S, Biradar P. Clinical Outcome of Floating Hip with Irreducible Hip Dislocation and Distal Third Femur Fracture with Intra-Articular Extension: A Rare Case Report. *Journal of Orthopaedic Case Reports* 2024 August;14(8): 20-24.