



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Letter to the Editor

## Multiple parameters required for diagnosis of COVID-19 in clinical practice



Dear Editor,

We read with interest the recent papers in this Journal by Hao who described clinical features of atypical 2019 novel coronavirus pneumonia with an initially negative RT-PCR assay.<sup>1</sup> We would like to share our opinions for diagnostic approach of COVID-19 infection, the diagnostic approach for COVID-19 infection should be made comprehensive analysis.

Since December 2019, a series of patients with unknown cause pneumonia had been reported in Wuhan of China. Recently, study on the early transmission dynamics had been reported that human-to-human was the epidemiologic characteristics for COVID-19 infection.<sup>2</sup> Therefore, it is very essential to diagnose precisely the patients suspected with COVID-19 infection for opportune isolation or treatment.

**Chest CT and RT-PCR for detection of COVID-19 infection.** Currently, the real-time reverse transcriptase polymerase chain reaction (RT-PCR) amplification of the viral DNA is considered as the “gold standard”. However, initial RT-PCR is not always positive in the patients with COVID-19 infection.<sup>3,4</sup> In that situation, chest computed tomographic (CT) images could be played an important role to detect the lesions in the pulmonary parenchyma in the patients suspected with COVID-19 infection. But it doesn't mean that the abnormalities of CT images could be observed in the COVID-19 infection while the initial RT-PCR is positive or negative.<sup>3–6</sup> Therefore, even though chest CT plays a key role in detection or diagnosis of COVID-19 infection, however, chest CT examination and RT-PCR results should be mutual verification for precise diagnosis in the patient suspected COVID-19 infection.

**Clinical characteristics.** Beside CT examination results, initial symptoms were helpful in screening. For clinical manifestation, fever, dyspnea, chest tightness, cough, sputum, weakness, vomiting, diarrhea, etc. were observed in the patients with COVID-19 infection.<sup>5,7</sup> Of these clinical characteristics, fever and cough were the most frequency percentage of initial symptoms, and the other manifestations should be taken consideration.

**Laboratory results.** Laboratory results were considered as auxiliary information to diagnose COVID-19 infection.<sup>5,7</sup> In our hospital, there were 14 cases confirmed COVID-19 infection. Of 14 patients, white-cell count was tended to be normal in 12 patients (85.71%), lymphocyte (8/14, 57.14%) and monocyte (9/14, 64.29%) percentage had a tendency to decrease. Significant statistical differences were observed in lymphocyte percentage decreased and C-reactive protein elevated (all  $P=0.015$ ) in the patients with COVID-19 infection between initial positive chest CT results (10/14) and negative chest CT results (4/14). Thus, knowing these

clinical laboratory results was helpful for the doctor to diagnose the current novel coronavirus infection.

**Exposure history.** Recent studies reported that most of the patients with COVID-19 infection had the exposure history to the source of transmission within past 14 days. Approximately 13 patients (2%) had the history of direct contact with wildlife, 75% cases in the epidemic area or contact with resident of epidemic area.<sup>5,7</sup> However, 28 cases (35%) with COVID-19 infections were no obvious history of exposure,<sup>5</sup> which needs to pay more attention in this situation.

In conclusion, even though chest CT has played a key role in detection or diagnosis of COVID-19 infection with some typical CT features while the initial RT-PCR result is negative. However, not all the cases had the initial abnormality chest CT results or positive RT-PCR in the patients with COVID-19 infection. Consequently, RT-PCR results, chest CT features, clinical manifestation, laboratory results, and exposure history should be made a comprehensive analysis to diagnose COVID-19 infection for the clinical decisions beyond clinical and radiological features.

## References

- Hao W. Clinical features of atypical 2019 novel coronavirus pneumonia with an initially negative RT-PCR assay. *J Infect* 2020. doi:10.1016/j.jinf.2020.02.008.
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, china, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020. doi:10.1056/NEJMoa2001316.
- Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, et al. Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology* 2020:200642. doi:10.1148/radiol.2020200642.
- Fang Y, Zhang H, Xie J, Lin M, Ying L, Pang P, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology* 2020:200432. doi:10.1148/radiol.2020200432.
- Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, et al. Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, china: a descriptive study. *Lancet Infect Dis* 2020. doi:10.1016/S1473-3099(20)30086-4.
- Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al. Time course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) pneumonia. *Radiology* 2020:200370. doi:10.1148/radiol.2020200370.
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020. doi:10.1056/NEJMoa2002032.

Pinggui Lei\*<sup>1</sup>

Department of Radiology, the Affiliated Hospital of Guizhou Medical University, Guiyang 550004, China

Bing Fan<sup>1</sup>

Department of Radiology, Jiangxi Provincial People's Hospital, Nanchang 330006, China

Jujiang Mao

*Department of Radiology, the Affiliated Hospital of Guizhou Medical  
University, Guiyang 550004, China*

Pingxian Wang

*Department of Medical Insurance, the Affiliated Hospital of Guizhou  
Medical University, Guiyang 550004, China*

\*Corresponding author.

E-mail address: [pingguilei@foxmail.com](mailto:pingguilei@foxmail.com) (P. Lei)

<sup>1</sup> Equal contributors