

Interaction between IP₃ receptors and BK channels in arterial smooth muscle: non-canonical IP₃ signaling at work

Patricio E. Mujica^{1,2} and Francisco G. González²

¹Graduate School of Biomedical Sciences, Newark Division, and ²Department of Pharmacology and Physiology, New Jersey Medical School, University of Medicine and Dentistry of New Jersey, Newark, NJ 07101

The dynamic regulation of blood flow in the peripheral circulation comprises multiple mechanisms that work in concert to provide adequate control of blood supply to the various vascular beds that feed organ systems. Blood vessels actively respond to different stimuli to adapt vessel diameter according to the continuously changing metabolic requirements of such organ systems. Among these vasoactive stimuli, shear stress as well as endocrine and paracrine factors play a substantial role in the maintenance of vascular homeostasis and the balance between vasoconstriction and vasodilation by acting on the different cell types that form the vascular wall, such as vascular endothelial and smooth muscle cells (SMCs) (Patterson et al., 2002).

Intracellular calcium concentration ($[Ca^{2+}]_i$) is critical for vascular smooth muscle function, as increased $[Ca^{2+}]_i$ triggers SMC contraction. In response to vasoactive ligands that bind to G_q/phospholipase C-coupled receptors, inositol 1,4,5-trisphosphate (IP₃) is produced. IP₃ binds to its receptor (IP₃R) in the membrane of the SMC SR, which facilitates Ca²⁺ release from the SR, raising $[Ca^{2+}]_i$ and inducing cell contraction. SMCs can also respond to membrane depolarization via voltage-dependent Ca²⁺ channels (VDCCs) that mediate Ca²⁺ entry from the extracellular milieu independently of SR Ca²⁺ release (Xi et al., 2008; Zhao et al., 2008), triggering vasoconstriction. Moreover, vascular SMCs express plasma membrane large-conductance, Ca²⁺-activated K⁺ (BK_{Ca}) channels, which are gated by Ca²⁺ with relatively low (micromolar) affinity. As a consequence of a large increase in $[Ca^{2+}]_i$ after IP₃-mediated depletion of internal stores, a BK_{Ca}-dependent hyperpolarizing K⁺ current will induce inactivation of VDCCs, thereby preventing vasoconstriction (Patterson et al., 2002).

The interplay between the different cellular mechanisms that increase $[Ca^{2+}]_i$ is important for the control of vascular tone. In this context, the report we review here (Zhao et al., 2010) sheds an interesting light onto how a negative feedback mechanism involving type 1 IP₃

receptors (IP₃R1) and BK_{Ca} channels is established in cerebral artery SMCs. The findings of Zhao et al. (2010) bring to our attention the possibility that IP₃ may further regulate vascular tone by increasing the apparent Ca²⁺ sensitivity of the BK_{Ca} channel through a mechanism involving an interaction between the channel and IP₃R1, independently of SR Ca²⁺ release. Using electrophysiological approaches on freshly isolated cerebral artery SMCs, the authors found that IP₃ increased the open probability (P_O) of BK_{Ca} channels in both cell-attached and excised inside-out membrane patches in a dose-dependent manner. Neither fast Ca²⁺ buffering with BAPTA nor elevated pipette Ca²⁺ concentration (2 mM) altered the ability of IP₃ to increase P_O of BK_{Ca} channels, suggesting that the observed phenomenon was independent of both intracellular calcium release and plasma membrane calcium influx. Moreover, when the authors examined the Ca²⁺ sensitivity of BK_{Ca} channels in inside-out patches at a physiological membrane potential of -40 mV, they found that 10 μM IP₃ induced a decrease of the mean apparent dissociation constant (K_d) for Ca²⁺ of the channel from ~20 to ~12 μM.

Other reports have suggested that IP₃ may regulate vasoconstriction by inducing a direct coupling between IP₃R1 and canonical transient receptor potential (TRPC) type 3 (TRPC3) channels (Xi et al., 2008; Adebisi et al., 2010), raising the question of whether IP₃ regulates BK_{Ca} activity by means of IP₃R1. Indeed, Zhao et al. (2010) found that heparin, an IP₃R blocker as well as a monoclonal antibody against the cytosolic C terminus of IP₃R1, prevented the IP₃-mediated increase in BK_{Ca} channel P_O. Moreover, experiments performed on cerebral artery SMCs isolated from IP₃R1 knockout mice showed that IP₃ is unable to increase BK_{Ca} channel P_O in the absence of the IP₃R. This requirement for the IP₃R suggests that the receptor could interact directly with the BK_{Ca} channel, a hypothesis that was tested by coimmunoprecipitation and immunofluorescence resonance energy transfer (immuno-FRET) studies. Coimmunoprecipitation experiments revealed that both

Correspondence to Patricio E. Mujica: mujicape@umdnj.edu

Abbreviations used in this paper: BK_{Ca}, large-conductance, Ca²⁺-activated K⁺; $[Ca^{2+}]_i$, intracellular calcium concentration; immuno-FRET, immunofluorescence resonance energy transfer; IP₃, inositol 1,4,5-trisphosphate; IP₃R, IP₃ receptor; SMC, smooth muscle cell; TRPC, canonical transient receptor potential; VDCC, voltage-dependent Ca²⁺ channel.

© 2011 Mujica and González. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

α and $\beta 1$ subunits of the BK_{Ca} channel are precipitated by an anti- IP_3R1 antibody, suggesting that the three polypeptides coexist in a higher-order macromolecular complex, whereas immuno-FRET showed that BK_{Ca} and IP_3R1 localized in close spatial proximity. The data suggest that IP_3 can modulate vascular tone by regulating BK_{Ca} channel activity independently of the canonical SR Ca^{2+} release pathway, and highlight that this phenomenon requires an interaction between the BK_{Ca} channel and IP_3R1 .

The paper by Zhao et al. (2010) advances our knowledge of the regulation of vascular function by revealing a novel molecular signaling mechanism for IP_3 , a ubiquitous second messenger that mediates the actions of many G_q /phospholipase C-coupled vasoconstrictor agonists such as endothelin-1 and angiotensin II (Patterson et al., 2002; Adebisi et al., 2010). The same group has contributed to the understanding of this non-canonical IP_3 signaling by providing evidence that IP_3Rs can directly couple to plasma membrane ion channels such as TRPC3, thereby modifying their activity (Xi et al., 2008; Adebisi et al., 2010). In arterial myocytes, both endothelin-1 and IP_3 activate a nonselective cation current (I_{Cat}) that depends on the functional coupling of TRPC3 and IP_3R1 , an interaction that proved to be isoform selective and independent of SR Ca^{2+} release. Although TRPC3-mediated I_{Cat} would cause membrane depolarization and activation of VDCCs, thereby inducing vasoconstriction, the novel finding that BK_{Ca} channels are also activated by IP_3 in an IP_3R1 -dependent manner raises an interesting question regarding the balance between depolarizing and hyperpolarizing currents in arterial myocytes, which could play a significant role in the control of basal vascular tone and of other active vascular processes such as the myogenic response or endothelium-mediated SMC hyperpolarization. In fact, BK_{Ca} channels have been implicated in the control of myogenic tone as a hemodynamic autoregulatory mechanism (Hill et al., 2010), and the fact that IP_3 is able to simultaneously regulate TRPC3 and BK_{Ca} channels provides insight into the role that paracrine factors may play in the modulation of the myogenic response and the local regulation of blood flow. The role of TRPC channels in the control of vasoconstriction has been addressed before, and it has been suggested that the main isoform involved in the myogenic response (i.e., vasoconstriction in response to increased transmural pressure) is TRPC6, whereas TRPC3 would be implicated in the response to vasoconstrictor agents that elevate IP_3 (Welsh et al., 2002; Adebisi et al., 2010). Beyond local regulation, it will be interesting to probe the importance of this specific coupling between IP_3R1 and BK_{Ca} channels in the control of blood flow distribution and its role in other vascular beds.

Despite the compelling evidence discussed above, some questions still remain. For instance, all IP_3 stimulations in inside-out patch clamp experiments were

performed at concentrations of 10 μM IP_3 or higher, with the support of an IP_3 dose-response curve that yielded maximal stimulations at 10 μM and an apparent K_d for IP_3 of $\sim 4.1 \pm 1.3 \mu M$. Although the authors' discussion about arterial myocyte global $[IP_3]_i$ is enlightening, we agree that more accurate measurements of basal and agonist-stimulated arterial myocyte $[IP_3]_i$ are required to obtain a clearer picture of the physiologically relevant IP_3 levels that will induce IP_3R1 -mediated activation of BK_{Ca} channels. To this effect, electrophysiological experiments performed at $\sim 4 \mu M$ IP_3 (the apparent K_d indicated by the data) may further clarify the relevance of submaximal IP_3 levels that may be encountered in a physiological context. It was reported previously that Bt- IP_3 , a cell-permeable IP_3 analogue, could induce TRPC3-mediated depolarization in endothelium-denuded, SR Ca^{2+} -depleted arteries pressurized at 20 mmHg at concentrations as low as 1 μM (Xi et al., 2008), whereas the data provided by Zhao et al. (2010) show no increase in the mean P_O of BK_{Ca} channels at such IP_3 levels. This discrepancy may be a result of the experimental approach, but we cannot rule out the possibility that the very nature of the interaction between IP_3R1 and plasma membrane ion channels may be different. IP_3Rs have multiple IP_3 -binding sites with varying affinities (Foskett et al., 2007), and receptor-activating conformational changes induced by IP_3 binding at different sites could favor interactions with different proteins.

We think that elucidating whether the IP_3 -induced shift in Ca^{2+} sensitivity of the BK_{Ca} channel (from ~ 20 to $\sim 12 \mu M$) is relevant for its biophysical properties, as well as for its physiological role, would be an important step in the clarification of the general picture this paper conveys. However, it remains unsolved whether this effect is in fact a result of a direct molecular interaction between IP_3R1 and the BK_{Ca} channel. We think that the authors do not provide a conclusive demonstration for the occurrence of a direct interaction, despite that evidence in favor of the requirement of IP_3R1 for IP_3 -mediated activation of BK_{Ca} channels is presented. Heparin, a general IP_3R blocker, and an antibody against IP_3R1 both abolish IP_3 -induced BK_{Ca} activation in inside-out membrane patches. The effect of these IP_3R1 -blocking strategies would require an intact and functional receptor (i.e., able to bind IP_3 and activate BK_{Ca}) excised with the patch. In other work by this group, it was shown that a synthetic peptide corresponding to the N-terminal sequence of the IP_3R containing the TRPC3-binding region activated a Gd^{3+} -sensitive, nonselective cation current (I_{Cat}). Also, membrane-permeant (TAT-conjugated) synthetic peptides were used to probe the coupling between IP_3R1 and TRPC3. Data revealed the presence of a calmodulin- and IP_3R -binding domain in TRPC3 channels that supports the occurrence of the proposed interaction (Adebisi et al., 2010). To our knowledge, the presence of calmodulin- and IP_3R -binding domains

has not been described in BK_{Ca} channels. The use of these experimental approaches could prove helpful in the validation of the authors' hypothesis by establishing the actual domains that may be mediating the interaction between IP₃R1 and BK_{Ca} channels.

IP₃R1 was also shown to localize in close proximity to BK_{Ca} channels by immuno-FRET. As opposed to the standard FRET approach, which makes use of genetically encoded fluorophores directly coupled to the proteins of interest, immuno-FRET is based on the energy transfer between fluorophores coupled to the secondary antibodies of a standard indirect immunofluorescence. In both methodologies, resonance energy transfer occurs between donor and acceptor molecules located within a specific distance (the Förster radius, R₀) characteristic of each FRET pair, and it is unlikely to occur when the distance between fluorophores is >100 Å (Pietraszewska-Bogiel and Gadella, 2011). Positive immuno-FRET signals are a good indicator of the interaction between IP₃R1 and BK_{Ca} if the distance between donor and acceptor represents the same separation between these proteins, but this is hard to determine because the three-dimensional structure of the immunocomplex is unknown. Moreover, antibodies are large proteins with a molecular weight of ~150 kD. Analysis of the crystal structure of various immunoglobulins has shown that the distance between antigen-binding sites can vary between 118 and 185 Å (Saphire et al., 2002).

Because R₀ for the FRET pair used by the authors (Cy2/Cy3) is ~50–60 Å (Zhao et al., 2010), it is possible that the actual distance between IP₃R1 and BK_{Ca} may be >100 Å, suggesting that energy transfer could be a consequence of the interaction between components of the immunocomplex, regardless of whether or not IP₃R1 and BK_{Ca} interact and/or localize nearby.

The hypothesis that IP₃R1 and BK_{Ca} channels are found in close proximity is also supported by coimmunoprecipitation experiments showing that an IP₃R1 antibody is able to pull down both subunits of the BK_{Ca} channel. Although it is not conclusive evidence of direct interaction, coimmunoprecipitation data clearly suggests the existence of a macromolecular complex that comprises both entities. Despite this attractive inference, Zhao et al. (2010) do not provide a control reverse immunoprecipitation to strengthen their conclusions. Also, probing the presence of molecular chaperones that may stabilize these interactions could provide insight on the trafficking and signaling properties of the IP₃R1–BK_{Ca} complex, as well as providing explanation for the intracellular normalized FRET data the authors obtained. However, the fact that the regulatory β1 subunit of the channel was coimmunoprecipitated is particularly important; its role as a crucial regulator of channel activity and Ca²⁺ sensitivity (Brenner et al., 2000; Patterson et al., 2002) could provide insight on the mechanisms of BK_{Ca} modulation by IP₃ and IP₃R1. Experiments aimed




Figure 1. IP₃-induced interactions between IP₃Rs and ion channels in arterial SMCs. Induction of Ca²⁺ release from the SR via IP₃R (solid orange arrows) is the best-known mechanism of action of IP₃. Non-canonical IP₃ signaling is independent of SR Ca²⁺ release and involves an IP₃-induced interaction of IP₃R1 with plasma membrane channels such as TRPC3 and BK_{Ca} (red arrowheads), possibly localized in membrane caveolae. TRPC3 activation by IP₃R1 induces a nonselective cation current (I_{cat}; solid green arrow) that depolarizes the cell membrane, activating VDCC and BK_{Ca} channels (dashed green arrows). BK_{Ca} channels are also activated by Ca²⁺ (dashed orange arrows). Whereas VDCCs mediate Ca²⁺ influx through the plasma membrane (purple arrow) leading to contraction, K⁺ efflux through BK_{Ca} channels (blue arrow) leads to hyperpolarization of the cell membrane, which closes VDCCs, leading to SMC relaxation.

at discriminating which BK_{Ca} subunit mediates the interaction between the channel and the IP₃R or other components of the immunoprecipitated complex will help clarify this issue.

It has been shown previously that TRPC3 and TRPC6 channels can associate with BK_{Ca} channels in differentiated cells of a podocyte cell line (Kim et al., 2009), whereas recent work shows that, in arterial SMCs, caveolin-1, a lipid raft scaffolding protein, assembles IP₃R1 and TRPC3 channels to signaling complexes (Adebiyi et al., 2011). Moreover, it has been shown that BK_{Ca} channels are localized to lipid rafts in glioma cells, an interaction that places BK_{Ca} channels in close proximity to IP₃Rs (Weaver et al., 2007). The involvement of lipid rafts and caveolin-1 in the architecture of these macromolecular signaling complexes brings a bridging element between the different components that constitute them, and paves the road to further inquiry regarding the physiological role of such complexes and the potential interaction with other elements known to localize to plasma membrane microdomains. However, other potential interactions may take place between plasma membrane ion channels and SR membrane components. The STIM/Orai family of proteins has been shown to mediate store-operated Ca²⁺ entry in many different tissues including vascular smooth muscle, and they have been proposed to interact with members of the TRPC channel family to regulate Ca²⁺ entry at cellular domains where the plasma membrane is in close proximity to the SR membrane (Wang et al., 2008). Therefore, BK_{Ca} channels, IP₃, and IP₃Rs may also be subject to interaction with these ion channel families in the control of SMC Ca²⁺ signals, cell contractility, and vascular tone. Fig. 1 summarizes the potential interactions discussed above.

The evidence regarding the regulation of vascular homeostasis and, in particular, arterial smooth muscle physiology by non-canonical IP₃ signaling suggests that macromolecular signaling complexes that physically and functionally link the plasma and SR membranes may be an important common theme in terms of the fine modulation of signals that establish vascular SMC excitability and vascular tone, thereby controlling active processes such as the myogenic response. At the same time, they hint at the specific possibility that IP₃-mediated regulation of BK_{Ca} channels may not only involve the IP₃R but also other molecules present in these signaling microdomains. The recent determination of the crystal structure of the cytoplasmic Ca²⁺-sensing domain of the BK_{Ca} channels (Wu et al., 2010; Yuan et al., 2010) will certainly provide further insight to this matter by means of structure-driven mutational analysis to elucidate the specific regions of BK_{Ca} channels that interact with IP₃R1. Careful dissection of their components and integration of their independent and interdependent activities in a physiologically sound model of vascular

function will be the next step in the advancement of this field.

Please participate in a discussion of this Journal Club article on the *JGP* Facebook page (www.facebook.com/JGenPhysiol).

The authors would like to thank Drs. Jorge E. Contreras, Walter N. Durán and Martha C. Nowycky, Mrs. Ammy M. Santiago, Mr. Rafael Contreras, and Mr. Jorge Cantero for their helpful comments and suggestions.

The authors are supported by National Institutes of Health grants 5R01 HL070634 and 5R01 HL088479.

Jorge E. Contreras served as faculty advisor.

Lawrence G. Palmer served as editor.

REFERENCES

- Adebiyi, A., G. Zhao, D. Narayanan, C.M. Thomas-Gatewood, J.P. Bannister, and J.H. Jaggar. 2010. Isoform-selective physical coupling of TRPC3 channels to IP₃ receptors in smooth muscle cells regulates arterial contractility. *Circ. Res.* 106:1603–1612. doi:10.1161/CIRCRESAHA.110.216804
- Adebiyi, A., D. Narayanan, and J.H. Jaggar. 2011. Caveolin-1 assembles type 1 inositol 1,4,5-trisphosphate receptors and canonical transient receptor potential 3 channels into a functional signaling complex in arterial smooth muscle cells. *J. Biol. Chem.* 286:4341–4348. doi:10.1074/jbc.M110.179747
- Brenner, R., G.J. Peréz, A.D. Bonev, D.M. Eckman, J.C. Kosek, S.W. Wiler, A.J. Patterson, M.T. Nelson, and R.W. Aldrich. 2000. Vasoregulation by the β 1 subunit of the calcium-activated potassium channel. *Nature.* 407:870–876. doi:10.1038/35038011
- Foskett, J.K., C. White, K.-H. Cheung, and D.-O.D. Mak. 2007. Inositol trisphosphate receptor Ca²⁺ release channels. *Physiol. Rev.* 87:593–658. doi:10.1152/physrev.00035.2006
- Hill, M.A., Y. Yang, S.R. Ella, M.J. Davis, and A.P. Braun. 2010. Large conductance, Ca²⁺-activated K⁺ channels (BKCa) and arteriolar myogenic signaling. *FEBS Lett.* 584:2033–2042. doi:10.1016/j.febslet.2010.02.045
- Kim, E.Y., C.P. Alvarez-Baron, and S.E. Dryer. 2009. Canonical transient receptor potential channel (TRPC)3 and TRPC6 associate with large-conductance Ca²⁺-activated K⁺ (BKCa) channels: role in BKCa trafficking to the surface of cultured podocytes. *Mol. Pharmacol.* 75:466–477. doi:10.1124/mol.108.051912
- Patterson, A.J., J. Henrie-Olson, and R. Brenner. 2002. Vasoregulation at the molecular level: a role for the β 1 subunit of the calcium-activated potassium (BK) channel. *Trends Cardiovasc. Med.* 12:78–82. doi:10.1016/S1050-1738(01)00146-3
- Pietraszewska-Bogiel, A., and T.W.J. Gadella. 2011. FRET microscopy: from principle to routine technique in cell biology. *J. Microsc.* 241:111–118. doi:10.1111/j.1365-2818.2010.03437.x
- Saphire, E.O., R.L. Stanfield, M.D. Crispin, P.W. Parren, P.M. Rudd, R.A. Dwek, D.R. Burton, and I.A. Wilson. 2002. Contrasting IgG structures reveal extreme asymmetry and flexibility. *J. Mol. Biol.* 319:9–18. doi:10.1016/S0022-2836(02)00244-9
- Wang, Y., X. Deng, T. Hewavitharana, J. Soboloff, and D.L. Gill. 2008. Stim, ORAI and TRPC channels in the control of calcium entry signals in smooth muscle. *Clin. Exp. Pharmacol. Physiol.* 35:1127–1133. doi:10.1111/j.1440-1681.2008.05018.x
- Weaver, A.K., M.L. Olsen, M.B. McFerrin, and H. Sontheimer. 2007. BK channels are linked to inositol 1,4,5-trisphosphate receptors via lipid rafts: a novel mechanism for coupling [Ca²⁺]_i to ion channel activation. *J. Biol. Chem.* 282:31558–31568. doi:10.1074/jbc.M702866200

- Welsh, D.G., A.D. Morielli, M.T. Nelson, and J.E. Brayden. 2002. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circ. Res.* 90:248–250. doi:10.1161/hh0302.105662
- Wu, Y., Y. Yang, S. Ye, and Y. Jiang. 2010. Structure of the gating ring from the human large-conductance Ca(2+)-gated K(+) channel. *Nature.* 466:393–397. doi:10.1038/nature09252
- Xi, Q., A. Adebisi, G. Zhao, K.E. Chapman, C.M. Waters, A. Hassid, and J.H. Jaggar. 2008. IP₃ constricts cerebral arteries via IP₃ receptor-mediated TRPC3 channel activation and independently of sarcoplasmic reticulum Ca²⁺ release. *Circ. Res.* 102:1118–1126. doi:10.1161/CIRCRESAHA.108.173948
- Yuan, P., M.D. Leonetti, A.R. Pico, Y. Hsiung, and R. MacKinnon. 2010. Structure of the human BK channel Ca²⁺-activation apparatus at 3.0 Å resolution. *Science.* 329:182–186. doi:10.1126/science.1190414
- Zhao, G., A. Adebisi, E. Blaskova, Q. Xi, and J.H. Jaggar. 2008. Type 1 inositol 1,4,5-trisphosphate receptors mediate UTP-induced cation currents, Ca²⁺ signals, and vasoconstriction in cerebral arteries. *Am. J. Physiol. Cell Physiol.* 295:C1376–C1384. doi:10.1152/ajpcell.00362.2008
- Zhao, G., Z.P. Neeb, M.D. Leo, J. Pachuau, A. Adebisi, K. Ouyang, J. Chen, and J.H. Jaggar. 2010. Type 1 IP₃ receptors activate BK_{Ca} channels via local molecular coupling in arterial smooth muscle cells. *J. Gen. Physiol.* 136:283–291. doi:10.1085/jgp.201010453