



RESEARCH

Open Access

Fine mapping of the hereditary haemorrhagic telangiectasia (*HHT*)₃ locus on chromosome 5 excludes VE-Cadherin-2, Sprouty4 and other interval genes

Fatima S Govani, Claire L Shovlin*

Abstract

Background: There is significant interest in new loci for the inherited condition hereditary haemorrhagic telangiectasia (HHT) because the known disease genes encode proteins involved in vascular transforming growth factor (TGF)- β signalling pathways, and the disease phenotype appears to be unmasked or provoked by angiogenesis in man and animal models. In a previous study, we mapped a new locus for HHT (*HHT3*) to a 5.7 Mb region of chromosome 5. Some of the polymorphic markers used had been uninformative in key recombinant individuals, leaving two potentially excludable regions, one of which contained loci for attractive candidate genes encoding VE Cadherin-2, Sprouty4 and FGF1, proteins involved in angiogenesis.

Methods: Extended analyses in the interval-defining pedigree were performed using informative genomic sequence variants identified during candidate gene sequencing. These variants were amplified by polymerase chain reaction; sequenced on an ABI 3730xl, and analysed using FinchTV V1.4.0 software.

Results: Informative genomic sequence variants were used to construct haplotypes permitting more precise citing of recombination breakpoints. These reduced the uninformative centromeric region from 141.2-144 Mb to between 141.9-142.6 Mb, and the uninformative telomeric region from 145.2-146.9 Mb to between 146.1-146.4 Mb.

Conclusions: The *HHT3* interval on chromosome 5 was reduced to 4.5 Mb excluding 30% of the coding genes in the original *HHT3* interval. Strong candidates VE-cadherin-2 and Sprouty4 cannot be *HHT3*.

Background

Transforming growth factor (TGF)- β superfamily signalling is of fundamental importance to developmental and physiological regulation. In these pathways (reviewed in [1,2]), ligands such as TGF- β s, bone morphogenetic proteins (BMP)s, activins, nodals, growth/differentiation factors (GDF)s and inhibins bind to receptor complexes of paired type I and type II transmembrane receptor serine/threonine kinases. Activated type I receptors (ALKs 1-7) phosphorylate receptor-associated Smad proteins in complex-specific patterns [3-5]. There is increasing recognition of the role of alternative signalling pathways for particular ligands within designated cell types. In

endothelial cells (ECs), signalling through the TGF- β type II receptor, T β RII, can be propagated not only through ALK-5 via SMAD2/3 as in other cell types, but also through ALK-1 via SMAD1/5/8, providing two mutually antagonistic pathways [6,7]. The transmembrane glycoprotein endoglin is an accessory TGF- β receptor, highly expressed on ECs, and is one factor modulating the balance between ALK-1 and ALK-5 pathways [8].

The inherited vascular condition hereditary haemorrhagic telangiectasia (HHT) [9] is of significant relevance to TGF- β signalling because the genes for endoglin, ALK-1 and SMAD4 (a co-Smad and downstream effector of the TGF- β signalling pathway), are mutated in different HHT families [10-12]. HHT is transmitted as an autosomal dominant trait due to a single mutation in

* Correspondence: c.shovlin@imperial.ac.uk
NHLI Cardiovascular Sciences, Imperial College London, UK, Hammersmith Campus, Du Cane Rd, London W12 0NN, UK

either *ENG* encoding endoglin (HHT type 1); *ACVRL1* encoding ALK-1 (HHT type 2) or *SMAD4* (HHT in association with juvenile polyposis). Perturbation of TGF- β signalling pathways is therefore implicated in HHT pathogenesis.

HHT serves not only as a vascular model of aberrant TGF- β superfamily signalling, but also as a model of aberrant angiogenesis [13,14]. The abnormal blood vessels develop only in selected vascular beds (telangiectasia particularly in mucocutaneous and gastrointestinal sites; arteriovenous malformations (AVMs) most commonly in pulmonary, hepatic and cerebral circulations) [9,15]. At each site, only a small proportion of vessels are abnormal. The context in which HHT mutations are deleterious, when allowing apparently normal endothelial function for most vessels, now appear to be angiogenic in origin. Early studies modelling HHT in transgenic animals provided evidence of aberrant angiogenesis. Heterozygous mice developed HHT-like features; *Eng* and *Alk1* null mice died by E11.5 with normal vasculogenesis but abnormal angiogenesis [7,16-20]. The zebrafish *violet beaugarde* (*vbg*), an *Alk1* mutant, was also homozygous embryonic lethal, with mutant embryos displaying dilated cranial vessels attributed to an increased number of endothelial cells [21]. More recent studies have demonstrated that an *Alk1* deletion in adult mouse subdermal blood vessels resulted in AVM formation in wounded areas displaying angiogenesis [22]; that angiogenic stimuli promoted AVM formation in endothelial specific endoglin knockout mice, accompanied by an abnormal increase in EC proliferation [23] which was also observed in *Eng*^{-/-} mouse embryonic ECs [8], and that *Alk1* knockout mice had defective smooth muscle differentiation and recruitment and excessive angiogenesis [7]. These data from animal models have been accompanied by clinical reports that Bevacizumab, an antibody against vascular endothelial growth factor (VEGF)-A, and thalidomide, appear to have efficacy in treating clinical manifestations of HHT in man [24,25].

A current model to explain these observations, discussed in more detail in [26], is based on the EC-mural cell axis defined by Sato and Rifkin [27]. In angiogenesis, HHT mutations (endoglin and ALK-1) appear to impair recruitment of mural cells to the angiogenic sprout [7,28] at least in part via reduced EC secretion of TGF- β 1 [29,30] and/or reduced TGF- β 1 induced responses [7,29] resulting in defective mural cell stabilisation of the nascent vessel and persistent, excessive, EC proliferation. Thalidomide, which induced vessel maturation in *Eng*^{+/-} mice which normally suffer from excessive angiogenesis, appears to target mural cell recruitment, by increasing endothelial expression of PDGF-B at the endothelial tip cell, thus facilitating recruitment of

pericytes that express PDGFR-b, associated with increasing pericyte proliferation [25].

Further HHT genes were therefore predicted to identify new components or regulators of TGF- β /BMP signalling pathways of particular relevance to angiogenesis. More than 80% of HHT patients carry a sequence variation in *endoglin* or *ALK-1* [31], most (but not all [32,33]) disease-causing, whilst 1-2% carry a mutation in *SMAD4* [34]. HHT patients in whom no mutations have been detected in *endoglin*, *ALK-1* or *SMAD4* may have mutations in unsequenced intronic regions, or undetected large rearrangements of these genes, but importantly, they may have a mutation in a different gene. A third gene for pure HHT, *HHT3*, was mapped in our laboratory, using the pedigree illustrated in Figure 1A, to chromosome 5q [35] (Figure 1B), and a fourth, *HHT4*, to chromosome 7p [36].

The published chromosome 5 *HHT3* interval which contains the *HHT3* gene, is 5.7 Mb long, between microsatellite markers *D5S2011* and *D5S2490* (Figure 1B) [35]. This interval contains 38 genes according to the Ensembl genome browser <http://www.ensembl.org>. There were two groups of potentially very attractive candidate genes. The first encode proteins involved in angiogenesis: *PCDH12* and *SPRY4* (discussed further below), and *FGF1* encoding the proangiogenic fibroblast growth factor 1 [37]. A second group of interval genes encode proteins involved in signalling by serine/threonine kinases: *PPP2R2B* which encodes the B regulatory subunit of protein phosphatase 2A [38], and *STK32A*, encoding a serine/threonine kinase [39]. While these genes had been sequenced with no disease-causing mutations found ([35], Govani and Shovlin unpublished), this did not exclude the genes as disease-causing, due to the possibility of undetected intronic mutations. Importantly *PCDH12* (141.3 Mb), *SPRY4* (141.6 Mb) and *FGF1* (142 Mb) were at the centromeric extreme of the *HHT3* interval, and *PPP2R2B* (146.4 Mb) and *STK32A* (146.6 Mb) at the telomeric extreme, all within potentially excludable regions of the interval [35].

Excluding any of these genes from the *HHT3* interval was important not only to assist the identification of *HHT3* itself, but also to inform scientists working on the relevant proteins, particularly VE-Cadherin-2 (encoded by *PCDH12*) and Sprouty4 (encoded by *SPRY4*): VE-Cadherin-2, primarily expressed on ECs [40], was shown to be down-regulated in HHT1 and HHT2 blood outgrowth ECs [41] with a role in angiogenesis [42]. Furthermore, a close family member (VE-cadherin), associates with endoglin and ALK-1 in cell surface complexes, and promotes TGF- β 1 signalling by facilitating the recruitment of TGF- β receptor II (T β RII) into the active signalling complex [43]. Sprouty family members are involved in branching morphogenesis [44].




Figure 1 Previous *HHT3* interval. **A)** *HHT3* linked Family S pedigree (black symbols represent HHT-affected individuals; white symbols represent unaffected individuals; squares represent males and circles represent females. Individuals are denoted by their generation number, such that the oldest male is I.1, oldest female I.2). **B)** Published *HHT3* interval on chromosome 5 between microsatellite markers *D5S2011* and *D5S2490* [35], identifying the region shared by all affected individuals (open interval between *D5S402* and *D5S436*); regions where the designated individuals had inherited their affected parent's non disease-gene bearing allele (black bars), and regions in which the transition from the non disease-gene bearing allele (black bars) to disease-gene bearing allele (open interval) had occurred, but the exact sites of the meiotic recombination event was not definable using the markers studied in [35] (grey bars; uninformative markers). It was possible that the interval could be reduced to a minimum of *D5S402*-*D5S436*, according to where the recombination breakpoints had occurred in individuals II.1, II.3 and II.9. cen: centromere; tel: telomere; Mb: mega bases.

Overexpression of *Spry4* in ECs of embryonic day (E)9 mice inhibited angiogenesis [45], whereas surviving *Spry4* knockout mice [46] demonstrated enhanced angiogenesis [47].

In this study we report the fine mapping of the published *HHT3* interval using genomic sequence variants detected during candidate gene sequencing, and exclusion of key genes.

Methods

Pedigree

Ethics approval was obtained from the Multicentre Research Ethics Committee for Scotland (MREC/98/0/42; 07/MRE00/19), by the Hammersmith, Queen Charlotte's, Chelsea, and Acton Hospital Research Ethics Committee (LREC 99/5637M) and in 2007 by the Hammersmith Hospitals Trust (SHOV1022, 2007). The

study is registered on <http://www.clinicaltrials.gov> (NCT00230620). The pedigree (Family S, Figure 1A) and DNA extractions were as described in [35].

PCR and sequencing

Oligonucleotide primers were ordered from Eurofins MWG-Biotech and amplified in informative members of Family S by polymerase chain reaction (PCR). Genomic primers spanning exons and > 30 bp of flanking intronic sequences were amplified by PCR using *Phusion Hot-start* (New England Biolabs) or *AmpliTaq Gold* (Applied Biosystems) DNA polymerases. Reactions were performed as recommended by the manufacturers using 35-40 cycles of denaturation, annealing (at primer specific Ta) and extension. PCR products were treated with an ExoSAP mix containing SAP enzyme and buffer (Promega) and ExoI enzyme (New England Biolabs) as

previously described [48]; gel purified with Wizard SV gel and PCR clean-up system (Promega) following the manufacturer's protocol; or gel purified and ethanol precipitated using Costar Spin-X column centrifuge tubes [49] followed by ethanol precipitation [50]. Purified PCR products were sequenced in 10 µl reactions with 25 ng (\approx 3.2 pM) of the sequencing primer on an ABI 3730xl DNA analyzer (MRC CSC DNA core laboratory, Imperial College London, Hammersmith). The results were analysed using FinchTV V1.4.0 software (Geospiza, Inc).

Fine mapping

Non-disease causing genomic sequence variants found during *HHT3* candidate gene sequencing were used to fine map the interval, by providing further polymorphisms to track known recombination events through the uninformative part of the interval. Previously studied microsatellite polymorphisms had not been able to distinguish between disease and non-disease alleles (Figure 1B; grey bars), predominantly due to homozygosity in individual I.2 (Figure 1A). New sequence variants where individual I.2 was heterozygous (Figure 2) were sequenced in the interval-defining members of the pedigree (Figure 3). Adjacent genes (such as *NR3C1*, mutations in which cause glucocorticoid resistance [51]), were also sequenced to identify further polymorphisms to define the precise centromeric recombination event. The alleles inherited by each pedigree member were then assessed and assigned to maternal or paternal origin using Mendelian principles. Haplotypes were then constructed to minimise the number of recombination events that would be required to generate the segments of DNA in each offspring. NCBI dbSNP <http://www.ncbi.nlm.nih.gov/projects/SNP/> and SNPbrowser™ V3.5 (Applied Biosystems) were used to determine if the genomic sequence variants were previously known (when they were allocated the relevant *rs* dbSNP RefSNP label), or novel.

For confirmations of genotypes, SNPs *rs153148*, *rs34074000*, *rs164075*, *rs164077*, *rs6879694*, *rs12110022* and *rs11958187* were sequenced in two separate PCR reactions in two or more key individuals.

Genome analysis

Chromosome 5 positions of genomic sequence variants are based on NCBI Reference build 36 (hg18) assembly.

Results

The *HHT3* interval [35] was defined by recombination events in three affected individuals of Family S, II.1, II.3, and II.9 (Figure 1). At each end of the interval, there were regions illustrated by grey bars, in which it was not possible to state which maternal chromosome 5 allele sequences these individuals had inherited, due to homozygosity in individual I.2. New non-microsatellite genomic sequence variants were identified during candidate gene sequencing and analysed to assess if they could be used to track the inheritance of the sequences derived from the two different alleles of chromosome 5 from individual I.2 in the regions between *D5S2011* and *D5S402*, and between *D5S436* and *D5S2490*. A total of 89 genetic sequence variations were found (75 SNPs, 4 indels and 10 repeats) of which 73 were present on dbSNP (2010, build 131). Eighteen potentially informative genomic sequence variants where I.2 was a heterozygote (seventeen SNPs; one triplet repeat) were identified within the regions of interest (Figure 2).

Figure 3 tracks the inheritance of alleles at these SNPs and triplet repeat in the centromeric (Figure 3A), and telomeric (Figure 3B) extremes of the interval. Although I.2 shared three genotypes with I.1, it was possible to definitely state which of I.2's alleles had been inherited by each of their children at all sites except *rs164077*, *rs252108* and *rs252109* where genotypes were shared. Sites where the inheritance of the affected parent's alleles could definitely be determined were used to generate haplotypes.




Figure 2 Informative genomic sequence variants used in fine mapping. Locations of genomic variants where individual I.2 was heterozygous, potentially allowing her disease associated and non-disease associated alleles to be tracked. Sequence variants are illustrated in relation to candidate gene exons (grey boxes) and original microsatellite markers. By chance, all five genes are on the reverse strand of chromosome 5, designated by reverse arrows. *rs*; as on NCBI dbSNP.





Figure 3 Genomic variant haplotypes in interval-defining members of the pedigree. A) Allele inheritance and haplotypes for the inheritance of the key microsatellite markers, and new genomic sequence variants, across *PCDH12*, *SPRY4*, *FGF1* and *NR3C1* genes in eight members of Family S. For each sequence variant, offspring inherit one allele from their mother and one from their father. Occasional alleles where it was not possible to determine parental origin (and which were therefore not used for haplotype construction) are denoted in non-bold italics. Individual II.5 demonstrates the disease haplotype across the interval, a haplotype which is now shared by II.1, but not II.3 and her descendants until *rs17209237* in *NR3C1*. Therefore II.3 excludes the centromeric region between *rs153148* and the novel SNP in *FGF1*. **B)** Haplotypes for the inheritance of the new genomic sequence variants across the *PPP2R2B* gene in nine members of Family S. Affected individuals II.1, II.3, II.5, III.3 and III.4 demonstrate the disease haplotype which is not shared by II.9 at *rs11958187*, but is shared at *rs12110022*, *rs6879694*, and the remainder of the *HHT3* interval. Therefore II.9 excludes *rs11958187* and beyond from the telomeric extreme of the interval]: uninformative regions; bp: base pairs.

As shown in Figure 3A, sequencing 15 of these markers allowed better definition of the centromeric regions where II.1 and II.3 had inherited the disease-gene associated allele, and indicated that the site of the recombination breakpoint differed in the two individuals. In individual II.1, the breakpoint had occurred close to *D552011* such that he had inherited the maternal disease allele from *rs153148* (within *PCDH12*) at 141.3 Mb and onwards. However, in individual II.3, the recombination event had occurred further into the *HHT3* interval: II.3 continued to inherit a different maternal allele to her affected siblings from *rs153148* to a novel SNP (within *FGF1*) at 141.9 Mb. By 142.6 Mb (*rs17209237* and *rs4986593* within *NR3C1*), II.3 had inherited the same maternal allele as her affected siblings, siting the

recombination event between 141.9 Mb and 142.6 Mb (novel SNP and *rs17209237*)

These data in individual II.3 indicated that in addition to the definite *D55402-D55436* region (Figure 1B), the maximal centromeric extent of the disease gene-bearing chromosome 5 that could have been inherited by all affected individuals in the pedigree extended not from *D552011* at 141.2 Mb, but from the *FGF* SNP at 141.9 Mb (Figure 4). This reduction of the uninformative centromeric region by 0.7 Mb excluded 9 Ensembl interval genes (Figure 4). *PCDH12* and *SPRY4* were amongst the group of genes excluded (Figure 4).

The telomeric border of the *HHT3* interval was defined by a recombination event in affected individual II.9. This individual had inherited a non-disease allele at



microsatellite markers up to and including *D5S2490* (146.9 Mb), and was uninformative at microsatellite markers up to *D5S436* (145.2 Mb) [35] (Figure 1B). Figure 3B demonstrates the results of sequencing genomic sequence variants within genes in the potentially excludable telomeric extreme of the interval. II.9 had inherited the maternal disease allele from *rs6879694* to *rs12110022*, but a different maternal allele to his affected siblings at *rs11958187* (within *PPP2R2B*). This sited the recombination event between 146.1-146.4 Mb, excluding a further 3 Ensembl interval genes; *JAKMIP2*, *DPYSL*, and *STK32A* (Figure 4).

Discussion

Identification of further HHT genes is predicted to provide new insights into TGF- β /BMP signalling and angiogenesis pathways. The previously published *HHT3* interval included several highly attractive candidate

genes. As further evidence accumulated regarding the roles of particularly VE-cadherin and Sprouty family members, the possibility that one of these may be mutated in HHT increased their interest to endothelial cell biologists. While these genes had been sequenced with no disease-causing mutations found in the coding or untranslated regions in affected members of the chromosome 5 linked family ([35], Govani and Shovlin unpublished), such sequencing did not exclude the genes as disease-causing, due to the possibility of undetected mutations in intronic or poorly characterised regulatory regions.

It was recognised that fine mapping of the genomic variants found during candidate gene sequencing could help define the critical recombination events and exclude certain genes, including those for VE-cadherin-2 and Sprouty4. Hence candidate gene sequence variants were examined in the interval-defining members of the

Family S pedigree that included the crucial individuals whose recombination events had defined the centromeric and telomeric borders of the *HHT3* interval.

To optimise PCR fidelity, candidate genes and genomic sequence variants were amplified with *AmpliTaq Gold* or *Phusion Hotstart* DNA polymerases, enzymes that are inactive until they undergo heat activation to generate polymerase activity, thus providing an automated 'hot start' [52-54]. To further confirm key genotypes, SNPs *rs153148*, *rs34074000*, *rs164075*, *rs164077*, *rs6879694*, *rs12110022* and *rs11958187* were sequenced in two separate PCR reactions in two or more key individuals.

Confirmed genotypes allowed construction of haplotypes across the centromeric and telomeric extremes of the *HHT3* interval. The recombination events were then sited more precisely, leading to findings which excluded 12 interval genes such as strong candidates *PCDH12*, *SPRY4* and *STK32A*. *FGF1* and *PPP2R2B* could not be fully excluded as the centromeric and telomeric defining recombination events (SNPs 'novel' and *rs11958187*) are within the introns of both genes. It should be noted that the siting of a recombination breakpoint in a single affected member of the pedigree does not carry any pathogenic implications for the genes in question, unlike disease-causing chromosomal rearrangements when the causative rearrangement is shared by all of the affected individuals in the pedigree.

The original interval contained 38 genes according to the Ensembl genome browser. Twelve have been excluded in this study. The 26 genes that remain include 10 pseudogenes or processed transcripts with no known protein products; two genes of potential functional relevance to angiogenesis or TGF- β signalling (*FGF1* and *PPP2R2B*); three protein coding genes associated with other genetic disease (*ARHGAP26*; *NR3C1* and *POU4F3*) and 11 protein coding genes not associated with a genetic disease but whose gene product had no obvious function in TGF- β signalling and/or angiogenesis (*HMHB1*; *YIPF5*; *KCTD16*; *PRELID2*; *LARS*; *TCERG1*; *SH3RF2*; *GRXCR2*; *RBM27*; *PLAC8L1* and *GPR151*).

Conclusions

Further examination of key recombination events in a known *HHT3* pedigree has reduced the published *HHT3* gene interval. Crucially, the *PCDH12* and *SPRY4* genes were amongst twelve genes no longer in the reduced interval. Therefore VE-cadherin-2 and Sprouty4, proteins of significant interest to angiogenesis researchers due to their respective roles in pathophysiological angiogenesis and relationship to known interactors with the HHT gene products, are not mutated in *HHT3* [55].

Reduction of the interval has facilitated *HHT3* gene identification studies.

Abbreviations

HHT: hereditary haemorrhagic telangiectasia; TGF- β : transforming growth factor-beta; ALK: activin receptor-like kinase; AVMS: arteriovenous malformations; SNP: single nucleotide polymorphism; PCR: polymerase chain reaction; ECS: endothelial cells.

Acknowledgements

This work was funded by the British Heart Foundation. We are also grateful for support from the NIHR Biomedical Research Centre Funding Scheme. The study sponsors played no part in study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication. The authors thank Dr Gillian Wallace and Dr Stewart Cole for DNA extractions; Dr Mick Jones and Dr Carol Shoulders for helpful discussions; and the family for their willing participation in these studies.

Authors' contributions

CLS and FSG designed the study. All experimental work was performed by FSG, with advice from CLS. The manuscript was co-written by CLS and FSG; both authors approved the final version.

Competing interests

The authors declare that they have no competing interests.

Received: 8 June 2010 Accepted: 11 August 2010

Published: 11 August 2010

References

1. Massague J, Gomis RR: The logic of TGFbeta signaling. *FEBS Lett* 2006, **580**(12):2811-2820.
2. Piek E, Heldin CH, Ten Dijke P: Specificity, diversity, and regulation in TGF-beta superfamily signaling. *FASEB J* 1999, **13**(15):2105-2124.
3. Daly AC, Randall RA, Hill CS: Transforming growth factor beta-induced Smad1/5 phosphorylation in epithelial cells is mediated by novel receptor complexes and is essential for anchorage-independent growth. *Mol Cell Biol* 2008, **28**(22):6889-6902.
4. Park SO, Lee YJ, Seki T, Hong KH, Fliess N, Jiang Z, Park A, Wu X, Kaartinen V, Roman BL, Oh SP: ALK5- and TGFBR2-independent role of ALK1 in the pathogenesis of hereditary hemorrhagic telangiectasia type 2. *Blood* 2008, **111**(2):633-642.
5. Upton PD, Davies RJ, Trembath RC, Morrell NW: Bone morphogenetic protein (BMP) and activin type II receptors balance BMP9 signals mediated by activin receptor-like kinase-1 in human pulmonary artery endothelial cells. *J Biol Chem* 2009, **284**(23):15794-15804.
6. Goumans MJ, Valdimarsdottir G, Itoh S, Lebrin F, Larsson J, Mummery C, Karlsson S, ten Dijke P: Activin receptor-like kinase (ALK)1 is an antagonistic mediator of lateral TGFbeta/ALK5 signaling. *Mol Cell* 2003, **12**(4):817-828.
7. Oh SP, Seki T, Goss KA, Imamura T, Yi Y, Donahoe PK, Li L, Miyazono K, ten Dijke P, Kim S, Li E: Activin receptor-like kinase 1 modulates transforming growth factor-beta 1 signaling in the regulation of angiogenesis. *Proc Natl Acad Sci USA* 2000, **97**(6):2626-2631.
8. Pece-Barbara N, Vera S, Kathirkamathamby K, Liebner S, Di Guglielmo GM, Dejana E, Wrana JL, Letarte M: Endoglin null endothelial cells proliferate faster and are more responsive to transforming growth factor beta1 with higher affinity receptors and an activated Alk1 pathway. *J Biol Chem* 2005, **280**(30):27800-27808.
9. Govani FS, Shovlin CL: Hereditary haemorrhagic telangiectasia: a clinical and scientific review. *Eur J Hum Genet* 2009, **17**(7):860-871.
10. McAllister KA, Grogg KM, Johnson DW, Gallione CJ, Baldwin MA, Jackson CE, Helmbold EA, Markel DS, McKinnon WC, Murrell J, McCormick MK, Pericak-Vance MA, Heutink P, Oostra BA, Haitjema T, Westermann CJJ, Porteous ME, Guttmacher AE, Letarte M, Marchuk DA: Endoglin, a TGF-beta binding protein of endothelial cells, is the gene for hereditary haemorrhagic telangiectasia type 1. *Nat Genet* 1994, **8**(4):345-351.

11. Johnson DW, Berg JN, Baldwin MA, Gallione CJ, Marondel I, Yoon SJ, Stenzel TT, Speer M, Pericak-Vance MA, Diamond A, Guttmacher AE, Jackson CE, Attisano L, Kucherlapati R, Porteous ME, Marchuk DA: **Mutations in the activin receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia type 2.** *Nat Genet* 1996, **13**(2):189-195.
12. Gallione CJ, Repetto GM, Legius E, Rustgi AK, Schelley SL, Tejpar S, Mitchell G, Drouin É, Westermann CJJ, Marchuk DA: **A combined syndrome of juvenile polyposis and hereditary haemorrhagic telangiectasia associated with mutations in MADH4 (SMAD4).** *The Lancet* 2004, **363**(9412):852-859.
13. ten Dijke P, Goumans MJ, Pardali E: **Endoglin in angiogenesis and vascular diseases.** *Angiogenesis* 2008, **11**(1):79-89.
14. David L, Feige JJ, Bailly S: **Emerging role of bone morphogenetic proteins in angiogenesis.** *Cytokine Growth Factor Rev* 2009, **20**(3):203-212.
15. Shovlin CL, Guttmacher AE, Buscarini E, Faughnan ME, Hyland RH, Westermann CJJ, Kjeldsen AD, Plauchu H: **Diagnostic Criteria for Hereditary Hemorrhagic Telangiectasia (Rendu-Osler-Weber Syndrome).** *American Journal of Medical Genetics* 2000, **91**(1):66-67.
16. Bourdeau A, Dumont DJ, Letarte M: **A murine model of hereditary hemorrhagic telangiectasia.** *J Clin Invest* 1999, **104**(10):1343-1351.
17. Li DY, Sorensen LK, Brooke BS, Urness LD, Davis EC, Taylor DG, Boak BB, Wendel DP: **Defective angiogenesis in mice lacking endoglin.** *Science* 1999, **284**(5419):1534-1537.
18. Arthur HM, Ure J, Smith AJ, Renforth G, Wilson DL, Torsney E, Charlton R, Parums DV, Jowett T, Marchuk DA, Burn J, Diamond AG: **Endoglin, an ancillary TGFbeta receptor, is required for extraembryonic angiogenesis and plays a key role in heart development.** *Dev Biol* 2000, **217**(1):42-53.
19. Urness LD, Sorensen LK, Li DY: **Arteriovenous malformations in mice lacking activin receptor-like kinase-1.** *Nat Genet* 2000, **26**(3):328-331.
20. Srinivasan S, Hanes MA, Dickens T, Porteous ME, Oh SP, Hale LP, Marchuk DA: **A mouse model for hereditary hemorrhagic telangiectasia (HHT) type 2.** *Hum Mol Genet* 2003, **12**(5):473-482.
21. Roman BL, Pham VN, Lawson ND, Kulik M, Childs S, Lekven AC, Garrity DM, Moon RT, Fishman MC, Lechleider RJ, Weinstein BM: **Disruption of acvrl1 increases endothelial cell number in zebrafish cranial vessels.** *Development* 2002, **129**(12):3009-3019.
22. Park SO, Wankhede M, Lee YJ, Choi EJ, Fliess N, Choe SW, Oh SH, Walter G, Raizada MK, Sorg BS, Oh SP: **Real-time imaging of de novo arteriovenous malformation in a mouse model of hereditary hemorrhagic telangiectasia.** *J Clin Invest* 2009, **119**(11):3487-3496.
23. Mahmoud M, Allinson KR, Zhai Z, Oakenfull R, Ghandi P, Adams RH, Fruttiger M, Arthur HM: **Pathogenesis of arteriovenous malformations in the absence of endoglin.** *Circ Res* 2010, **106**(8):1425-1433.
24. Mitchell A, Adams LA, MacQuillan G, Tibballs J, vanden Driesen R, Delriviere L: **Bevacizumab reverses need for liver transplantation in hereditary hemorrhagic telangiectasia.** *Liver Transpl* 2008, **14**(2):210-213.
25. Lebrin F, Srun S, Raymond K, Martin S, van den Brink S, Freitas C, Breant C, Mathivet T, Larrivee B, Thomas JL, Arthur HM, Westermann CJ, Disch F, Mager JJ, Snijder RJ, Eichmann A, Mummery CL: **Thalidomide stimulates vessel maturation and reduces epistaxis in individuals with hereditary hemorrhagic telangiectasia.** *Nat Med* 2010, **16**(4):420-428.
26. Shovlin CL: **Hereditary haemorrhagic telangiectasia: Pathogenesis, diagnosis and treatment.** *Blood Reviews* 2010.
27. Sato Y, Rifkin DB: **Inhibition of endothelial cell movement by pericytes and smooth muscle cells: activation of a latent transforming growth factor-beta 1-like molecule by plasmin during co-culture.** *J Cell Biol* 1989, **109**(1):309-315.
28. Mancini ML, Terzic A, Conley BA, Oxburgh LH, Nicola T, Vary CP: **Endoglin plays distinct roles in vascular smooth muscle cell recruitment and regulation of arteriovenous identity during angiogenesis.** *Dev Dyn* 2009, **238**(10):2479-2493.
29. Carvalho RL, Itoh F, Goumans MJ, Lebrin F, Kato M, Takahashi S, Ema M, Itoh S, van Rooijen M, Bertolino P, Ten Dijke P, Mummery CL: **Compensatory signalling induced in the yolk sac vasculature by deletion of TGFbeta receptors in mice.** *J Cell Sci* 2007, **120**(Pt 24):4269-4277.
30. Letarte M, McDonald ML, Li C, Kathirkamathambay K, Vera S, Pece-Barbara N, Kumar S: **Reduced endothelial secretion and plasma levels of transforming growth factor-beta1 in patients with hereditary hemorrhagic telangiectasia type 1.** *Cardiovasc Res* 2005, **68**(1):155-164.
31. Richards-Yutz J, Grant K, Chao EC, Walther SE, Ganguly A: **Update on molecular diagnosis of hereditary hemorrhagic telangiectasia.** *Hum Genet* 2010, **128**(1):61-77.
32. Bayrak-Toydemir P, McDonald J, Mao R, Phansalkar A, Gedge F, Robles J, Goldgar D, Lyon E: **Likelihood ratios to assess genetic evidence for clinical significance of uncertain variants: hereditary hemorrhagic telangiectasia as a model.** *Exp Mol Pathol* 2008, **85**(1):45-49.
33. McDonald J, Gedge F, Burdette A, Carlisle J, Bukjok CJ, Fox M, Bayrak-Toydemir P: **Multiple sequence variants in hereditary hemorrhagic telangiectasia cases: illustration of complexity in molecular diagnostic interpretation.** *J Mol Diagn* 2009, **11**(6):569-575.
34. Gallione C, Aylsworth AS, Beis J, Berk T, Bernhardt B, Clark RD, Clericuzio C, Danesino C, Drautz J, Fahl J, Fan Z, Faughnan ME, Ganguly A, Garvie J, Henderson K, Kini U, Leedom T, Ludman M, Lux A, Maisenbacher M, Mazzucco S, Olivieri C, Ploos van Amstel JK, Prigoda-Lee N, Peyeritz RE, Reardon W, Vandezande K, Waldman JD, White RJ, Williams CA, Marchuk DA: **Overlapping spectra of SMAD4 mutations in juvenile polyposis (JP) and JP-HHT syndrome.** *Am J Med Genet A* 2010, **152A**(2):333-339.
35. Cole SG, Begbie ME, Wallace GMF, Shovlin CL: **A new locus for hereditary haemorrhagic telangiectasia (HHT3) maps to chromosome 5.** *Journal of Medical Genetics* 2005, **42**:577-582.
36. Bayrak-Toydemir P, McDonald J, Akarsu N, Toydemir RM, Calderon F, Tuncali T, Tang W, Miller F, Mao R: **A fourth locus for hereditary hemorrhagic telangiectasia maps to chromosome 7.** *Am J Med Genet A* 2006, **140**(20):2155-2162.
37. Hayrabedian S, Kyurkchiev S, Kehayov I: **FGF-1 and S100A13 possibly contribute to angiogenesis in endometriosis.** *J Reprod Immunol* 2005, **67**(1-2):87-101.
38. Mayer RE, Hendrix P, Cron P, Matthies R, Stone SR, Goris J, Merlevede W, Hofsteenge J, Hemmings BA: **Structure of the 55-kDa regulatory subunit of protein phosphatase 2A: evidence for a neuronal-specific isoform.** *Biochemistry* 1991, **30**(15):3589-3597.
39. Manning G, Whyte DB, Martinez R, Hunter T, Sudarsanam S: **The protein kinase complement of the human genome.** *Science* 2002, **298**(5600):1912-1934.
40. Ludwig D, Lorenz J, Dejana E, Bohlen P, Hicklin DJ, Witte L, Pytowski B: **cDNA cloning, chromosomal mapping, and expression analysis of human VE-Cadherin-2.** *Mamm Genome* 2000, **11**(11):1030-1033.
41. Fernandez LA, Garrido-Martin EM, Sanz-Rodriguez F, Pericacho M, Rodriguez-Barbero A, Eleno N, Lopez-Novoa JM, Duwell A, Vega MA, Bernabeu C, Botella LM: **Gene expression fingerprinting for human hereditary hemorrhagic telangiectasia.** *Hum Mol Genet* 2007, **16**(13):1515-1533.
42. Wallez Y, Vilgrain I, Huber P: **Angiogenesis: the VE-cadherin switch.** *Trends Cardiovasc Med* 2006, **16**(2):55-59.
43. Rudini N, Felici A, Giampietro C, Lampugnani M, Corada M, Swirsding K, Garre M, Liebner S, Letarte M, ten Dijke P, Dejana E: **VE-cadherin is a critical endothelial regulator of TGF-beta signalling.** *Embo J* 2008, **27**(7):993-1004.
44. Cabrita MA, Christofori G: **Sprouty proteins: antagonists of endothelial cell signaling and more.** *Thromb Haemost* 2003, **90**(4):586-590.
45. Lee SH, Schloss DJ, Jarvis L, Krasnow MA, Swain JL: **Inhibition of angiogenesis by a mouse sprouty protein.** *J Biol Chem* 2001, **276**(6):4128-4133.
46. Taniguchi K, Ayada T, Ichiyama K, Kohno R, Yonemitsu Y, Minami Y, Kikuchi A, Maehara Y, Yoshimura A: **Sprouty2 and Sprouty4 are essential for embryonic morphogenesis and regulation of FGF signaling.** *Biochem Biophys Res Commun* 2007, **352**(4):896-902.
47. Taniguchi K, Ishizaki T, Ayada T, Sugiyama Y, Wakabayashi Y, Sekiya T, Nakagawa R, Yoshimura A: **Sprouty4 deficiency potentiates Ras-independent angiogenic signals and tumor growth.** *Cancer Sci* 2009, **100**(9):1648-1654.
48. Kim JB, Blackshaw S: **One-step enzymatic purification of PCR products for direct sequencing.** *Curr Protoc Hum Genet* 2001, Chapter 11, Unit 11.16.
49. Schwarz H, Whitton JL: **A rapid, inexpensive method for eluting DNA from agarose or acrylamide gel slices without using toxic or chaotropic materials.** *Biotechniques* 1992, **13**(2):205-206.

50. Moore D: Purification and concentration of DNA from aqueous solutions. *Curr Protoc Immunol* 2001, Chapter 10, Unit 10 11.
51. Bray PJ, Cotton RG: Variations of the human glucocorticoid receptor gene (NR3C1): pathological and in vitro mutations and polymorphisms. *Hum Mutat* 2003, **21**(6):557-568.
52. Moretti T, Koons B, Budowle B: Enhancement of PCR amplification yield and specificity using AmpliTaq Gold DNA polymerase. *Biotechniques* 1998, **25**(4):716-722.
53. Phusion™ Hot Start High-Fidelity DNA Polymerase (manual). [<http://www.neb.com/nebecomm/ManualFiles/manualF-540.pdf>].
54. Chou Q, Russell M, Birch DE, Raymond J, Bloch W: Prevention of pre-PCR mis-priming and primer dimerization improves low-copy-number amplifications. *Nucleic Acids Res* 1992, **20**(7):1717-1723.
55. Govani FS: Regulation of endothelial cell gene expression: insights from HHT3 candidate gene studies. *PhD thesis* Imperial College London, NHLI Cardiovascular Sciences department 2009.

doi:10.1186/2040-2384-2-15

Cite this article as: Govani and Shovlin: Fine mapping of the hereditary haemorrhagic telangiectasia (*HHT*)3 locus on chromosome 5 excludes *VE-Cadherin-2*, *Sprouty4* and other interval genes. *Journal of Angiogenesis Research* 2010 **2**:15.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

