

# Ultrasound Guided Low Approach Interscalene Brachial Plexus Block for Upper Limb Surgery

Department of Anesthesiology and Pain Medicine, Jeju National University Hospital, Jeju,  
\*Ulsan University Hospital, University of Ulsan College of Medicine, Ulsan, Korea

Sun Kyung Park, Min Ha Sung\*, Hae Jin Suh, and Yun Suk Choi

**Background:** The interscalene brachial plexus block is widely used for pain control and anesthetic purposes during shoulder arthroscopic surgeries and surgeries of the upper extremities. However, it is known that interscalene brachial plexus block is not appropriate for upper limb surgeries because it does not affect the lower trunk (C8–T1, ulnar nerve) of the brachial plexus.

**Methods:** A low approach, ultrasound-guided interscalene brachial plexus block (LISB) was performed on twenty-eight patients undergoing surgery of the upper extremities. The patients were assessed five minutes and fifteen minutes after the block for the degree of block in each nerve and muscle as well as for any complications.

**Results:** At five minutes and fifteen minutes after the performance of the block, the degree of the block in the ulnar nerve was found to be  $2.8 \pm 2.6$  and  $1.1 \pm 1.8$ , respectively, based on a ten-point scale. Motor block occurred in the median nerve after fifteen minutes in 26 of the 28 patients (92.8%), and in all of the other three nerves in all 28 patients. None of the patients received additional analgesics, and none experienced complications.

**Conclusions:** The present study confirmed the achievement of an appropriate sensory and motor block in the upper extremities, including the ulnar nerve, fifteen minutes after LISB, with no complications. (Korean J Pain 2016; 29: 18-22)

**Key Words:** Complication; Interscalene brachial plexus block; Nerve stimulator; Sensory block; Ultrasound-guided; Upper limb surgery.

## INTRODUCTION

The brachial plexus runs from the C5–T1 ventral rami, forms the superior, middle, and inferior trunks, divides under the clavicle, leads to the lateral, posterior, and medial

cords, and finally forms the peripheral nerves running to the arms. The brachial plexus block is popular for anesthetic and pain control purposes in the upper limbs. There are a few approaches to the block, including the interscalene approach, supraclavicular approach, infraclavicular approach, and axillary approach.

Received September 3, 2015. Revised October 8, 2015. Accepted October 16, 2015.

Correspondence to: Yun Suk Choi

Department of Anesthesiology and Pain Medicine, Jeju National University Hospital, 15 Aran 13-gil, Jeju 63241, Korea

Tel: +82-64-717-2025, Fax: +82-64-717-2042, E-mail: solafide5@yahoo.co.kr

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


Copyright © The Korean Pain Society, 2016

vicular approach, and the axillary approach. The classic interscalene approach has been performed at the C6 level located in the cricoid cartilage and is useful in controlling pain after shoulder surgeries. However, this approach spares the lower trunk of the brachial plexus (C8–T1, ulnar nerve), thus it is not appropriate for some upper limb surgeries [1]. Some studies have reported findings regarding the efficacy of the interscalene brachial plexus block with approaches from lower regions in certain upper limb surgeries; however, there have been no studies that have reported this method being used with ultrasound. Therefore, the present study first divided the distance between C6 and the clavicle into three sections. Then we performed a low approach, ultrasound-guided interscalene brachial plexus block (LISB) on the interscalene groove that is located at a site about two-thirds of the distance caudally from C6 to examine the efficacy of the nerve and motor block at five and fifteen minutes after the block, and monitored for any signs of complications.

## MATERIALS AND METHODS


The present study was conducted on twenty-eight patients who were classified as physical status 1 or 2 by the American Society of Anesthesiologists (ASA) and who were scheduled to receive an upper limb surgery at our hospital between May 2013 and April 2014. We excluded patients with coagulation disorders, those who were under the age of 18 or over the age of 75, those who weighed less than 50 kg or more than 100 kg, patients with any kind of neurological deficit, or patients with surgical site infections.

After approval from the clinical research ethical review board at our hospital, we explained to the patients the objective of the study as well as potential risks and complications of the procedure. The study proceeded following consent from the patients (IRB number: 2013-02-013). The patients were not premedicated prior to receiving anesthesia. After they arrived in the operating room, patients were connected to a non-invasive blood pressure manometer, pulse oximeter, and electrocardiogram to monitor their vital signs every three minutes. The patients were in the supine position with their heads facing away from the side of the block. The region was prepped with betadine and the linear probe of the ultrasound (SonoSite M-Turbo, SonoSite, Inc., Bothell, WA, USA) was placed on the interscalene groove, which is located at about two-thirds of the distance caudally from C6 when the distance between C6 and the clavicle is divided into three sections, as suggested by Kim et al. [2] (Fig. 1). Sterilized plastic wrap and gels, and 22G, 50 mm needles (UniPlex NanoLine facet tip UP 3/50, Pajunk Medical Produkte GmbH, Geisingen, Germany) were used. With the help of a nerve stimulator (Stimuplex-S, B. Braun Melsungen AG, Melsungen, Germany) at 0.7 mA, the contraction of each muscle (pectoralis, deltoid, arm, forearm or hand) was confirmed (Fig. 2). For the local anesthetic, 40 ml of 1.5% lidocaine, including 1 : 200,000 epinephrine was used. At five and fifteen minutes after the injection of the local anesthetic, we confirmed the sensory block using alcohol wipes on the musculocutaneous nerve, median nerve, radial nerve, and ulnar nerve with a scale ranging from 0 (no sensation) to 10 (normal sensation) [2]. We also checked



**Fig. 1.** Anatomical landmark (A) and ultrasound probe and block needle position during ultrasound-guided low approach interscalene brachial plexus block (B).

for muscular contractions by assessing flexion of the elbow (musculocutaneous nerve), extension of the elbow and wrist (radial nerve), pronation of the arm and flexion of the wrist (median nerve), and flexion and opposition of the fourth and fifth fingers toward the thumb (ulnar nerve), and considered signs of paralysis (loss of contraction) to



**Fig. 2.** Ultrasound image showing low approach interscalene brachial plexus block. Needle pathway (arrows), SCM: sternocleidomastoid muscle, IJV: internal jugular vein, ST: superior trunk of brachial plexus, MSM: middle scalene muscle.

**Table 1.** Patient Demographic Data

	n = 28
Age (yr)	47.8 ± 14.6
Height (cm)	160.0 ± 8.6
Weight (kg)	59.7 ± 10.6
Sex (M/F)	10/18
Operation time (min)	59.6 ± 33.0
Block performance time (sec)	341.7 ± 59.2

Values are mean ± standard deviation or number of patients.

**Table 2.** Type of Surgery

Type of surgery	N=28
Elbow curettage & drilling	8
Distal radius ORIF	6
Finger ORIF	3
Ulnar metal removal	3
Ganglion excision	3
Tendon reposition	3
Arthroscopic debridement, wrist	1
Tenosynovectomy	1

Values are number of patients. ORIF: open reduction & internal fixation.

indicate a successful motor block [3]. One anesthesiologist performed the LISB procedure and one orthopedist performed the surgery. We confirmed cases of hemidiaphragmatic paralysis after the surgery by performing a chest X-ray and consulting a radiologist regarding the results.

## RESULTS

Among the 28 subjects of this study, 10 were male and 18 were female. The patients' demographic and clinical data including age, body weight, height, gender, surgery length, and type of surgery are illustrated in **Tables 1 and 2**. At five and fifteen minutes after the block procedure, the degree of sensory block in the ulnar nerve was found to be  $2.8 \pm 2.6$  and  $1.1 \pm 1.8$ , respectively, on a scale of ten. Muscular block occurred in the median nerve after fifteen minutes in 26 of the 28 patients (92.8%), and in all of the other three nerves in all 28 patients (**Tables 3 and 4**). None of the patients received additional analgesics after the surgery, and there were no abnormalities during the surgeries. In addition, there were no signs of complica-

**Table 3.** Characteristics of the Sensory Block Using a Low Approach Interscalene Brachial Plexus Block

	5 min after injection	15 min after injection
Sensory block (0-10)		
Musculocutaneous nerve	0.7 ± 1.3	0.2 ± 0.6
Median nerve	1.6 ± 2.3	0.9 ± 1.9
Radial nerve	0.5 ± 1.1	0.1 ± 0.3
Ulnar nerve	2.8 ± 2.6	1.1 ± 1.8

Values are mean ± standard deviation. Sensory block (0-10); 0: loss of sensation, 10: normal sensation.

**Table 4.** Characteristics of the Motor Block Using a Low Approach Interscalene Brachial Plexus Block

	5 min after injection No. (%)	15 min after injection No. (%)
Musculocutaneous nerve	26 (92.8%)	28 (100%)
Median nerve	20 (71.4%)	26 (92.8%)
Radial nerve	25 (89.2%)	28 (100%)
Ulnar nerve	23 (82.1%)	28 (100%)

Values are number of patients (percentage).

**Table 5.** Analgesic Requirement and Complications after a Low Approach Interscalene Brachial Plexus Block

	N = 28
Patients requiring analgesics during operation	0
Patients with complications	
Nausea	0
Horner syndrome	0
Dyspnea	0

Values are number of patients.

tions, such as dyspnea or Horner syndrome, during the surgery, in the recovery room, or in the wards (**Table 5**).

## DISCUSSION

This study confirmed that an appropriate sensory and motor block was achieved in the upper extremities, including the ulnar nerve, fifteen minutes after LISB, and that there were no complications associated with the block.

Two methods of LISB have been introduced, namely the anatomical landmark approach and the ultrasound-guided approach. There are three approaches to the anatomical landmark method: first, it can be performed in between the cricoid cartilage and the clavicle. Second, it can be performed 2 cm above the clavicle, and third, it can be performed on the interscalene groove, which is located at about two-thirds of the distance caudally from C6 after dividing the distance between C6 and the clavicle into three sections. There are two methods for the ultrasound-guided approach: first, it can be performed on the superior trunk where C5 and C6 are combined. Second, the injection can be performed on the caudal side of the C6 nerve root [2,4,5]. Although different studies define and name the procedures slightly differently (low approach, lower interscalene approach, or superior trunk approach), these procedures are identical in terms of using an approach through the lower regions of the C6 level compared to the existing ISB.

Owing to the advances in procedural techniques and the application of ultrasound technology, several studies have reported that the use of local anesthetics can be reduced when performing ISB. However, most of these studies limit their scope to shoulder surgeries, which are irrelevant to the block of the inferior trunk (C8–T1, ulnar nerve) [6–8]. It has been known that the ulnar nerve is not af-

ected in about 30–50% of ISBs performed with the classic approach [1]. Furthermore, it has been reported that the ulnar nerve was not blocked in about 7–33.3% of the cases that took the classic approach ISB and used 30 ml of local anesthetics on the caudal side of the C6 nerve root [9,10]. Hence, in the present study, we used 40 ml of local anesthetics to ensure a quick onset and complete block of the ulnar nerve.

LISB is known to involve a short effect distance (from the C5 nerve root to the C8 nerve root) and to diffuse local anesthetics via the deep cervical fascia. In addition, LISB has been reported to bring about appropriate sensory and motor blocks required for upper limb surgeries even with a single injection [10]. Against this backdrop, the present study was planned and conducted.

Moreover, according to Plante et al., [9] who compared two groups of patients who were injected with local anesthetics either in the upper region of the C5 nerve root or the lower region of the C6 nerve root during an ultrasound-guided interscalene brachial plexus block for analgesia in arthroscopic shoulder surgeries, the group of patients who received local anesthetics in the lower region of the C6 nerve root had appropriate sensory blocks in all of the nerves; they reported that the sensory and motor blocks were especially noticeable in the ulnar nerve and that there was a rapid onset in the ulnar nerve. The results of this study showed that the sensory block in the ulnar nerve was  $2.8 \pm 2.6$  on a 10-point scale at five minutes after LISB and  $1.1 \pm 1.8$  at fifteen minutes after LISB. About 82.1% of motor neurons were blocked at five minutes, but 100% were blocked after fifteen minutes. In addition, there were no additional injections of analgesics during the surgery. Therefore, it can be said that the ulnar nerve was appropriately blocked via LISB.

Meanwhile, ISB is known to induce a temporary paralysis in the ipsilateral hemidiaphragm due to phrenic nerve palsy. The phrenic nerve is located within 2 mm of the brachial plexus of the cricoid cartilage and divides 3 mm per 1 cm as it descends caudally. Thus, it can be predicted that the incidence of phrenic nerve palsy-induced hemidiaphragmatic paralysis can be reduced if ISB is performed more caudal to the C6 level or on the superior trunk [11, 12]. In the present study, there were no signs of dyspnea or hemidiaphragmatic paralysis. In addition, LISB is known to reduce the damage to the dorsal scapular and long thoracic nerves, both of which split from the C5 nerve root

[4]. Although we did not assess whether any such damages occurred in the present study, none of the patients experienced any such problems.

As mentioned above, there were no complications in the current study. We presume that we were able to reduce the risk of complications, such as vascular injection or nerve injury, by using an ultrasound nerve stimulator in addition to the inherent merits of the LISB method.

In the present study, the motor block in the median nerve was shown to be about 71.4% at five minutes after the procedure was performed. The block increased to 92.8% at fifteen minutes, and there were no additional analgesics injected and no additional block was performed. Other studies have also reported similarly slow blocks in the median nerve within fifteen minutes [9], which is thought to be due to the fact that the median nerve innervates all of C5, C6, C7, and T1.

There are a few limitations to this study. First, we did not have a large pool of subjects. Second, we could not observe the diffusion of local anesthetics through injecting contrast medium. Third, we did not compare the procedure of interest with other approaches. Thus far, studies on LISB are only in the form of case reports or brief reports; hence, in the future, LISB should be compared with other approaches, and cases of LISB using different doses of local anesthetics should be compared as well.

In conclusion, the present study confirmed that the nerves in the upper extremities, including the ulnar nerve, were appropriately blocked fifteen minutes after the performance of LISB, and that there were no complications induced by the block.

## ACKNOWLEDGEMENTS

This work was supported by the research grant of Jeju National University in 2013.

## REFERENCES

1. Neal JM, Gerancher JC, Hebl JR, Ilfeld BM, McCartney CJ, Franco CD, et al. Upper extremity regional anesthesia: essentials of our current understanding, 2008. *Reg Anesth Pain Med* 2009; 34: 134–70.
2. Kim JH, Chen J, Bennett H, Lesser JB, Resta-Flarer F, Barczewska-Hillel A, et al. A low approach to interscalene brachial plexus block results in more distal spread of sensory–motor coverage compared to the conventional approach. *Anesth Analg* 2011; 112: 987–9.
3. Liu FC, Liou JT, Tsai YF, Li AH, Day YY, Hui YL, et al. Efficacy of ultrasound–guided axillary brachial plexus block: a comparative study with nerve stimulator–guided method. *Chang Gung Med J* 2005; 28: 396–402.
4. Burckett–St Laurent D, Chan V, Chin KJ. Refining the ultrasound–guided interscalene brachial plexus block: the superior trunk approach. *Can J Anaesth* 2014; 61: 1098–102.
5. Gadsden JC, Tsai T, Iwata T, Somasundaram L, Robards C, Hadzic A. Low interscalene block provides reliable anesthesia for surgery at or about the elbow. *J Clin Anesth* 2009; 21: 98–102.
6. Renes SH, Rettig HC, Gielen MJ, Wilder–Smith OH, van Geffen GJ. Ultrasound–guided low–dose interscalene brachial plexus block reduces the incidence of hemidiaphragmatic paresis. *Reg Anesth Pain Med* 2009; 34: 498–502.
7. Gautier P, Vandepitte C, Ramquet C, DeCoopman M, Xu D, Hadzic A. The minimum effective anesthetic volume of 0.75% ropivacaine in ultrasound–guided interscalene brachial plexus block. *Anesth Analg* 2011; 113: 951–5.
8. Fredrickson MJ, Smith KR, Wong AC. Importance of volume and concentration for ropivacaine interscalene block in preventing recovery room pain and minimizing motor block after shoulder surgery. *Anesthesiology* 2010; 112: 1374–81.
9. Plante T, Rontes O, Bloc S, Delbos A. Spread of local anesthetic during an ultrasound–guided interscalene block: does the injection site influence diffusion? *Acta Anaesthesiol Scand* 2011; 55: 664–9.
10. Bharti N, Bhardawaj N, Wig J. Comparison of ultrasound–guided supraclavicular, infraclavicular and below–C6 interscalene brachial plexus block for upper limb surgery: a randomised, observer–blinded study. *Anaesth Intensive Care* 2015; 43: 468–72.
11. Nadeau MJ, Lévesque S, Dion N. Ultrasound–guided regional anesthesia for upper limb surgery. *Can J Anaesth* 2013; 60: 304–20.
12. Kessler J, Schafhalter–Zoppoth I, Gray AT. An ultrasound study of the phrenic nerve in the posterior cervical triangle: implications for the interscalene brachial plexus block. *Reg Anesth Pain Med* 2008; 33: 545–50.