

Towards a Reconceptualization of Striatal Interactions Between Glutamatergic and Dopaminergic Neurotransmission and Their Contribution to the Production of Movements

Hélène N. David*

NNOXe Pharmaceuticals, 3107 Avenue des Hôtels, Suite 18C, Québec, QC, G1W 4W5, Canada

Abstract: According to the current model of the basal ganglia organization, simultaneous activation of the striato-nigral direct pathway by glutamatergic and dopaminergic neurotransmission should lead to a synergistic facilitatory action on locomotor activity, while in contrast activation of the indirect pathway by these two neurotransmissions should lead to antagonistic effects on locomotor activity. Based on published data, as a break with the current thinking, we propose a reconceptualization of functional interactions between dopaminergic and glutamatergic neurotransmission. In this model, dopaminergic neurotransmission is seen as a motor pacemaker responsible for the basal and primary activation of striatal output neurons and glutamate as a driver providing a multiple combination of tonic, phasic, facilitatory and inhibitory influxes resulting from the processing of environmental, emotional and mnemonic stimuli. Thus, in the model, glutamate-coded inputs would allow tuning the intrinsic motor-activating properties of dopamine to adjust the production of locomotor activity into goal-oriented movements.

Key Words: Dopamine-glutamate striatal interactions, locomotor activity, goal-oriented movements, nucleus accumbens, striatum.

1. FUNCTIONAL NEUROANATOMY BETWEEN GLUTAMATERGIC AND DOPAMINERGIC NEUROTRANSMISSION IN THE STRIATUM COMPLEX (FIGS. 1 & 2)

The striatum including its dorsal and ventral parts, respectively the caudate-putamen and the nucleus accumbens, is the major input structure of the basal ganglia, which are a set of subcortical structures that is now widely accepted to contribute not only to the control of motor activity and movements but also to the processing of cognitive and 'limbic' (emotional and motivational) functions. While the motor system directly mediates the production of movements, the limbic system is involved in the elaboration of emotions and motivations, which basic drives can influence the production of movements and therefore the pertinence and efficacy of actions and goal-oriented behaviors [39,42]. Within the framework of interactions between the limbic system and the motor system, the nucleus accumbens, which corresponds to the limbic striatum, is considered to be a critical interface between the motor system and the limbic system in the brain. While the shell of the nucleus accumbens is generally thought to be associated more particularly with visceral responses, the core of the nucleus accumbens, which possesses similarities with the dorsal striatum (i.e. the caudate-putamen) both in terms of motor functions and connectivity, is believed to be allied more with motor functions [8,42].

From an anatomofunctional point of view, the striatum complex receives neuronal projections from all areas of the cerebral cortex and from limbic and other brain structures.

These inputs originating from the cerebral cortex, the centro-median and parafascicularis nucleus of the thalamus, the hippocampus, and the amygdala are excitatory and use glutamate as a neurotransmitter. These inputs converge to the limbic and motor striatum where they make synaptic contacts at pre- and postsynaptic levels with striatal spiny output neurons [29,32,69,70] and striatal aspiny GABAergic and cholinergic interneurons [47,52]. In addition to these excitatory inputs, further inputs from outside and from within the striatum provide additional neuronal influxes, the function of which is thought to modify the responsiveness of striatal projection neurons to glutamatergic inputs. Among the influxes originating from outside the striatum are dopaminergic inputs from the substantia nigra pars compacta, the ventral tegmental area, and the retrorubral nucleus [31,61], as well as serotonergic inputs from the raphe nuclei [79]. Further evidence supporting such a dual regulation of striatal projection neurons is that glutamatergic and dopaminergic terminals directly converge on the dendritic spines of the same striatal projection neurons [13,69]. Like striatal output neurons, striatal interneurons are quite evenly distributed within the striatum [41]. Most of them make synapses with striatal projection neurons and thus may be viewed as intermediate relays between neuronal inputs from outside the striatum and striatal projection neurons. In turn, the striatum complex projects back upon the frontal lobe of the cortex *via* direct and indirect topographically organized pathways those which pass through the thalamus to organize goal-oriented movements [3-7,24,67]. Both the direct and indirect pathways utilize the same neuronal cell type, namely the striatal spiny projection neuron that represents approximately 95 % of the neuronal population of the striatum [10,26,44,85]. Spiny output neurons are quite evenly distributed within the striatum complex and are both the major input targets, although not exclusively, and the major output neurons of the

*Address correspondence to this author at NNOXe Pharmaceuticals, 3107 Avenue des Hôtels, Suite 18C, Québec, QC, G1W 4W5, Canada; E-mail: hdavid@cyceron.fr; hn.david@nnoxepharmaceuticals.com




Fig. (1). Basal Ganglia - Dorsal Pathways.

The dorsal striatum receives cortical informations and transfer them to the output structures of the basal ganglia: the entopeduncular nucleus (EN) and the substantia nigra *pars reticulata* (SNR) through the direct and indirect pathway. Then, the information is sent to the frontal cortex through the thalamus. STN: subthalamic nucleus; NAcc: Nucleus Accumbens; SNC: substantia nigra pars compacta; Thal: Thalamus; RR: retrorubral nucleus; GP: Globus Pallidus; VTA: ventral tegmental area. voie glutamatergique ; voie GABAergique; voie dopaminergique.

striatum. Their connections are thus the major determinant of the functional organization of the striatum. Striatal projection neurons all use the inhibitory neurotransmitter γ -aminobutyric acid (GABA). Striatal output neurons of the “direct pathway” project directly from the striatum to the substantia nigra pars reticulata. In contrast, striatal output neurons of the “indirect pathway” connect the striatum to the substantia nigra pars reticulata *via* intermediate synaptic connections in the globus pallidus and the subthalamic nucleus. Projections from the striatum to the globus pallidus and from the globus pallidus to the subthalamic nucleus are GABAergic. Neurons originating from the subthalamic nucleus are glutamatergic, and project to the substantia nigra pars reticulata. Finally, the

substantia nigra pars reticulata, the major output structure of the basal ganglia in rodents, sends GABAergic fibers to the thalamus, which in turn projects back to the motor cortex and other cortical area of the frontal lobe through excitatory glutamatergic fibers.

Despite the dual regulation of striatal output neurons by dopaminergic and glutamatergic inputs, there are very few axo-axonic synapses between glutamatergic and dopaminergic terminals throughout the entire striatum complex [13,31, 66,69,77]. Although this lack of axo-axonic synapses between glutamatergic and dopaminergic terminals limits the possibility of presynaptic interactions in a classical manner,




Fig. (2). Basal Ganglia - Ventral Pathways.

The nucleus accumbens receives cortical informations and transfer them to the output structures of the basal ganglia: the entopeduncular nucleus (EN) and the substantia nigra *pars reticulata* (SNR) through the direct and indirect pathway. Then, the information is sent to the frontal cortex through the thalamus. STN: subthalamic nucleus; NAcc: Nucleus Accumbens; SNC: substantia nigra pars compacta; Thal: Thalamus; RR: retrorubral nucleus; GP: Globus Pallidus; VTA: ventral tegmental area. voie glutamatergique ; voie GABAergique; voie dopaminergique.

dopamine and glutamate spilled over from the synaptic cleft modulate each other's release through diffusion of neurotransmitters away from the synapse, leading to subsequent activation of extrasynaptic heteroreceptors through the so-called volume neurotransmission [69,77], and further interact by modulating in concert the activity of striatal output neurons.

2. STRIATAL MOTOR INTERACTIONS BETWEEN GLUTAMATERGIC AND DOPAMINERGIC NEUROTRANSMISSION: CURRENT THINKING

The current thinking on striatal motor interactions between glutamatergic and dopaminergic neurotransmission has come mainly from the work of Albin *et al.* [3] and DeLong [25] who first described a scheme, known as the model of basal ganglia, which was originally proposed to allow structuring glutamatergic and dopaminergic interactions in the dorsal striatum, i.e. the caudate and the putamen, as regards to the production of locomotor activity and movements. Due to its considerable success, this model has been adopted as a general principle and mode of reasoning and thinking as regards to glutamatergic and dopaminergic striatal motor interactions.

According to the current thinking as regards to striatal motor function, activation by glutamate (Fig. 3) of the striatal GABAergic output neurons that project directly to the substantia nigra pars reticulata (the striato-nigral output neurons that form the direct pathway) is thought to provide a direct inhibitory action on the substantia nigra pars reticulata GABAergic neurons, the inhibition of which in turn leads to a disinhibition of the glutamatergic neurons involved in motor activities that project from the thalamus to the cortex; behaviorally, this allows the production of locomotor activity

and movements. On the contrary, activation of the striato-pallidal GABAergic output neurons, which project indirectly to the substantia nigra pars reticulata through a trisynaptic link and form the so-called indirect pathway, is believed to lead to an inhibition of the GABAergic neurons of the globus pallidus, thereby leading to a disinhibition of the subthalamic nucleus glutamatergic neurons and as a consequence to an activation of the substantia nigra pars reticulata GABAergic neurons that project to the thalamus; overall, this provides an inhibitory effect on thalamo-cortical glutamatergic neurons involved in motor activities and thereby reduces the production of locomotor activity and movements. Although the output activity of the basal ganglia is influenced by the opposing effects of glutamatergic neurotransmission at the direct and indirect pathways, it is assumed that the direct pathway exerts a predominant control on locomotor activity.

Support for this model has been obtained from experiments in non-human primates that have demonstrated that the activation of striato-nigral and striato-pallidal output neurons, respectively, facilitate and suppress the production of motor activity and movements [4,5,25]. In addition, in rodents, further support for the predominant role of striato-nigral output neurons in the control of motor activity has been obtained from pharmacological studies that have consistently shown that an intrastriatal injection of glutamate or glutamate receptor agonists in the caudate-putamen or the nucleus accumbens induces an increase in locomotor activity [11,12,14,15,20,22,23,27,34,36,48,64,68,71,72,75,82,86 but 45,49,65]. However, leading to the conclusion that the role in the control of motor activity of glutamate inputs within the striatum is probably much more complex than that proposed by the model above, other studies have shown that the intrastriatal injection of glutamate receptor antagonists in the




Fig. (3). Current Model of Basal Ganglia - Glutamatergic Control.

caudate-putamen or the nucleus accumbens also results in an increase in locomotor activity [12,15,20,23,27,36,43,50,59, 65,68,75,86 but 18].


The model of basal ganglia organization and motor function also proposes that dopamine modulates glutamatergic inputs by exerting a dual effect on striatal GABAergic output neurons (Fig. 4): exciting striatal output neurons of the direct pathway, which mainly express D1-like receptors, and inhibiting striatal output neurons of the indirect pathway, which mainly express D2-like receptors. The segregation of D1-like- and of D2-like postsynaptic receptors has provided the anatomical basis for the D1-like receptor-mediated excitatory effect of dopamine on striato-nigral neurons that form the direct pathway and the D2-like receptor-mediated inhibitory action of dopamine on striato-pallidal neurons that constitute the first synaptic link of the indirect pathway. According to the anatomical organization of the basal ganglia from the striatum complex to the motor cortex, the model predicts that activation of either D1-like receptors or D2-like postsynaptic receptors leads to the production of locomotor activity as follows: activation by dopamine of the striato-nigral GABAergic output neurons is thought to provide a direct inhibitory action on the substantia nigra pars reticulata GABAergic neurons. This inhibition in turn leads to a disinhibition of the glutamatergic neurons involved in motor activities that project from the thalamus to the cortex and thereby to the production of locomotor activity and movements. Alternatively, activation of the striato-pallidal GABAergic output neurons is believed to result in a disinhibition of the globus pallidus GABAergic neurons, leading to a subsequent inhibition of the subthalamic nucleus glutamatergic neurons and to a reduction in activity of the substantia nigra pars reticulata GABAergic neurons that project to the thalamus. This inhibition in turn finally leads to a disin-

hibition of the glutamatergic neurons involved in motor activities that project from the thalamus to the cortex and thereby to the production of locomotor activity and movements.

A large body of evidence supports a functional synergy between D1-like and D2-like postsynaptic receptors that are mainly located on striato-nigral- and striato-pallidal output neurons, respectively [1,2,46,80,81,84]. At other times, activation of each dopaminergic receptor subtype produces antagonistic effects [57]. Accordingly, the very vast majority of behavioral motor studies has shown: (i) activation by specific agonists of D1-like receptors results in an increase in locomotor activity [18,20-23,28,33,53,54,73,87 but 55,60]; (ii) co-activation of D1-like- and D2-like receptors – a pharmacological condition shown to be required for the full expression of the postsynaptic effects of D2-like receptor agonists because of the enabling role of D1-like receptors on D2-like postsynaptic receptor activation – results in synergistic motor effects leading to a locomotor activity much more greater than that produced by the activation of D1-like receptors alone, thought to be mainly mediated by D2-like postsynaptic receptors [16,18,19,23,28,30,33,37,40,60] and reported as such in the text below; (iii) activation in the caudate-putamen or the nucleus accumbens of D2-like receptors alone – a pharmacological condition thought to reflect mainly D2-like presynaptic receptor mechanisms [38,76,83] results at a very few exceptions [24] in a decrease in locomotor activity [16,18,20-24,33,55,63,71,73 but 51,60]. In addition, support for the model of basal ganglia has been obtained from studies that have consistently shown that blockade of dopaminergic receptors by intrastriatal infusion of D1-like and/or D2-like receptor antagonists resulted in a decrease in locomotor activity [35,53,58,60,78].

Excitatory through the direct pathway

D1R



Excitatory through the indirect pathway

Postsynaptic D2R




Fig. (4). Current Model of Basal Ganglia - Dopaminergic Control.

According to this model, simultaneous activation of glutamatergic receptors and D1-like receptors within the striatum complex should lead to a synergistic facilitatory action on locomotor activity through the direct striato-nigral pathway, while activation of glutamatergic receptors and D2-like receptors should lead to antagonistic effects on locomotor activity through the indirect pathway. However, it is clear from the data summarized in Table 1, obtained from *focal* injection studies performed in the caudate-putamen or the nucleus accumbens of “intact” animals, that an attempt to compel behavioral motor interactions between glutamatergic and dopaminergic neurotransmission into the current model of basal ganglia organization, and an exclusively excitatory role of glutamatergic neurotransmission, would fail to provide a reliable framework of functional interactions between both neurotransmitter systems in the striatum complex.

3. TOWARDS A RECONCEPTUALIZATION OF FUNCTIONAL STRIATAL INTERACTIONS BETWEEN GLUTAMATERGIC AND DOPAMINERGIC NEUROTRANSMISSION AND THEIR ROLE IN THE PRODUCTION OF MOVEMENT

In this section, we will develop particular points that we believe to be important to layout by which glutamatergic and

dopaminergic neurotransmissions may interact within the striatum complex to modulate striatal projection neurons and thereby the production of locomotor activity and movements in “intact” animals. Our analysis is based on an overview of previous published papers that have investigated in non-lesioned animals functional interactions between glutamatergic and dopaminergic neurotransmission in the caudate-putamen and the core of the nucleus accumbens using focal (intra-striatal) injection of glutamate and dopamine receptor ligands. Behavioral systemic injection studies with glutamatergic and dopaminergic neurotransmission were omitted intentionally because this technique is not without drawbacks since the pharmacological effects of systemic injection are diffuse, thereby altering many other brain structures that are involved in motor control throughout the brain.

Modulation by glutamatergic neurotransmission of locomotor responses produced by activation of dopaminergic postsynaptic receptors is in fact very heterogeneous and contrasts with the current model of basal ganglia striatal motor function, which assumes that glutamate exerts a facilitatory and an inhibitory action on locomotor activity through striato-nigral and striato-pallidal projection neurons respectively. Based on the behavioral motor findings obtained from focal injection studies that are summarized in Table 1 and

Table 1. The Nucleus Accumbens: When Modelization Meets Experimental Research

Effect of Glutamate Receptor Ligands on Locomotor Activity Produced by Dopamine Receptor Activation	As Predicted by the bg Model	Experimental Results	References
NMDA receptor activation on D1-like receptor activation	+	+	[23]
NMDA receptor blockade on D1-like receptor activation	-	n.e.	[86]
		-	[23]
NMDA receptor activation on D2-like receptor activation	-	-	[23]
NMDA receptor blockade on D2-like receptor activation	+	-	[23]
Non-NMDA receptor activation on D1-like receptor activation	+	-	[23]
Non-NMDA receptor blockade on D1-like receptor activation	-	-	[23]
		n.e.	[18]
Non-NMDA receptor activation on D2-like receptor activation	-	+	[23]
Non-NMDA receptor blockade on D2-like receptor activation	+	-	[23;40]
Group I mGlu receptor activation on D1-like receptor activation	+	+	[63]
Group I mGlu receptor blockade on D1-like receptor activation	-	-	[20]
Group I mGlu receptor activation on D2-like receptor activation	-	-	[63]
Group I mGlu receptor blockade on D2-like receptor activation	+	n.e.	[20]
Group II mGlu receptor activation on D1-like receptor activation	+	+	[21]
Group II mGlu receptor blockade on D1-like receptor activation	-	n.e.	[21]
Group II mGlu receptor activation on D2-like receptor activation	-	n.e.	[21]
Group II mGlu receptor blockade on D2-like receptor activation	+	-	[21]
Group III mGlu receptor activation on D1-like receptor activation	+	-	[22]
Group III mGlu receptor blockade on D1-like receptor activation	-	+	[63]
Group III mGlu receptor activation on D2-like receptor activation	-	+	[22]
Group III mGlu receptor blockade on D2-like receptor activation	+	+	[63]

from other electrophysiological investigations [17], a more reliable and productive approach may be to fit the effects of glutamatergic neurotransmission into a modulatory, either facilitatory or inhibitory, role in a way that depends on the type of receptor involved. This proposal clearly breaks away from the current model of striatal motor function and the current theory about the role and function of glutamate, which is viewed as an excitatory neurotransmitter that produces activation of striato-nigral and striato-pallidal projection neurons. Though such a modulatory function of glutamatergic neurotransmission within the striatum complex may lead to heated debates and conflicts of opinion, there is now evidence that, depending on the type of receptor involved, activation of glutamatergic receptors can produce either a facilitatory or an inhibitory action (or have no action) on the locomotor responses produced by activation of dopaminergic postsynaptic receptors. Reciprocally, there is now a growing body of evidence that not only dopaminergic neurotransmission modulates glutamate release in the striatum, but that glutamatergic neurotransmission, depending on the type of receptor involved, also exerts a modulatory, either facilitatory or inhibitory, action on striatal dopamine release in *ex vivo* and *in vivo* models [24].

If glutamate is considered as a neurotransmitter which activation can positively and negatively control both striato-nigral and striato-pallidal output neurons, and *in fine* facilitate or inhibit the production of locomotor activity and movements mediated by D1-like receptor- or D2-like postsynaptic receptor activation, it could be predicted that the types of receptors and glutamatergic pathway (e.g. from the prefrontal cortex, the hippocampus or the amygdala) activated would determine the direction in which functional interactions between glutamatergic and dopaminergic neurotransmission occur. However, the mechanisms by which couplings of specific glutamatergic and dopaminergic receptor subtypes occur cannot be foreseen with enough accuracy

at this time, and only hypotheses can be drawn. Coupling possibilities may involve intracellular interactions on common neurons as well as spatial interactions through trans (poly)synaptic mechanisms. For example, co-activation of NMDA and D1-like receptors produces behavioral and electrophysiological synergistic effects while co-activation of NMDA and D2-like receptors produces antagonistic effects. Consistent with intracellular mechanisms, it is well known that NMDA receptors and D1-like receptors tap into the same transduction system that is the adenosine monophosphate cyclic – phosphokinase A transduction system, while D2-like receptors inhibit it. Likewise, also consistent with intracellular mechanisms, activation of group III mGlu receptors inhibits the adenosine monophosphate cyclic – phosphokinase A transduction system and behaviorally opposes and favors the locomotor-activating properties of D1-like receptor activation and D2-like postsynaptic receptor activation, respectively. Another possibility is spatial interactions including volume neurotransmission and trans(poly)synaptic mechanisms, in which glutamatergic and dopaminergic receptors interact with each other distally. This possibility (Fig. 5) should be considered with particular interest when functional interactions between glutamatergic and dopaminergic neurotransmission do not fit with compatible intracellular mechanisms, even if they do fit with the current model of striatal motor function. For example, co-activation of D1-like receptors and group II mGlu receptors produces behavioral motor synergistic effects as predicted by the current model of basal ganglia motor function, but are respectively positively and negatively linked to adenylate cyclase. Therefore, intracellular interactions between these receptors on common neurons do not appear tenable and spatial interactions through volume neurotransmission mechanisms should have to be considered. Consistent with such mechanisms, ultrastructural studies have shown that group II, in contrast with group I and group III, mGlu receptors have no




Fig. (5). What can emerge from functional studies?

close apposition with synapses [74]. Likewise, co-activation of non-NMDA and D1-like receptors produces motor antagonistic effects while co-activation of non-NMDA and D2-like receptors produces synergistic effects. These effects do not fit with the current model of striatal motor function and the fact that non-NMDA, like NMDA, receptors produce postsynaptic excitatory currents at striatal output neurons [56]; therefore, the possibility that glutamatergic and dopaminergic neurotransmission interact spatially through trans (poly)synaptic mechanisms to modulate striatal projection neurons must be considered. The same principles of data analysis and interpretation are also applicable to blockade of glutamatergic receptors. In that way, the importance of investigating the effects of receptor blockade clearly has to be emphasized. Indeed, it is obvious from the data summarized in Table 1 that the effects produced by the activation or blockade of glutamatergic receptors on D1-like receptor- and D2-like pre- and postsynaptic receptor-mediated locomotor responses are not always reciprocal. This may be due to the fact that, for a given subtype of receptors, certain receptors are tonically activated while others are “silent” and can be phasically activated. Indeed, if a receptor is tonically activated, it cannot be further activated by a receptor agonist but can be blocked by a receptor antagonist; likewise, if a receptor is silent, it cannot be further blocked by a receptor antagonist but can be activated phasically by a receptor agonist. We propose that this may provide informations about the neuro-anatomical localization within the striatum complex of the glutamatergic receptors studied, either on phasically activated neurons (PANs), such as the spiny output neurons that exhibit increases in firing in relation to movement, or on tonically activated neurons (TANs) such as the cholinergic interneurons that spike in a tonic and irregular fashion [9].

Because all glutamatergic and dopaminergic receptor subtypes are expressed in striatal projection neurons, we

believe that priority of interpretation must be given to intracellular mechanisms. However, spatial mechanisms must also be considered whether or not intracellular mechanisms explain functional interactions between glutamatergic and dopaminergic neurotransmission. Doubtlessly, such a complex combination of modulatory properties can no longer be regarded as a bulk system where glutamatergic inputs are all considered as excitatory. As a basis for an extended, but not yet complete, model of glutamatergic innervation of striatal output neurons, we propose that depending on their anatomical origin glutamatergic terminals may tap onto specific, and not all, glutamatergic receptor subtypes. Support for this are recent studies [62] that have shown that a reversible inhibition by the local anesthetic lidocaine of the prefrontal cortex, the hippocampus complex or the basolateral amygdala (which are brain structures that project glutamatergic fibers toward the striatum complex) has differential facilitatory and inhibitory effects on hyperlocomotion induced by activation in the nucleus accumbens of D1-like receptors or D2-like postsynaptic receptors that mimic those of glutamatergic receptor activation or blockade.

In addition, according to the current view as regards to dopamine-glutamate interactions in the striatum complex, glutamate is often seen as an excitatory neurotransmitter that is modulated by dopamine. However, in addition of the experimental evidence summarized in Table 1, there are neuro-anatomical arguments to suggest that glutamate may be viewed on the contrary as a modulator, or better a *driver*, of dopaminergic neurotransmission (Fig. 6). Glutamatergic pathways that project to the striatum complex mainly originate from the whole cortex and limbic structures, such as the amygdala and the dorsal and ventral hippocampus. The cortex and limbic structures are well known to play a key role in the processing of extrinsic (environmental) and intrinsic (emotional and mnemonic) stimuli. Dopaminergic pathways that




Fig. (6). Glutamate-coded inputs would allow tuning the intrinsic motor-activating properties of dopamine to adjust the production of locomotor activity into goal-oriented movements.

project to the dorsal and ventral striatum mainly originate from the substantia nigra pars compacta, the ventral tegmental area and retrorubral nucleus, which deeper brain structures are thought to be only slightly involved in the processing of extrinsic and intrinsic stimuli compared to the cortex and limbic structures. Therefore, in contrast with the current thinking of a dopaminergic modulation of glutamatergic neurotransmission, we propose that dopaminergic neurotransmission should be seen as a subcortical-infralimbic motor pacemaker responsible for the basal and primary activation of striatal output neurons, and glutamate as a driver providing facilitatory and inhibitory inputs resulting from the processing of extrinsic and intrinsic stimuli by the cortical areas and limbic structures. These glutamate-mediated facilitatory and inhibitory inputs would allow modulating the motor-activating properties of dopamine at striatal output neurons in order to transform and adjust the production of motor activity into goal-oriented movements. As a result, the relevance and accuracy of goal-oriented movements depend on the appropriate processing of extrinsic (environmental) and intrinsic (emotional and mnemonic) stimuli; and this is the reason why, for instance, strategic processes, perceptive awareness, and learning influence goal-oriented movements. However, because infusion in the striatum complex of either glutamate receptor agonists or antagonists consistently results in an increase of locomotor activity (see above), thereby indicating phasic and tonic facilitatory properties of glutamatergic neurotransmission *per se*, it is likely that the direction of the glutamatergic modulation of dopaminergic neurotransmission would depend on the level of the dopaminergic tone within the striatum complex. In other words, glutamatergic inputs would adapt their inputs as a function of outer and inner factors to the body to modulate dopaminergic neurotransmission and to produce accurate goal-oriented movements.

CONCLUSION

The principal function of oncoming glutamatergic and dopaminergic fibers that innervate the striatum complex is to modulate the activity and responsiveness of striatal output neurons. The present article emphasizes from a behavioral motor point of view the complexity of interactions between glutamatergic and dopaminergic neurotransmission, and further propose a reconceptualization of these interactions in order to induce debate. Clearly, in the future, a great amount of research will be necessary to better define and understand intrastriatal interactions between these neurotransmitter systems. However, we believe that there is enough theoretical and empirical evidence to propose an extended, receptor and pathway-based, model of basal ganglia motor function, in which striatal dopaminergic neurotransmission can be seen as a motor pacemaker responsible for the basal and primary activation of striatal output neurons, and glutamate as a driver providing a multiple combination of tonic, phasic, facilitatory and inhibitory influxes resulting from the processing of extrinsic environmental and intrinsic emotional and mnemonic stimuli. In this framework, and in contrast with the current thinking as regards to glutamate function, glutamate-coded inputs would allow adjusting the intrinsic motor-activating properties of dopamine in order to fine-tune the production of motor activity into goal-oriented movements.

REFERENCES

- [1] Abbraini, J.H., Fechtali, T., Rostain, J-C. (1992) Pressure reversed extracellular striatal dopamine decrease produced by D1 receptor agonist SKF 38393, and D2 receptor agonist LY 171555, but failed to change the effect of the activation of both D1 and D2 receptors. *Neuroscience*, **50**, 395-402.
- [2] Abbraini, J.H., Tomei, C., Rostain, J-C. (1991) Role of dopamine receptors in the occurrence of the behavioral motor disturbances in rats exposed to high pressure. *Pharmacol. Biochem. Behav.*, **39**, 773-779.
- [3] Albin, R.L., Young, A.B., Penney, J.B. (1989) The functional anatomy of basal ganglia disorders. *Trends Neurosci.*, **12**, 366-375.
- [4] Alexander, G.E., Crutcher, M.D. (1990a) Preparation for movement: neural representations of intended direction in three motor areas of the monkey. *J. Neurophysiol.*, **64**, 133-150.
- [5] Alexander, G.E., Crutcher, M.D. (1990b) Neural representations of the target (goal) of visually guided arm movements in three motor areas of the monkey. *J. Neurophysiol.*, **64**, 164-178.
- [6] Alexander, G.E., Crutcher, M.D., DeLong, M.R. (1990) Basal ganglia-thalamocortical circuits: parallel substrates for motor, oculomotor, "prefrontal" and "limbic" functions. *Prog. Brain Res.*, **85**, 119-146.
- [7] Alexander, G.E., DeLong, M.R., Crutcher, M.D. (1986) Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.*, **9**, 357-381.
- [8] Alheid, G.F., Heimer, L. (1988) New perspectives in basal forebrain organization of special relevance for neuropsychiatric disorders: the striatopallidum, amygdaloid, and corticopetal components of substantia innominata. *Neuroscience*, **27**, 1-39.
- [9] Bennett, B.D., Wilson, C.J. (2003) TANs, PANs and STANs. In: Graybiel, A.M., DeLong, M.R., Kitai, S.T. Eds, *The Basal Ganglia VI*, Springer, pp. 225-236.
- [10] Bishop, G.A., Chang, H.T., Kitai, S.T. (1982) Morphological and physiological properties of neostriatal neurons: an intracellular horseradish peroxidase study in the rat. *Neuroscience*, **7**, 179-191.
- [11] Boldry, R.C., Uretsky, N.J. (1988) The importance of dopaminergic neurotransmission in the hypermotility response produced by the administration of N-methyl-D-aspartic acid into the nucleus accumbens. *Neuropharmacology*, **27**, 569-577.
- [12] Boldry, R.C., Willins, D.L., Wallace, L.J., Uretsky, N.J. (1991) The role of endogenous dopamine in the hypermotility response to intra-accumbens AMPA. *Brain Res.*, **559**, 100-108.
- [13] Bouyer, J.J., Park, D.H., Joh, T.H., Pickel, V.M. (1984) Chemical and structural analysis of the relation between cortical inputs and tyrosine hydroxylase-containing terminals in rat neostriatum. *Brain Res.*, **302**, 267-275.
- [14] Breyse, N., Risterucci, C., Amalric, M. (2002) D1 and D2 dopamine receptors contribute to the locomotor response induced by Group II mGluRs activation in the rat nucleus accumbens. *Pharmacol. Biochem. Behav.*, **73**, 347-357.
- [15] Burns, L.H., Everitt, B.J., Kelley, A.E., Robbins, T.W. (1994) Glutamate-dopamine interactions in the ventral striatum: role in locomotor activity and responding with conditioned reinforcement. *Psychopharmacology (Berl)*, **115**, 516-528.
- [16] Canales, J.J., Iversen, S.D. (2000) Dynamic dopamine receptor interactions in the core and shell of nucleus accumbens differentially coordinate the expression of unconditioned motor behaviors. *Synapse*, **36**, 297-306.
- [17] Cepeda, C., Levine, M.S. (1998) Dopamine and N-methyl-D-aspartate receptor interactions in the neostriatum. *Dev. Neurosci.*, **20**, 1-18.
- [18] Choi, K.H., Zarandi, B., Todd, K.G., Biondo, A.M., Greenshaw, A.J. (2000) Effects of AMPA/kainite receptor blockade on responses to dopamine receptor agonists in the core and shell of the rat nucleus accumbens. *Psychopharmacology (Berl)*, **150**, 102-111.
- [19] Dalia, A., Uretsky, N.J., Wallace, L.J. (1998) Dopaminergic agonists administered into the nucleus accumbens: effects on extracellular glutamate and on locomotor activity. *Brain Res.* **788**, 111-117.
- [20] David, H.N., Abbraini, J.H. (2001a) The group I metabotropic glutamate receptor antagonist S-4-CPG modulates the locomotor response produced by the activation of D1-like, but not D2-like, dopamine receptors in the rat nucleus accumbens. *Eur. J. Neurosci.*, **13**, 2157-2164.

- [21] David, H.N., Abiraini, J.H. (2001b) Differential modulation of the D1-like- and D2-like dopamine receptor-induced locomotor responses by group II metabotropic glutamate receptors in the rat nucleus accumbens. *Neuropharmacology*, **41**, 454-463.
- [22] David, H.N., Abiraini, J.H. (2002) Group III metabotropic glutamate receptors and D1-like and D2-like dopamine receptors interact in the rat nucleus accumbens to influence locomotor activity. *Eur. J. Neurosci.*, **15**, 869-875.
- [23] David, H.N., Sissaoui K., Abiraini J.H. (2004) Modulation of the locomotor responses induced by D(1)-like and D(2)-like dopamine receptor agonists and D-amphetamine by NMDA and non-NMDA glutamate receptor agonists and antagonists in the core of the rat nucleus accumbens. *Neuropharmacology*, **46**, 179-191.
- [24] David, H.N., Anseau, M., Abiraini, J.H. (2005) Dopamine-glutamate reciprocal modulation of release and motor responses in the rat caudate-putamen and nucleus accumbens of "intact" animals. *Brain Res. Rev.*, **50**, 336-360.
- [25] DeLong, M.R. (1990) Primate models of movement disorders of basal ganglia origin. *Trends Neurosci.*, **13**, 281-285.
- [26] DiFiglia, M., Pasik, P., Pasik, T. (1976) A golgi study of neuronal types in the neostriatum of monkeys. *Brain Res.*, **114**, 245-256.
- [27] Donzanti, B.A., Uretsky, N.J. (1983) Effects of excitatory amino acids on locomotor activity after bilateral microinjection into the rat nucleus accumbens: possible dependence on dopaminergic mechanisms. *Neuropharmacology*, **22**, 971-981.
- [28] Dreher, J.K., Jackson, D.M. (1989) Role of D1 and D2 dopamine receptors in mediating locomotor activity elicited from the nucleus accumbens of rats. *Brain Res.*, **487**, 267-277.
- [29] Dube, L., Smith, A.D., Bolam, J.P. (1988) Identification of synaptic terminals of thalamic or cortical origin in contact with distinct medium-size spiny neurons in the rat neostriatum. *J. Comp. Neurol.*, **267**, 455-471.
- [30] Essman, W.D., McGonigle, P., Lucki, I. (1993) Anatomical differentiation within the nucleus accumbens of the locomotor stimulatory actions of selective dopamine agonists and d-amphetamine. *Psychopharmacology (Berl)*, **112**, 233-241.
- [31] Freund, T.F., Powell, J.F., Smith, A.D. (1984) Tyrosine hydroxylase-immunoreactive boutons in synaptic contact with identified striatonigral neurons, with particular reference to dendritic spines. *Neuroscience*, **13**, 1189-1215.
- [32] Frotscher, M., Rinne, U., Hassler, R., Wagner, A. (1981) Termination of cortical afferents on identified neurons in the caudate nucleus of the cat. A combined Golgi-EM degeneration study. *Exp. Brain Res.*, **41**: 329-337.
- [33] Gong, W., Neill, D.B., Lynn, M., Justice, J.B. Jr. (1999) Dopamine D1/D2 agonists injected into nucleus accumbens and ventral pallidum differentially affect locomotor activity depending on site. *Neuroscience*, **93**, 1349-1358.
- [34] Hamilton, M.H., De Bellerocche, J.S., Gardiner, I.M., Herberg, L.J. (1986) Stimulatory effect of N-methyl aspartate on locomotor activity and transmitter release from rat nucleus accumbens. *Pharmacol. Biochem. Behav.*, **25**, 943-948.
- [35] Hauber, W., Münkler, M. (1997) Motor depressant effects mediated by dopamine D2 and adenosine A2A receptors in the nucleus accumbens and the caudate-putamen. *Eur. J. Pharmacol.*, **323**, 127-131.
- [36] Honore, T., Davies, S.N., Drejer, J., Fletcher, E.J., Jacobsen, P., Lodge, D., Nielsen, F.E. (1988) Quinoxalinediones: potent competitive non-NMDA glutamate receptor antagonists. *Science*, **241**, 701-703.
- [37] Ikemoto, S. (2002) Ventral striatal anatomy of locomotor activity induced by cocaine, D-amphetamine, dopamine and D1/D2 agonists. *Neuroscience*, **113**, 939-955.
- [38] Imperato, A., Di Chiara, G. (1988) Effects of locally applied D-1 and D-2 receptor agonists and antagonists studied with brain dialysis. *Eur. J. Pharmacol.*, **156**, 385-393.
- [39] Joel, D., Weiner, I. (2000) The connections of the dopaminergic system with the striatum in rats and primates: an analysis with respect to the functional and compartmental organization of the striatum. *Neuroscience*, **96**, 451-474.
- [40] Kaddis, F.G., Wallace, L.J., Uretsky, N.J. (1993) AMPA/kainate antagonists in the nucleus accumbens inhibit locomotor stimulatory response to cocaine and dopamine agonists. *Pharmacol. Biochem. Behav.*, **46**, 703-708.
- [41] Kawaguchi, Y. (1997) Neostriatal cell subtypes and their functional roles. *Neurosci. Res.*, **27**, 1-8.
- [42] Kelley, A.E. (1999) Neural integrative activities of nucleus accumbens subregions in relation to learning and motivation. *Psychobiology*, **27**, 198-213.
- [43] Kelley, A.E., Throne, L.C. (1992) NMDA receptors mediate the behavioral effects of amphetamine infused into the nucleus accumbens. *Brain Res. Bull.*, **29**, 247-254.
- [44] Kemp, J.M., Powell, T.P.S. (1971) The structure of the caudate nucleus of the cat: light and electron microscopic study. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.*, **262**, 383-401.
- [45] Kronthaler, U.O., Schmidt, W.J. (2000) Activation of striatal group II metabotropic glutamate receptors has a differential effect on dopamine-D1 and -D2 receptor antagonist-induced hypokinesia in the rat. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **361**, 289-297.
- [46] LaHoste, G.F., Yu, J., Marshall, J.F. (1993) Striatal Fos expression is indicative of dopamine D1/D2 synergism and receptor supersensitivity. *Proc. Natl. Acad. Sci. USA*, **90**, 7451-7455.
- [47] Lapper, S.R., Bolam, J.P. (1992) Input from the frontal cortex and the parafascicular nucleus to cholinergic interneurons in the dorsal striatum of the rat. *Neuroscience*, **51**, 533-545.
- [48] Mao, L., Wang, J.Q. (2000a) Motor stimulation following bilateral injection of the group-I metabotropic glutamate receptor agonist into the dorsal striatum of rats: evidence against dependence on ionotropic glutamate receptors. *Psychopharmacology (Berl)*, **148**, 367-373.
- [49] Mao, L., Wang, J.Q. (2000b) Distinct inhibition of acute cocaine-stimulated motor activity following microinjection of a group III metabotropic glutamate receptor agonist into the dorsal striatum of rats. *Pharmacol. Biochem. Behav.*, **67**, 93-101.
- [50] Mao, L., Wang, J.Q. (1999) Protection against acute amphetamine-induced behavior by microinjection of a group II metabotropic glutamate receptor agonist into the dorsal striatum of rats. *Neurosci. Lett.*, **270**, 103-106.
- [51] McPherson, R.J., Marshall, J.F. (1996) Intra-striatal AP5 differentially affects behaviors induced by local infusions of D1 vs. D2 dopamine agonists. *Brain Res.*, **739**, 19-25.
- [52] Meredith, G.E., Wouterlood, F.G. (1990) Hippocampal and midline thalamic fibers and terminals in relation to the choline acetyltransferase-immunoreactive neurons in nucleus accumbens of the rat: a light and electron microscopic study. *J. Comp. Neurol.*, **296**, 204-221.
- [53] Meyer, M.E. (1993) Effects of intraaccumbens dopamine agonist SK&F38393 and antagonist SCH23390 on locomotor activities in rats. *Pharmacol. Biochem. Behav.*, **45**, 843-847.
- [54] Meyer, M.E., Van Hartesveldt, C., Potter, T.J. (1993) Locomotor activity following intra-accumbens microinjections of dopamine D1 agonist SK&F 38393 in rats. *Synapse*, **13**, 310-314.
- [55] Mogenson, G.J., Wu, M. (1991) Effects of administration of dopamine D2 agonist quinpirole on exploratory locomotion. *Brain Res.*, **551**, 216-220.
- [56] Mori, A., Takahashi, T., Miyashita Y., Kasai, H. (1994) Two distinct glutamatergic synaptic inputs to striatal medium spiny neurons of neonatal rats and paired-pulse depression. *J. Physiol.*, **476**, 217-228.
- [57] Nestler, E.J. (1994) Hard target: Understanding dopaminergic neurotransmission. *Cell*, **79**, 923-926.
- [58] Ossowska, K., Karcz, M., Wardas, J., Wolfarth, S. (1990) Striatal and nucleus accumbens D1/D2 dopamine receptors in neuroleptic catalepsy. *Eur. J. Pharmacol.*, **182**, 327-334.
- [59] Ouagazzal, A., Amalric, M. (1995) Competitive NMDA receptor antagonists do not produce locomotor hyperactivity by a dopamine-dependent mechanism. *Eur. J. Pharmacol.*, **294**, 137-146.
- [60] Plaznik, A., Stefanski, R., Kostowski, W. (1989) Interaction between accumbens D1 and D2 receptors regulating rat locomotor activity. *Psychopharmacology (Berl)*, **99**, 558-562.
- [61] Pickel, V.M., Chan, J., Sesack, S.R. (1992) Cellular basis for interactions between catecholaminergic afferents and neurons containing Leu-enkephalin-like immunoreactivity in rat caudate-putamen nuclei. *J. Neurosci. Res.*, **31**, 212-230.
- [62] Rouillon, C., Abiraini, J.H., David, H.N. (2007) Hippocampal modulation of locomotor activity induced by focal activation of postsynaptic dopamine receptors in the core of the nucleus accumbens. *Hippocampus*, **17**, 1028-1036.

- [63] Rouillon, C., Degoulet, M., Chevallier, K., Abirami, J.H., David, H.N. (2008) Modulation by group I mGlu receptor activation and group III mGlu receptor blockade of locomotor responses induced by D1-like and D2-like receptor agonists in the nucleus Accumbens. *Brain Res.*, **1198**, 44-54.
- [64] Sadri-Vakili, G., Johnson, D.W., Janis, G.C., Gibbs, T.T., Pierce, R.C., Farb, D.H. (2003) Inhibition of NMDA-induced striatal dopamine release and behavioral activation by the neuroactive steroid 3alpha-hydroxy-5beta-pregnan-20-one hemisuccinate. *J. Neurochem.*, **86**, 92-101.
- [65] Schmidt, W.J., Bury, D. (1988) Behavioural effects of N-methyl-D-aspartate in the anterodorsal striatum of the rat. *Life Sci.*, **43**, 545-549.
- [66] Sesack, S.R., Pickel, V.M. (1990) In the rat medial nucleus accumbens, hippocampal and catecholaminergic terminals converge on spiny neurons and are in apposition to each other. *Brain Res.*, **527**, 266-279.
- [67] Shink, E., Bevan, M.D., Bolam, J.P., Smith, Y. (1996) The subthalamic nucleus and the external pallidum: two tightly interconnected structures that control the output of the basal ganglia in the monkey. *Neuroscience*, **73**, 335-357.
- [68] Shreve, P.E., Uretsky, N.J. (1988) Role of quisqualic acid receptors in the hypermotility response produced by the injection of AMPA into the nucleus accumbens. *Pharmacol. Biochem. Behav.*, **30**, 379-384.
- [69] Smith, Y., Bennett, B.D., Bolam, J.P., Parent, A., Sadikot, A.F. (1994) Synaptic relationships between dopaminergic afferents and cortical or thalamic input in the sensorimotor territory of the striatum in monkey. *J. Comp. Neurol.*, **344**, 1-19.
- [70] Somogyi, P., Bolam, J.P., Smith, A.D. (1981) Monosynaptic cortical input and local axon collaterals of identified striatonigral neurons. A light and electron microscopic study using the Golgi-peroxidase transport-degeneration procedure. *J. Comp. Neurol.*, **195**, 567-584.
- [71] Swanson, C.J., Kalivas, P.W. (2000) Regulation of locomotor activity by metabotropic glutamate receptors in the nucleus accumbens and ventral tegmental area. *J. Pharmacol. Exp. Ther.*, **292**, 406-414.
- [72] Swanson, C.J., Baker, D.A., Carson, D., Worley, P.F., Kalivas, P.W. (2001) Repeated cocaine administration attenuates group I metabotropic glutamate receptor-mediated glutamate release and behavioral activation: a potential role for Homer. *J. Neurosci.*, **21**, 9043-9052.
- [73] Swanson, C.J., Heath, S., Stratford, T.R., Kelley, A.E. (1997) Differential behavioral responses to dopaminergic stimulation of nucleus accumbens subregions in the rat. *Pharmacol. Biochem. Behav.*, **58**, 933-945.
- [74] Tamaru, Y., Nomura, S., Mizuno N., Shigemoto, R. (2001) Distribution of metabotropic glutamate receptor mGluR3 in the mouse CNS: differential location relative to pre- and postsynaptic sites. *Neuroscience*, **106**, 481-503.
- [75] Thanos, P.K., Jhamandas, K., Beninger, R.J. (1992) N-methyl-D-aspartate unilaterally injected into the dorsal striatum of rats produces contralateral circling: antagonism 2-amino-7-phosphonoheptanoic acid and cis-fluoxetine. *Brain Res.*, **589**, 55-61.
- [76] Timmerman, W., De Vries J.B., Westerink, B.H. (1990) Effects of D-2 agonists on the release of dopamine: localization of the mechanism of action. *Naunyn Schmiedebergs Arch. Pharmacol.*, **342**, 650-654.
- [77] Totterdell, S., Smith, A.D. (1989) Convergence of hippocampal and dopaminergic input onto identified neurons in the nucleus accumbens of the rat. *J. Chem. Neuroanat.*, **2**, 285-298.
- [78] Van den Boss, R., Cools, A.R., Ogren, S.O. (1988) Differential effects of the selective D2-antagonist raclopride in the nucleus accumbens of the rat on spontaneous and d-amphetamine-induced activity. *Psychopharmacology (Berl)*, **95**, 447-451.
- [79] Van de Kar, L.D., Lorens, S.A. (1979) Differential serotonergic innervation of individual hypothalamic nuclei and other forebrain regions by the dorsal and median midbrain raphe nuclei. *Brain Res.*, **162**, 45-54.
- [80] Wachtel, S.R., Hu, X.T., Galloway, M.P., White, F.J. (1989) D1 dopamine receptor stimulation enables the postsynaptic, but not autoreceptor, effects of D2 dopamine agonists in nigrostriatal and mesoaccumbens dopamine systems. *Synapse*, **4**, 327-346.
- [81] Walters, J.R., Bergstrom, D.A., Carlson, J.H., Chase, T.N., Braun, A.R. (1987) D1 dopamine receptor activation required for postsynaptic expression of D2 agonist effects. *Science*, **236**, 719-722.
- [82] Wang, J.Q., Mao, L. (2000) Sustained behavioral stimulation following selective activation of group I metabotropic glutamate receptors in rat striatum. *Pharmacol. Biochem. Behav.*, **65**, 439-447.
- [83] Westerink, B.H., De Boer, P., Timmerman, W., De Vries, J.B. (1990) *In vivo* evidence for the existence of autoreceptors on dopaminergic, serotonergic, and cholinergic neurons in the brain. *Ann. N. Y. Acad. Sci.*, **604**, 492-504.
- [84] White, F.J. (1987) D-1 dopamine receptor stimulation enables the inhibition of nucleus accumbens neurons by a D-2 receptor agonist. *Eur. J. Pharmacol.*, **135**, 101-105.
- [85] Wilson, C.J., Groves, P.M. (1980) Fine structure and synaptic connections on the common spiny neuron of the rat neostriatum: a study employing intracellular injection of horseradish peroxidase. *J. Comp. Neurol.*, **194**, 599-615.
- [86] Wu, M., Brudzynski, S.M., Mogenson, G.J. (1993a) Functional interactions of dopamine and glutamate in the nucleus accumbens in the regulation of locomotion. *Can. J. Physiol. Pharmacol.*, **71**, 407-413.
- [87] Wu, M., Brudzynski, S.M., Mogenson, G.J. (1993b) Differential effects of quinpirole in the nucleus accumbens depending on the initial level of locomotor activity. *Brain Res. Bull.*, **32**, 395-398.