



Note

Re-epitypification of *Neofusicoccum loricinum*

Yukako Hattori^{a*}, Chiharu Nakashima^b & Hayato Masuya^a^aDepartment of Mushroom Science and Forest Microbiology, Forestry and Forest Products Research Institute (FFPRI), 1 Matsunosato, Tsukuba, Ibaraki, 305-8687 Japan^bGraduate School of Bioresources, Mie University, 1577 Kurima-machiya, Tsu, Mie, 514-8507 Japan

ABSTRACT

The lectotype and epitype of *Neofusicoccum loricinum* proposed in previous studies failed to meet the requirements of the International Code of Nomenclature for Algae, Fungi, and Plants. Here, we propose a valid lectotype and epitype for this species that meet the requirements of the Code.

Keywords: *Botryosphaeriaceae*, *Botryosphaeriales*, epitype, taxonomy

Article history: Received 6 October 2023, Revised 21 November 2023, Accepted 23 November 2023, Available online 6 March 2024.

Neofusicoccum loricinum (Sawada) Hattori & C. Nakash. inhabits Far East Asia and is the cause of shoot blight of larch (EPPO datasheets on pests recommended for regulation, <https://gd.eppo.int/>; Sep 2023). This species was established in 1950 by Sawada as *Physalospora loricina* Sawada. Although Sawada (1950) described only the sexual state, subsequent research on shoot blight revealed that the asexual state of this species, identified as *Macrophoma* sp., causes the disease, and elucidated its life history (Uozumi, 1961; Yokota, 1962, 1963). Yamamoto (1961) proposed that the causal fungus *P. loricina* should be treated as the genus *Guignardia* based on the morphological characteristics of its asexual state, and a new combination should be proposed as *Guignardia loricina* (Sawada) W. Yamam. & Kaz. Itô. Subsequently, Shang (1987) transferred *G. loricina* to the genus *Botryosphaeria* based on the morphological characteristics of its sexual state noted by Luttrell (1973). Hattori et al. (2021) reexamined the specimens of *G. loricina*, including the syntype specimens of *P. loricina* (IUM-FS515, IUM-FS516, IUM-FS517, IUM-FS518, IUM-FS519, and IUM-FS520). In their study, a multi-locus molecular phylogenetic analysis using ITS, rpb2, tef1- α , and tub2 sequences was conducted with newly isolated strains of *G. loricina* to resolve taxonomic problems associated with this species. According to the results of phylogenetic analysis, *B. loricina* was transferred to the genus *Neofusicoccum*, and epitypification for the species was proposed based on their newly materials. However, the treatment failed to meet the requirements of the International Code of Nomenclature for Algae, Fungi, and Plants (Shenzhen Code; the identifier was not given (Art. F.5.4), and designation of an epitype is not effected unless the lectotype that the epitype supports is explicitly cited (Art. 9.9)). Recently, Zhao et al. (2021) proposed a lectotype based on the illustrations in Sawada

(1950) despite the existence of a syntype. Due to the presence of the syntype, this proposal was also not compliant with the Code (Art. 9.12). Therefore, we propose a lectotype and an epitype compliant with the requirements of the Code.

Taxonomy

Neofusicoccum loricinum (Sawada) Y. Hattori & C. Nakash., Mycoscience 62: 252, 2021. MycoBank no.: MB 837720.

≡ *Physalospora loricina* Sawada, Bull. Gov. Forest Exp. Sta. 46: 126, 1950.

≡ *Guignardia loricina* (Sawada) W. Yamam. & Kaz. Itô, Sci. Rep. Hyogo Univ. Agric. 5: 9, 1961.

≡ *Botryosphaeria loricina* (Sawada) Y.Z. Shang, Acta Microbiol. Sin. 6: 249, 1987.

Lectotype: JAPAN, Aomori, Sanbongi, Fukamochi, on *Larix kaempferi*, 27 Sep 1949, collected by K. Sawada, Syntype of *Physalospora loricina*, IUM-FS515, designated here as lectotype (MBT 10015084).

Epitype: JAPAN, Ibaraki, Mito, on *Larix decidua*, 14 Jun 1973, collected by H. Kondo, TFM: FPH-4038, designated here (MBT 10015083), ex-epitype culture FFPRI 411215 = MUCC 2662.

Disclosure

The authors declare no conflict of interest. All the experiments undertaken as part of this study comply with the current laws of the country in which they were performed.

Acknowledgments

We are grateful to Dr. Konstanze Bensch of Westerdijk Fungal Biodiversity Institute for advice on typification. This study was supported, in part, by a JSPS KAKENHI grant (no. 20J12025) and Research Grant #202303 of the Forestry and Forest Products Re-

* Corresponding author.

Department of Mushroom Science and Forest Microbiology, FFPRI, 1 Matsunosato, Tsukuba, Ibaraki, 305-8687 Japan

E-mail address: hattori31@ffpri.affrc.go.jp



This is an open-access paper distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivative 4.0 international license (CC BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>).

search Institute.

References

- European and Mediterranean Plant Protection Organization (EPPO). (2023). *Neofusicoccum laricinum*. EPPO datasheets on pests recommended for regulation. <https://gd.eppo.int/> (accessed 2023-09-12)
- Hattori, Y., Ando, Y., & Nakashima C. (2021). Taxonomical re-examination of the genus *Neofusicoccum* in Japan. *Mycoscience*, 62, 250–259. <https://doi.org/10.47371/mycosci-2021-03-008>.
- Luttrell, E. S. (1973). Loculoascomycetes. In: G. C. Ainsworth, F. K. Sparrow, & A. S. Sussman (Eds.): *The Fungi*, Vol. IVa. pp. 135–219. Academic Press, New York and London.
- Sawada, K. (1950). Fungi inhabiting conifers in the Tohoku district. II. Fungi on various conifers except ‘Sugi.’ *Bulletin of the Government Forest Experimental Station Meguro*, 46, 111–150.
- Shang, Y. Z. (1987). Taxonomic study on the pathogen fungus of shoot blight of larch (in Chinese). *Acta Mycologia Sinica*, 6, 248–249.
- Uozumi, T. (1961). Studies on the shoot blight disease of larch. With special reference to the life history of the causal fungus, *Physalospora laricina* Sawada. *Bulletin of the Government Forest Experimental Station Tokyo*, 132, 47–54.
- Yamamoto, W. (1961). Species of the genera of *Glomerella* and *Guignardia* with special reference to their imperfect stage (in Japanese with English resume). *Science Reports of the Hyogo University of Agriculture. Series, Agricultural Biology*, 5, 1–12.
- Yokota, S. (1962). Studies on Shoot-blight Disease of Larch Trees II. The distribution, morphology and life history of the causal fungus (in Japanese with English resume). *Bulletin of the Government Forest Experimental Station Tokyo*, 142, 173–202.
- Yokota, S. (1963). Studies on Shoot-blight Disease of Larch Trees III. Expulsion and dissemination of spores and their germinations (in Japanese with English resume). *Bulletin of the Government Forest Experimental Station Tokyo*, 151, 1–44.
- Zhao, P., Crous, P. W., Hou, L. W., Duan, W. J., Cai, L., Ma, Z. Y., & Liu, F. (2021). Fungi of quarantine concern for China I: Dothideomycetes. *Persoonia*, 47, 45–105. <https://doi.org/10.3767/persoonia-2021-47-02>.