



# Defining competencies in robotic thyroidectomy: development of a model assessing an expert operator's intraoperative performance skills and cognitive strategies

Hyejin Kim<sup>1#</sup>, Hyeong Won Yu<sup>2#^</sup>, Jong-hyuk Ahn<sup>3</sup>, Tae Seon Lee<sup>4,5</sup>, Kyu Eun Lee<sup>6^</sup>

<sup>1</sup>Department of Public Health, Graduate School, Yonsei University, Seoul, Republic of Korea; <sup>2</sup>Department of Surgery, Seoul National University Bundang Hospital, Seongnam, Republic of Korea; <sup>3</sup>Department of Surgery, Seoul National University Hospital, Seoul, Korea; <sup>4</sup>Department of Neurosurgery, Severance Hospital & Yonsei University College of Medicine, Seoul, Republic of Korea; <sup>5</sup>Center for Medical Education, Graduate School of Medicine, Nagoya University, Nagoya, Japan; <sup>6</sup>Department of Surgery, Seoul National University Hospital and College of Medicine, Seoul, Republic of Korea

**Contributions:** (I) Conception and design: TS Lee, KE Lee; (II) Administrative support: HW Yu, KE Lee; (III) Provision of study materials or patients: HW Yu, JH Ahn, KE Lee; (IV) Collection and assembly of data: H Kim; (V) Data analysis and interpretation: H Kim, HW Yu, TS Lee; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

<sup>#</sup>These authors contributed equally to this work.

**Correspondence to:** Kyu Eun Lee, MD, PhD. Department of Surgery, Seoul National University Hospital and College of Medicine, 101 Daehak-ro, Jongno-gu, Seoul 110-744, Republic of Korea. Email: kyueunlee@snu.ac.kr; Tae Seon Lee, PhD. Department of Neurosurgery, Severance Hospital & Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, Republic of Korea; Center for Medical Education, Graduate School of Medicine, Nagoya University, Nagoya, Japan. Email: tseonlee@yonsei.ac.kr.

**Background:** The changing medical education environment emphasizes the need for time efficiency, increasing the demand for competency-based medical education to improve trainees' learning strategies. This study was performed to determine the competencies required for successful performance of robotic thyroidectomy (RT) and to determine the consensus of experts for performance of RT.

**Methods:** Data were collected through 12 semi-structured interviews with RT experts and 11 field observations. Cognitive task analysis was performed to determine the competencies required for experts to perform RT. A modified Delphi methodology was used to determine how 20 experts rated the importance of each item of RT performance on a Likert 7-point scale. The criteria for the Delphi consensus were set at a Cronbach's  $\alpha \geq 0.80$  with two survey rounds.

**Results:** After 11 field observations and 12 semi-structured interviews, 89 items were identified within six modules. These items were grouped into sub-modules according to their theme. The modified Delphi survey, involving 21 experts, reached the consensus standard during the second round (Cronbach's  $\alpha=0.954$ ), enabling the identification of the 64 most important items within six modules related to RT performance: midline incision to isthmectomy (MID module; n=8), lateral dissection (LAT module; n=7), preservation of inferior parathyroid glands (INF module; n=16), preservation of recurrent laryngeal nerve and dissection of the ligament of Berry (BER module; n=21), dissection of the thyroid upper pole (SUP module; n=10), and specimen removal and closure (END module; n=2).

**Conclusions:** This mixed-method study combining qualitative and quantitative methodology identified modules of core competencies required to perform RT. These modules can be used as a standard and objective guide to train surgeons to perform RT and evaluate outcomes.

**Keywords:** Thyroidectomy; clinical competence; robotics; surgical procedures; reference standards

<sup>^</sup> ORCID: Hyeong Won Yu, 0000-0001-7338-5157; Kyu Eun Lee, 0000-0002-2354-3599.

Submitted Nov 16, 2023. Accepted for publication Feb 25, 2024. Published online Mar 22, 2024.

doi: 10.21037/gS-23-467

View this article at: <https://dx.doi.org/10.21037/gS-23-467>

## Introduction

Recently, the educational paradigm in surgery has changed from the traditional apprenticeship model to competency-based medical education, which emphasizes the acquisition of standard technical and cognitive proficiency for performing surgery safely and effectively (1-3). To incorporate intraoperative decision-making ability into the standard surgical training curriculum, many studies have attempted to identify cognitive components associated with the success of various types of surgery, such as conventional open thyroidectomy, management of trauma patients, laparoscopic transabdominal adrenalectomy, and flexible pharyngo-laryngoscopy (3-6). Research on the techniques required for open thyroidectomy was utilized to develop high-definition video teaching modules and interactive web-based educational platforms (7,8). Although modules and platforms have been developed for open and laparoscopic surgery, surgeons are still being trained to perform robotic surgery through apprenticeship, without a standardized educational curriculum (9).

Robotic thyroidectomy (RT) is remote-access surgery and results in higher cosmetic satisfaction than conventional surgical methods (10). These advantages

have increased demands for RT worldwide, increasing the number of training programs for RT. Previously, Madani and colleagues defined a model of competencies for successful open thyroidectomy performed by expert endocrine surgeons to use it as references for trainees in the context of competency-based medical education (3). RT, however, requires advanced surgical skills and utilizes different surgical instruments and approaches including behavioral and cognitive strategies compared with open thyroidectomy (11). Thus, this study was designed to organize key tasks and decision-making procedures for RT by modules and to evaluate their importance in determining the competencies of expert performance and cognitive strategies required for RT.

## Methods

### *Setting and participants*

This study recruited surgeons who perform RT with the bilateral axillo-breast approach (BABA). We defined subject matter experts (SMEs) as individuals who had conducted 40 or more BABA RT operations (11-14). Moreover, we included individuals with a minimum of 5 years of clinical experience to guarantee proficiency in BABA-RT. We used convenience sampling to enlist 12 SMEs for the qualitative cognitive task analysis (CTA) via email. Additionally, we recruited 21 surgeons through email for the modified Delphi survey. The purpose and method of the study were explained to the recruited surgeons, and they provided informed consent before participating in the study via telephone or email. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study protocol was approved by the Institutional Review Board of Seoul National University Hospital (No. H-1912-116-1090).

### *BABA RT*

BABA RT is a surgical method in which the thyroid gland is dissected using a da Vinci robot inserted through small wounds on both sides of the axilla and breast (15). BABA RT differs from conventional open thyroidectomy, in that it

### Highlight box

#### Key findings

- This study organized the key tasks and decision-making procedures for robotic thyroidectomy (RT) into modules and assessed their significance in delineating the competencies for expert performance and the cognitive strategies essential for RT.

#### What is known and what is new?

- While RT demands advanced surgical skills and employs various surgical instruments and approaches, including behavioral and cognitive strategies, there is a lack of research defining the competencies necessary for successful RT performance when compared to conventional thyroidectomy.
- We identified 89 items within six modules that define the competencies for RT.

#### What is the implication, and what should change now?

- These modules can be used as a standard and objective guide to train surgeons to perform RT and evaluate outcomes.

**Table 1** Sample questions used for semi-structured interviews

Structure	Questions
General questions	Describe the tasks required to perform BABA RT in key steps
Questions by module	What is the purpose of this task?
	What is the sequence of actions necessary to complete this task?
	What conditions must be present before starting this task?
	What decisions have to be made during this task, including the various options and criteria to choose among options?
	What errors can occur and what tips/tricks can be used to avoid such errors during this task?
	What performance standards or quality indicators are used to ensure successful completion of this task?

BABA RT, bilateral axillo-breast approach robotic thyroidectomy.

starts with the lower pole of the thyroid gland and continues to the upper pole. BABA RT allows thyroid lobectomy, total thyroidectomy, and total thyroidectomy with modified lateral neck dissection (16,17). This study, however, was restricted to surgery involving a midline incision to remove the dissected specimen and midline closure. The RT procedures were divided into six modules, as described (15).

#### **Qualitative study: CTA**

This study was of mixed-method design, combining qualitative and quantitative methods. To collect qualitative data, one of the researchers observed the performance of RT by study participants, focusing mainly on communication with colleagues, emergency situations and coping methods, and changes in decisions during surgery. Each RT was video recorded and used as an adjunct in the subsequent semi-structured interview. The interviewee asked prescribed questions about the intent, cautions, and decision-making during each step of RT (*Table 1*) (3). The interviewer was careful not to interrupt the interviewee and to not ask close-ended questions, facilitating open and unbiased answers. Each interview was recorded, and each interview took roughly 1 hour. The recorded interviews were transcribed verbatim.

The qualitative data were analyzed by CTA, a robust method that systematically captures automated cognitive tasks performed by experts. CTA was based on naturalistic decision-making studies and has been used extensively to explore cognitive processes in medical settings (18). In this study, each surgical procedure performed by the surgeon was recorded and reviewed several times, and the cognitive factors of the task were analyzed by two task analysis experts

(Ph.D. in cognitive psychology and medical education and two clinical health psychologists). To ensure the saturation of data, surgery textbooks that described the steps for open thyroidectomy and RT were reviewed to add any missing information (15,19-21).


The contents were subsequently reviewed and modified by two expert surgeons to develop a structured framework, completing it as modules in structured stages for the entire operation. This method defined the explicit and implicit knowledge and skills required, without bias, for surgical judgment and decision making.

#### **Quantitative study: modified Delphi survey**

The modified Delphi method, which is effective in drawing consensus from experts through iterative survey rounds and has been recommended for solving problems in clinical practice that lack scientific evidence (22), was utilized for quantitative analysis. Survey items were constructed from the data obtained from CTA. During the first round of the Delphi survey, participants were asked to anonymously rate the importance of each item, ranging from 1 (not at all important) to 7 (very important). During the second round, 1 week later, participants were asked to again rate the importance of each item. Rounds one and two included 21 and 20 RT experts, respectively, of various backgrounds and experience. The design of this study is shown in *Figure 1*.

#### **Statistical analysis**

To quantitatively verify the results obtained from the qualitative study, the responses to each survey round were reported as the means and variances. A consensus among



**Figure 1** Flowchart of the study design.

experts was defined as a Cronbach's  $\alpha \geq 0.80$  for the reliability of the participants' responses to each item. Items evaluated by over 80% of respondents as  $\geq 5$  on the Likert scale were selected as the key component in implementing RT. The results were reported as the mean (standard deviation), median (range), and N (%). All statistical analyses were performed using Microsoft Excel<sup>®</sup> software. This mixed methodology study design has been used in various studies that define key operative competencies (2,3,6).

## Results

### *Qualitative study: CTA*

CTA analysis of the data collected by the interviews and the review of video recordings of RT identified 89 meaningful items, which were divided into six modules based on the order of the tasks: midline incision to isthmectomy (MID, n=18), lateral dissection (LAT, n=16), preservation of inferior parathyroid glands (INF, n=18), preservation of recurrent laryngeal nerve (RLN) and dissection of the ligament of Berry (BER, n=22), dissection of the thyroid upper pole (SUP, n=12), and specimen removal and closure (END, n=3). The items within each module were categorized based on structural similarities and arranged as submodules (Table 2; for the full list, see Table S1).

### *Quantitative study: modified Delphi survey*

In the quantitative study, the participants rated the 89 identified Delphi items by their importance. Twenty-seven RT experts, including participants in the first phase, were asked to participate in the Delphi survey, with 21 SMEs participating in the first round and 20 in the second round. The concordance of the responses among experts met the criteria for consensus set in the study, with the Cronbach's  $\alpha$  values of 0.91 in round 1 and 0.954 in round 2.

The most important of the 89 items was determined by selecting the items (n=64) that received scores  $\geq 5$  from >80% of respondents, with these items regarded as the critical behavioral and cognitive competencies for RT. Figure 2 shows the results of this study.

Table 3 shows the 10 most important items required for safe and efficient RT, as determined by experts. The five top-ranked items of each of the six modules are graphically shown in Figure 3. The order of these modules is not necessarily linear and can be altered at the discretion of the surgeon (for the full list, see Table S2).

Additional comments regarding the MID module included: (I) the need for preoperative evaluation of the possibility of isthmectomy, as cancer on the isthmus itself may alter the isthmectomy position; and (II) that care should be taken not to injure the trachea. An additional

Table 2 Items synthesized through the cognitive task analysis

MID module (n=18)	LAT module (n=16)	INF module (n=18)	BER module (n=22)	SUP module (n=12)	END module (n=3)
<i>Identification of the midline</i>	<i>Dissection of the surgical plane between the thyroid and the strap muscle</i>	<i>Identification of the RLN</i>	<i>Dissection between the medial thyroid and trachea</i>	<i>Dissection in the upward direction</i>	<i>Specimen out</i>
<ul style="list-style-type: none"> <li>Find the correct midline location on the strap muscles</li> </ul>	<ul style="list-style-type: none"> <li>Separate the strap muscles and thyroid gland from cranial to caudal</li> </ul>	<ul style="list-style-type: none"> <li>Identify the RLN between the central lymph nodes</li> </ul>	<ul style="list-style-type: none"> <li>Separate the trachea and cricothyroid muscle from the thyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>To prevent EBSLN injury, attach as much to the thyroid as possible and simultaneously ligate the upper thyroid artery well.</li> </ul>	<ul style="list-style-type: none"> <li>If the thyroid is too large to remove, expand the trocar tunnel site sufficiently</li> </ul>
<i>Retract the midline bilaterally with both graspers</i>	<ul style="list-style-type: none"> <li>Dissect the surgical plane as close as possible to the surface of the thyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Ensure safe distances based on the range of heat conduction to prevent thermal injury</li> </ul>	<i>Dissection between the thyroid gland and fascia</i>	<ul style="list-style-type: none"> <li>Avoid EBSLN injury</li> </ul>	<ul style="list-style-type: none"> <li>Use a surgical lap bag to safely remove the specimen from the operative field to prevent metastasis to other sites</li> </ul>
<ul style="list-style-type: none"> <li>When retracting the midline with graspers on both sides, provide proper symmetrical tension and cautions to prevent muscle tearing</li> </ul>	<ul style="list-style-type: none"> <li>Be careful when dissecting between the thyroid gland and strap muscle</li> </ul>	<ul style="list-style-type: none"> <li>Avoid retracting the thyroid gland excessively, as it may cause mechanical injury to the RLN</li> </ul>	<ul style="list-style-type: none"> <li>Finish the lateral dissection on the lateral side of the thyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Map the course of the EBSLN using nerve monitoring</li> </ul>	<i>Use of hemostatic dressing and anti-adhesion adjuvant</i>
<i>Midline incision</i>	<ul style="list-style-type: none"> <li>Determine whether the strap muscle is injured</li> </ul>	<ul style="list-style-type: none"> <li>Identify the course and the location of the RLN</li> </ul>	<i>Thyroid retraction—Zuckerkanndl</i>	<i>Identification and preservation of the superior parathyroid glands</i>	<ul style="list-style-type: none"> <li>Sew the strap muscles with running sutures during midline closure (cranial to caudal)</li> </ul>
<ul style="list-style-type: none"> <li>Follow the surgical plane well and dissect it</li> </ul>	<i>Lateral retraction of the strap muscle</i>	<i>Identification of the inferior parathyroid gland</i>	<ul style="list-style-type: none"> <li>Retract the thyroid to enable entering the harmonic</li> </ul>	<ul style="list-style-type: none"> <li>Avoid bleeding during the ligation of the superior thyroid artery because the operative field is narrow</li> </ul>	<i>Drain insertion and midline closure</i>
<ul style="list-style-type: none"> <li>Identify the sternothyroid and sternohyoid muscles</li> </ul>	<ul style="list-style-type: none"> <li>Sufficiently separate the thyroid from the strap muscle</li> </ul>	<ul style="list-style-type: none"> <li>Recognize the typical location and shape of the parathyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Avoid RLN injury caused by traction</li> </ul>	<ul style="list-style-type: none"> <li>Determine which blood vessels to leave</li> </ul>	
<ul style="list-style-type: none"> <li>Ensure sufficient incision to Delphian lymph node</li> </ul>	<ul style="list-style-type: none"> <li>Be careful not to pull the strap muscle excessively, as it can tear and bleed</li> </ul>	<ul style="list-style-type: none"> <li>Identify the color of the parathyroid gland</li> </ul>	<i>Preservation of the RLN</i>	<ul style="list-style-type: none"> <li>Avoid upper parathyroid injury</li> </ul>	
<ul style="list-style-type: none"> <li>Avoid muscle injury when making the midline incision</li> </ul>	<i>Identification of the common carotid artery</i>	<ul style="list-style-type: none"> <li>Identify the anatomical variations in the location of the parathyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Consider the various shapes of the RLN</li> </ul>	<i>Identification and preservation of EBSLN</i>	
<ul style="list-style-type: none"> <li>Incise from the thyroid cartridge to the suprasternal notch (or the location at which central node dissection is possible)</li> </ul>	<ul style="list-style-type: none"> <li>Identify the correct depth and course of the common carotid artery</li> </ul>	<ul style="list-style-type: none"> <li>Identify the blood stream distribution and blood vessel travel of the parathyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Continue to check the course of the RLN from view to view</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether the EBSLN is functional</li> </ul>	
<i>Identification of the trachea</i>	<ul style="list-style-type: none"> <li>Avoid the blood vessels around the common carotid artery</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether to leave the parathyroid or perform auto-transplantation after removal</li> </ul>	<ul style="list-style-type: none"> <li>Distinguish the artery from the RLN</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether the signals come from the EBSLN using nerve monitoring</li> </ul>	
<ul style="list-style-type: none"> <li>Avoid injury to the trachea</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether the common carotid artery moves well according to the heartbeat</li> </ul>	<ul style="list-style-type: none"> <li>If it is difficult to distinguish between the lymph nodes and parathyroid, determine whether to leave some or remove all depending on the cancer stage</li> </ul>	<ul style="list-style-type: none"> <li>Predict RLN location and angle</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether the cricothyroid muscle twitches</li> </ul>	
<ul style="list-style-type: none"> <li>Expose the trachea as much as possible</li> </ul>	<ul style="list-style-type: none"> <li>Determine whether the common carotid artery is well exposed along the thyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Avoid damage to the parathyroid gland and blood vessels leading to the parathyroid</li> </ul>	<ul style="list-style-type: none"> <li>Determine the intensity of pulling when the Berry ligament and RLN are adjacent</li> </ul>	<i>Ligation of the superior thyroid artery and vein</i>	
<i>Identification of the isthmus</i>	<i>Thyroid retraction—lower</i>	<i>Preservation of the blood stream of the parathyroid</i>	<ul style="list-style-type: none"> <li>Dissect the RLN while protecting it by covering it with a gauze ball to prevent thermal or mechanical injury</li> </ul>	<ul style="list-style-type: none"> <li>Adjust robotic arms for better visibility</li> </ul>	
<ul style="list-style-type: none"> <li>Avoid injury to the trachea</li> </ul>	<ul style="list-style-type: none"> <li>Accurately locate the parathyroid gland and RLN</li> </ul>	<ul style="list-style-type: none"> <li>Preserve blood vessels that affect the parathyroid</li> </ul>	<ul style="list-style-type: none"> <li>Strong retraction of the thyroid may damage the RLN</li> </ul>	<ul style="list-style-type: none"> <li>Rapidly expose and ligate the superior thyroid artery</li> </ul>	
<ul style="list-style-type: none"> <li>Confirm the location of the isthmus carefully</li> </ul>	<ul style="list-style-type: none"> <li>Avoid bleeding in the thyroid capsule</li> </ul>	<ul style="list-style-type: none"> <li>Ensure safe distances considering the range of heat conduction to prevent thermal injury</li> </ul>	<ul style="list-style-type: none"> <li>Avoid thermal injury</li> </ul>	<i>Other items related to dissection of the thyroid upper pole</i>	
<ul style="list-style-type: none"> <li>Determine whether the isthmus was visible as soon as the midline was opened from the sternothyroid muscle</li> </ul>	<ul style="list-style-type: none"> <li>Expose the lower pole and part of the upper part of the thyroid gland</li> </ul>	<ul style="list-style-type: none"> <li>Identify the inferior and middle thyroidal veins</li> </ul>	<ul style="list-style-type: none"> <li>Occasionally, a non-recurrent laryngeal nerve is present that drives directly into the vagus nerve from the upper part of the subclavian artery and enters the larynx</li> </ul>	<ul style="list-style-type: none"> <li>Use nerve monitoring to identify the vagus nerve (located close to the carotid artery)</li> </ul>	

Table 2 (continued)

Table 2 (continued)

MID module (n=18)	LAT module (n=16)	INF module (n=18)	BER module (n=22)	SUP module (n=12)	END module (n=3)
<i>Isthmectomy</i>	<ul style="list-style-type: none"> <li>Use a switching motion to support and lift the thyroid gland to check the tissue around the common carotid artery</li> </ul>	<ul style="list-style-type: none"> <li>Preserve the parathyroid as much as possible</li> </ul>	<ul style="list-style-type: none"> <li>Nerve monitoring determining the amplitude of the nerves when initially stimulated (whether the signal has been reduced by more than 50%)</li> </ul>		
<ul style="list-style-type: none"> <li>Preserve the inferior thyroid vein on the non-operative side</li> <li>Consider the location of the isthmus</li> <li>Avoid vessel injury (such as thyroid ima)</li> <li>Avoid injury to the cricoid cartilage</li> <li>Determine whether the left and right sides of the thyroid are separated</li> </ul>	<p><i>Other items related to lateral dissection</i></p> <ul style="list-style-type: none"> <li>Determine whether the central lymph nodes are removed cleanly along the thyroid gland</li> <li>Determine whether the middle thyroid vein is exposed and properly ligated</li> </ul>	<ul style="list-style-type: none"> <li>Avoid injury to the parathyroid and parathyroid feeding vessels</li> </ul> <p><i>Other items associated with the preservation of the inferior parathyroid glands</i></p> <ul style="list-style-type: none"> <li>Avoid retracting the parathyroid directly to prevent damage to the parathyroid</li> <li>If inevitable, retract tissues around the parathyroid or grab the blood vessels going to the parathyroid</li> </ul>	<p><i>Dissection of the ligament of Berry</i></p> <ul style="list-style-type: none"> <li>Expose Berry ligament sufficiently</li> <li>Determine the intensity of pulling when the Berry ligament and RLN are adjacent</li> <li>Dissect the thyroid gland below the Berry ligament</li> <li>Check the cricothyroid muscle in the upper area</li> </ul>		
<p><i>Other items related to midline incision and isthmectomy</i></p> <ul style="list-style-type: none"> <li>Determine preoperatively whether the isthmectomy is possible (if there is cancer on the isthmus itself, the isthmectomy position might have to be changed)</li> </ul>			<ul style="list-style-type: none"> <li>Minimize residual thyroid tissue, as microscopic amounts of thyroid tissue may remain when the thyroid and the RLN are attached, or when the thyroid tissue covers the RLN, similar to the ears</li> <li>Hemostasis is difficult if bleeding occurs in the Berry ligament</li> <li>Determine whether the Berry ligament is well removed while protecting the RLN</li> </ul> <p><i>Other items related to preservation of the RLN, dissection of the ligament of Berry</i></p> <ul style="list-style-type: none"> <li>Use a compression method with energy or a gauze ball for hemostasis</li> </ul>		

MID, midline incision to isthmectomy; LAT, lateral dissection; INF, preservation of inferior parathyroid glands; BER, preservation of RLN and dissection of the ligament of Berry; SUP, dissection of the thyroid upper pole; END, specimen removal and closure; RLN, recurrent laryngeal nerve; EBSLN, external branch of superior laryngeal nerve.



MID (n=8; M=5.83)	LAT (n=7; M=5.56)	INF (n=16; M=5.99)	BER (n=21; M=5.94)	SUP (n=10; M=5.83)	END (n=2; M=6.15)
Identification of midline (n=0) Retract midline bilaterally with both graspers (n=0) Midline incision (n=2; M=5.45) Identification of trachea (n=2; M=5.98) Identification of isthmus (n=2; M=5.85) Isthmectomy (n=1; M=5.55) Other items related to midline incision and isthmectomy (n=1; M=6.5)	Dissection of surgical plane between thyroid and strap muscle (n=0) Lateral retraction of the strap muscle (n=2; M=5.45) Identification of common carotid artery (n=1; M=5.2) Thyroid retraction—lower (n=3; M=5.9) Other items related to lateral dissection (n=1; M=5.15)	Identification of RLN (n=4; M=6.38) Identification of inferior parathyroid gland (n=7; M=5.75) Preservation of the blood stream of parathyroid (n=4; M=6.19) Other items to preservation of inferior parathyroid glands (n=1; M=5.3)	Dissection between medial thyroid and trachea (n=1; M=5.5) Dissection between thyroid gland and fascia (n=0) Thyroid retraction—Zuckermandl (n=2; M=6) Preservation of RLN (n=10; M=6.13) Dissection of ligament of Berry (n=7; M=5.77) Other items related to preservation of RLN, dissection of the ligament of Berry (n=1; M=5.5)	Dissection in the upward direction (n=2; M=6.45) Identification and preservation of superior parathyroid glands (n=3; M=5.93) Identification and preservation of EBSLN (n=3; M=5.5) Ligation of superior thyroid artery and vein (n=5.7) Other items related to dissection of the thyroid upper pole (n=0)	Specimen out (n=2; M=6.15) Use of hemostatic dressing and anti-adhesion adjuvant (n=0) Drain insertion and midline closure (n=0)

**Figure 2** Modules of RT. n, the number of items; M, mean of importance scores; MID, midline incision to isthmectomy; LAT, lateral dissection; INF, preservation of inferior parathyroid glands; BER, preservation of RLN and dissection of the ligament of Berry; SUP, dissection of the thyroid upper pole; END, specimen removal and closure; RLN, recurrent laryngeal nerve; RT, robotic thyroidectomy.

comment regarding the LAT module consisted of accurate determination of the locations of the parathyroid gland and RLN, whereas additional comments regarding the INF module included (I) identifying the course and location of the RLN, (II) ensuring safe distances from the heat source based on the range of heat conduction thereby preventing thermal injury, (III) not retracting the thyroid gland excessively as it may cause mechanical injury to the RLN. Additional comments regarding the BER module included (I) checking the course of the RLN from view to view, (II) avoiding thermal injury, and (III) avoiding RLN injury caused by traction, whereas an additional comment regarding the SUP module noted that to prevent external branch of superior laryngeal nerve (EBSLN) injury, dissection should be made as close to the thyroid gland

as possible while ligating the upper thyroid artery well (Table S3).

## Discussion

The present study utilized CTA to identify 89 items within six modules for RT, with the modified Delphi survey identifying the 64 items finally in these six modules most important for RT performance, including 8, 7, 16, 21, 10, and 2 items in the MID, LAT, INF, BER, SUP, and END modules, respectively. The core performance skills and cognitive strategies required to perform BABA RT were determined using a mixed research method.

The list of procedural tasks and non-technical skills defined throughout this study may function as the basis

**Table 3** Results of modified Delphi consensus on items required to perform robotic thyroidectomy (top 10)

Final rank	Module	Items	Round 1			Round 2	
			Mean (SD)	% rating over 5	Rank	Mean (SD)	% rating over 5
1	BER	Continue to check the course of the RLN from view to view	6.67 (0.64)	100	3	6.85 (0.65)	95
2	LAT	Accurately locate the parathyroid gland and RLN	6.76 (0.61)	100	2	6.8 (0.68)	95
2	INF	Identify the course and location of the RLN	6.81 (0.50)	100	1	6.8 (0.51)	100
4	INF	Ensure safe distances based on the range of heat conduction to prevent thermal injury	6.67 (0.56)	100	3	6.7 (0.56)	100
5	SUP	To prevent EBSLN injury, dissect as close to the thyroid as possible, and at the same time, ligate the upper thyroid artery well	6.52 (0.59)	100	6	6.55 (0.50)	100
6	MID	Determine whether isthmectomy is possible preoperatively (if there is a cancer on the isthmus itself, the isthmectomy position might be changed)	6.38 (0.79)	100	12	6.5 (0.81)	95
6	BER	Avoid thermal injury	6.52 (0.66)	100	6	6.5 (0.50)	100
8	MID	Avoid injury to the trachea when identifying the isthmus	6.43 (0.85)	100	11	6.45 (0.67)	100
8	INF	Excess retraction of the thyroid gland may cause mechanical injury to the RLN	6.19 (0.91)	90	20	6.45 (0.86)	95
8	BER	Avoid RLN injury caused by traction	6.48 (0.73)	100	10	6.45 (0.74)	100

SD, standard deviation; BER, preservation of RLN and dissection of the ligament of Berry; RLN, recurrent laryngeal nerve; LAT, lateral dissection; INF, preservation of inferior parathyroid glands; SUP, dissection of the thyroid upper pole; EBSLN, external branch of superior laryngeal nerve; MID, midline incision to isthmectomy.


for developing a standardized and valid training program, which can enhance surgeon proficiency in surgical skills and ultimately ensure patient safety. In particular, we attempted to evaluate the factors associated with competency-based medical education, including knowledge, skills, values, and attitudes, which could be reflected in both the behavioral tasks and cognitive schema of RT (4,23). The skills learned in the operative field are based on both situational and practice-based learning, both of which could affect the development of professional identity throughout the proper training and evaluation as a thyroid surgeon.

Previous studies have attempted to define the competencies required for each targeted medical treatment, including open thyroidectomy, radiation protection, advanced care planning, endoscopic submucosal dissection, and laparoscopic transabdominal adrenalectomy (2-4,24-26). In some of these studies, when the researchers developed the items for the Delphi consensus, they only used literature review or task analyses of their own performances (24-26).

In addition, two of these studies analyzed the performance of multiple SMEs, but did not apply modified Delphi methodology for validation (2,4). This study adopted a mixed-method approach, differentiating this study from previous research, enabling the collection of as much unbiased information as possible and determining the most important competencies for RT, as assessed by various experts, thereby improving its validity (3).

This study, however, did not assess pre-operative steps, such as adjusting the settings on the robot. Although intraoperative patient care includes pre-operative preparation and post-operative management, this study assessed operative skills and techniques with the surgical robot. Pre-operative preparation for RT may include steps associated with the ease of tool usage, as settings for the surgical robot before the operative procedure can be important factors. Another consideration is that our study is demographically homogeneous. Since BABA RT is a surgical method that originated in Korea, this initial





**Figure 3** Top 5 items of each modules. MID, midline incision to isthmectomy; LAT, lateral dissection; INF, preservation of inferior parathyroid glands; BER, preservation of RLN and dissection of the ligament of Berry; SUP, dissection of the thyroid upper pole; END, specimen removal and closure; RLN, recurrent laryngeal nerve; EBSLN, external branch of superior laryngeal nerve.

Delphi study had to focus on expert surgeons in Korea. However, presently, the same surgery is being conducted on individuals of different races in various countries worldwide. With the insights gained from this study, it will be possible to develop surgical guidelines from a global perspective in the future. Nevertheless, this study focused on the surgical process and analyzed the surgeon's cognitive behavior when performing RT, resulting in meaningful results, regarded as offsetting the preoperative process. The unique features of RT that is distinct from open thyroidectomy identified in the present study were the proficient use of robotic instruments and the method of

securing the operative field visually. These characteristics may be regarded as results associated with differences in surgical tools.

Surgeons should be mindful that, despite their proficiency in the surgical skills we provide, complications may arise in rare cases. A recent analysis of a large-scale robotic surgery study revealed rare complications in 60 out of 5,011 patients. These rare complications comprised hematoma in 4 cases (0.44%), chyle leakage in 15 cases (0.3%), flap injury in 4 cases (0.08%), RLN injury in 7 cases (0.14%), open conversion in 8 cases (0.16%), and pneumothorax in 4 cases (0.08%) (27).

## Conclusions

Because of the lack, to date, of step-by-step surgical procedure guidelines for BABA-RT, the results of this study may be used to develop standardized educational criteria for training novice surgeons sufficiently to perform operations independently. These systematic and evidence-based procedures for RT could reduce the quality gap in accordance with training settings, and may contribute to the long-term narrowing of medical gaps among communities and surgeons. It might also help RT trainees to form accurate mental representations of successful performance and improve their surgical skills based on more detailed feedback.

## Acknowledgments

The authors thank all expert panel members participating in the Delphi process.

*Funding:* This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Science, ICT & Future Planning, Republic of Korea (No. 2018R1A1A1A05077361); the Korea Medical Device Development Fund grant funded by the Korea Government (the Ministry of Science and ICT, the Ministry of Trade, Industry and Energy, the Ministry of Health & Welfare, the Ministry of Food and Drug Safety) (No. RS-2020-KD000146); the MSIT (Ministry of Science and ICT), Korea, under the ITRC (Information Technology Research Center) support program (No. IITP-2021-2018-0-01833) supervised by the IITP (Institute for Information & Communications Technology Planning & Evaluation).

## Footnote

*Data Sharing Statement:* Available at <https://gs.amegroups.com/article/view/10.21037/gS-23-467/dss>

*Peer Review File:* Available at <https://gs.amegroups.com/article/view/10.21037/gS-23-467/prf>

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://gs.amegroups.com/article/view/10.21037/gS-23-467/coif>). The authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all

aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study protocol was approved by the Institutional Review Board of Seoul National University Hospital (No. H-1912-116-1090) and informed consent was taken from all the participants.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Ryan MS, Lomis KD, Deiorio NM, et al. Competency-Based Medical Education in a Norm-Referenced World: A Root Cause Analysis of Challenges to the Competency-Based Paradigm in Medical School. *Acad Med* 2023;98:1251-60.
2. Madani A, Grover K, Kuo JH, et al. Defining the competencies for laparoscopic transabdominal adrenalectomy: An investigation of intraoperative behaviors and decisions of experts. *Surgery* 2020;167:241-9.
3. Madani A, Watanabe Y, Vassiliou M, et al. Defining competencies for safe thyroidectomy: An international Delphi consensus. *Surgery* 2016;159:86-94, 96-101.
4. Madani A, Vassiliou MC, Watanabe Y, et al. What Are the Principles That Guide Behaviors in the Operating Room?: Creating a Framework to Define and Measure Performance. *Ann Surg* 2017;265:255-67.
5. Melchioris J, Henriksen MJV, Dijkers FG, et al. Diagnostic flexible pharyngo-laryngoscopy: development of a procedure specific assessment tool using a Delphi methodology. *Eur Arch Otorhinolaryngol* 2018;275:1319-25.
6. Madani A, Gips A, Razek T, et al. Defining and Measuring Decision-Making for the Management of Trauma Patients. *J Surg Educ* 2018;75:358-69.
7. Madani A, Gornitsky J, Watanabe Y, et al. Measuring

- Decision-Making During Thyroidectomy: Validity Evidence for a Web-Based Assessment Tool. *World J Surg* 2018;42:376-83.
8. Chorath KT, Luu NN, Douglas JE, et al. Assessment of YouTube as an educational tool in teaching thyroidectomy and parathyroidectomy. *J Laryngol Otol* 2022;136:952-60.
  9. Larkins K, Khan M, Mohan H, et al. A systematic review of video-based educational interventions in robotic surgical training. *J Robot Surg* 2023;17:1329-39.
  10. Li C, Gao Y, Zhou P, et al. Comparison of the Robotic Bilateral Axillo-Breast Approach and Conventional Open Thyroidectomy in Pediatric Patients: A Retrospective Cohort Study. *Thyroid* 2022;32:1211-9.
  11. Fassari A, Gurrado A, Iossa A, et al. Definition of learning curve for thyroidectomy: systematic review on the different approaches. *Gland Surg* 2023;12:989-1006.
  12. Liu SY, Kim JS. Bilateral axillo-breast approach robotic thyroidectomy: review of evidences. *Gland Surg* 2017;6:250-7.
  13. Lee J, Yun JH, Choi UJ, et al. Robotic versus Endoscopic Thyroidectomy for Thyroid Cancers: A Multi-Institutional Analysis of Early Postoperative Outcomes and Surgical Learning Curves. *J Oncol* 2012;2012:734541.
  14. Kim WW, Jung JH, Park HY. The Learning Curve for Robotic Thyroidectomy Using a Bilateral Axillo-Breast Approach From the 100 Cases. *Surg Laparosc Endosc Percutan Tech* 2015;25:412-6.
  15. Youn YK, Lee KE, Choi JY. Robotic Thyroidectomy: Bilateral Axillo-Breast Approach (BABA). In: Youn YK, Lee KE, Choi JY, editors. *Color Atlas of Thyroid Surgery: Open, Endoscopic and Robotic Procedures*. Berlin, Heidelberg: Springer; 2014:83-125.
  16. You JY, Kim HK, Kim HY, et al. Bilateral axillo-breast approach robotic thyroidectomy: review of a single surgeon's consecutive 317 cases. *Gland Surg* 2021;10:1962-70.
  17. Kang YJ, Stybayeva G, Hwang SH. Surgical safety and effectiveness of bilateral axillo-breast approach robotic thyroidectomy: a systematic review and meta-analysis. *Braz J Otorhinolaryngol* 2023. [Epub ahead of print]. doi: 10.1016/j.bjorl.2023.101376.
  18. Swaby L, Shu P, Hind D, et al. The use of cognitive task analysis in clinical and health services research - a systematic review. *Pilot Feasibility Stud* 2022;8:57.
  19. Korean Association of Thyroid and Endocrine Surgeons. *Textbook of Endocrine Surgery*. Seoul, Republic of Korea: Koonja Publisher; 2018.
  20. Zollinger R, Ellison E, Pawlik T, et al. *Zollinger's Atlas of Surgical Operations*. 11th edition. McGraw Hill; 2021.
  21. Persemlidis D, Inabnet WB 3rd, Gagner M. *Endocrine Surgery*. 2nd edition. Boca Raton: CRC Press; 2016.
  22. Nasa P, Jain R, Juneja D. Delphi methodology in healthcare research: How to decide its appropriateness. *World J Methodol* 2021;11:116-29.
  23. The Accreditation Council for Graduate Medical Education. *Surgery Milestones*. 2019.
  24. Doyen B, Maurel B, Cole J, et al. Defining the Key Competencies in Radiation Protection for Endovascular Procedures: A Multispecialty Delphi Consensus Study. *Eur J Vasc Endovasc Surg* 2018;55:281-7.
  25. Sudore RL, Heyland DK, Lum HD, et al. Outcomes That Define Successful Advance Care Planning: A Delphi Panel Consensus. *J Pain Symptom Manage* 2018;55:245-255.e8.
  26. Takao M, Bilgic E, Waschke K, et al. Defining competencies for endoscopic submucosal dissection (ESD) for gastric neoplasms. *Surg Endosc* 2019;33:1206-15.
  27. Kwak J, Yu HW, Ahn JH, et al. A Time Trend Analysis of 5,000 Robotic Thyroidectomies via Bilateral Axillo-Breast Approach. *World J Surg* 2023;47:403-11.

**Cite this article as:** Kim H, Yu HW, Ahn JH, Lee TS, Lee KE. Defining competencies in robotic thyroidectomy: development of a model assessing an expert operator's intraoperative performance skills and cognitive strategies. *Gland Surg* 2024;13(3):340-350. doi: 10.21037/gs-23-467