

CORRIGENDUM

The clavipectoral fascia as the unique anatomical criteria for distinguishing breast parenchymal lesions from axillary lymph node metastasis

Antonios Patrinos^{1,2,*}, Maria Zarokosta^{1,2}, John Tsiaoussis³, George Noussios⁴, Theodoros Piperos^{1,2}, Menelaos Zoulamoglou¹, Ioannis Flessas^{1,2}, and Theodoros Mariolis-Sapsakos^{1,2}

¹General and Oncologic Hospital of Kifissia 'Agioli Anargyroi', Athens, Greece ²Anatomy and Histology Laboratory, Nursing School of the University of Athens, Greece ³Laboratory of Anatomy-Histology-Embryology Medical School of Heraklion, University of Crete, Greece, and ⁴Department of Physical Education and Sport Science, 'Aristoteles University' of Thessaloniki, Serres, Greece

*Correspondence address. Anatomy and Histology Laboratory, Nursing School, National and Kapodistrian University of Athens, Greece, Papadiamantopoulou 123, Goudi, Athens, Greece, P.C:15773. Tel/Fax: 210-7461483; E-mail: antopatrinis@gmail.com

Journal of Surgical Case Reports, 2019;5, 1–3; doi: 10.1093/jscr/rjz135

In the original version of this report the affiliation of Dr. John Tsiaoussis was incorrectly listed. This has now been corrected and the author apologises for the error.