

Assessment of Changes in Body Composition After 3 Months of Dulaglutide Treatment [Response to Letter]

Shuqin Chen^{1,*}, Xuepeng Wang^{2,*}, Yong Jin³, Xueqin Chen⁴, Qifa Song⁵, Gang Wei⁶, Li Li¹

¹Department of Endocrinology and Metabolism, The First Affiliated Hospital of Ningbo University, Ningbo, People's Republic of China; ²Department of Infectious Disease, The First Affiliated Hospital of Ningbo University, Ningbo, People's Republic of China; ³Department of Internal Medicine, Ningbo Yinzhou No. 2 Hospital, Ningbo, People's Republic of China; ⁴Department of Traditional Medicine, The First Affiliated Hospital of Ningbo University, Ningbo, People's Republic of China; ⁵Medical Data Center, The First Affiliated Hospital of Ningbo University, Ningbo, People's Republic of China; ⁶Beijing Diabetes Institute, Beijing Key Laboratory of Diabetes Research and Care, Department of Endocrinology, Beijing Tongren Hospital, Capital Medical University, Beijing, People's Republic of China

*These authors contributed equally to this work

Correspondence: Li Li, Department of Endocrinology and Metabolism, The First Affiliated Hospital of Ningbo University, Ningbo, 31500, People's Republic of China, Email lilyningbo@163.com; Gang Wei, Beijing Diabetes Institute, Beijing Key Laboratory of Diabetes Research and Care, Department of Endocrinology, Beijing Tongren Hospital, Capital Medical University, Beijing, 100000, People's Republic of China, Email gangwei_2013@163.com

Dear editor

We thank Panjaitan et al for their thoughtful and constructive comments on our recent report about dulaglutide.¹ We really appreciate the interesting highlights they have summarized.

We agree with the author that a dose-escalation experimental design is needed to allow patients to tolerate the adverse effects of the drug and to compare the efficacy of the drug at different doses. The Food and Drug Administration (FDA) has approved 4 doses (0.75 mg, 1.5 mg, 3 mg, and 4.5 mg) of dulaglutide for the treatment of type 2 diabetes in adults. Now dulaglutide (Eli Lilly) is available in 0.75 mg and 1.5 mg doses in China. And only 0.5 mL: 1.5 mg dulaglutide (Trulicity, Eli Lilly) is currently available at the subject's hospital. There is no 0.75 mg or other high-dose formulations available at the subject's hospital. Side effects, especially gastrointestinal side effects, are relatively rare and tolerated by patients in the practical use of this size of drug (0.5 mL: 1.5 mg, Trulicity, Eli Lilly) in the subject's hospital. A previous study also showed that dose titration of dulaglutide (LY2189265) did not result in a clear reduction in gastrointestinal side effects, as measured by the Gastroparesis Cardinal Symptom Index.² Therefore, 1.5 mg dulaglutide without dose titration was used in our study.

It is generally accepted that dulaglutide has a dose-dependent therapeutic effect, whether for glucose lowering or weight loss. In the AWARD-CHN2 study, the 1.5 mg dose of dulaglutide appears to be more effective in glycemic control than the 0.75 mg dose.³ A recent systematic review and meta-analysis conducted by Li et al showed that higher dose of dulaglutide have greater effects on body weight and waist circumference reduction.⁴ It would be interesting to have the opportunity to investigate whether the effect on body composition is also dose dependent.

We agree with the reviewer's concern that a 3-month follow-up period is relatively short. The appropriate follow-up period was an important issue to consider when we designed the study. The choice of a 3-month follow-up period was a reasonable choice based on our preliminary review of the relevant literature. Data from previous studies of dulaglutide showed that the fastest period of weight change after the use of 1.5 mg dulaglutide was the first 3 months (12 weeks), after which weight change entered a slow period.⁵ The aim of our study was to find out whether the weight change was accompanied by bone and muscle loss. Taking all factors into account, 3 months was chosen as the follow-up period for this study.

We are very happy that the readers are interested in our work. However, our current research work is still insufficient to explain many important issues. These questions include: 1. The reason for the change in body composition by dulaglutide is not clear. 2. Is there a rebound of body weight and corresponding changes in body composition after stopping the use of dulaglutide? As mentioned by the reviewers in the letter, large clinical trials with long-term follow-up are needed to address these questions.

Again, we appreciate the constructive feedback from the reviewers and look forward to more interesting and exciting results in the future.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Chen S, Wang X, Jin Y, et al. Assessment of changes in body composition after 3 months of dulaglutide treatment. *Diabetes Metab Syndr Obes.* 2024;17:1301–1308. doi:10.2147/DMSO.S443631
2. Umpierrez GE, Blevins T, Rosenstock J, Cheng C, Anderson JH, Bastyr EJ. The effects of LY2189265, a long-acting glucagon-like peptide-1 analogue, in a randomized, placebo-controlled, double-blind study of overweight/obese patients with type 2 diabetes: the EGO study. *Diabetes Obes Metab.* 2011;13(5):418–425. doi:10.1111/j.1463-1326.2011.01366.x
3. Li Y, Li L, De Peng Y, et al. Efficacy and safety of dulaglutide versus insulin glargine in Chinese T2DM patients: a subgroup analysis of a randomized trial (AWARD-CHN2). *Diabetes Ther.* 2019;10(4):1435–1452. doi:10.1007/s13300-019-0646-y
4. Li Y, Gong X, Găman M-A, Hernández-Wolters B, Velu P, Li Y. The effect of subcutaneous dulaglutide on weight loss in patients with type 2 diabetes mellitus: systematic review and meta-analysis of randomized controlled trials. *Eur J Clin Invest.* 2024;54(4):e14125. doi:10.1111/eci.14125
5. Frias JP, Bonora E, Nevarez Ruiz L, et al. Efficacy and safety of dulaglutide 3.0 mg and 4.5 mg versus dulaglutide 1.5 mg in metformin-treated patients with type 2 diabetes in a randomized controlled trial (AWARD-11). *Diabetes Care.* 2021;44(3):765–773. doi:10.2337/dc20-1473

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Diabetes, Metabolic Syndrome and Obesity 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Diabetes, Metabolic Syndrome and Obesity editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>

<https://doi.org/10.2147/DMSO.S473832>