

POSTER PRESENTATION

Open Access

# Modulation of anti-tumor lymphocyte function by neurotransmitter glutamate

Thomas Hodo<sup>1</sup>, Menaka Thounaojam<sup>1</sup>, Anil Shanker<sup>2\*</sup>

From Society for Immunotherapy of Cancer 29th Annual Meeting  
National Harbor, MD, USA. 6-9 November 2014

Most research to date pertaining to neural influence on immune response involves immunosuppression via the anti-inflammatory pathway. However, there is emerging evidence indicating that neurotransmitters have the ability to promote immune activation. We are investigating whether neurotransmitters can modulate and/or activate T cell function in situations where immunosuppression is prevalent such as in the tumor microenvironment. Published work suggests that glutamate, serotonin, dopamine, and Substance P trigger immune responses such as cytokine secretion, integrin expression, and chemotaxis. We saw that both CD4 and CD8 T cells express high surface protein levels of glutamate receptor AMPA iGluR3, which is able to import  $\text{Ca}^{2+}$  and  $\text{Na}^+$ . We also found that mGluR1 is significantly upregulated on lymphocytes upon activation. Our data further show that T cells in the tumor-draining LN and tumor-infiltrating lymphocytes have upregulated expression of iGluR3 and mGluR1. Treatment with glutamate or its receptor agonist augmented T cell proliferation following CD3-CD28-mediated TCR stimulation. Thus, modulation of glutamate receptor signaling can be useful for enhancing anti-tumor T cell immunity such as inhibition of AICD, enhancement of proliferation, and increased cytokine production [1,2]. Indeed, overactivation of lymphocytes in multiple sclerosis is closely tied to the overexpression of AMPA GluR3 on T cells [3]. Experiments are under way to dissect in an adoptive transfer set up whether glutamate-modulated immune effector function involves specific activation of anti-tumor lymphocytes to elicit cytolytic response that is needed to cause tumor cell death. Our findings will help identify novel neuro-immune modulators that may serve to enhance anti-tumor T cell response.

## Authors' details

<sup>1</sup>Meharry Medical College School of Medicine, Nashville, TN, United States.

<sup>2</sup>Meharry Medical College School of Medicine / Vanderbilt-Ingram Cancer Center, Nashville, TN, United States.

Published: 6 November 2014

## References

1. Chiocchetti A, Miglio G, Mesturini R, Varsaldi F, Mocellin M, Orilieri E, Dianzani C, Fantozi R, Dianzani U, Lombardi G: Group I mGlu receptor stimulation inhibits activation-induced cell death of human T lymphocytes. *British journal of pharmacology* 2006, **148**:760-768.
2. Pacheco R, Oliva H, Martinez-Navio JM, Climent N, Ciruela F, Gatell JM, Gallart T, Mallol J, Lluis C, Franco R: Glutamate released by dendritic cells as a novel modulator of T cell activation. *J Immunol* 2006, **177**:6695-6704.
3. Sarchielli P, Di Filippo M, Candeliere A, Chiasserini D, Mattioni A, Tenaglia S, Bonucci M, Calabresi P: Expression of ionotropic glutamate receptor GLUR3 and effects of glutamate on MBP- and MOG-specific lymphocyte activation and chemotactic migration in multiple sclerosis patients. *Journal of neuroimmunology* 2007, **188**:146-158.

doi:10.1186/2051-1426-2-S3-P38

Cite this article as: Hodo et al.: Modulation of anti-tumor lymphocyte function by neurotransmitter glutamate. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P38.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>2</sup>Meharry Medical College School of Medicine / Vanderbilt-Ingram Cancer Center, Nashville, TN, United States  
Full list of author information is available at the end of the article