

CORRECTION

Open Access

Correction to: Advanced imaging in adult diffusely infiltrating low-grade gliomas



Nail Bulakbaşı^{1*} and Yahya Paksoy²

Correction to: Insights Imaging

<https://doi.org/10.1186/s13244-019-0793-8>

The original article "Advanced imaging in adult diffusely infiltrating low-grade gliomas" contains errors in Table 1 in rows k_{trans} and V_e ; the correct version of Table 1 can be viewed in this Correction article.

The original article can be found online at <https://doi.org/10.1186/s13244-019-0793-8>.

* Correspondence: nail.bulakbasi@kyrenia.edu.tr

¹Medical Faculty, University of Kyrenia, Sehit Yahya Bakir Street, Karakum, Mersin-10, Kyrenia, Turkish Republic of Northern Cyprus, Turkey
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Table 1 Radiomic data for differential diagnosis of low-grade vs high-grade gliomas

Parameters	Low-grade glioma	High-grade glioma	Cut of value ^{Ref} (Range) ^{Ref}
rCBV _{max}	Low	High	1.76 [10, 19] (0.94–3.34) [9–11, 19–21]
rCBV _{MD}	Low	High	1.44 [22] (1.08–1.81) [22]
nADC _{min}	High	Low	1.07 × 10 ⁻³ mm ² /s [24] (0.31–1.31) [23–28]
Cho/Cr ratio	Low	High	1.56 [19] (1.3–2.04) [29–31]
MK _{MD}	Low	High	0.17 [28] (0.11–0.28) [28]
FA _{TC}	Low	High	0.3 [25] (0.14–0.63) [25]
MD _{min}	High	Low	0.98 mm ² /s [25] (0.76–0.91) [25]
k _{trans}	Low	High	1.18 [22] (0.91–1.45) [22]
V _e	Low	High	1.43 [22] (1.06–1.80) [22]
ITTS grade	1.2	2.6	NA [9]
APT signal (%)	Low	High	2.23% [15] (1.53%–3.70%) [15]

rCBV_{max} maximum relative cerebral blood volume, *rCBV_{MD}* standardized mean difference of *rCBV_{max}*, *nADC_{min}* Normalized minimum apparent diffusion coefficient, *Cho/Cr* Choline/Creatine, *MK_{MD}* mean difference in mean kurtosis, *FA_{TC}* Odds ratio of fractional anisotropy in the tumor core, *MD_{min}* Minimum mean diffusivity, *k_{trans}* standardized mean difference of volume transfer coefficient, *V_e* standardized mean difference of volume fraction of extravascular extracellular space, *ITTS* Intratumoral susceptibility score, *APT* Percent amide proton transfer signal

Author details

¹Medical Faculty, University of Kyrenia, Sehit Yahya Bakir Street, Karakum, Mersin-10, Kyrenia, Turkish Republic of Northern Cyprus, Turkey. ²Selcuk University, Konya, Turkey.

Published online: 22 April 2020

Reference

- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK (2016) WHO classification of tumors of the central nervous system, revised 4th edition. IARC, Lyon.
- Sahm F, Reuss D, Koelsche C et al (2014) Farewell to oligoastrocytoma: in situ molecular genetics favor classification as either oligodendrogloma or astrocytoma. Acta Neuropathol 128(4):551–559. <https://doi.org/10.1007/s00401-014-1326-7>
- Bready D, Placantonakis DG (2019) Molecular pathogenesis of low-grade glioma. Neurosurg Clin N Am 30(1):17–25. <https://doi.org/10.1016/j.nec.2018.08.011>
- Duffau H (2016) Long-term outcomes after supratotal resection of diffuse low-grade gliomas: a consecutive series with 11-year follow-up. Acta Neurochir (Wien) 158(1):51–58. <https://doi.org/10.1007/s00701-015-2621-3>
- National Comprehensive Cancer Network. Central nervous system cancers (Version 1.2019). https://www.nccn.org/professionals/physician_gls/pdf/cns.pdf. Accessed Mar 5 2019
- Olson JJ, Kalkanis SN, Ryken TC (2015) Evidence-based clinical practice parameter guidelines for the treatment of adults with diffuse low-grade glioma: introduction and methods. J Neurooncol 125(3):449–456. <https://doi.org/10.1007/s11060-015-1847-5>
- Fouke SJ, Benzinger T, Gibson D, Ryken TC, Kalkanis SN, Olson JJ (2015) Therole of imaging in the management of adults with diffuse low-grade glioma: a systematic review and evidence-based clinical practice guideline. J Neurooncol 125(3):457–479. <https://doi.org/10.1007/s11060-015-1908-9>
- Caulo M, Panara V, Tortora D et al (2014) Data-driven grading of brain gliomas: a multiparametric MR imaging study. Radiology 272(2):494–503. <https://doi.org/10.1148/radiol.14132040>
- Li X, Zhu Y, Kang H et al. (2015) Glioma grading by microvascular permeability parameters derived from dynamic contrast-enhanced MRI and intratumoral susceptibility signal on susceptibility weighted imaging. Cancer Imaging 15:4. <https://doi.org/10.1166/s40644-015-0039-z>
- Delgado AF, Delgado AF (2017) Discrimination between glioma grades I and III using dynamic susceptibility contrast perfusion MRI: a meta-analysis. AJNR Am J Neuroradiol 38(7):1348–1355. <https://doi.org/10.3174/ajnr.A5218>
- Kong L, Chen H, Yang Y, Chen L (2017) A meta-analysis of arterial spin labeling perfusion values for the prediction of glioma grade. Clin Radiol 72(3):255–261. <https://doi.org/10.1016/j.crad.2016.10.016>
- ACR Committee on Drugs and Contrast Media (2018) ACR manual on contrast media (Version 10.3, 2018). https://www.acr.org/-/media/ACR/Files/Clinical-Resources/Contrast_Media.pdf. Accessed 2018.
- Choi C, Ganji SK, DeBerardinis RJ et al. (2012) 2-hydroxyglutarate detection by magnetic resonance spectroscopy in IDH-mutated patients with gliomas. Nat Med 18(4):624–629. <https://doi.org/10.1038/nm.2682>
- Tietze A, Choi C, Mickey B et al. (2018) Noninvasive assessment of isocitrate dehydrogenase mutation status in cerebral gliomas by magnetic resonance spectroscopy in a clinical setting. J Neurosurg 128(2):391–398. <https://doi.org/10.3171/2016.10.JNS161793>
- Suh CH, Park JE, Jung SC, Choi CG, Kim SJ, Kim HS (2019) Amide proton transfer-weighted MRI in distinguishing high- and low-grade gliomas: a systematic review and meta-analysis. Neuroradiology 61(5):525–534. <https://doi.org/10.1007/s00234-018-02152-2>
- Park JE, Kim HS, Park KJ, Choi CG, Kim SJ (2015) Histogram analysis of amide proton transfer imaging to identify contrast-enhancing low-grade brain tumor that mimics high-grade tumor: increased accuracy of MR perfusion. Radiology 277(1):151–161. <https://doi.org/10.1148/radiol.2015142347>
- Biller A, Badde S, Nagel A et al. (2016) Improved brain tumor classification by sodium MR imaging: prediction of IDH mutation status and tumor progression. AJNR Am J Neuroradiol 37(1):66–73. <https://doi.org/10.3174/ajnr.A4493>
- Pepin KM, McGee KP, Arani A et al. (2018) MR elastography analysis of glioma stiffness and IDH1-mutation status. AJNR Am J Neuroradiol 39(1):31–36. <https://doi.org/10.3174/ajnr.A5415>
- Law M, Yang S, Wang H et al. (2003) Glioma grading: sensitivity, specificity, and predictive values of perfusion MR imaging and proton MR spectroscopic imaging compared with conventional MR imaging. AJNR Am J Neuroradiol 24(10):1989–1998
- Bulakbasi N, Kocaoglu M, Farzaliyev A, Tayfun C, Ucoz T, Somuncu I (2005) Assessment of diagnostic accuracy of perfusion MR imaging in primary and metastatic solitary malignant brain tumors. AJNR Am J Neuroradiol 26(9):2187–2199
- Anzalone N, Castellano A, Cadioli M et al. (2018) Brain gliomas: multicenter standardized assessment of dynamic contrast-enhanced and dynamicsusceptibility contrast MR images. Radiology 287(3):933–943. <https://doi.org/10.1148/radiol.2017170362>
- Liang J, Liu D, Gao P et al. (2018) Diagnostic values of DCE-MRI and DSC-MRI for differentiation between high-grade and low-grade gliomas: a comprehensive meta-analysis. Acad Radiol 25(3):338–348. <https://doi.org/10.1016/j.acra.2017.10.001>
- Bulakbasi N, Guvenç I, Onguru O, Erdogan E, Tayfun C, Ucoz T (2004) The added value of the apparent diffusion coefficient calculation to magnetic resonance imaging in the differentiation and grading of malignant brain tumors. J Comput Assist Tomogr 28(6):735–746
- Server A, Kulle B, Gadmar ØB, Josefson R, Kumar T, Nakstad PH (2011) Measurements of diagnostic examination performance using quantitative apparent diffusion coefficient and proton MR spectroscopic imaging in the preoperative evaluation of tumor grade in cerebral gliomas. Eur J Radiol 80(2):462–470. <https://doi.org/10.1016/j.ejrad.2010.07.017>

