



Alteration of interoceptive sensitivity: expanding the spectrum of behavioural disorders in amyotrophic lateral sclerosis

Pasquale Moretta¹ · Myriam Spisto² · Francesco Pio Ausiello² · Rosa Iodice² · Natascia De Lucia² · Gabriella Santangelo³ · Luigi Trojano³ · Elena Salvatore² · Raffaele Dubbioso² 

Received: 2 April 2022 / Accepted: 17 June 2022 / Published online: 25 June 2022
© The Author(s) 2022

Abstract

Background Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disorder with progressive loss of upper and lower motor neurons. Non-motor-symptoms, such as cognitive, emotional, autonomic, and somatosensory alterations, have been also described. Interoception represents the link between the body and brain, since it refers to the ability to consciously perceive the physical condition of the inner body, including one's heartbeat (i.e., interoceptive sensitivity, IS).

Objectives To evaluate IS in ALS patients by means of a well-established task: the heartbeat perception task. Moreover, we evaluated possible correlations between IS and neuropsychological, affective, and disease-related characteristics.

Methods Fifty-five ALS patients (mean-age = 60.3 ± 12.5 years; mean disease-duration = 20.9 ± 18.8 months) and 41 caregivers (CG) underwent the heartbeat perception task and an extensive evaluation of motor, cognitive, body awareness, affective, and emotion domains.

Results ALS patients showed lower IS than CG (0.68 ± 0.24 vs 0.82 ± 0.16 ; $p = 0.003$). Significant correlations were found between IS and self-reported measures of alexithymia (subscale of Toronto Alexithymia scale-20 “difficulties in describing feelings”; $\rho = -.391$, $p = .003$) and interoceptive awareness (subscale of Multidimensional assessment of interoceptive awareness “not worrying about pain”; $\rho = .405$, $p = .002$). No significant differences were found on questionnaires for depression and anxiety between patients with ALS and their caregivers ($p > .05$).

Conclusions ALS patients show reduced interoceptive sensitivity that is associated with poorer ability to describe feelings and with lower focalization on pain, regardless of cognitive and motor impairment. Alteration of interoception may represent a specific behavioural sign within the spectrum of emotion processing deficits described in ALS patients.

Keywords Interoception · Emotion · Behavioural · Cognition · HRV · Rehabilitation

Introduction

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease mainly affecting upper and lower motor neurons, with consequent motor impairment. In the last two decades,

non-motor symptoms, in the cognitive, behavioural, and emotional domains, have been identified as well [1–4].

Cognitive impairment is reported in approximately 30–50% of patients with ALS [5] and may negatively impact on survival [6, 7] and caregiver burden [8]. Among patients with cognitive impairment, a relatively small proportion (about 10%) presents frontotemporal dementia, mostly the behavioural variant [5]. Cognitive deficits have been shown in different domains. Most consistently, executive dysfunction and verbal fluency deficits are reported, but recently impairments in language and memory have been identified in patients with ALS [4, 9]. Moreover, facial emotion recognition, judgments of emotional valence, and social cognition skills such as decision-making and theory of mind (ToM) may also be impaired [10]. About 30% of patients show behavioural alterations, mainly characterized by apathy and irritability [11]. The prevalence of

✉ Raffaele Dubbioso
raf dubbioso@gmail.com

¹ Istituti Clinici Scientifici Maugeri IRCCS, Neurological Rehabilitation Unit of Telesse Terme Institute, 82037 Telesse Terme, Benevento, Italy

² Department of Neurosciences, Reproductive Sciences and Odontostomatology, University of Naples Federico II, Via Sergio Pansini, 5, 80131 Naples, Italy

³ Department of Psychology, University of Campania Luigi Vanvitelli, Naples, Italy

depression in ALS varies considerably ranging from 5 to 30%, whilst anxiety occurs in 20–35% of patients [12].

Structural and functional neuroimaging studies showed significant cortical thinning and reduced functional activation in the pre-central gyrus and supplementary motor area [13–15] as well as in extra-motor regions such as the prefrontal and temporal cortices [15, 16]. Interestingly, the involvement of limbic areas, including the insula, has been recently described in the neurodegenerative process of the ALS [10, 17, 18].

These areas, and the insula in particular, are thought to be key regions for interoception [19–21] that is the perception of sensations from inside the body, related to the function of internal organs [22]. Moreover, it has been proposed that subjective “feeling states” are dependent on the process of interoception: the representation and contextualisation of somatic and visceral responses elicited by emotional stimuli. In ALS, many components of the emotional processing are altered [23], and the insula has been suggested as a key anatomical region involved in several non-motor symptoms, including depression, anxiety, apathy, anhedonia, and fatigue [24]. However, besides interoceptive brain centres, viscerosensory afferent pathways, mainly from the vagus nerve, play an important role in modulating interoception [25, 26]

Accuracy in interoception is expressed as interoceptive sensitivity (IS). IS can be measured by using the heartbeat detection task, that is the ability to detect sensations from one’s heart.

Impairment of the heartbeat detection task, and therefore of IS, has been found in neurological disorders characterized by altered emotion processing, such as Parkinson disease [27], vascular focal brain lesions [28], and multiple sclerosis [29]. To the best of our knowledge, no study has systematically evaluated IS in patients with ALS and its correlation with cognitive and behavioural aspect of disease.

On these bases, the present study had three main aims: (1) to assess interoceptive sensitivity in patients affected by ALS, by comparing scores obtained on the heartbeat perception task in patients and in their caregivers; (2) to ascertain possible relations of IS with behavioural and cognitive measures in patients with ALS; (3) to explore whether IS difficulties are related with severity of clinical disability and/or the rate of disease progression. We enrolled patients’ caregivers as the control sample to rule out possible confounding factors related to interoception assessment, such as depression and anxiety [30], since similar levels of psychological distress have been described in both groups [31, 32].

Materials and methods

Participants

This study included patients with ALS consecutively admitted to the ALS centre of the University Hospital Federico II of

Naples in the period June 2020 to June 2021. Patients aged 18 years or more and met the “probable,” “probable laboratory-supported,” or “definite” diagnostic categories as per the revised El Escorial criteria for ALS [33]. Patients with history of neurologic disorders affecting cognition (major stroke, severe head injuries, mental retardation), alcohol dependence or drug dependence, severe mental illness, or use of high-dose psychoactive medications were not included in data analysis. In addition, patients with a diagnosis of frontotemporal dementia (FTD) [44], unable to communicate adequately, either verbally or by writing, were excluded.

Functional assessment of patients was performed by the Revised Amyotrophic Lateral Sclerosis Functional Rating Scale (ALSFRS-R). Disease onset (i.e., spinal vs bulbar onset) as well as disease phenotype, according to Chiò classification [34], were also recorded.

Moreover, we evaluated respiratory function, through spirometry performed with the patient sitting upright. Results for forced vital capacity (FVC) were expressed as a percentage of predicted value, from an average of three trials [35].

Genetic analysis was performed in all patients, exploring *C9orf72* repeat expansion and mutations of *SOD1*, *TARDBP*, and *FUS* genes. Routine magnetic resonance imaging (MRI) with a 3-T scanner was obtained for all patients.

All patients completed the cognitive, and psychological assessment specified below, the heartbeat counting task, and an assessment of interoceptive awareness.

We also recruited patients’ caregivers as control’s group. Only patient’s main informal caregivers were enrolled; professional caregivers were not included in the study. We excluded caregivers affected by any neurological, psychiatric, or other relevant clinical condition.

Neuropsychological assessment and cognitive classification

To assess cognitive and behavioural profile, neuropsychologists with specific expertise in ALS assessment (M.S., F.P.A.) administered a multi-domain battery to all participants. For assessing global cognitive functioning both ALS patients and caregivers underwent the Italian versions of the Mini-Mental State Examination (MMSE) [36] and of the Edinburgh Cognitive and Behavioural ALS Screen (ECAS) [37], a rapid screening test (15–20 min), including an ALS-specific section (assessing executive functions, social cognition, verbal fluency and language; 0–100 points), and a non-ALS-specific section (that assesses memory and visuospatial abilities; 0–36 points). ECAS total score ranges from 0 (worst performance) to 136 (best performance). Moreover, a brief caregiver interview provides an assessment of behaviour changes (Behavioural Disinhibition, Apathy/Inertia, Loss of Sympathy/Empathy, Perseverative/Stereotype,

Change in Eating Behaviour; from 0 to 10) and psychotic symptoms (from 0 to 3) usually associated with ALS [37].

In addition, patients underwent a battery of neuropsychological tests to assess specific cognitive domains: executive, attentive, memory, and visuospatial functions.

Executive functions were assessed by means of Wisconsin-Card-Sorting-Test (WCST) [38], the Stroop test [39], and the phonemic and semantic fluency tests [40]. Verbal short-term memory was evaluated by means of Digit Span forward test [41], and long-term verbal and visuo-spatial memory by means of: Rey Auditory Verbal Learning Test (RAVLT) [40] and Rey-Osterrieth Complex Figure Test Different Recall [42].

Non-verbal intelligence was assessed by means of the Raven's coloured progressive matrices [40], and visuospatial functions by means of Clock Drawing Test [26] and Rey-Osterrieth Complex Figure Test Copy [42].

Clinically relevant depression and anxiety symptoms were evaluated through the Hamilton Depression Scale (HDS) [43] and Beck Anxiety Inventory (BAI) [44]. The raw data of the neuropsychological tests were adjusted for age and years of education, according to Italian normative data [40, 45]. Adjusted scores were considered below the cut-off threshold (indicating deficit in cognitive performance) when they were below the fifth percentile from the Italian reference population's mean.

The cognitive status of the patients was classified according to diagnostic criteria published by Strong et al. [46] into the following categories: (i) ALS with normal cognition (ALS-nci); (ii) ALS with behavioural impairment (ALS-bi); (iii) ALS with cognitive impairment (ALS-ci); (iv) ALS with cognitive and behavioural impairment (ALS-cbi). As said above, patients that fulfilled the criteria for ALS with FTD (ALS-FTD) were not included in the study.

Assessment of interoceptive sensitivity

The IS was measured by the heartbeat perception task, a validated and reliable task (Cronbach's $\alpha = 0.69\text{--}0.90$) assessing interoceptive sensitivity [47, 48]. In the heartbeat perception task, the participants are required to count, in their mind or whispering, how many beats of their heart they perceive in a specific time frame (2×35 s, 2×25 s, and 2×45 s; trials were randomized across subjects), while being at rest in a comfortable sitting position. The number of beats provided by each participant was compared with the number of beats recorded by an ECG trace (Natus Dantec, Keypoint G4, Planegg, Germany) during the same time interval [47, 48]. Before starting the real task, participants performed a 12-s practice trial. The participants were not aware of the duration of the interval that was going to be presented, or of their accuracy; moreover, during task execution, they could not use strategies, such as taking the beat from their own wrist or chest. Accuracy of heartbeat perception was calculated

as the mean score of three heartbeat perception intervals according to the following formula: $1/6 \sum [(1 - ((\text{recorded heartbeats} - \text{counted heartbeats}) / \text{recorded heartbeats}))]$ [48]. Using this transformation, the IS score can vary between 0 and 1, with higher scores indicating smaller differences between recorded and perceived heartbeats (i.e., higher accuracy corresponds to higher interoceptive sensitivity).

Heart rate variability

To account for the possible contribution of autonomic afference to IS, participants' vagally mediated heart rate variability (HRV) was determined during the second part of the ECG recording session. Specifically, we asked the participants to lay still and to stay awake during a 300-s lasting time interval. The recordings were detrended (smooth priors: $\lambda = 500$), visually inspected and artefact corrected (adaptive filtering: cubic spline interpolation) before they were subjected to a time-domain analysis. The time-domain analysis was used for the determination of two vagally mediated HRV indices: the root mean square of successive differences between consecutive heartbeats (RMSSD) and the standard deviation of RR intervals (SDRR) [49].

Evaluation of interoception awareness, alexithymia, and apathy

To assess the interoceptive awareness, all participants underwent the Self Awareness Questionnaire (SAQ) [50], a self-report questionnaire evaluating awareness related to visceral sensations (F1) and to somatosensory sensations (F2). It includes 35 items rated on a 5-point Likert scale (0 = never; 1 = sometimes; 2 = often; 3 = very often; 4 = always); score ranges from 0 to 140, with higher score indicating greater subjective awareness of internal states of own body.

We also employed the Italian version of Multidimensional Assessment of Interoceptive Awareness (MAIA), to assess multiple dimensions of interoception [51, 52]. The MAIA is a self-report questionnaire including 32 items on a 6-points Likert scale, in which the participant must rate "how often each statement applies to you generally in daily life," with ordinal responses coded from 0 ("never") to 5 ("always"). This multidimensional tool embeds eight scales: (1) noticing, the awareness of one's body sensations; (2) not-distracting, the tendency not to ignore or distract oneself from sensations of pain or discomfort; (3) not-worrying, the tendency not to experience emotional distress or worry with sensations of pain or discomfort; (4) attention regulation, the ability to sustain and control attention to body sensation; (5) emotional awareness, the awareness of the connection between body sensations and emotional states; (6) self-regulation, the ability to regulate psychological distress by

attention to body sensations; (7) body listening, the tendency to actively listen to the body for insight; and (8) trusting, the experience of one's body as safe and trustworthy.

To assess the ability to identify and describe emotions, we used the Italian version of the Toronto Alexithymia Scale-20 items (TAS-20) [53], the most widely used self-report tool to assess the Alexithymia construct. The 20 items explore three factors reflecting the main aspects of the alexithymia: difficulty in identifying feelings; difficulty in describing feelings; externally oriented thinking. Each item must be rated on a 5-point Likert scale (from 1 = “completely agree” to 5 = “strongly disagree”). The total score ranges 20–100, with higher scores indicating higher levels of alexithymia. The Italian version of TAS-20 has been demonstrated to show good test–retest reliability (0.86) and adequate internal consistency (Cronbach's alpha: 0.75) in a wide sample of healthy adults and of medical and psychiatric outpatients (Bressi et al., 1996).

After completing the above tests and scales, all participants fulfilled the Italian version of the self-report version of Apathy Evaluation Scale (AES-S) [54, 55], a questionnaire including 18 items concerning behavioural (items 2, 6, 10, 11, 12), cognitive (items 1, 3, 4, 5, 7, 9, 13, 16), emotional (items 8, 14), and other (items 15, 17, 18) aspects of apathy. All items are scored on 4-point Likert scale (to mean “not at all true,” “slightly true”), “somewhat true” or “very true.” The total score ranges from 18 to 72 points, and higher scores indicate more severe apathy. Patients achieving an AES score ≤ 37 are classified as ALS-apathetic; patients with AES score > 37 as ALS-no-apathetic.

Statistical analysis

Continuous data were expressed as mean \pm standard deviation (SD). Categorical variables were summarized as relative frequencies. We compared the neuropsychological scores achieved by each participant with normative values, to assess prevalence and clinical relevance of cognitive impairment. We compared IS scores of patients and caregivers and IS scores of patients with apathy (ALS-apathetic) or without apathy (ALS-no-apathetic), by Mann–Whitney U test, with alpha level set at $p=0.05$. As a last step, we compared IS as a function of cognitive status defined based on Strong et al. (2017) criteria by means of Kruskal–Wallis test, with alpha level set at $p=0.05$.

Spearman's correlations were computed to examine the possible relationship between IS and clinical, psychological, and cognitive variables. The clinical variables were disease duration, disease severity (ALSFRS-R score), disease progression rate [(48—ALSFRS-R score at clinical examination)/disease duration (months)] [56], HRV measures (i.e., RMSSD and SDRR), functional independence (Activities of daily living—ADL, Instrumental activities of daily living—IADL scores); the psychological variables were depression (Hamilton), anxiety (BAI), apathy (AES), interoceptive awareness (MAIA), and alexithymia (TAS-20); cognitive variables

were ECAS scores, cognitive global functioning (MMSE). Spearman's correlations were conducted on the ALS patients' group only. Significance level was set at $p < 0.05$. All analyses were performed using Statistical Package for Social Sciences (SPSS) version 24 (IBM Corp., Armonk, NY, USA). Graphs were elaborated by means of GraphPad Prism version 8.4.2 for Windows (GraphPad Software, La Jolla California USA).

Results

Fifty-five ALS patients (36 males, mean age 60.3 ± 12.5 years) and 41 caregivers (CG) (17 males, mean age 56.5 ± 10.7) were included in the study. Genetic analysis was negative in all patients but two harbouring the hexanucleotide repeat expansion of the *C9orf72* gene. Routine MRI imaging did not disclose any significant abnormality in ALS patients.

Demographic and clinical data of the study populations are shown in Table 1. Table 2 and Table S1 summarize the results of the cognitive tests.

ALS patients obtained significantly lower scores on IS than CG (0.68 ± 0.24 vs 0.82 ± 0.16 , Mann–Whitney $U = 746.00$, $p = 0.003$) (Fig. 1). No statistically significant differences were found in self-reported measures of anxiety (BAI), depression (Hamilton), apathy (AES), Alexithymia (TAS-20), and interoceptive awareness (MAIA) between two groups (all $p > 0.05$; Table S1). No differences on IS scores were found between ALS-apathetic and ALS-no-apathetic groups.

Moreover, comparisons of IS scores performed by means of non-parametric Kruskal–Wallis test for multiple independent groups as function of Strong classification did not show significant differences among categories (see Table S2).

Lastly, Spearman's correlation in ALS group (Fig. 2) showed a significant negative association between IS and Alexithymia (score on the *difficulty in describing of feelings subscale* of TAS-20; $\rho = -0.391$, $p = 0.003$), a positive association with interoceptive awareness (score on *tendency of worrying about pain subscale* of MAIA; $\rho = 0.405$, $p = 0.002$). No correlations were found between IS and measures of disease severity (ALSFRS-R) and disease progression rate (ALSFRS-R rate). Moreover, no correlations were found between IS and scores of anxiety (BAI), depression (Hamilton), apathy (AES), cognitive tests, and of subscores of ECAS (all $p > 0.05$). Lastly, no significant correlation was observed between IS and HRV measures (all $p > 0.05$). Significant correlations are shown in Fig. 2.

Discussion

This study represents the first attempt to investigate the interoceptive sensitivity, measured by means of the heartbeat perception task, in patients with ALS. Our findings showed that

Table 1 Demographic and clinical characteristics of ALS patients

Sample size (<i>n</i>)	55
Age at onset	60.3 (12.5)
Gender (F/M)	19/36
Education (years)	10.6 (4.7)
Disease duration (in months)	20.9 (18.8)
Onset region (spinal/bulbar)	47/8
Disease phenotype	
Classic	20/55
Bulbar	6/55
Flail-arm	8/55
Flail-leg	9/55
Pyramidal	7/55
Pure lower motor neuron	4/55
Respiratory	1/55
ALSFRS-R	34.5 (6.8)
ALSFRS-R-rate of progression	0.89 (0.73)
FVC (%)	79.25 (25.1)
ADL	4.7 (1.6)
IADL	5.3 (1.9)

Continuous data are presented as mean (standard deviation) unless otherwise indicated

Abbreviations: ALSFRS-R Amyotrophic Lateral Sclerosis Functional Rating Scale-Revised, FVC forced vital capacity, ADL activities of daily life, IADL instrumental activities of daily life

patients were less accurate on IS task compared to caregivers; altered IS was not associated with disease severity or disease progression rate. Moreover, we found no correlation between IS and psychological self-report measure of anxiety and depression. Instead, we found some correlations with self-report measures of alexithymia and interoceptive awareness. Specifically, in our sample of patients with ALS, low IS was associated with a low ability in describing feelings and a higher tendency to not experience emotional distress or worry with sensations of pain or discomfort. This result is of particular interest because it could reflect a general reduction of interoception mediated by the insula.

This process represents the body-to-brain axis of sensation concerning the state of the internal body and the organs [20]. The central representation and perception of changes in bodily physiology are the basis for emotional feeling states [57]. In this process, the insula is thought to play a key role. Indeed, the connections and activation profile of the insula suggest that it integrates visceral and somatic input and forms a representation of the state of the body [22].

The deficit of IS found in patients with ALS might be an epiphenomenon of a degenerative process involving the insula and a specific behavioural symptom. Indeed, we did not find any correlation between the degree of deficit in IS and the severity of affective symptoms such as apathy, and other psychological symptoms. Similar findings have

Table 2 Comparisons of clinical of neuropsychological measures and questionnaires between patients with amyotrophic lateral sclerosis (ALS) and caregivers (CG)

	Total	ALS	CG	<i>P</i> -value
Sample size (<i>n</i>)	96	55	41	-
Females/males	42/54	19/36	23/18	-
Age (years)	58.9 (10.9)	60.1 (11.1)	57 (10.4)	0.117
Education (years)	11.4 (5.1)	10.4 (4.5)	12.7 (5.5)	0.064
ECAS Total	92.6 (21.8)	88.3 (24.2)	99.5 (15.1)	0.027
ECAS Total ALS	71.2 (15.8)	68.2 (17.5)	75.3 (12.3)	0.098
ECAS Total No ALS	21.6 (5.9)	20.3 (6.4)	23.8 (4.5)	0.003
ECAS language	22.2 (5.3)	21.7 (5.9)	23.1 (4.1)	0.547
ECAS fluency	16.7 (5.6)	16.3 (6.2)	17.4 (4.4)	0.658
ECAS executive functions	31.9 (9.1)	29.9 (9.7)	35.2 (6.8)	0.023
ECAS memory	11.1 (4.8)	9.8 (4.8)	12.9 (4.3)	0.001*
ECAS visuospatial	10.7 (1.8)	10.6 (1.9)	10.8 (1.4)	0.713
MMSE	28.1 (1.9)	27.8 (2.1)	28.5 (1.6)	0.113
ADL	5.1 (1.3)	4.7 (1.6)	5.6 (0.5)	0.026
IADL	5.9 (1.3)	5.3 (1.9)	6.7 (1.9)	0.001*
HDS	6.4 (4.8)	6.9 (4.7)	5.8 (5.1)	0.176
BAI	8.9 (7.8)	9.4 (8.1)	8.3 (7.5)	0.442
AES	28.9 (8.3)	30.2 (9.3)	27 (6.2)	0.114
TAS-20	45.7 (13.3)	46.2 (12.9)	44.9 (14.1)	0.470
MAIA	23.3 (6.2)	24.1 (5.9)	22.6 (6.5)	0.414
SAQ	17.3 (10.8)	17.6 (10.8)	16.8 (10.9)	0.708

Continuous data are presented as mean and standard deviation unless otherwise indicated

Abbreviations: ECAS Edinburgh Cognitive and Behavioural ALS Screen, MMSE Mini Mental Examination State, ADL activities of daily life, IADL instrumental activities of daily life, BAI Beck Anxiety Inventory, HDS Hamilton Depression Scale, AES Apathy Evaluation Scale, TAS-20 Toronto Alexithymia Scale, MAIA Multidimensional Assessment of Interoceptive Awareness, SAQ Self Awareness Questionnaire

*Significant differences according to Bonferroni correction ($p=0.002$; number of comparisons = 19)

been described in Parkinson disease, where poor IS on the heartbeat count task did not correlate with non-motor symptoms such as depression, anxiety, apathy, and cognitive functions [58]. Ricciardi et al. [58] suggested that the lack of correlation between self-reported measures of affective status and bodily awareness might reflect separate functional roles for the anterior and posterior insula, being the former involved in cognitive/affective functions and the latter in viscerosensory/somatosensory awareness [58, 59]. In this perspective, our results might be ascribed to a dissociation between an impairment of IS, as measured by the heartbeat perception task, and a relative sparing of interoceptive processing, more closely related to affective/emotional function, in ALS.




Fig. 1 Differences in Interoceptive Sensitivity (IS) during heartbeat count task between ALS patients and caregivers (CG)

However, our findings also showed the significant association of IS with some dimensions of alexithymia and interoceptive awareness explored by self-report questionnaires. Thus, from a different perspective, this result would demonstrate the significant functional interaction between the two components of interoception mediated by insula (cognitive/affective and viscerosensory/somatosensory) [20]. In addition, the lack of any correlation between HRV parameters and IS suggested that the impairment




Fig. 2 Correlations between interoceptive sensitivity (IS) during heartbeat count task with alexithymia (subscale of Toronto Alexithymia Scale, TAS-20, “difficulties in describing feelings”), left panel,


of interoception does not seem to be underpinned by an alteration of the afferent autonomic pathway but rather to a degeneration of the central brain networks.

Lastly, we demonstrated that IS is impaired in ALS, regardless of cognitive impairment, or general attentional disturbances, pointing out that behavioural interoceptive alteration is a specific signature of emotion processing deficit. This is in line with other emotion processing disturbances described in ALS, such as recognition of facial expressions of anger, sadness, and disgust that are impaired even when the cognition is preserved [60].

This study has some limitations. First, the relatively small sample size reduced the possibility to explore the association between ALS phenotypes and IS, especially when data were stratified in sub-categories according to the Strong criteria. Second, the lack of longitudinal observations did not allow to verify the reliability of the differences in IS measures between ALS and CG. Lastly, we did not assess variables potentially influencing IS such as personality traits and emotional status.

Conclusions

In conclusion, our data demonstrated a reduction of interoceptive sensitivity in patients with ALS. This deficit was related neither to presence and severity of cognitive and behavioural symptoms nor to motor disability. Our findings suggested that altered interoceptive accuracy may represent a specific behavioural sign within the spectrum of emotion processing deficits described in ALS. We speculated that impaired interoceptive processing might be linked to insular degeneration rather than to altered autonomic afferences, but this issue remains to be investigated by neuroimaging studies.



and interoceptive awareness (subscale of Multidimensional Assessment of Interoceptive Awareness, MAIA, “not worrying about pain”), right panel

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10072-022-06231-4>.

Author contribution All authors contributed to the study conception and design. Raffaele Dubbioso, Pasquale Moretta, and Myriam Spisto: conceptualization, methodology, and writing the final draft. Francesco Pio Ausiello, Rosa Iodice, and Natascia De Lucia: data curation, draft preparation. Gabriella Santangelo, Luigi Trojano, and Elena Salvatore: supervision.

Funding Open access funding provided by Università degli Studi di Napoli Federico II within the CRUI-CARE Agreement.

Declarations

Ethical approval and Informed consent Written consent was obtained from all patients and caregivers; the study was approved by the local Ethics Committee of University of Naples Federico II (N. 100/17).

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abrahams S, Leigh PN, Harvey A et al (2000) Verbal fluency and executive dysfunction in amyotrophic lateral sclerosis (ALS). *Neuropsychologia* 38:734–747. [https://doi.org/10.1016/S0028-3932\(99\)00146-3](https://doi.org/10.1016/S0028-3932(99)00146-3)
- Beeldman E, Raaphorst J, Twenenaar MK et al (2016) The cognitive profile of ALS: a systematic review and meta-analysis update. *J Neurol Neurosurg Psychiatry* 87:611–619. <https://doi.org/10.1136/jnnp-2015-310734>
- De Lucia N, Ausiello FP, Spisto M et al (2020) The emotional impact of COVID-19 outbreak in amyotrophic lateral sclerosis patients: evaluation of depression, anxiety and interoceptive awareness. *Neurol Sci* 41:2339–2341. <https://doi.org/10.1007/s10072-020-04592-2>
- Girardi A, MacPherson SE, Abrahams S (2011) Deficits in emotional and social cognition in amyotrophic lateral sclerosis. *Neuropsychology* 25:53–65. <https://doi.org/10.1037/a0020357>
- van Es MA, Hardiman O, Chio A et al (2017) Amyotrophic lateral sclerosis. *Lancet* 390:2084–2098
- Montuschi A, Iazzolino B, Calvo A et al (2015) Cognitive correlates in amyotrophic lateral sclerosis: a population-based study in Italy. *J Neurol Neurosurg Psychiatry* 86:168–173. <https://doi.org/10.1136/jnnp-2013-307223>
- Chiò A, Moglia C, Canosa A et al (2019) Cognitive impairment across ALS clinical stages in a population-based cohort. *Neurology* 93:E984–E994. <https://doi.org/10.1212/WNL.0000000000008063>
- Chiò A, Vignola A, Mastro E et al (2010) Neurobehavioral symptoms in ALS are negatively related to caregivers' burden and quality of life. *Eur J Neurol* 17:1298–1303. <https://doi.org/10.1111/j.1468-1331.2010.03016.x>
- Bora E (2017) Meta-analysis of social cognition in amyotrophic lateral sclerosis. *Cortex* 88:1–7. <https://doi.org/10.1016/j.cortex.2016.11.012>
- Lulé DE, Aho-Özhan HE, Vázquez C et al (2019) Story of the ALS-FTD continuum retold: Rather two distinct entities. *J Neurol Neurosurg Psychiatry* 90:586–589. <https://doi.org/10.1136/jnnp-2018-318800>
- Zucchi E, Ticozzi N, Mandrioli J (2019) Psychiatric symptoms in amyotrophic lateral sclerosis: beyond a motor neuron disorder. *Front Neurosci* 13:1–11. <https://doi.org/10.3389/fnins.2019.00175>
- Kawada T, Thakore NJ, Piro EP (2016) Depression in ALS in a large self-reporting cohort. *Neurology* 87:1631–1632. <https://doi.org/10.1212/01.wnl.0000503501.75283.f1>
- Mezzapesa DM, D'Errico E, Tortelli R et al (2013) Cortical thinning and clinical heterogeneity in amyotrophic lateral sclerosis. *PLoS ONE* 8:1–8. <https://doi.org/10.1371/journal.pone.0080748>
- Zhang J, Yin X, Zhao L et al (2014) Regional alterations in cortical thickness and white matter integrity in amyotrophic lateral sclerosis. *J Neurol* 261:412–421. <https://doi.org/10.1007/s00415-013-7215-5>
- Proudfoot M, Bede P, Turner MR (2019) Imaging cerebral activity in amyotrophic lateral sclerosis. *Front Neurol* 9:1148. <https://doi.org/10.3389/fneur.2018.01148>
- D'Ambrosio A, Gallo A, Trojsi F et al (2014) Frontotemporal cortical thinning in amyotrophic lateral sclerosis. *Am J Neuroradiol* 35:304–310. <https://doi.org/10.3174/ajnr.A3753>
- Benbrika S, Doidy F, Carlier L et al (2021) Longitudinal study of cognitive and emotional alterations in amyotrophic lateral sclerosis: clinical and imaging data. *Front Neurol* 12:1–11. <https://doi.org/10.3389/fneur.2021.620198>
- Ishaque A, Ta D, Khan M et al (2021) Distinct patterns of progressive gray and white matter degeneration in amyotrophic lateral sclerosis. *Human Brain Mapping* 43:1519–1534. <https://doi.org/10.1002/hbm.25738>
- Pollatos O, Schandry R, Auer DP, Kaufmann C (2007) Brain structures mediating cardiovascular arousal and interoceptive awareness. *Brain Res* 1141:178–187. <https://doi.org/10.1016/j.brainres.2007.01.026>
- Quadt L, Critchley HD, Garfinkel SN (2018) The neurobiology of interoception in health and disease. *Ann N Y Acad Sci* 1428:112–128. <https://doi.org/10.1111/nyas.13915>
- Wang X, Wu Q, Egan L et al (2019) Anterior insular cortex plays a critical role in interoceptive attention. *Elife* 15(8):e42265. <https://doi.org/10.1101/464867>
- Craig AD (2009) How do you feel—now? The anterior insula and human awareness. *Nat Rev Neurosci* 10:59–70
- Carelli L, Solca F, Tagini S et al (2021) Emotional processing and experience in amyotrophic lateral sclerosis: A systematic and critical review. *Brain Sci* 11:1–15. <https://doi.org/10.3390/brainsci11101356>
- Zhang T, Bai T, Xie W et al (2021) Abnormal connectivity of anterior-insular subdivisions and relationship with somatic symptom in depressive patients. *Brain Imaging Behav* 15:1760–1768. <https://doi.org/10.1007/s11682-020-00371-x>
- Lischke A, Pahnke R, Mau-Moeller A, Weippert M (2021) Heart rate variability modulates interoceptive accuracy. *Front Neurosci* 14:612445. <https://doi.org/10.3389/fnins.2020.612445>
- Quadt L, Critchley HD, Garfinkel SN (2018) The neurobiology of interoception in health and disease. *Ann N Y Acad Sci* 1428:112–128. <https://doi.org/10.1111/NYAS.13915>

27. Santangelo G, Vitale C, Baiano C et al (2018) Interoceptive processing deficit: a behavioral marker for subtyping Parkinson's disease. *Parkinsonism Relat Disord* 53:64–69. <https://doi.org/10.1016/j.parkrelidis.2018.05.001>
28. Raimo S, Boccia M, Di Vita A et al (2020) Interoceptive awareness in focal brain-damaged patients. *Neurol Sci* 41:1627–1631. <https://doi.org/10.1007/s10072-019-04172-z>
29. Gonzalez Campo C, Salamone PC, Rodríguez-Arriagada N et al (2020) Fatigue in multiple sclerosis is associated with multimodal interoceptive abnormalities. *Mult Scler J* 26:1845–1853. <https://doi.org/10.1177/1352458519888881>
30. Conroy É, Kennedy P, Heverin M et al (2021) Informal caregivers in amyotrophic lateral sclerosis: a multi-centre, exploratory study of burden and difficulties. *Brain Sci* 11:1–16. <https://doi.org/10.3390/brainsci11081094>
31. Chiò A, Gauthier A, Calvo A et al (2005) Caregiver burden and patients' perception of being a burden in ALS. *Neurology* 64:1780–1782. <https://doi.org/10.1212/01.WNL.0000162034.06268.37>
32. Vignola A, Guzzo A, Calvo A et al (2008) Anxiety undermines quality of life in ALS patients and caregivers. *Eur J Neurol* 15:1231–1236. <https://doi.org/10.1111/j.1468-1331.2008.02303.x>
33. Brooks BR, Miller RG, Swash M, Munsat TL (2000) El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 1:293–299. <https://doi.org/10.1080/146608200300079536>
34. Chiò A, Calvo A, Moglia C et al (2011) Phenotypic heterogeneity of amyotrophic lateral sclerosis: a population based study. *J Neurol Neurosurg Psychiatry* 82:740–746. <https://doi.org/10.1136/jnnp.2010.235952>
35. Czaplinski A, Yen AA, Appel SH (2006) Forced vital capacity (FVC) as an indicator of survival and disease progression in an ALS clinic population. *J Neurol Neurosurg Psychiatry* 77:390–392. <https://doi.org/10.1136/jnnp.2005.072660>
36. Folstein M, Folstein S, McHugh P (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 12:189–198. [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6)
37. Poletti B, Solca F, Carelli L et al (2016) The validation of the Italian Edinburgh Cognitive and Behavioural ALS Screen (ECAS). *Amyotroph Lateral Scler Front Degener* 17:489–498. <https://doi.org/10.1080/21678421.2016.1183679>
38. Laiacona M, Inzaghi MG, De Tanti A, Capitani E (2000) Wisconsin card sorting test: a new global score, with Italian norms, and its relationship with the Weigl sorting test. *Neurol Sci* 21:279–291. <https://doi.org/10.1007/s100720070065>
39. Barbarotto R, Laiacona M, Frosio R et al (1998) A normative study on visual reaction times and two Stroop colour-word tests. *Neurol Sci* 19:161–170. <https://doi.org/10.1007/bf00831566>
40. Carlesimo C, Caltagirone C, Gainotti G (1996) The Mental Deterioration Battery: normative data, diagnostic reliability and qualitative analyses of cognitive impairment. The Group for the Standardization of the Mental Deterioration Battery. *Eur Neurol* 36:378–384. <https://doi.org/10.1159/000117297>
41. Orsini A, Grossi D, Capitani E et al (1987) Verbal and spatial immediate memory span: normative data from 1355 adults and 1112 children. *Italian J Neurol Sci* 8:537–548. <https://doi.org/10.1007/BF02333660>
42. Caffarra P, Vezzadini G, Dieci F et al (2002) Rey-Osterrieth complex figure: Normative values in an Italian population sample. *Neurol Sci* 22:443–447. <https://doi.org/10.1007/s100720200003>
43. Hamilton M (1960) A rating scale for depression. *J Neurol Neurosurg Psychiatry* 23:56–62. <https://doi.org/10.1136/jnnp.23.1.56>
44. Beck AT, Epstein N, Brown G, Steer RA (1988) An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 56:893–897. <https://doi.org/10.1037/0022-006X.56.6.893>
45. Spinnler H, Tognoni G (1987) Standardizzazione e taratura italiana di test neuropsicologici. *Italian J Neurol Sci Suppl* 8:1–120
46. Strong MJ, Abrahams S, Goldstein LH et al (2017) Amyotrophic lateral sclerosis - frontotemporal spectrum disorder (ALS-FTSD): revised diagnostic criteria. *Amyotroph Lateral Scler Front Degener* 18:153–174. <https://doi.org/10.1080/21678421.2016.1267768>
47. Dunn BD, Stefanovitch I, Buchan K et al (2009) A reduction in positive self-judgment bias is uniquely related to the anhedonic symptoms of depression. *Behav Res Ther* 47:374–381. <https://doi.org/10.1016/j.brat.2009.01.016>
48. Pollatos O, Traut-Mattausch E, Schandry R (2009) Differential effects of anxiety and depression on interoceptive accuracy. *Depress Anxiety* 26:167–173. <https://doi.org/10.1002/da.20504>
49. Shaffer F, Ginsberg JP (2017) An overview of heart rate variability metrics and norms. *Front Public Health* 5:258. <https://doi.org/10.3389/fpubh.2017.00258>
50. Longarzo M, D'Olimpio F, Chiavazzo A et al (2015) The relationships between interoception and alexithymic trait. The Self-Awareness Questionnaire in healthy subjects. *Front Psych* 6:1149. <https://doi.org/10.3389/fpsyg.2015.01149>
51. Mehling WE, Price C, Daubenmier JJ et al (2012) The Multidimensional Assessment of Interoceptive Awareness (MAIA). *PLoS ONE* 7:e48230. <https://doi.org/10.1371/journal.pone.0048230>
52. Cali G, Ambrosini E, Picconi L et al (2015) Investigating the relationship between interoceptive accuracy, interoceptive awareness, and emotional susceptibility. *Front Psychol* 6:1–13. <https://doi.org/10.3389/fpsyg.2015.01202>
53. Bressi C, Taylor G, Parker J et al (1996) Cross validation of the factor structure of the 20-item Toronto Alexithymia Scale: An Italian multicenter study. *J Psychosom Res* 41:551–559. [https://doi.org/10.1016/S0022-3999\(96\)00228-0](https://doi.org/10.1016/S0022-3999(96)00228-0)
54. Raimo S, Trojano L, Spitaleri D et al (2014) Apathy in multiple sclerosis: a validation study of the apathy evaluation scale. *J Neurol Sci* 347:295–300. <https://doi.org/10.1016/j.jns.2014.10.027>
55. Santangelo G, Barone P, Cuoco S et al (2014) Apathy in untreated, de novo patients with Parkinson's disease: validation study of Apathy Evaluation Scale. *J Neurol* 261:2319–2328. <https://doi.org/10.1007/s00415-014-7498-1>
56. Kimura F, Fujimura C, Ishida S et al (2006) Progression rate of ALS-FRS-R at time of diagnosis predicts survival time in ALS. *Neurology* 66:265–267. <https://doi.org/10.1212/01.wnl.0000194316.91908.8a>
57. Lange CG, James W (1967) The emotions. Hafner
58. Ricciardi L, Ferrazzano G, Demartini B et al (2016) Know thyself: exploring interoceptive sensitivity in Parkinson's disease. *J Neurol Sci* 364:110–115. <https://doi.org/10.1016/j.jns.2016.03.019>
59. Chang LJ, Yarkoni T, Khaw MW, Sanfey AG (2013) Decoding the role of the insula in human cognition: Functional parcellation and large-scale reverse inference. *Cereb Cortex* 23:739–749. <https://doi.org/10.1093/cercor/bhs065>
60. Sedda A (2014) Disorders of emotional processing in amyotrophic lateral sclerosis. *Curr Opin Neurol* 27:659–665

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.