

# Negative selection, not receptor editing, is a physiological response of autoreactive thymocytes

Taras Kreslavsky,<sup>1</sup> Hye-Jung Kim,<sup>1</sup> Sergei B. Koralov,<sup>2</sup> Dvora Ghitza,<sup>2</sup> Thorsten Buch,<sup>3</sup> Harvey Cantor,<sup>1</sup> Klaus Rajewsky,<sup>2</sup> and Harald von Boehmer<sup>1</sup>

<sup>1</sup>Department of Cancer Immunology and AIDS, Dana-Farber Cancer Institute; and <sup>2</sup>Program in Cellular and Molecular Medicine, Children's Hospital, and Immune Disease Institute, Harvard Medical School, Boston, MA 02115

<sup>3</sup>Institute for Medical Microbiology, Immunology, and Hygiene, Technische Universität München, 81545 Munich, Germany

**Antigen receptor editing—a process of secondary rearrangements of antigen receptor genes in autoreactive lymphocytes—is a well-established tolerance mechanism in B cells, whereas its role in T cells remains controversial. Here, we investigated this issue using a novel *Tcra* knock-in locus, which ensured appropriate timing of TCR $\alpha$  expression and allowed secondary rearrangements. Under these conditions the only response to self-antigen that could be unambiguously identified was negative selection of CD4/CD8 double positive thymocytes. No evidence could be obtained for antigen-induced TCR editing, whereas replacement of the transgenic TCR $\alpha$  chain by ongoing gene rearrangement occurred in some cells irrespective of the presence or absence of self-antigen.**

## CORRESPONDENCE

Harald von Boehmer:  
harald\_von\_boehmer@  
dfci.harvard.edu

Abbreviations used: CD8SP, CD8 single positive; DN, double negative; DP, double positive; ES, embryonic stem.

The diversity of antigen receptors in the adaptive immune system of vertebrates is generated by a random recombinatorial process, and thus receptors recognizing self-antigens are constantly generated, creating a risk of autoimmunity. Therefore, the immune system requires an array of mechanisms to disarm autoreactive lymphocytes. This includes negative selection, induction of anergy, diversion to lineages with regulatory properties, and antigen receptor editing.

Receptor editing is a process of secondary rearrangements of antigen receptor genes in response to recognition of self-antigen. This phenomenon is well described for B cells (Nemazee and Hogquist, 2003); however, its role in T cell tolerance remains controversial (Mostoslavsky and Alt, 2004), in part because of the lack of an appropriate mouse model. Indirect experiments using mice expressing TCR transgenes that were not contained within endogenous TCR loci suggested that editing may exist because

T cells with receptors specific for peptide ligands expressed in the thymic cortex appeared to undergo rearrangements of the endogenous *Tcra* locus (McGargill et al., 2000). In this setting, endogenous TCR $\alpha$  chains have to compete with the transgenic TCR $\alpha$  for pairing with TCR $\beta$  chains. The varying efficiency of such competition for different TCR transgenes may explain the fact that OT-I, but not HY or 2C TCRs, allowed for the appearance of transgene-negative, TCR-positive cells in the presence of antigen (Mayerova and Hogquist, 2004). Other studies have suggested that editing may even take place after thymic egress and peripheral antigens could induce TCR $\beta$  rearrangements in peripheral T cells specific for these antigens (McMahan and Fink, 1998). More direct experiments in which a rearranged V $\alpha$ J $\alpha$  exon of the HY TCR $\alpha$  chain was knocked-in into the *Tcra* locus in a way that mimics the product of physiological V $\alpha$ J $\alpha$  rearrangement (HY-I mouse), and thus could be deleted by secondary rearrangements, indicated, however, an absence of TCR editing

T. Kreslavsky's present address is the Research Institute of Molecular Pathology, Vienna, Austria.

S.B. Koralov's present address is the Dept. of Pathology, NYU Langone Medical Center, New York, NY.

K. Rajewsky's present address is the Max-Delbrück-Center for Molecular Medicine, Berlin, Germany.

© 2013 Kreslavsky et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

(Buch et al., 2002). Secondary rearrangements did occur to some extent in this system in the absence of the self-antigen, but were not increased in its presence. These secondary rearrangements, also observed in another *Tcra* knock-in system (Wang et al., 1998), are thought to be a mechanism that enhances the generation of TCRs suitable for positive selection in WT mice. Unexpectedly, however, the knock-in TCR $\alpha$  allele constructed by Buch et al. (2002) was expressed prematurely, in CD4/CD8 double-negative 3 (DN3) thymocytes (Croxford et al., 2008), whereas under physiological conditions *Tcra* is rearranged only at the CD4/CD8 double positive (DP) stage. Thus expression of the TCR in this system could occur too early, at a point in time when editing may not yet be possible. Indeed, rearrangement of antigen receptors takes place only in nonproliferating populations, as Rag2 is unstable in dividing cells (Li et al., 1996; Lee and Desiderio, 1999), whereas expression of the TCR $\beta$  chain in DN3 cells initiates a burst of proliferation that is likely to interfere with V(D)J recombination. Although, in B cells, premature expression of an autoreactive BCR does not interfere with editing, which still occurs after a wave of proliferation (Pelanda et al., 1997), similar rules do not have to apply to T cells. Moreover, premature TCR $\alpha\beta$  expression is known to divert cells to  $\gamma\delta$ -like lineage (Bruno et al., 1996; Terrence et al., 2000; Baldwin et al., 2005; Egawa et al., 2008) and it is unclear whether that is compatible with editing. Some in vitro experiments were performed to address this issue, and appeared to contradict this possibility (Buch et al., 2002); however, this system did not allow tackling the issue properly in the in vivo setting. Thus, a model with correct timing of TCR expression is required to address the impact of TCR $\alpha$  editing under physiological conditions.

A successful approach to correct the timing of TCR $\alpha$  transgene expression was developed by Baldwin et al. (2005). The authors placed a floxed STOP cassette between the promoter and the HY TCR $\alpha$  transgene, thus preventing premature TCR $\alpha$  expression. The STOP cassette was then excised by CD4-Cre that is expressed from the DP stage, ensuring proper timing of TCR $\alpha$  expression (HY<sup>cd4</sup> mouse). However, the transgene used could not be deleted by secondary rearrangements, and thus *Tcra* editing could not be addressed in this system. We adopted this approach to correct the timing of TCR $\alpha$  expression in the HY-I system. In the resulting HY-STOP mice, the onset of HY TCR $\alpha$  expression took place physiologically, at the DP stage. Correction of the timing of TCR expression rescued the abnormalities in the gross thymic phenotype observed in HY-I mice. However, no evidence for TCR $\alpha$  editing was found in this system, and autoreactive cells instead were eliminated by negative selection.

## RESULTS AND DISCUSSION

### Generation of HY-STOP mice


We aimed to design a *Tcra* locus whose expression was only initiated at the physiologically appropriate nonproliferative DP stage, but that would allow for secondary rearrangements potentially induced by self-antigen encounter. To this end, we

chose to modify the previously reported HY $\alpha^{1A}$  *Tcra* allele of HY-I mouse (Buch et al., 2002). This allele was designed to mimic the product of V $\alpha$ J $\alpha$  recombination: the rearranged V $\alpha$ J $\alpha$  segment of the HY TCR $\alpha$  chain was inserted into the TCR $\alpha$  locus followed by deletion of the TCR $\delta$  locus (Buch et al., 2002). The H2D<sup>b</sup>-restricted HY TCR recognizes Y-chromosome-encoded antigen, and thus is self-reactive in males but not females. However, the HY $\alpha^{1A}$  locus suffered from premature expression at the DN stages of T cell development (Buch et al., 2002; Fig. 1 B). To correct this abnormality, we generated and retargeted embryonic stem (ES) cells from HY-I mice to introduce a floxed STOP cassette with two additional polyadenylation signal sequences (Lakso et al., 1992; Buch et al., 2005) between rearranged HY V $\alpha$ J $\alpha$  and the first downstream J segment (Fig. 1 A). As polyadenylation signals in the STOP cassette lead to termination of transcription, the truncated mRNA should include a rearranged V $\alpha$ J $\alpha$  exon, but not a constant region of the TCR $\alpha$ , and thus the TCR $\alpha$  protein should not be produced unless the STOP cassette is removed. As the HY $\alpha^{1A}$  allele contains a “leftover” loxP site upstream of HY V $\alpha$ J $\alpha$  (Buch et al., 2002), we used loxP2272 sites that can recombine with each other but not with conventional loxP sites (Siegel et al., 2001; Luche et al., 2007) to flank the STOP cassette. Generation of the ES cells, targeting, ES cell screening and blastocyst injections, were performed as described in the Material and methods.

The resulting *Tcra*<sup>HY-STOP</sup> chimeras were bred to *Tcra*<sup>-/-</sup> females containing the TCR HY $\beta$  (Uematsu et al., 1988) and Roryt-Cre (Eberl and Littman, 2004) transgenes. *Tcra*<sup>HY-STOP/+</sup> HY $\beta$  Roryt-Cre animals, henceforth referred to as HY-STOP mice, were used in all further analyses. The TCR HY $\beta$  transgene was required to produce a male-autoreactive HY TCR once HY $\alpha$  is expressed. Expression of Roryt-Cre is initiated at the DP stage (Eberl and Littman, 2004)—the time in T cell development when rearrangement and expression of TCR $\alpha$  take place physiologically, and thus the STOP cassette should be removed from the *Tcra*<sup>HY-STOP</sup> locus. In *Tcra*<sup>HY-STOP/+</sup> mice, the only functional copy of the *Tcra* locus is the one containing prerearranged HY $\alpha$  and the STOP cassette, and thus expression of any TCR $\alpha$  other than HY $\alpha$  is indicative of secondary rearrangements in the locus.

### Thymic phenotype of HY-STOP mice

We first compared the timing of TCR $\alpha\beta$  expression in HY-I and HY-STOP female mice. Unlike HY-I animals, where the majority of DN thymocytes were TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>+</sup>, virtually no such cells could be detected in HY-STOP animals. DP thymocytes, however, did initiate the expression of HY $\alpha$  (Fig. 1 B). Importantly, only a fraction of DP cells were positive for HY $\alpha$ , mimicking the physiological situation in WT mice where TCR $\alpha$  rearrangement takes place during transition through the DP stage and only a fraction of DP thymocytes expresses the  $\alpha\beta$ TCR. Thus, as expected, insertion of the STOP cassette interfered with premature expression of TCR $\alpha$ , and its removal by Roryt-Cre ensured a physiological onset of TCR $\alpha$  expression at the DP stage.



**Figure 1. Corrected timing of TCR $\alpha$  expression and normal gross thymic phenotype of HY-STOP mice.** (A) Targeting strategy for generation of the *TcrA*<sup>HY-STOP</sup> locus. Original HY-I knock-in locus (top), targeting vector (center), and targeted locus with a STOP cassette are shown. HY V $\alpha$ J $\alpha$ , recombined V $\alpha$ J $\alpha$  element of HY TCR; Neo, neomycin resistance gene; frt, frt sites; pA, polyadenylation signal; STOP, transcriptional STOP cassette; loxP, leftover loxP site in HY-I locus; loxP2272, alternative loxP sites; TK, thymidine kinase gene; Amp, ampicillin resistance gene. (B) Expression of surface TCR $\beta$  and HY $\alpha$  on CD4<sup>-</sup>CD8<sup>-</sup> (DN), CD4<sup>+</sup>CD8<sup>+</sup> (DP), and CD4<sup>-</sup>CD8<sup>+</sup> (CD8SP) thymocytes from the HY-I (*TcrA*<sup>HY-I/HY-I</sup> HY $\beta$ ) and HY-STOP (*TcrA*<sup>HY-STOP/HY-I</sup> HY $\beta$  Ror $\gamma$ t-Cre) female mice. Representative results of at least three independent experiments are shown. (C) Total thymocytes numbers and numbers of TCR<sup>-</sup> DP thymocytes in male and female HY-I and HY-STOP mice. \*, P < 0.5; \*\*, P < 0.01 (n = 3-5); student's t test. Error bars represent the SD. Representative results of two independent experiments are shown. (D) Comparison of CD4/CD8 profiles of HY-I and HY-STOP thymocytes from male and female mice. Representative results of at least three independent experiments are shown.

As reported previously (Groettrup and von Boehmer, 1993), we could detect some level of surface expression of transgenic TCR $\beta$  on DP cells even in the absence of TCR $\alpha$  (e.g., on a *Tcr $\alpha$ <sup>-/-</sup>* background; Fig. S1 A). Thus, in experiments that required strict gating on TCR $\alpha\beta$ -negative cells, the gate was applied as shown in Fig. S1A.

Unlike the original HY-I mice, which had a strong decrease in the DP compartment and a relative increase in DN cells even in female mice because of premature TCR expression, both female and male HY-STOP mice had a normal frequency of DP and DN thymocytes (Fig. 1 D). Thymic cellularity, which was severely reduced in male HY-I mice due to premature deletion of DP cells, was consistently increased in male HY-STOP mice. However, male HY-STOP mice still had an approximately twofold decrease in total thymocyte numbers when compared with HY-STOP females (Fig. 1 C). This yet to be explained phenomenon is not caused by premature deletion of H-Y-specific cells, as TCR<sup>-</sup> DP cells are also affected (Fig. 1 C), and thus seems to represent a bystander effect. We conclude that corrected timing of TCR expression largely rescued the severe abnormalities found in the gross thymic phenotype of HY-I mice. This conclusion concurs with earlier studies in the HY<sup>cd4</sup> model (Baldwin et al., 2005).

We next tested whether the HY-STOP locus is functional in terms of V $\alpha$ J $\alpha$  recombination. It was reported for HY-I mice that even in the absence of self-antigen, their TCR $\alpha$  locus underwent some level of V $\alpha$ -J $\alpha$  rearrangement resulting in deletion of the HY $\alpha$  V $\alpha$ J $\alpha$  in a fraction of thymocytes, a process that in normal mice is believed to enhance the formation of receptors suitable for positive selection. Consistently, in HY-STOP female mice, some thymocytes were HY TCR<sup>-</sup> but expressed high levels of surface TCR $\beta$ . These cells expressed V $\alpha$ J $\alpha$  joins other than that of the HY TCR, as staining was detectable with V $\alpha$ 2 and V $\alpha$ 3.2 antibodies (Fig. 2 A). These non-HY $\alpha$  chains can only be a product of rearrangements in the *Tcr $\alpha$ <sup>HYSTOP</sup>* allele, as *Tcr $\alpha$ <sup>HYSTOP/-</sup>* animals were used. Therefore, insertion of the STOP cassette did not disrupt the *Tcr $\alpha$*  locus architecture and did not interfere with its ability to undergo secondary rearrangements. However, these rearrangements were not induced by antigen, as they occurred in female mice. These TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup> cells differentiated toward both CD4 and CD8 lineages and, as expected, were uniformly HY $\beta$ <sup>+</sup> as judged by V $\beta$ 8.1/8.2 antibody staining. This demonstrates that, as reported previously (Uematsu et al., 1988), the transgenic TCR $\beta$  chain was able to support positive selection in the context of TCR $\alpha$  chains other than HY $\alpha$  (Fig. 2 B).


In summary, HY-STOP mice had a grossly normal thymic phenotype, their TCR $\alpha$  expression timing was physiological, and their *Tcr $\alpha$*  locus was capable of secondary rearrangements. These properties, not found together in any other model system, make HY-STOP mice suitable to definitively address the question of TCR editing.

#### Lack of evidence for TCR editing in HY-STOP animals

We next sought to find evidence for or against TCR editing in this system. To this end, we compared the frequencies

and numbers of clonotype-positive and negative TCR $\alpha\beta$ -expressing cells in male and female HY-STOP mice. BM chimeras with sex-matched donors and recipients were used in these experiments to increase the numbers of animals analyzed in parallel.

In female HY-STOP mice, the majority of TCR<sup>+</sup> cells express the clonotypic HY TCR, whereas a minority undergoes secondary rearrangement and are TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup> (Fig. 1 B). Two scenarios are conceivable in male mice. If self-reactive thymocytes would undergo TCR-editing, cells that were HY $\alpha$ -positive in females would convert to an HY $\alpha$ -negative state in males. Alternatively, if negative selection is the main tolerogenic mechanism, these cells would be eliminated by apoptosis. Both scenarios would predict a proportional decrease in HY $\alpha$ <sup>+</sup> cells among TCR<sup>+</sup> cells (Fig. 3 A). However, in the case of TCR editing, this would be accompanied by a corresponding increase in absolute numbers of TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup> thymocytes (converted from autoreactive HY $\alpha$ <sup>+</sup> cells), whereas if the predominant mechanism of tolerance is negative selection of TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>+</sup> cells, no increase in TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup>




**Figure 2. HY-STOP *Tcr $\alpha$*  locus is functional and can undergo antigen-independent secondary rearrangements.** (A) Female HY-STOP thymocytes were stained for surface TCR $\beta$ , HY $\alpha$ , and V $\alpha$ 2 or V $\alpha$ 3.2. Note the presence of TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup> cells (left), a fraction of which can be stained with V $\alpha$  antibodies (center), and the lack of binding of V $\alpha$  antibodies to TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>+</sup> cells (right). Representative results of two independent experiments are shown. (B) Female HY-STOP thymocytes were stained for surface TCR $\beta$ , HY $\alpha$ , TCR $\beta$ 8.1/8.2, CD4, and CD8 $\alpha$ . Staining for TCR $\beta$  and TCR $\beta$ 8.1/8.2 on HY $\alpha$ <sup>-</sup> cells (left) and expression of CD4 and CD8 $\alpha$  by TCR $\beta$ <sup>+</sup>HY $\alpha$ <sup>-</sup> cells (right) is shown. Representative staining for one of three analyzed animals is shown.

thymocytes would be expected (Fig. 3 A). Consistent with both scenarios, a strong decrease in  $\text{HY}\alpha^+$  thymocytes was observed in male HY-STOP mice as compared to females (Fig. 3, B and C). However, this decrease was not accompanied by any measurable increase in absolute numbers of  $\text{TCR}\beta^+\text{HY}\alpha^-$  DP thymocytes. Indeed, male mice had a decrease in the numbers of  $\text{TCR}\beta^+\text{HY}\alpha^-$  cells (Fig. 3 C). As male HY-STOP mice have a twofold reduced overall thymic cellularity (that also affects  $\text{TCR}^-$  DP precursors), the frequency of  $\text{TCR}\beta^+\text{HY}\alpha^-$  cells among total thymocytes may be a fairer measure, as it takes in account this reduction. This approach again failed to reveal a significant increase in  $\text{TCR}\beta^+\text{HY}\alpha^-$  cells in male mice (Fig. 3 C). Thus, autoreactive cells disappeared from male mice without a sizable contribution to the pool of nonautoreactive T cells. Expression of recombination activating genes is rapidly turned off in the DP thymocytes upon TCR signaling. TCR editing would require prolongation or reinduction of *Rag* expression in autoreactive thymocytes. We thus measured *Rag1* and *Rag2* expression in  $\text{TCR}\beta^-$  and  $\text{TCR}\beta^+\text{HY}\alpha^+$  DP cells

from male and female HY-STOP mice by qPCR (Fig. 3 D). As expected, expression of HY TCR led to a 5–10 fold decrease in *Rag* gene expression in female DP thymocytes. However, even a more prominent 55–170 fold decrease was observed in male  $\text{HY}\alpha^+$  DP cells, suggesting that the stronger TCR signal from an autoreactive TCR was repressing *Rag* expression more efficiently rather than prolonging it. We conclude that no evidence for antigen-induced TCR editing can be found in the HY-STOP system.


### Autoreactive T cells in HY-STOP mice undergo negative selection

We next compared the phenotype of clonotype-positive cells in male and female HY-STOP mice. As expected, no  $\text{CD4}^+\text{CD8}^-$  cells were found among  $\text{HY}\alpha^+$  cells (Fig. 4 A), once again confirming appropriate timing of  $\text{TCR}\alpha$  expression.  $\text{CD8SP}$   $\text{HY}\alpha^+$  cells were detected in female but not male mice, and  $\text{HY}\alpha^+$  DP cells were present both in males and females. However, the levels of co-receptor expression were different



**Figure 3. Presence of self-antigen does not enhance secondary rearrangements.** (A) Schematic representation of the predicted effects of deletion and TCR-editing scenarios on numbers of clonotype-positive and -negative thymocytes. (B) Expression of surface  $\text{TCR}\beta$  and  $\text{HY}\alpha$  on DP and CD8SP thymocytes from HY-STOP male and female BM chimeras. Representative results of at least five independent experiments are shown. (C) Absolute numbers (left) and percentage from total thymocytes (right) of  $\text{TCR}\beta^+\text{HY}\alpha^+$  and  $\text{TCR}\beta^+\text{HY}\alpha^-$  DP cells in male and female HY-STOP BM chimeras. \*\*,  $P < 0.01$ ; NS, not significant ( $P > 0.05$ ); student's *t* test;  $n = 5$ . Error bars represent the SD. Representative results of two independent experiments are shown. (D) Expression of *Rag1* and *Rag2* was analyzed by TaqMan qPCR in sorted  $\text{TCR}\beta^-$  and  $\text{TCR}\beta^+\text{HY}\alpha^+$  DP thymocytes from HY-STOP male and female BM chimeras. Mean relative expression normalized against *Gapdh* expression is shown. Error bars represent SD (individual mice;  $n = 4$  for each group from two independent experiments). \*\*\*,  $P < 0.001$ ; \*\*,  $P < 0.01$ ; student's *t* test; NS, not significant ( $P > 0.05$ ).

between males and females. Whereas all DP cells in females expressed high levels of CD4 and CD8, a large fraction of male  $\text{HY}\alpha^+$  DP cells had a  $\text{CD4}^{\text{low}}\text{CD8}^{\text{low}}$  phenotype (Fig. 4 A). This  $\text{DP}^{\text{dull}}$  phenotype was previously reported to be associated with negative selection (Page et al., 1993). Likewise, male but not female cells expressed high levels of the co-inhibitory receptor PD-1 and the transcription factor Helios (both known to be up-regulated in cells undergoing negative selection (Baldwin and Hogquist, 2007; Daley et al., 2013; Mingueau



**Figure 4.  $\text{HY}\alpha^+$  cells in male  $\text{HY-STOP}$  mice are negatively selected.** (A) Expression of CD4 and CD8 co-receptors by male and female  $\text{HY}\alpha^+$  thymocytes. Representative results of at least five independent experiments are shown. Note co-receptor dulling and lack of  $\text{CD4}^{\text{low}}\text{CD8}^{\text{low}}$  cells among male clonotype-positive cells. (B) Expression of negative selection-associated markers PD-1 and Helios by total  $\text{HY}\alpha^+$  DP thymocytes from male and female  $\text{HY-STOP}$  mice and by  $\text{DP}^{\text{dull}}$  cells from male mice. Representative results of three independent experiments are shown. (C) Detection of cleaved caspase-3 in male and female  $\text{HY}\alpha^+$  thymocytes from  $\text{HY-I}$  (top) and  $\text{HY-STOP}$  (bottom) mice. Representative results of two independent experiments are shown.

et al., 2013), with the male  $\text{DP}^{\text{dull}}$  population expressing the highest levels of both markers (Fig. 4 B). To assess directly whether clonotype-positive cells in males undergo apoptosis, we performed staining for the active form of the “executioner” caspase-3 (act-Casp3). Male, but not female,  $\text{HY}\alpha^+$  DP cells had a detectable population of act-Casp3<sup>+</sup> cells (Fig. 4 C) at a frequency close to that reported previously for the  $\text{HY}^{\text{cd4}}$  model, where TCR editing is not possible (McCaughy et al., 2008). We conclude that, under conditions permissive for TCR editing, encounter of self-antigen in the thymus results in apoptosis rather than secondary rearrangement of the *Tcr* locus.

Interestingly, in male  $\text{HY-I}$  mice (i.e., the experimental system previously used by Buch et al. [2002]) the frequency of act-Casp3<sup>+</sup>  $\text{HY}\alpha^+$  thymocytes was approximately fivefold lower than in  $\text{HY-STOP}$  males (Fig. 4 C). Thus, premature TCR expression partially interferes with negative selection—possibly due to rescue of some of the autoreactive thymocytes by their diversion to  $\gamma\delta$ -like lineage. This result emphasizes the importance of models with physiological timing of TCR expression for studies of T cell tolerance.

In spite of the virtual nonexistence of  $\text{HY}\alpha^+$  CD8SP thymocytes in male mice, few such cells escaped to the periphery and acquired a  $\text{CD8}\alpha\beta^+\text{CD122}^+\text{CD44}^+$  phenotype. These cells expressed lower levels of TCR and CD8 than clonotype-positive cells in females, and many of them up-regulated Ly49, a phenotype previously reported for CD8 T cells with regulatory properties (Kim et al., 2010, 2011; not depicted).

Although antigen receptor editing is a well-established tolerance mechanism in B cells, studies in T cells provided conflicting results. In the two systems where TCR editing was suggested to occur, it would have to take place under non-physiological circumstances. In the OT-I system, endogenous TCR $\alpha$  chains had to compete with the transgenic TCR $\alpha$  expressed by all thymocytes (McGargill et al., 2000). Another study used a knock-in into the *Tcr* locus that contained a PGK promoter-driven neomycin resistance gene and an Ig heavy chain enhancer in addition to the rearranged  $\text{V}\alpha\text{J}\alpha$  (Wang et al., 1998). These elements may well influence the accessibility of the locus to the recombination machinery. Moreover, this locus did not mimic a normal  $\text{V}\alpha\text{J}\alpha$  recombination product in that it contained intact *Tcrd* gene segments. Experiments in a more physiological *Tcr* knock-in allele that closely mimicked a product of  $\text{V}\alpha\text{J}\alpha$  recombination failed to find evidence for TCR editing (Buch et al., 2002). However, all these models suffered from premature expression of TCR $\alpha$  that is known to alter T cell development. Here, we revisited the problem of TCR editing with a new *Tcr* knock-in allele that ensured physiological timing of TCR $\alpha$  expression and was capable of secondary rearrangements. No evidence for TCR editing was found in this system and autoreactive thymocytes were instead eliminated by negative selection.

It remains theoretically possible that TCR editing can still contribute to tolerance under certain circumstances—for example for MHC class II-restricted TCRs, when antigen is expressed in a particular subcompartment of the thymus (i.e., the medulla rather than the cortex) or by a particular type of

antigen-presenting cell. However, both TCRs for which editing was suggested to take place were MHC class I restricted (Wang et al., 1998; McGargill et al., 2000). Moreover, initial evidence for TCR editing in the OT-I model was provided in a system where antigen expression was restricted to thymic cortex (McGargill et al., 2000) and later extended to a situation when it was expressed ubiquitously (Mayerova and Hogquist, 2004). This pattern of antigen expression is thus very similar to the situation in male HY-STOP mice, where ubiquitously expressed antigen is first encountered by autoreactive thymocytes in the cortex, and therefore cannot explain the observed differences.

It was suggested that TCRs have certain intrinsic differences determining their ability or inability to induce editing (Mayerova and Hogquist, 2004). An obvious candidate for such a difference is the affinity of the TCR for self-ligand. Both TCRs that were suggested to induce TCR editing were relatively inefficient in induction of negative selection, as judged by the high frequency of TCR-transgenic DP thymocytes in the presence of antigen (Wang et al., 1998; McGargill et al., 2000), whereas negative selection by the HY TCR is highly efficient. Thus, it is conceivable that only more weakly autoreactive TCRs mediate TCR editing. Editing, however, would require prolongation or reinduction of *Rag* expression in autoreactive thymocytes. We demonstrate here that both weak (female HY mice) and strong (male HY mice) TCR signals repress *Rag* expression. It does not seem probable, although remains formally possible, that a TCR signal of intermediate strength would have an exclusive ability to induce *Rag*. Along these lines, McGargill et al. (2000) reported less efficient *Rag* down-regulation in unfractionated OT-I DP thymocytes when antigen was present. However, as the DP compartment includes cells both before and after TCR signaling, this result can be explained by selective survival of *Rag*-high preselection DP thymocytes.

Antigen-independent secondary rearrangements reported previously in other models (Wang et al., 1998; Buch et al., 2002) also took place in the HY-STOP system. Such secondary rearrangements together with the fact that TCR editing was suggested only for systems where negative selection was inefficient, may explain the seeming discrepancy between the present and some earlier results. Indeed, as suggested previously (Buch et al., 2002), the observed relative increase in clone-type-negative cells interpreted as resulting from TCR editing may be simply explained by selective survival of the cells that rearranged their *Tcr* loci before antigen exposure and therefore escaped negative selection.

In conclusion, the experiments in this novel physiological knock-in model do not provide any evidence for self-antigen-induced TCR editing, although unequivocally showing negative selection by deletion of autoreactive thymocytes.

## MATERIALS AND METHODS

**Generation of HY-I (*Tcr*<sup>tm1.1(Tcr)</sup>*Rsk*) ES cells.** All ES cell work was performed in ES cell medium/DMEM containing 15% fetal calf serum, 2 mM L-glutamine, 1 mM sodium pyruvate, 0.1 mM MEM-non essential amino acids, 2,000 U/ml LIF, and 0.1 mM β-mercaptoethanol. C57BL/6

females were superovulated and mated with HY-I males. 3.5 d after coitus, blastocysts were isolated and cultured on embryonic fibroblast feeders in ES cell medium supplemented with PD98059 MEK Inhibitor (final concentration 13.5 μg/ml). On day 4 or 5 after plating, the blastocysts, outgrowing inner cell mass-derived clumps, were trypsinized and plated on fresh feeder cells. Presence of HY-I allele in the resulting ES cell lines was confirmed by PCR.

**Generation of HY-STOP mice.** Targeting pC3T1STOP construct was derived from pC3T1 vector used for generation of the original HY-I mouse (Buch et al., 2002) by removal of elements upstream of HY VαJα and insertion of floxed STOP cassette (Buch et al., 2005) downstream of it. Conventional LoxP sites flanking the STOP cassette were replaced with loxP2272 sites that can only recombine with each other (Siegel et al., 2001; Luche et al., 2007). Linearized pC3T1STOP vector was electroporated into HY-I ES cells. ES cells were selected with G418 and ganciclovir. ES cell clones were screened by PCR with primers upstream of short arm of homology (5'-AAGGCTGTTGTGATAAATGTGC-3') and in the STOP cassette (5'-CGGAATAGGAACTTCGTCGAG-3'; Fig. 1 A). Integration site was confirmed by sequencing the PCR product. The fact that HY-I and not WT allele was targeted was confirmed by PCR that detects disruption of the original HY-I locus by STOP cassette insertion. Two clones satisfying these criteria were injected into C57BL/6 albino blastocysts to generate chimeric males that were directly bred to *Tcr*<sup>-/-</sup> females containing TCR HYβ and Rorγt-Cre transgenes to generate mice with *Tcr*<sup>HY-STOP/-</sup>HYβ Rorγt-Cre genotype referred as HY-STOP mice. All mice were bred and maintained in the specific pathogen-free animal facilities of the Dana-Farber Cancer Institute (DFCI). All animal procedures were done in compliance with the guidelines of the DFCI Animal Resources Facility, which operates under regulatory requirements of the U.S. Department of Agriculture and the Association for Assessment and Accreditation of Laboratory Animal Care.

**BM chimeras.** In some experiments BM chimeras were used to amplify mice with *Tcr*<sup>HY-STOP/-</sup>HYβ Rorγt-Cre genotype. To this end, C57BL/6 males and females were lethally irradiated (1,000 rad, with a γ-cell 40 irradiator with a cesium source) and injected with 2–8 × 10<sup>6</sup> T cell-depleted, sex-matched BM cells from *Tcr*<sup>HY-STOP/-</sup>HYβ Rorγt-Cre animals. Mice were analyzed 4–10 wk after the transfer.

**Flow cytometry.** mAbs specific for CD4 (RM4-5), CD8a (53–6.7), CD69, TCRβ (H57-597), TCR Vβ8.1/8.2 (KJ16–133.18), HYα (T3.70), Vα2 (B20.1), Vα3.2 (RR3–16), PD-1 (29F.1A12), and Helios (22.F6) were purchased from BD, eBioscience, or BioLegend and were used as biotin, FITC, phycoerythrin (PE), peridinin chlorophyll protein (PerCP), PerCP-Cy5.5, PE-Cy7, allophycocyanin (APC), APC-Cy7, or Pacific blue conjugates. Fluorochrome-conjugated streptavidin was used to reveal staining with biotinylated mAb. Intracellular staining for Helios was performed using the Foxp3 staining buffer set (eBioscience). Intracellular staining for cleaved caspase-3 was performed using 5A1E rabbit monoclonal antibody, Alexa Fluor 647-labeled anti-rabbit F(ab')<sub>2</sub> fragment (both from Cell Signaling Technology) and the Cytotfix/Cytoperm kit (BD) according to the manufacturer's instructions. Flow cytometry was performed on FACSria (BD). Data were analyzed with FlowJo software (Tree Star).

**Real-time PCR.** Indicated populations were sorted, total RNA was prepared with RNeasy Plus kit (QIAGEN) with gDNA Eliminator columns. cDNA was synthesized using Superscript II reverse transcription (Invitrogen) according to the manufacturer's recommendations. Real-time RT-PCR was performed on CFX Connect cyclor (Bio-Rad Laboratories) using TaqMan PCR master mix (Bioline). Mm01270936\_m1 (*Rag1*), Mm00501300\_m1 (*Rag2*), and Mm99999915\_g1 (*Gapdh*) TaqMan Gene Expression Assays (Applied Biosystems) were used. No *Rag1* or *Rag2* signals were detected in the absence of reverse transcription. Standard curve method was used for data analysis.

**Online supplemental material.** Fig. S1 shows TCRβ and HYα staining of WT, *Tcr*<sup>-/-</sup>, *Tcr*<sup>-/-</sup> HYβ, and HY-STOP DP thymocytes, isotype controls for

this staining, and gating strategy for TCR $\beta$ <sup>-</sup> cells. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20130876/DC1>.

We would like to thank S. Schlenner and M. Gleimer for helpful discussions. The authors are grateful to V. Schmidt and G. Singh for technical assistance and to the staff of Brigham & Women's Hospital Transgenic Core Facility for ES cell transfection and microinjection.

These studies were supported by National Institutes of Health grants R01 A145846 and R01 A151378 to Harald von Boehmer and R37 AI054636 to Klaus Rajewsky, and Swiss National Fund SNF 310030\_132713 grant to Thorsten Buch.

The authors declare no competing financial interests.

Submitted: 29 April 2013

Accepted: 30 July 2013

## REFERENCES

- Baldwin, T.A., and K.A. Hogquist. 2007. Transcriptional analysis of clonal deletion in vivo. *J. Immunol.* 179:837–844.
- Baldwin, T.A., M.M. Sandau, S.C. Jameson, and K.A. Hogquist. 2005. The timing of TCR $\alpha$  expression critically influences T cell development and selection. *J. Exp. Med.* 202:111–121. <http://dx.doi.org/10.1084/jem.20050359>
- Bruno, L., H.J. Fehling, and H. von Boehmer. 1996. The alpha beta T cell receptor can replace the gamma delta receptor in the development of gamma delta lineage cells. *Immunity.* 5:343–352. [http://dx.doi.org/10.1016/S1074-7613\(00\)80260-5](http://dx.doi.org/10.1016/S1074-7613(00)80260-5)
- Buch, T., F. Rieux-Laucat, I. Förster, and K. Rajewsky. 2002. Failure of H-Y-specific thymocytes to escape negative selection by receptor editing. *Immunity.* 16:707–718. [http://dx.doi.org/10.1016/S1074-7613\(02\)00312-6](http://dx.doi.org/10.1016/S1074-7613(02)00312-6)
- Buch, T., F.L. Heppner, C. Tertilt, T.J. Heinen, M. Kremer, F.T. Wunderlich, S. Jung, and A. Waisman. 2005. A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation after toxin administration. *Nat. Methods.* 2:419–426. <http://dx.doi.org/10.1038/nmeth762>
- Croxford, A.L., O. Akilli-Ozturk, F. Rieux-Laucat, I. Förster, A. Waisman, and T. Buch. 2008. MHC-restricted T cell receptor signaling is required for alpha beta TCR replacement of the pre T cell receptor. *Eur. J. Immunol.* 38:391–399. <http://dx.doi.org/10.1002/eji.200737054>
- Daley, S.R., D.Y. Hu, and C.C. Goodnow. 2013. Helios marks strongly autoreactive CD4<sup>+</sup> T cells in two major waves of thymic deletion distinguished by induction of PD-1 or NF- $\kappa$ B. *J. Exp. Med.* 210:269–285. <http://dx.doi.org/10.1084/jem.20121458>
- Eberl, G., and D.R. Littman. 2004. Thymic origin of intestinal alphabeta T cells revealed by fate mapping of ROR $\gamma$ mat<sup>+</sup> cells. *Science.* 305:248–251. <http://dx.doi.org/10.1126/science.1096472>
- Egawa, T., T. Kreslavsky, D.R. Littman, and H. von Boehmer. 2008. Lineage diversion of T cell receptor transgenic thymocytes revealed by lineage fate mapping. *PLoS ONE.* 3:e1512. <http://dx.doi.org/10.1371/journal.pone.0001512>
- Groettrup, M., and H. von Boehmer. 1993. T cell receptor  $\beta$  chain dimers on immature thymocytes from normal mice. *Eur. J. Immunol.* 23:1393–1396. <http://dx.doi.org/10.1002/eji.1830230633>
- Kim, H.J., B. Verbinnen, X. Tang, L. Lu, and H. Cantor. 2010. Inhibition of follicular T-helper cells by CD8(+) regulatory T cells is essential for self tolerance. *Nature.* 467:328–332. <http://dx.doi.org/10.1038/nature09370>
- Kim, H.J., X. Wang, S. Radfar, T.J. Sproule, D.C. Roopenian, and H. Cantor. 2011. CD8<sup>+</sup> T regulatory cells express the Ly49 Class I MHC receptor and are defective in autoimmune prone B6-Yaa mice. *Proc. Natl. Acad. Sci. USA.* 108:2010–2015. <http://dx.doi.org/10.1073/pnas.1018974108>
- Lakso, M., B. Sauer, B. Mosinger Jr., E.J. Lee, R.W. Manning, S.H. Yu, K.L. Mulder, and H. Westphal. 1992. Targeted oncogene activation by site-specific recombination in transgenic mice. *Proc. Natl. Acad. Sci. USA.* 89:6232–6236. <http://dx.doi.org/10.1073/pnas.89.14.6232>
- Lee, J., and S. Desiderio. 1999. Cyclin A/CDK2 regulates V(D)J recombination by coordinating RAG-2 accumulation and DNA repair. *Immunity.* 11:771–781. [http://dx.doi.org/10.1016/S1074-7613\(00\)80151-X](http://dx.doi.org/10.1016/S1074-7613(00)80151-X)
- Li, Z., D.I. Dordai, J. Lee, and S. Desiderio. 1996. A conserved degradation signal regulates RAG-2 accumulation during cell division and links V(D)J recombination to the cell cycle. *Immunity.* 5:575–589. [http://dx.doi.org/10.1016/S1074-7613\(00\)80272-1](http://dx.doi.org/10.1016/S1074-7613(00)80272-1)
- Luche, H., O. Weber, T. Nageswara Rao, C. Blum, and H.J. Fehling. 2007. Faithful activation of an extra-bright red fluorescent protein in “knock-in” Cre-reporter mice ideally suited for lineage tracing studies. *Eur. J. Immunol.* 37:43–53. <http://dx.doi.org/10.1002/eji.200636745>
- Mayerova, D., and K.A. Hogquist. 2004. Central tolerance to self-antigen expressed by cortical epithelial cells. *J. Immunol.* 172:851–856.
- McCaughy, T.M., T.A. Baldwin, M.S. Wilken, and K.A. Hogquist. 2008. Clonal deletion of thymocytes can occur in the cortex with no involvement of the medulla. *J. Exp. Med.* 205:2575–2584. <http://dx.doi.org/10.1084/jem.20080866>
- McGargill, M.A., J.M. Derbinski, and K.A. Hogquist. 2000. Receptor editing in developing T cells. *Nat. Immunol.* 1:336–341. <http://dx.doi.org/10.1038/79790>
- McMahan, C.J., and P.J. Fink. 1998. RAG reexpression and DNA recombination at T cell receptor loci in peripheral CD4<sup>+</sup> T cells. *Immunity.* 9:637–647. [http://dx.doi.org/10.1016/S1074-7613\(00\)80661-5](http://dx.doi.org/10.1016/S1074-7613(00)80661-5)
- Mingueneau, M., T. Kreslavsky, D. Gray, T. Heng, R. Cruse, J. Ericson, S. Bendall, M.H. Spitzer, G.P. Nolan, K. Kobayashi, et al; Immunological Genome Consortium. 2013. The transcriptional landscape of  $\alpha\beta$  T cell differentiation. *Nat. Immunol.* 14:619–632. <http://dx.doi.org/10.1038/ni.2590>
- Mostoslavsky, R., and F.W. Alt. 2004. Receptor revision in T cells: an open question? *Trends Immunol.* 25:276–279. <http://dx.doi.org/10.1016/j.it.2004.04.001>
- Nemazee, D., and K.A. Hogquist. 2003. Antigen receptor selection by editing or downregulation of V(D)J recombination. *Curr. Opin. Immunol.* 15:182–189. [http://dx.doi.org/10.1016/S0952-7915\(03\)00008-6](http://dx.doi.org/10.1016/S0952-7915(03)00008-6)
- Page, D.M., L.P. Kane, J.P. Allison, and S.M. Hedrick. 1993. Two signals are required for negative selection of CD4<sup>+</sup>CD8<sup>+</sup> thymocytes. *J. Immunol.* 151:1868–1880.
- Pelanda, R., S. Schwers, E. Sonoda, R.M. Torres, D. Nemazee, and K. Rajewsky. 1997. Receptor editing in a transgenic mouse model: site, efficiency, and role in B cell tolerance and antibody diversification. *Immunity.* 7:765–775. [http://dx.doi.org/10.1016/S1074-7613\(00\)80395-7](http://dx.doi.org/10.1016/S1074-7613(00)80395-7)
- Siegel, R.W., R. Jain, and A. Bradbury. 2001. Using an in vivo phagemid system to identify non-compatible loxP sequences. *FEBS Lett.* 505:467–473. [http://dx.doi.org/10.1016/S0014-5793\(01\)02806-X](http://dx.doi.org/10.1016/S0014-5793(01)02806-X)
- Terrence, K., C.P. Pavlovich, E.O. Matechak, and B.J. Fowlkes. 2000. Premature expression of T cell receptor (TCR) $\alpha\beta$  suppresses TCR $\gamma\delta$  gene rearrangement but permits development of  $\gamma\delta$  lineage T cells. *J. Exp. Med.* 192:537–548. <http://dx.doi.org/10.1084/jem.192.4.537>
- Uematsu, Y., S. Ryser, Z. Dembić, P. Borgulya, P. Krimpenfort, A. Berns, H. von Boehmer, and M. Steinmetz. 1988. In transgenic mice the introduced functional T cell receptor beta gene prevents expression of endogenous beta genes. *Cell.* 52:831–841. [http://dx.doi.org/10.1016/0092-8674\(88\)90425-4](http://dx.doi.org/10.1016/0092-8674(88)90425-4)
- Wang, F., C.Y. Huang, and O. Kanagawa. 1998. Rapid deletion of rearranged T cell antigen receptor (TCR) Valpha-Jalpha segment by secondary rearrangement in the thymus: role of continuous rearrangement of TCR alpha chain gene and positive selection in the T cell repertoire formation. *Proc. Natl. Acad. Sci. USA.* 95:11834–11839. <http://dx.doi.org/10.1073/pnas.95.20.11834>