


RESEARCH ARTICLE

Glaucoma and Alzheimer: Neurodegenerative disorders show an adrenergic dysbalance

Bettina Hohberger^{1*}, Harald Prüss², Christian Mardin¹, Robert Lämmer¹, Johannes Müller³, Gerd Wallukat³

1 Department of Ophthalmology, Universität of Erlangen-Nürnberg, Friedrich-Alexander-University of Erlangen-Nürnberg, Erlangen, Germany, **2** Department of Neurology, Charité-Universitätsmedizin Berlin, Berlin, Germany, **3** Berlin Cures GmbH, Berlin, Germany

* bettina.hohberger@uk-erlangen.de



Abstract

Glaucoma disease is characterized by an increased intraocular pressure (IOP), glaucomatous alterations of the optic disc and corresponding visual field defects. Even lowering the main risk factor IOP until an individual target level does not prevent this neurodegenerative disorder from proceeding. Several autoimmune mechanisms were discovered, partly showing a functionality. One of these autoimmune phenomena targets the β 2-adrenergic receptor (β 2-AR; i.e. agonistic autoantibodies; β 2-agAAb) and is linked to an elevated IOP and an impaired retinal microcirculation. As neurodegenerative disorder, Alzheimer's Disease (AD) is postulated to share a common molecular mechanism with glaucoma. In the present study we investigated autoimmune phenomena targeting the β 2-AR in patients with AD. Sera of the patients were analyzed in a rat cardiomyocyte bioassay for the presence of functional autoantibodies against β 2-AR. In addition, different species of amyloid beta ($A\beta$) monomers were tested ($A\beta$ 1-14, $A\beta$ 10-25, $A\beta$ 10-37, $A\beta$ 1-40, $A\beta$ 1-42, $A\beta$ 28-40, and $A\beta$ -[Pyr]3-43). Our results demonstrate that none of the short-chain $A\beta$ ($A\beta$ 1-14, $A\beta$ 10-25, or $A\beta$ 28-40) showed any agonistic or inhibitory effect on β 2-AR. Contrary, long-chain $A\beta$ -[Pyr]3-43, representing a major neurogenic plaque component, exerted an activation that after blocking by the β 2-AR antagonist ICI118.551, could be identified as that the effect was realized via the β 2-AR. Moreover, the long chain $A\beta$ 1-40, $A\beta$ 1-42, and $A\beta$ 10-37, yet not the short-chain $A\beta$ peptides prevented the clenbuterol induced desensitization of the β 2-AR. In addition, we identified functional autoantibodies in the sera of AD patients, activating the β 2-AR, like the β 2-agAAb found in patients with glaucoma. As autoimmune mechanisms were reportedly involved in the pathogenesis of glaucoma and Alzheimer's Disease, we postulate that overstimulation of the β 2-AR pathway can induce an adrenergic overdrive, that may play an important role in the multifactorial interplay of neurodegenerative disorders.

OPEN ACCESS

Citation: Hohberger B, Prüss H, Mardin C, Lämmer R, Müller J, Wallukat G (2022) Glaucoma and Alzheimer: Neurodegenerative disorders show an adrenergic dysbalance. PLoS ONE 17(10): e0272811. <https://doi.org/10.1371/journal.pone.0272811>

Editor: Madepalli K. Lakshmana, Torrey Pines Institute for Molecular Studies, UNITED STATES

Received: February 11, 2022

Accepted: July 26, 2022

Published: October 6, 2022

Copyright: © 2022 Hohberger et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data are within the paper and its [Supporting information](#) files.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Glaucoma disease is a worldwide burden, being the second leading cause of blindness in the industrial nations. This neurodegenerative disease is characterized by an increased intraocular

pressure (IOP), a glaucomatous optic disc and visual field defects, going along with disease progression. Up to now the exact pathomechanisms are still elusive. An elevated IOP, the main risk factor, is the target of conservative, laser and surgical therapy. Yet, despite lowering IOP, glaucoma disease proceeds, as no causal therapy is available until now [1]. Neurodegeneration in glaucoma is featured by apoptosis of the retinal ganglion cells (RGC). Immune and autoimmune mechanisms can interplay in these molecular interactions of loss of RGC as recent studies suggested. Data of mice models yielded a significant RGC loss mediated by T-lymphocytes via heat-shock-proteins in the presence of only slightly elevated IOP [2]. Changed cytokine levels [3] and microglial activation were observed in a mice model of ocular hypertension [4]. Even normotensive eyes were observed to show an apoptosis of RGC when the contralateral eye was hypertensive (mice model). This effect was attributed to lymphocytes as well [5]. Apoptosis itself, a vascular dysregulation or other external factors (e.g. nutrition, alteration of levels of trace elements) are able to trigger immune and autoimmune mechanisms. Thus, it is not surprising that diverse autoantibodies were detected in sera of patients with glaucoma. In this context, autoantibodies with a functional consequence are of special interest. Recent studies presented autoantibodies targeting the adrenergic β_2 -receptor (β_2 -AR) in sera of patients with open-angle glaucoma. These β_2 -autoantibodies activate their target (β_2 -agAAb) with consecutive overstimulation and desensitization of the receptor contrary to physiological agents. *In vivo* data suggested a link of the β_2 -agAAb to the increased IOP and impaired retinal microcirculation [6,7]. We propose that a chronic overstimulation of the β_2 -AR can result in a dysregulated homeostatic adrenergic balance. Receptor desensitization and internalization might be only one of the molecular features [8]. If this adrenergic disbalance has sufficient 'power' on cellular level, clinical characteristics occur in the patients (*imbalanced autonomic theory*; see also [9]). This adrenergic disbalance is certainly not only a local problem (in the eye). We assume that adrenergic alterations might be even in extraocular tissue of patients with neurodegenerative diseases.

Patients with Alzheimer's Disease (AD) show amyloid beta ($A\beta$) depositions being proposed to be the major mediator of neurotoxicity in the brain [10]. $A\beta$ is assumed to be one crosslink between glaucoma and AD. RGC express $A\beta$ [11,12]. Especially, intravitreal injection of $A\beta_{1-42}$ was observed to induce RGC loss [13]. $A\beta$ is generated after cleavage of the amyloid precursor protein (APP). Ni et al. has shown that a β_2 -adrenergic stimulation that activates the gamma secretase is involved in the formation of $A\beta$ from the APP [14]. This cleavage can be mediated by vessel endothelial cell enzymes and platelets, especially when platelets contact the endothelium. Any thrombosis can enhance this $A\beta$ generation. Any thrombosis can enhance this $A\beta$ generation.

Glaucoma and AD disease were observed to share common features clinically and biochemically. Patients with open-angle glaucoma are at higher risk [15] and showed a higher incidence of AD [16]. Vis-a-vis patients with AD are at higher risk of glaucoma [17]. Both, glaucoma and AD, were observed to be age-related disorders [18–22]. As the eye, especially retinal tissue, represents "cerebral tissue" considering embryology, it is not remarkable that neurodegeneration and amyloid- β plaques were seen within the retina in AD patients [23–25]. Since several years a common pathomechanism of glaucoma and AD has been suggested [26–28]. Patients with AD showed amyloid beta ($A\beta$) depositions being proposed to be the major mediator of neurotoxicity in the brain [10]. $A\beta$ is assumed to be one crosslink between glaucoma and AD. RGC express $A\beta$ [11,12]. Especially, intravitreal injection of $A\beta_{1-42}$ was observed to induce RGC loss [13]. Degeneration of RG and retrograde degeneration of axons in the lateral geniculate body are two findings, being present in AD patients [13,25,29,30]. Even amyloid- β can induce RGC loss within the retina itself [13]. These neurodegenerative features were also observed in glaucomatous eyes [31–34]. The diagnostic correlate of RGC loss can be seen in retinal nerve fiber layer thinning (RNFL) *in vivo*, visualized by magnet

resonance tomography and optical coherence tomography (OCT) [31,33]. Both, patients with glaucoma and AD, showed a RNFL thinning especially in the superior and/or inferior retina [35–38]. Thus, it can be hypothesized that there is common pathway, linking both neurodegenerative disorders (glaucoma and AD). Functional AAb against the β 2-AR were not only seen in the sera of patients with glaucoma but also in sera of patients with AD symptoms [39]. These β 2-agAAb but also the long chain A β peptides prevent the desensitization of the β 2AR and induce a permanent stimulation of this adrenergic receptor. As β 2-agAAb show a link to an impaired microcirculation (glaucoma) and A β expression is increased during impaired microcirculation (i.e. during and after thrombosis; AD), we hypothesize that there is a common molecular mechanisms between adrenergic dysfunction (β 2-agAAb) and A β . The aim of the study was to investigate sera of patients with AD and glaucoma for the presence of autoantibodies against β 2-AR, and the effect of fragments and the complete A β peptides on the β 2-AR and whose desensitization.

Material and methods

Patients and controls

Sera of 11 subjects with AD (3 female, 8 male, age: 40–87 years) and 18 subjects with primary open-angle glaucoma (7 female, 11 male, age: 42–76 years) were collected for the present study. Glaucoma patients were recruited at the Department of Ophthalmology, University of Erlangen-Nürnberg, Friedrich-Alexander-University Erlangen-Nürnberg [Erlangen Glaucoma Registry, ISSN 2191-5008, CS-2011; NTC00494923] [1]. Patients with AD were recruited at the Department of Neurology, Charité'-Universitätsmedizin Berlin. Blood samples were collected and centrifuged, and sera were stored at -20°C. All individuals declared informed consent. The study was performed according to the tenets of the Declaration of Helsinki and approved by the ethic committee of the university of Erlangen, the institutional Review Board of Charité'-Universitätsmedizin Berlin and the Max-Delbrück-Centrum for Molecular Medicine, Berlin-Buch (Y 9004/19, 02.11.2020).

Primary Open-Angle Glaucoma (POAG) patients

The criteria of POAG were: an open anterior chamber angle, IOP>21 mmHg (confirmed at least once, measured by Goldmann applanation tonometry), and a glaucomatous optic disc, classified after Jonas [40]. Functional loss of perimetric field was measured and had to be confirmed at least once according to the following criteria: (I) at least three adjacent test points having a deviation \geq 5 dB and with one test point with a deviation >10 dB lower than normal, or (II) at least two adjacent test points with a deviation \geq 10 dB, or (III) at least three adjacent test points with a deviation \geq 5 dB abutting the nasal horizontal meridian or (IV) a mean visual field defect of >2.6 dB.

Patients with AD

In the study we included 11 patients (3 female, 8 male) with diagnosed AD with a mean age of 70 years. All the investigated patients had hypertension and 3/11 patients had coronary heart disease. One patient was diseased with a peripheral artery disease, 7/11 patients were treated with antidementia drugs.

Cardiomyocyte bioassay

The cardiomyocyte bioassay was done as described previously (Junemann et al., 2018). Cell culture of cardiac myocytes of heart ventricle (1–3 day-old Sprague-Dawley rats) was used.

digested with a 0.25% solution of crude porcine trypsin (Serva, Germany). The cells were dispersed by trypsinization and suspended in a SM20-I medium (Biochrom, Germany), glutamine (Serva, Germany), containing streptomycin (HEFA Pharma; Germany), penicillin (Heyl, Germany), hydrocortisone (Merck, Germany), fluorodeoxyuridine (Serva, Germany), and 10% heat-inactivated neonatal calf serum (Gibco, Germany). The cells were seeded (field density of 160.000 cells/cm²) the culture medium was replaced after 24 hours. The cardiomyocytes were cultured for 3–4 days (37°C) before being used in the experiments. Firstly, the basal beating rate of the cardiomyocytes was measured for 15 seconds; at a heated stage of an inverted microscope at 37°C visually or with an ION OPTIX measurement place. Subsequently the immunoglobulin (Ig) fractions were added for 60 min in a dilution of 1:40. Next, the beating rates (BR) of the cardiomyocytes were counted again on 6 marked spontaneously beating cells or cell cluster. The data are expressed as “increase in number of beats/15sec”. The calculated cut off value was 1.83. Application of a specific β 2-blocker, i.e. ICI 118.551 (0.1 μ M) was done for identification of the receptor type.

Preparation of the serum immunoglobulins

Human IgG were prepared from each patients' sera by direct ammonium sulfate precipitation (final concentration of 40%; overnight at 4°C). The precipitates were centrifuged (4,000 \times g; 30 min) and the pellets were dissolved in dialysis buffer (154 mmol/l NaCl, 10 mmol/l Na₂HPO₄/NaH₂PO₄, pH 7.2). This procedure was repeated twice. Each sample was dialyzed against phosphate-buffered saline (4°C; 3 days). Storage of the samples was done at -20°C.

Desensitization experiments

In these experiments the cardiomyocytes were treated with 1 μ M clenbuterol with and without different A β -peptides and with AAb directed against the β 2-adrenoceptor. The incubation time was 120 min followed by a washing procedure and anew stimulation with clenbuterol at the same concentration. In this experiment we used AAb prepared from the sera of AD patients and the short- and long-chain A β -peptides: A β 1–14, A β 25–35, A β 28–40, A β 10–37, A β 1–40, and A β -1-42.

A β peptides

The fragmented A β peptides were synthesized by Dr. Beyermann (FMP Berlin). The long-chain A β peptides A β 1–40, A β 1–42 and also A β -[Pyr[3–43 were from Sigma-Aldrich.

Statistical analysis

Statistical analysis was done by SPSS (version 21) and Excel and the unpaired t-test. Data are shown as absolute value, mean and standard deviation.

Results

β 2-AAb in patients with glaucoma and Alzheimer's disease

AAbs prepared from patients with POAG showed a mean beat rate of 4.5 \pm 0.1 U (range: 3.3–5.2 U) that was realized via the β 2-AR. This increase in the beat rate could be blocked by the specific β 2-blocker ICI 118.551 (Fig 1). Immunoglobulins prepared from AD patient sera showed an increase of the mean beating rate of 4.5 \pm 0.4 U (range: 2.3–7.2 U). As observed for patients with POAG, this increase in the beat rate could be blocked by the specific β 2-blocker ICI 118.551 (Fig 1). These AAb recognize also the β 2-adrenoceptor. Thus, a seropositivity for β 2-AAb was observed for patients with POAG and AD. Healthy individuals, showing a




Fig 1. Comparison of the activity of the functional β 2-adrenoceptor autoantibodies (β 2-AAb) of patients with glaucoma (POAG) (n = 12) and Alzheimer's disease (AD) (n = 11). Both agonist-like effects were blocked by the β 2-adrenoceptor antagonist ICI 118.551 (0.1 μ M), ($p < 0.001$). The experiments with the AAb of the AD patients were done in the presence of the α 1- adrenoceptor antagonist urapidil to block the agAAb against the α 1-adrenoceptor that are also present in the sera of AD patients.

<https://doi.org/10.1371/journal.pone.0272811.g001>

seronegativity of G-Protein coupled receptors (GPCR)-AAb, have been tested in our previous works using the same assay [7,41].

In vitro analysis of the effects of amyloid β ($A\beta$) peptides

In order to identify a possible functional action of $A\beta$ peptides on the GPCR, different species of amyloid beta ($A\beta$) peptide were tested in the used bioassay. Following $A\beta$ peptides were investigated in a concentration of 0.1 μ M: $A\beta$ 1-14, $A\beta$ 10-37, $A\beta$ 25-35, $A\beta$ 28-40, $A\beta$ 1-40, $A\beta$ 1-42, and $A\beta$ -[Pyr]3-43). Our results demonstrate that only the $A\beta$ peptide $A\beta$ -[Pyr]3-43, but not the other $A\beta$ peptides or $A\beta$ fragments are able to activate the β 2 adrenoceptor similar as the classical β -adrenergic agonists. The long-chain $A\beta$ -[Pyr]3-43, representing a major neurogenic plaque component, exerted an activation of the β 2-AR blocked by the β 2-AR antagonist ICI 118.551 but not by the β 1-AR antagonists bisoprolol or the α 1- AR blocker urapidil (Fig 2).

Desensitization experiments

A long-term incubation of the cardiomyocytes with adrenergic agonists leads to desensitization of the adrenoceptors and to a resistant to the used drug. The incubation of the




Fig 2. Functional effect realized via the β 2 adrenoceptor by the truncated A β peptide A β -[PYR]3-43 (n = 5): This effect was blocked by ICI 118.551 (n-4, p<0.001), yet not by the β 1 adrenoceptor antagonist bisoprolol or the α 1-adrenergic antagonist urapidil.

<https://doi.org/10.1371/journal.pone.0272811.g002>

cardiomyocytes with the β 2-adrenergic agonist clenbuterol caused a time dependent reduction of the response to this agonist. A washing procedure removed this agonist completely and a new stimulation with 1 μ M clenbuterol generate only a strong reduced response (Fig 3a). Yet this protection mechanism of the cell can be disturbed by agAAb directed against these receptors and by long-chain amyloid peptides. Additionally, β 2-agAAb caused a positive chronotropic effect that could not be washed out by a washing procedure. Therefore, the agAAb against the β 2-adrenoceptor must be blocked and removed from the receptor by the specific




Fig 3. Desensitization of the β 2 adrenergic response by the β 2-adrenergic agonist clenbuterol (a). This receptor desensitization was missed if the cells were stimulated with the β 2-adrenergic AAb prepared from AD patients (b).

<https://doi.org/10.1371/journal.pone.0272811.g003>




Fig 4. Influence of the Aβ peptides on the clenbuterol induced β2-adrenoceptor desensitization: The desensitization of the β-AR response by clenbuterol, was not influenced by the short chain Aβ peptides (a). However, the long chain Aβ peptides Aβ 10–37, Aβ1–40, and Aβ1–42 prevent the desensitization and exert a permanent stimulation of the β2-AR signal cascade (b).

<https://doi.org/10.1371/journal.pone.0272811.g004>

β2-adrenoceptor antagonist ICI118.551. After the washing procedure the addition of clenbuterol led to a maximal stimulation. These data demonstrate that the β2-agAAb prevent the desensitization of the receptor (Fig 3b). Similar results were observed, if the long chain Aβ peptides plus clenbuterol were added to the cardiomyocytes. Under these conditions the β2-adrenoceptor was not desensitized and after the washing procedure the clenbuterol effect was not diminished (4b, Tab1). Contrary, in the presence of the short-chain Aβ peptides clenbuterol desensitized the β2 adrenoceptor like clenbuterol itself Fig 4a, Table 1).

Discussion

Glaucoma disease is one of the leading causes of irreversible blindness worldwide [42]. Its complex pathogenesis is known to be multifactorial with an immune and autoimmune involvement. Clinical studies observed that this neurodegenerative ocular disorder shows several common features with neurodegenerative disorders of the brain (e.g. AD) [26,27,43,44]. Thus, it is commonly accepted that both disease entities seem to have common molecular pathways and clinical characteristics. To the best of our knowledge the present study offers data of a common pathogenic involvement of the β2-adrenergic pathway both in AD and glaucoma. In sera of patients with AD and glaucoma a seropositivity for β2-agAAb was observed, respectively. In addition, the long-chain Aβ-[Pyr]3–42, representing the major neurogenic plaque component, showed a direct activation of the β2-AR that was blocked by the specific antagonist ICI118.551. This N-terminal truncated isoform of Aβ can trigger

Table 1. Influence of short and long chain Amyloid β peptides on the clenbuterol induced β2-adrenoceptor desensitization.

Time		5'	60'	120'	3 x wash	125'	Clenbuterol 1 μM	130'
Clenbuterol 1 μM	Clenbuterol 1μM	10.2±0.3	8.7±0.3	5.9±0.3		0.2±0.1		2.7±0.1
	+ Aβ 1–14	8.7±0.6	6.7±0.5	3.5±0.5		0.2±0.1		2.4±0.1
	+ Aβ 25–35	8.9±1.2	7.4±1.0	5.7±0.6		0.2±0.1		2.9±0.2
	+ Aβ 28–40	10.5±0.2	7.8±0.6	4.3±0.6		0.3±0.2		2.3±0.2
	+ Aβ 1–40	10.8±0.2	10.7±0.4	9.8±0.2		0.6±0.1		9.3±2.9
	+ Aβ 1–42	8.9±0.1	9.3±0.4	8.9±0.4		0.2±0.1		9.6±0.3
	+ Aβ 10–37	9.6±0.2	10.1±0.3	8.2±0.8		0.4±0.1		10.3±0.2

<https://doi.org/10.1371/journal.pone.0272811.t001>

neurodegeneration in a transgenic mouse model [45]) and developed a high neurotoxicity [46]. The other functional active A β peptides realize their effect in the used test model not via the β 2 adrenoceptor. Another aspect of the effects of the A β peptides exist in fact that the long chain A β 1–40, A β 1–42, A β 10–37 peptides prevented the clenbuterol induced desensitization of the β 2-AR similar as the β 2-adrenoceptor autoantibodies. All three mechanisms result finally in a boosted activation and permanent stimulation of the β 2-adrenergic pathways, potentially causing a dangerous adrenergic overdrive realized via the β 2-AR.

The autonomic system is a main actor in human body. It regulates important function and is the basis of human life. Any alterations in the sympathetic and/or parasympathetic pathways results in an impairment or overaction of the autonomic system. Adrenergic receptors are also present on pericytes thus it can be hypothesized that any alterations in adrenergic activity might influence and alter microcirculation. It is commonly accepted that an impaired microcirculation can induced amyloid plaques in AD [47–49] and is a very early pathogenetic alteration in glaucoma disease [50–52]. Up to now, it is still elusive which factors induce this impaired microcirculation in AD. Patients with AD showed an early involvement of the choroïdal microcirculation [53]. These clinical data were repeated in different murine models of AD (e.g. App NL-F) [54]. In addition, alterations in all retinal layers were observed, being dependent on whether patients have mild or moderate AD [55]. Interestingly, recent data showed that an impaired microcirculation is able to change the amyloid- β plaques in a mouse model [49]. Even in glaucoma pathogenesis, the exact pathogenesis is under investigation. There is evidence that an involvement of the autonomic system, especially the adrenergic β 2-mediated pathway might be present in dysregulation of IOP (glaucoma) [56] and capillary microcirculation (glaucoma [6] and AD), as patients with glaucoma and AD showed a seropositivity for β 2-AAb, respectively.

In patients with AD disease the β 2-adrenergic dysregulation might be mediated in three different ways. First is the direct activation of the β 2-AR by its classical natural agonist adrenaline or noradrenaline. Both compounds activate the β 1-, the β 2- and the β 3-AR. The stimulation with adrenaline, isoprenaline or the more β 2-AR specific agonist clenbuterol induced a positive chronotropic response in the cardiomyocyte bioassay, which could be blocked by the specific β 2-AR antagonist ICI118.551. It was shown that patients with AD develop functional active autoantibodies directed against the β 2-AR. These β 2-agAAb activate the β 2-AR in a similar manner as the β 2-adrenergic agonists [39], but in contrast to the agonists the β 2-agAAb prevent the desensitization of the corresponding receptor. This observation was also made in glaucoma patients [7]. The agonist like effect of this β 2-agaAAb was blocked by ICI118.551, yet not by the β 1-AR antagonist bisoprolol. Moreover, the agonistic effect of the β 2-agAAb was neutralized by a peptide corresponding to the second extracellular loop of this receptor [7]. However, the β 2-AR agAAB prepared from AD patient sera recognize the first extracellular loop of the β 2-AR [39]. In contrast to the classical β -AR agonists, β 2-agAAb activate the β 2-AR mediated signal cascade permanently. Moreover, the β 2-agAAb also prevent the desensitization, normally seen for the classical β 2-AR agonists, when agonists were added to β 2-agAAb pre-stimulated cardiomyocytes. This permanent stimulation without desensitization of the β 2-AR may represent a pathogenic factor that played an essential role in the pathogenesis of several diseases [57]. Another feature of these β 2-agAAb is the possible antagonistic action of these β 2-agAAb directed against the β 2-AR. It was shown by Mijares et al. that monovalent Fab fragments of monoclonal β 2-agAbbs exhibit antagonistic and not agonistic properties [58]. However, if these monovalent Fab fragments were crosslinked by an anti Fab antibody than the agonist-like effects were restored [59]. These data demonstrate that the β 2-agAAb also can act in an antagonistic manner under particular circumstances.

Secondly, it was shown that the activation of the β_2 -AR may play a role in the formation of amyloid β (A β). β_2 -AR agonists activate the gamma-secretase, forming A β from the amyloid precursor proteins (APP), resulting in an elevated accumulation of A β [14]. Moreover, a microperfusion of A β 1–40 in combination with the β_2 -adrenergic agonist clenbuterol into the hippocampus caused an augmented degradation of the A β 1:40 peptide in comparison to the controls without clenbuterol [60]. Some of these degradation products of the long-chain A β and A β 1–40 and A β 1–42 itself can stimulate different GPCR in a similar manner as the classical receptor agonists. For example, the A β -fragments A β 25–35 and A β 10–37 activate the α_1 -AR and this activation was abolished by the antagonist prazosin or urapidil. These A β -peptides recognize the first extracellular loop of the α_1 -adrenergic receptor and the A β 25–35 and A β 10–37 induced activation could be blocked by peptides corresponding to the first extracellular loop [61]. Such α_1 -adrenoceptor antibodies can influence moderately the contractility of the coronary and middle cerebral artery and strongly the renal artery of the rat [62]. Moreover, Tohda et al. demonstrated that A β 25–35 induced memory impairment, axonal atrophy and synaptic loss [63].

Wang and coworker observed in prefrontal cortical neurons that A β 1–40 and A β 1–42 can bind to the β_2 -AR, and activate the protein kinase A (PKA) in these [64,65]. This observation was not confirmed in our experiments using a functional cardiomyocyte assay. In our bioassay A β 1–40, A β 1–42 but also A β 1–14 activate the fast estradiol receptor GPR 30 (unpublished data). In contrast, the N-terminal truncated A β peptide A β -[Pyr]3–43 that representing a major A β plaque component was able to activate the β_2 -AR subtype in our cardiomyocyte model. This A β peptide, activate the β_2 -AR in similar manner as the agonist clenbuterol. This activation was neither influenced by the α_1 -adrenergic antagonist urapidil nor the β_1 -antagonist bisoprolol. Only the β_2 -antagonist ICI118,551 was able to block the activity induced by A β - [Pyr]3–43. Therefore, we assume that in the experiments of Wang on neuronal tissues the A β -peptides A β 1–40 and A β 1–42 were truncated by endogenous peptidases to A β -[Pyr]3–43 that can than recognize the β_2 -AR subtype.

A third possibility to influence the β_2 -AR activity can be seen in fact that the long-chain A β peptides A β 1–42, A β 1–40, and A β 10–37, yet not the short breaking products of the amyloid A β 1–14, A β 25–35 and A β 28–40 prevent desensitization of the β_2 -adrenergic agonist clenbuterol. The combination of the long chain A β peptides and clenbuterol induced a permanent activation of the β_2 -signal cascade similar to the response seen for the agonist-like β_2 -AABs. The β_2 -AABs induced their long-lasting effect via a cross-link between the two arms of the antibody molecule and two receptors. This cross-link stabilized the active conformation of the receptor and led to a permanent stimulation (as described above). Why the long-chain A β -peptides can prevent the desensitization of the adrenoceptors is an open question and under investigation. Yet, we were able to show that other proteins for example the C-reactive protein (CRP) can also prevent the desensitization of different GPCRs [66]. Moreover, the unspecific matrix-metalloprotease (MMP) inhibitor GM6001 prevented the desensitization of the adrenoceptors [67]. Therefore, is it assumable that proteases may play an important role in the process of desensitization. These observations indicate that the long chain A β can also contribute to the adrenergic overdrive.

We hypothesize that a dysregulated β_2 -mediated signal cascade might be a molecular mechanism of an autonomic dysregulation, both observed in neurodegenerative disorders, e.g. glaucoma and AD. A seropositivity for β_2 -AAB and the β -adrenergic agonists itself can enhance β_2 -AR activation. In addition to β_2 -AAB, the A β peptide A β -[Pyr]3–43 was also able to activate the β_2 -AR mediated signal cascade directly. Furthermore, the long chain A β peptides induced like the β_2 -AAB a permanent stimulation and deactivate the protection mechanism of

the cells against an adrenergic overstimulation. All these mechanisms may play an important role in the pathogenesis of glaucoma and AD.

Conclusion

Glaucoma and AD are known to be neurodegenerative disorders with an involvement of auto-immune mechanisms and impaired microcirculation. As both disease entities showed a seropositivity for β 2-AAbs we postulate that an overstimulated β 2-AR pathway can induce an adrenergic overdrive, which can be enhanced by amyloid- β peptides in patients with AD.

Supporting information

S1 File.
(DOCX)

Author Contributions

Conceptualization: Bettina Hohberger, Gerd Wallukat.

Data curation: Bettina Hohberger, Gerd Wallukat.

Investigation: Harald Prüss, Christian Mardin, Robert Lämmer, Gerd Wallukat.

Methodology: Gerd Wallukat.

Resources: Johannes Müller.

Supervision: Bettina Hohberger, Gerd Wallukat.

Visualization: Gerd Wallukat.

Writing – original draft: Bettina Hohberger, Gerd Wallukat.

Writing – review & editing: Harald Prüss, Christian Mardin, Robert Lämmer, Johannes Müller.

References

1. Hohberger B, Monczak E, Mardin CY. [26 Years of the Erlangen Glaucoma Registry: Demographic and Perimetric Characteristics of Patients Through the Ages]. *Klinische Monatsblätter für Augenheilkunde*. 2019; 236(5):691–8. Epub 2017/07/28. <https://doi.org/10.1055/s-0043-112856> PMID: 28750434.
2. Chen H, Cho KS, Vu THK, Shen CH, Kaur M, Chen G, et al. Commensal microflora-induced T cell responses mediate progressive neurodegeneration in glaucoma. *Nature communications*. 2018; 9(1):3209. <https://doi.org/10.1038/s41467-018-05681-9> PMID: 30097565.
3. Fernandez-Albarral JA, Salazar JJ, de Hoz R, Marco EM, Martin-Sanchez B, Flores-Salguero E, et al. Retinal Molecular Changes Are Associated with Neuroinflammation and Loss of RGCs in an Experimental Model of Glaucoma. *Int J Mol Sci*. 2021; 22(4). Epub 2021/03/07. <https://doi.org/10.3390/ijms22042066> PMID: 33669765.
4. Ramirez AI, de Hoz R, Fernandez-Albarral JA, Salobrar-Garcia E, Rojas B, Valiente-Soriano FJ, et al. Time course of bilateral microglial activation in a mouse model of laser-induced glaucoma. *Sci Rep*. 2020; 10(1):4890. Epub 2020/03/19. <https://doi.org/10.1038/s41598-020-61848-9> PMID: 32184450.
5. Gramlich OW, Godwin CR, Heuss ND, Gregerson DS, Kuehn MH. T and B Lymphocyte Deficiency in Rag1^{-/-} Mice Reduces Retinal Ganglion Cell Loss in Experimental Glaucoma. *Investigative ophthalmology & visual science*. 2020; 61(14):18. Epub 2020/12/16. <https://doi.org/10.1167/iovs.61.14.18> PMID: 33320171.
6. Hohberger B, Hosari S, Wallukat G, Kunze R, Krebs J, Muller M, et al. Agonistic autoantibodies against beta2-adrenergic receptor influence retinal microcirculation in glaucoma suspects and patients. *PloS one*. 2021; 16(5):e0249202. Epub 2021/05/08. <https://doi.org/10.1371/journal.pone.0249202> PMID: 33961631 EP1832600A1: Peptides against autoantibodies associated with glaucoma and use of these peptides; Kunze R, patent EP1832600A1: Peptides against autoantibodies associated with glaucoma

and use of these peptides; Krebs J, none; Muller M, none; Hennig T, none; Lammer R, none; Horn F, none; Munoz L.E., none; Herrmann M, patent EP1832600A1: Peptides against autoantibodies associated with glaucoma and use of these peptides; Mardin C, none. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

7. Junemann A, Hohberger B, Rech J, Sheriff A, Fu Q, Schlotzer-Schrehardt U, et al. Agonistic Autoantibodies to the beta2-Adrenergic Receptor Involved in the Pathogenesis of Open-Angle Glaucoma. *Frontiers in immunology*. 2018; 9:145. Epub 2018/02/28. <https://doi.org/10.3389/fimmu.2018.00145> PMID: 29483909.
8. Wallukat G. The beta-adrenergic receptors. *Herz*. 2002; 27(7):683–90. Epub 2002/11/20. <https://doi.org/10.1007/s00059-002-2434-z> PMID: 12439640.
9. Szentiványi A. The beta-adrenergic theory of the atopic abnormality in bronchial asthma. *J Allergy*. 1968; 42:203–32.
10. Wyss-Coray T. Inflammation in Alzheimer disease: driving force, bystander or beneficial response? *Nat Med*. 2006; 12(9):1005–15. Epub 2006/09/09.
11. Lee S, Jiang K, McIlmoyle B, To E, Xu QA, Hirsch-Reinshagen V, et al. Amyloid Beta Immunoreactivity in the Retinal Ganglion Cell Layer of the Alzheimer's Eye. *Frontiers in neuroscience*. 2020; 14:758. Epub 2020/08/28. <https://doi.org/10.3389/fnins.2020.00758> PMID: 32848548.
12. den Haan J, Morrema THJ, Verbraak FD, de Boer JF, Scheltens P, Rozemuller AJ, et al. Amyloid-beta and phosphorylated tau in post-mortem Alzheimer's disease retinas. *Acta Neuropathol Commun*. 2018; 6(1):147. Epub 2018/12/30. <https://doi.org/10.1186/s40478-018-0650-x> PMID: 30593285.
13. Simons ES, Smith MA, Dengler-Criss CM, Criss SD. Retinal ganglion cell loss and gliosis in the retinofugal projection following intravitreal exposure to amyloid-beta. *Neurobiol Dis*. 2021; 147:105146. Epub 2020/10/31. <https://doi.org/10.1016/j.nbd.2020.105146> PMID: 33122075.
14. Ni Y, Zhao X, Bao G, Zou L, Teng L, Wang Z, et al. Activation of beta2-adrenergic receptor stimulates gamma-secretase activity and accelerates amyloid plaque formation. *Nat Med*. 2006; 12(12):1390–6. Epub 2006/11/23. <https://doi.org/10.1038/nm1485> PMID: 17115048.
15. Lin IC, Wang YH, Wang TJ, Wang IJ, Shen YD, Chi NF, et al. Glaucoma, Alzheimer's disease, and Parkinson's disease: an 8-year population-based follow-up study. *PLoS one*. 2014; 9(9):e108938. Epub 2014/10/03. <https://doi.org/10.1371/journal.pone.0108938> PMID: 25275530.
16. Moon JY, Kim HJ, Park YH, Park TK, Park EC, Kim CY, et al. Association between Open-Angle Glaucoma and the Risks of Alzheimer's and Parkinson's Diseases in South Korea: A 10-year Nationwide Cohort Study. *Sci Rep*. 2018; 8(1):11161. Epub 2018/07/26. <https://doi.org/10.1038/s41598-018-29557-6> PMID: 30042382.
17. Lai SW, Lin CL, Liao KF. Glaucoma may be a non-memory manifestation of Alzheimer's disease in older people. *Int Psychogeriatr*. 2017:1–7. Epub 2017/05/30. <https://doi.org/10.1017/S1041610217000801> PMID: 28552085.
18. Guerreiro R, Bras J. The age factor in Alzheimer's disease. *Genome Med*. 2015; 7:106. Epub 2015/10/21.
19. Molinuevo JL, Casado-Naranjo I. Clinical profile of Alzheimer's disease: is the age of the patient a decisive factor? Results of the INFLUENCE study. *J Alzheimers Dis*. 2014; 39(2):227–32. Epub 2013/10/24. <https://doi.org/10.3233/JAD-131433> PMID: 24150103.
20. McMonnies CW. Glaucoma history and risk factors. *J Optim*. 2017; 10(2):71–8. Epub 2016/03/31. <https://doi.org/10.1016/j.optom.2016.02.003> PMID: 27025415.
21. Doucette LP, Rasnitsyn A, Seifi M, Walter MA. The interactions of genes, age, and environment in glaucoma pathogenesis. *Survey of ophthalmology*. 2015; 60(4):310–26. Epub 2015/04/25. <https://doi.org/10.1016/j.survophthal.2015.01.004> PMID: 25907525.
22. Klein R, Klein BE, Moss SE. Age-related eye disease and survival. The Beaver Dam Eye Study. *Archives of ophthalmology*. 1995; 113(3):333–9. Epub 1995/03/01. <https://doi.org/10.1001/archophth.1995.01100030089026> PMID: 7887847.
23. Whitehouse PJ, Price DL, Clark AW, Coyle JT, DeLong MR. Alzheimer disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol*. 1981; 10(2):122–6. Epub 1981/08/01. <https://doi.org/10.1002/ana.410100203> PMID: 7283399.
24. van de Nes JA, Nafe R, Schlote W. Non-tau based neuronal degeneration in Alzheimer's disease—an immunocytochemical and quantitative study in the supragranular layers of the middle temporal neocortex. *Brain research*. 2008; 1213:152–65. Epub 2008/05/06. <https://doi.org/10.1016/j.brainres.2008.03.043> PMID: 18455153.
25. Hinton DR, Sadun AA, Blanks JC, Miller CA. Optic-nerve degeneration in Alzheimer's disease. *The New England journal of medicine*. 1986; 315(8):485–7. Epub 1986/08/21. <https://doi.org/10.1056/NEJM198608213150804> PMID: 3736630.

26. Mancino R, Martucci A, Cesareo M, Giannini C, Corasaniti MT, Bagetta G, et al. Glaucoma and Alzheimer Disease: One Age-Related Neurodegenerative Disease of the Brain. *Curr Neuropharmacol*. 2018; 16(7):971–7. Epub 2017/12/07. <https://doi.org/10.2174/1570159X16666171206144045> PMID: 29210654.
27. Rawlyk B, Chauhan BC. Retinal ganglion cell loss in postmortem tissue of Alzheimer disease, glaucoma, and healthy normal subjects. *Canadian journal of ophthalmology Journal canadien d'ophtalmologie*. 2021; 56(2):e66–e70. Epub 2020/10/21. <https://doi.org/10.1016/j.jcjo.2020.09.018> PMID: 33080180.
28. Inyushin M, Zayas-Santiago A, Rojas L, Kucheryavykh Y, Kucheryavykh L. Platelet-generated amyloid beta peptides in Alzheimer's disease and glaucoma. *Histol Histopathol*. 2019; 34(8):843–56. <https://doi.org/10.14670/HH-18-111> PMID: 30945258
29. Sadun AA, Bassi CJ. Optic nerve damage in Alzheimer's disease. *Ophthalmology*. 1990; 97(1):9–17. Epub 1990/01/01. [https://doi.org/10.1016/s0161-6420\(90\)32621-0](https://doi.org/10.1016/s0161-6420(90)32621-0) PMID: 2314849.
30. Braak H, Braak E. Staging of Alzheimer-related cortical destruction. *Int Psychogeriatr*. 1997; 9 Suppl 1:257–61; discussion 69–72. Epub 1997/01/01. PMID: 9447446.
31. Kosior-Jarecka E, Pankowska A, Polit P, Stepniewski A, Symms MR, Koziol P, et al. Volume of Lateral Geniculate Nucleus in Patients with Glaucoma in 7Tesla MRI. *J Clin Med*. 2020; 9(8). Epub 2020/07/30. <https://doi.org/10.3390/jcm9082382> PMID: 32722571.
32. Lee JY, Jeong HJ, Lee JH, Kim YJ, Kim EY, Kim YY, et al. An investigation of lateral geniculate nucleus volume in patients with primary open-angle glaucoma using 7 tesla magnetic resonance imaging. *Investigative ophthalmology & visual science*. 2014; 55(6):3468–76. Epub 2014/04/12. <https://doi.org/10.1167/iov.14-13902> PMID: 24722700.
33. Schmidt MA, Knott M, Heidemann R, Michelson G, Kober T, Dorfler A, et al. Investigation of lateral geniculate nucleus volume and diffusion tensor imaging in patients with normal tension glaucoma using 7 tesla magnetic resonance imaging. *PloS one*. 2018; 13(6):e0198830. Epub 2018/06/08. <https://doi.org/10.1371/journal.pone.0198830> PMID: 29879191 adherence to PLOS ONE policies on sharing data and materials. Siemens Healthcare AG did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.
34. Russo R, Varano GP, Adornetto A, Nucci C, Corasaniti MT, Bagetta G, et al. Retinal ganglion cell death in glaucoma: Exploring the role of neuroinflammation. *Eur J Pharmacol*. 2016; 787:134–42. Epub 2016/04/06. <https://doi.org/10.1016/j.ejphar.2016.03.064> PMID: 27044433.
35. Blanks JC, Schmidt SY, Torigoe Y, Porrello KV, Hinton DR, Blanks RH. Retinal pathology in Alzheimer's disease. II. Regional neuron loss and glial changes in GCL. *Neurobiology of aging*. 1996; 17(3):385–95. Epub 1996/05/01. [https://doi.org/10.1016/0197-4580\(96\)00009-7](https://doi.org/10.1016/0197-4580(96)00009-7) PMID: 8725900.
36. Blanks JC, Torigoe Y, Hinton DR, Blanks RH. Retinal pathology in Alzheimer's disease. I. Ganglion cell loss in foveal/parafoveal retina. *Neurobiology of aging*. 1996; 17(3):377–84. Epub 1996/05/01. [https://doi.org/10.1016/0197-4580\(96\)00010-3](https://doi.org/10.1016/0197-4580(96)00010-3) PMID: 8725899.
37. Schrems WA, Schrems-Hoesl LM, Mardin CY, Laemmer R, Kruse FE, Horn FK. Can Glaucomatous Visual Field Progression be Predicted by Structural and Functional Measures? *Journal of glaucoma*. 2017; 26(4):373–82. Epub 2017/01/25. <https://doi.org/10.1097/JGJ.0000000000000628> PMID: 28118204.
38. Sehi M, Bhardwaj N, Chung YS, Greenfield DS, Advanced Imaging for Glaucoma Study G. Evaluation of baseline structural factors for predicting glaucomatous visual-field progression using optical coherence tomography, scanning laser polarimetry and confocal scanning laser ophthalmoscopy. *Eye*. 2012; 26(12):1527–35. Epub 2012/10/13. <https://doi.org/10.1038/eye.2012.203> PMID: 23060026.
39. Wallukat G, Pruss H, Muller J, Schimke I. Functional autoantibodies in patients with different forms of dementia. *PloS one*. 2018; 13(3):e0192778. Epub 2018/03/15. <https://doi.org/10.1371/journal.pone.0192778> PMID: 29538413.
40. Jonas JB, Gusek GC, Naumann GO. Optic disc morphometry in chronic primary open-angle glaucoma. I. Morphometric intrapapillary characteristics. *Graefes's archive for clinical and experimental ophthalmology = Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 1988; 226(6):522–30. <https://doi.org/10.1007/BF02169199> PMID: 3209079
41. Hohberger B, Kunze R, Wallukat G, Kara K, Mardin CY, Lammer R, et al. Autoantibodies Activating the beta2-Adrenergic Receptor Characterize Patients With Primary and Secondary Glaucoma. *Frontiers in immunology*. 2019; 10:2112. Epub 2019/10/22. <https://doi.org/10.3389/fimmu.2019.02112> PMID: 31632387.
42. Blindness GBD, Vision Impairment C, Vision Loss Expert Group of the Global Burden of Disease S. Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight: an analysis for the Global Burden of

- Disease Study. *Lancet Glob Health*. 2021; 9(2):e144–e60. Epub 2020/12/05. [https://doi.org/10.1016/S2214-109X\(20\)30489-7](https://doi.org/10.1016/S2214-109X(20)30489-7) PMID: 33275949.
43. Ramirez AI, de Hoz R, Salobar-Garcia E, Salazar JJ, Rojas B, Ajoy D, et al. The Role of Microglia in Retinal Neurodegeneration: Alzheimer's Disease, Parkinson, and Glaucoma. *Front Aging Neurosci*. 2017; 9:214. Epub 2017/07/22. <https://doi.org/10.3389/fnagi.2017.00214> PMID: 28729832.
 44. Criscuolo C, Fabiani C, Cerri E, Domenici L. Synaptic Dysfunction in Alzheimer's Disease and Glaucoma: From Common Degenerative Mechanisms Toward Neuroprotection. *Front Cell Neurosci*. 2017; 11:53. Epub 2017/03/16. <https://doi.org/10.3389/fncel.2017.00053> PMID: 28289378.
 45. Wirths O, Breyhan H, Cynis H, Schilling S, Demuth HU, Bayer TA. Intraneuronal pyroglutamate-Abeta 3–42 triggers neurodegeneration and lethal neurological deficits in a transgenic mouse model. *Acta Neuropathol*. 2009; 118(4):487–96. Epub 2009/06/24. <https://doi.org/10.1007/s00401-009-0557-5> PMID: 19547991.
 46. Tekirian TL, Yang AY, Glabe C, Geddes JW. Toxicity of pyroglutaminated amyloid beta-peptides 3(pE)-40 and -42 is similar to that of A beta1-40 and -42. *Journal of neurochemistry*. 1999; 73(4):1584–9. Epub 1999/09/29. <https://doi.org/10.1046/j.1471-4159.1999.0731584.x> PMID: 10501204.
 47. Ishimaru H, Ishikawa K, Haga S, Shoji M, Ohe Y, Haga C, et al. Accumulation of apolipoprotein E and beta-amyloid-like protein in a trace of the hippocampal CA1 pyramidal cell layer after ischaemic delayed neuronal death. *Neuroreport*. 1996; 7(18):3063–7. Epub 1996/11/25. <https://doi.org/10.1097/00001756-199611250-00054> PMID: 9116241.
 48. Armstrong RA. Plaques and tangles and the pathogenesis of Alzheimer's disease. *Folia Neuropathol*. 2006; 44(1):1–11. Epub 2006/03/28. PMID: 16565925.
 49. Zhang Y, Bander ED, Lee Y, Muoser C, Schaffer CB, Nishimura N. Microvessel occlusions alter amyloid-beta plaque morphology in a mouse model of Alzheimer's disease. *J Cereb Blood Flow Metab*. 2020; 40(10):2115–31. Epub 2019/11/21. <https://doi.org/10.1177/0271678X19889092> PMID: 31744388.
 50. Hohberger B, Lucio M, Schlick S, Wollborn A, Hosari S, Mardin C. OCT-angiography: Regional reduced macula microcirculation in ocular hypertensive and pre-perimetric glaucoma patients. *PloS one*. 2021; 16(2):e0246469. Epub 2021/02/12. <https://doi.org/10.1371/journal.pone.0246469> PMID: 33571215.
 51. Grieshaber MC, Mozaffarieh M, Flammer J. What is the link between vascular dysregulation and glaucoma? *Survey of ophthalmology*. 2007; 52 Suppl 2:S144–54. Epub 2007/12/06. <https://doi.org/10.1016/j.survophthal.2007.08.010> PMID: 17998040.
 52. Wey S, Amanullah S, Spaeth GL, Ustaoglu M, Rahmatnejad K, Katz LJ. Is primary open-angle glaucoma an ocular manifestation of systemic disease? Graefe's archive for clinical and experimental ophthalmology = *Albrecht von Graefes Archiv fur klinische und experimentelle Ophthalmologie*. 2019; 257(4):665–73. Epub 2019/01/16. <https://doi.org/10.1007/s00417-019-04239-9> PMID: 30643967.
 53. Salobar-Garcia E, Mendez-Hernandez C, Hoz R, Ramirez AI, Lopez-Cuenca I, Fernandez-Albarral JA, et al. Ocular Vascular Changes in Mild Alzheimer's Disease Patients: Foveal Avascular Zone, Choroidal Thickness, and ONH Hemoglobin Analysis. *J Pers Med*. 2020; 10(4). Epub 2020/11/19. <https://doi.org/10.3390/jpm10040231> PMID: 33203157.
 54. Salobar-Garcia E, Lopez-Cuenca I, Sanchez-Puebla L, de Hoz R, Fernandez-Albarral JA, Ramirez AI, et al. Retinal Thickness Changes Over Time in a Murine AD Model APP (NL-F/NL-F). *Front Aging Neurosci*. 2020; 12:625642. Epub 2021/02/06. <https://doi.org/10.3389/fnagi.2020.625642> PMID: 33542683.
 55. Salobar-Garcia E, de Hoz R, Ramirez AI, Lopez-Cuenca I, Rojas P, Vazirani R, et al. Changes in visual function and retinal structure in the progression of Alzheimer's disease. *PloS one*. 2019; 14(8):e0220535. Epub 2019/08/16. <https://doi.org/10.1371/journal.pone.0220535> PMID: 31415594.
 56. Hohberger B, Worn M, Lammer R, Mahajan A, Mardin C, Schotzer-Schrehardt U, et al. Agonistic beta2-Adrenergic Receptor Autoantibodies Characterize the Aqueous Humor of Patients With Primary and Secondary Open-Angle Glaucoma. *Frontiers in immunology*. 2021; 12:550236. Epub 2021/05/25. <https://doi.org/10.3389/fimmu.2021.550236> PMID: 34025634.
 57. Wallukat G, Morwinski M, Kowal K, Forster A, Boewer V, Wollenberger A. Autoantibodies against the beta-adrenergic receptor in human myocarditis and dilated cardiomyopathy: beta-adrenergic agonism without desensitization. *European heart journal*. 1991; 12 Suppl D:178–81. https://doi.org/10.1093/eurheartj/12.suppl_d.178 PMID: 1717272.
 58. Mijares A, Lebesgue D, Wallukat G, Hoebeke J. From agonist to antagonist: Fab fragments of an agonist-like monoclonal anti-beta(2)-adrenoceptor antibody behave as antagonists. *Mol Pharmacol*. 2000; 58(2):373–9. Epub 2000/07/25. <https://doi.org/10.1124/mol.58.2.373> PMID: 10908305.
 59. Wallukat G, Schimke I. Agonistic autoantibodies directed against G-protein-coupled receptors and their relationship to cardiovascular diseases. *Semin Immunopathol*. 2014; 36(3):351–63. Epub 2014/04/30. <https://doi.org/10.1007/s00281-014-0425-9> PMID: 24777744.

60. Richter R.M BM, Schulze W., Wallukat G. Impact of β 2- activation on the processing of amyloid- β peptides in the rat hippocampus. *Neuroscience Meeting Planner (San Diego)* 2010.
61. Haase N, Herse F, Spallek B, Haase H, Morano I, Qadri F, et al. Amyloid-beta peptides activate alpha1-adrenergic cardiovascular receptors. *Hypertension*. 2013; 62(5):966–72. Epub 2013/09/05. <https://doi.org/10.1161/HYPERTENSIONAHA.113.01348> PMID: 24001898.
62. Yan L, Xu Y, Yao H, Xue W, Tian J, Ren H, et al. The effects of autoantibodies against the second extracellular loop of alpha(1)-adrenoceptor on vasoconstriction. *Basic Res Cardiol*. 2009; 104(5):581–9. Epub 2009/03/31. <https://doi.org/10.1007/s00395-009-0020-8> PMID: 19330480.
63. Tohda C, Matsumoto N, Zou K, Meselhy MR, Komatsu K. A beta(25–35)-induced memory impairment, axonal atrophy, and synaptic loss are ameliorated by M1, a metabolite of protopanaxadiol-type saponins. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*. 2004; 29(5):860–8. <https://doi.org/10.1038/sj.npp.1300388> PMID: 15010693
64. Wang D, Yuen EY, Zhou Y, Yan Z, Xiang YK. Amyloid beta peptide-(1–42) induces internalization and degradation of beta2 adrenergic receptors in prefrontal cortical neurons. *The Journal of biological chemistry*. 2011; 286(36):31852–63. Epub 2011/07/16. <https://doi.org/10.1074/jbc.M111.244335> PMID: 21757762.
65. Wang D, Govindaiah G, Liu R, De Arcangelis V, Cox CL, Xiang YK. Binding of amyloid beta peptide to beta2 adrenergic receptor induces PKA-dependent AMPA receptor hyperactivity. *FASEB J*. 2010; 24(9):3511–21. Epub 2010/04/17. <https://doi.org/10.1096/fj.10-156661> PMID: 20395454.
66. Wallukat G, Mattheck S, Wenzel K, Schrod W, Vogt B, Brunner P, et al. C-Reactive Protein (CRP) Blocks the Desensitization of Agonistic Stimulated G Protein Coupled Receptors (GPCRs) in Neonatal Rat Cardiomyocytes. *J Clin Med*. 2022; 11(4). Epub 2022/02/26. <https://doi.org/10.3390/jcm11041058> PMID: 35207331.
67. Bacova BS, Radosinska J, Wallukat G, Barancik M, Wallukat A, Knezl V, et al. Suppression of beta1-Adrenoceptor Autoantibodies is Involved in the Antiarrhythmic Effects of Omega-3 Fatty Acids in Male and Female Hypertensive Rats. *Int J Mol Sci*. 2020; 21(2). Epub 2020/01/18. <https://doi.org/10.3390/ijms21020526> PMID: 31947691.