

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(2E)-1-(4-Chlorophenyl)-3-[4-(propan-2-yl)phenyl]prop-2-en-1-oneBadiadka Narayana,^a Vinutha V. Salian,^a Balladka K. Sarojini^b and Jerry P. Jasinski^{c*}^aDepartment of Studies in Chemistry, Mangalore University, Mangalagangothri 574 199, India, ^bDepartment of Studies in Chemistry, Industrial Chemistry Section, Mangalore University, Mangalagangothri 574 199, India, and ^cDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA
Correspondence e-mail: jjasinski@keene.edu

Received 26 May 2014; accepted 30 June 2014


Edited by G. Smith, Queensland University of Technology, Australia

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.093; wR factor = 0.286; data-to-parameter ratio = 15.7.

In the title compound, $\text{C}_{18}\text{H}_{17}\text{ClO}$, the dihedral angle between the benzene rings is $53.5(1)^\circ$. The mean plane of the prop-2-en-1-one group is twisted by $24.5(8)$ and $33.5(3)^\circ$ from the chloro- and propanyl-substituted rings, respectively.

Related literature

For the non-linear optical properties of the chalcones, see: Sarojini *et al.* (2006); Poornesh *et al.* (2009) and for their biological activity, see: Nielsen *et al.* (1998); Mai *et al.* (2014); Insuasty *et al.* (2013). For related structures, see: Jasinski *et al.* (2009, 2012); Butcher *et al.* (2007); Harrison *et al.* (2006). For standard bond lengths, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{18}\text{H}_{17}\text{ClO}$
 $M_r = 284.76$
Monoclinic, $P2_1/c$
 $a = 8.8547(5)$ Å
 $b = 5.8455(3)$ Å
 $c = 28.8034(17)$ Å
 $\beta = 97.396(6)^\circ$ $V = 1478.46(14)$ Å³
 $Z = 4$
Cu $K\alpha$ radiation
 $\mu = 2.21$ mm⁻¹
 $T = 173$ K
 $0.41 \times 0.32 \times 0.14$ mm

Data collection

Agilent Eos Gemini diffractometer
Absorption correction: multi-scan
CrysAlis PRO and *CrysAlis RED*
(Agilent, 2012)
 $T_{\min} = 0.370$, $T_{\max} = 1.000$
8687 measured reflections
2868 independent reflections
2269 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.093$
 $wR(F^2) = 0.286$
 $S = 1.04$
2868 reflections
183 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.87$ e Å⁻³
 $\Delta\rho_{\min} = -0.44$ e Å⁻³

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis RED* (Agilent, 2012); program(s) used to solve structure: *SUPERFLIP* (Palatinus *et al.*, 2012); program(s) used to refine structure: *SHELXL2012* (Sheldrick, 2008); molecular graphics: *OLEX2* (Dolomanov *et al.*, 2009); software used to prepare material for publication: *OLEX2*.

BN thanks the UGC for financial assistance through a BSR one-time grant for the purchase of chemicals. VVS thanks the DST for financial assistance through a PURSE grant. JPJ acknowledges the NSF-MRI program (grant No. CHE-1039027) for funds to purchase the X-ray diffractometer.

Supporting information for this paper is available from the IUCr electronic archives (Reference: ZS2302).

References

- Agilent (2012). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies, Yarnton, England.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Butcher, R. J., Yathirajan, H. S., Narayana, B., Mithun, A. & Sarojini, B. K. (2007). *Acta Cryst. E* **63**, o30–o32.
- Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
- Harrison, W. T. A., Yathirajan, H. S., Sarojini, B. K., Narayana, B. & Vijaya Raj, K. K. (2006). *Acta Cryst. E* **62**, o1578–o1579.
- Insuasty, B., Montoya, A., Becerra, D., Quiroga, J., Abonia, R., Robledo, S., Velez, I. D., Upegui, Y., Noguera, M. & Cobo, J. (2013). *Eur. J. Med. Chem.* **67**, 252–262.
- Jasinski, J. P., Butcher, R. J., Veena, K., Narayana, B. & Yathirajan, H. S. (2009). *Acta Cryst. E* **65**, o1965–o1966.
- Jasinski, J. P., Golen, J. A., Nayak, P. S., Narayana, B. & Yathirajan, H. S. (2012). *Acta Cryst. E* **68**, o366.
- Mai, C. W., Yaeghoobi, M., Abd-Rahman, N., Kang, Y. B. & Pichika, M. R. (2014). *Eur. J. Med. Chem.* **77**, 378–387.
- Nielsen, S. F., Christensen, S. B., Cruciani, G., Kharazmi, A. & Liljefors, T. (1998). *J. Med. Chem.* **41**, 4819–4832.
- Palatinus, L., Prathapa, S. J. & van Smaalen, S. (2012). *J. Appl. Cryst.* **45**, 575–580.
- Poornesh, P., Shettigar, S., Umesh, G., Manjunatha, K. B., Kamath, K. P., Sarojini, B. K. & Narayana, B. (2009). *Opt. Mater.* **31**, 854–859.
- Sarojini, B. K., Narayana, B., Ashalatha, B. V., Indira, J. & Lobo, K. G. (2006). *J. Cryst. Growth*, **295**, 54–59.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2014). E70, o855 [doi:10.1107/S1600536814015281]

(2E)-1-(4-Chlorophenyl)-3-[4-(propan-2-yl)phenyl]prop-2-en-1-one**Badiadka Narayana, Vinutha V. Salian, Balladka K. Sarojini and Jerry P. Jasinski****1. Comment**

Chalcones are an important class of natural compounds and have been widely applied as synthons in synthetic organic chemistry. The nonlinear optical properties of the different chalcone derivatives have been reported (Sarojini *et al.*, 2006; Poornesh *et al.*, 2009). These α,β -unsaturated ketones also possess a wide variety of biological activities, including anti-leishmanial (Nielsen *et al.*, 1998), anticancer (Mai *et al.*, 2014) and antitumor activity (Insuasty *et al.*, 2013). The crystal structures of some chalcone derivatives viz., a second polymorph of (2E)-1-(4-fluorophenyl)-3-(3, 4, 5-trimethoxyphenyl)prop-2-en-1-one, (2E)-1-(3,4-dichlorophenyl)-3-(2-hydroxyphenyl)prop-2-en-1-one (Jasinski *et al.*, 2009, 2012), (2E)-1-(2,4-dichlorophenyl)-3-[4-(methylsulfonyl)phenyl]prop-2-en-1-one (Butcher *et al.*, 2007) and 2-bromo-1-chlorophenyl-3-(4-methoxyphenyl)prop-2-en-1-one (Harrison *et al.*, 2006) have been reported. In view of the importance of chalcone derivatives, we report herein the crystal structure of the title compound, C₁₈H₁₇ClO.

In the title compound, the dihedral angle between the mean planes of the phenyl rings is 53.5 (1)°. The mean plane of the prop-2-en-1-one group (C1/C2/O1/C8) is twisted away from the two phenyl rings by 24.5 (8)° (C2–C7) and 33.5 (3)° (C10–C15) (Fig. 1). Bond lengths are in normal ranges (Allen *et al.*, 1987). No classical hydrogen bonds are observed.

2. Experimental

To a mixture of cuminaldehyde (1.5 mL, 0.01 mol) and 4-chloroacetophenone (1.3 mL, 0.01 mol) in ethanol (50 mL), 15 mL of 10 % sodium hydroxide solution was added and stirred at 273–278 K for 3 h (Fig. 2). The precipitate formed was collected by filtration. Single crystals were grown from ethanol by slow the evaporation method (m.p.: 343–345 K).

3. Refinement

All of the H atoms were placed in their calculated positions and then refined using the riding model with atom—H bond lengths of 0.95–1.00 Å or 0.98 Å (CH₃). Isotropic displacement parameters for these atoms were set to 1.2 (CH) or 1.5 (CH₃) times U_{eq} of the parent atom. The Me group was refined as an ideally rotating group. No twinning has been observed.




Figure 1

ORTEP drawing of $C_{18}H_{17}ClO$, showing the atom labeling scheme, with 30% probability displacement ellipsoids.




Figure 2

Synthesis of $C_{18}H_{17}ClO$.

(2E)-1-(4-Chlorophenyl)-3-[4-(propan-2-yl)phenyl]prop-2-en-1-one

Crystal data

$C_{18}H_{17}ClO$
 $M_r = 284.76$
 Monoclinic, $P2_1/c$
 $a = 8.8547$ (5) Å
 $b = 5.8455$ (3) Å
 $c = 28.8034$ (17) Å
 $\beta = 97.396$ (6)°
 $V = 1478.46$ (14) Å³
 $Z = 4$
 $F(000) = 600$

$D_x = 1.279$ Mg m⁻³
 Melting point = 343–345 K
 Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å
 Cell parameters from 2529 reflections
 $\theta = 4.6$ – 72.0 °
 $\mu = 2.21$ mm⁻¹
 $T = 173$ K
 Prism, colourless
 $0.41 \times 0.32 \times 0.14$ mm

Data collection

Agilent Eos Gemini
 diffractometer
 Radiation source: Enhance (Cu) X-ray Source
 Graphite monochromator
 Detector resolution: 16.0416 pixels mm⁻¹
 ω scans
 Absorption correction: multi-scan
CrysAlis PRO and *CrysAlis RED* (Agilent,
 2012)

$T_{\min} = 0.370$, $T_{\max} = 1.000$
 8687 measured reflections
 2868 independent reflections
 2269 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\max} = 72.1$ °, $\theta_{\min} = 5.0$ °
 $h = -10 \rightarrow 10$
 $k = -7 \rightarrow 6$
 $l = -35 \rightarrow 29$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.093$
 $wR(F^2) = 0.286$
 $S = 1.04$

2868 reflections
 183 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.1595P)^2 + 1.6733P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.87 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	-0.25150 (13)	0.8390 (2)	0.33097 (5)	0.0819 (5)
O1	0.2335 (4)	0.2755 (5)	0.48768 (11)	0.0695 (8)
C1	0.2184 (5)	0.4826 (7)	0.48178 (13)	0.0576 (10)
C2	0.1070 (4)	0.5752 (6)	0.44409 (14)	0.0550 (9)
C3	0.0602 (5)	0.4451 (7)	0.40431 (15)	0.0625 (10)
H3	0.1034	0.2981	0.4012	0.075*
C4	-0.0468 (5)	0.5249 (8)	0.36970 (16)	0.0675 (11)
H4	-0.0755	0.4358	0.3424	0.081*
C5	-0.1133 (5)	0.7368 (8)	0.37462 (16)	0.0638 (11)
C6	-0.0724 (5)	0.8693 (8)	0.41294 (17)	0.0691 (12)
H6	-0.1192	1.0137	0.4161	0.083*
C7	0.0386 (5)	0.7896 (7)	0.44713 (16)	0.0677 (12)
H7	0.0696	0.8833	0.4736	0.081*
C8	0.3018 (5)	0.6508 (8)	0.51427 (16)	0.0671 (11)
H8	0.3047	0.8073	0.5055	0.080*
C9	0.3719 (5)	0.5833 (8)	0.55524 (17)	0.0676 (11)
H9	0.3598	0.4270	0.5630	0.081*
C10	0.4664 (5)	0.7227 (8)	0.59013 (16)	0.0652 (11)
C11	0.5252 (5)	0.9341 (8)	0.57874 (15)	0.0676 (11)
H11	0.5026	0.9925	0.5478	0.081*
C12	0.6155 (5)	1.0579 (7)	0.61201 (14)	0.0612 (10)
H12	0.6546	1.2016	0.6038	0.073*
C13	0.6505 (4)	0.9774 (6)	0.65722 (13)	0.0529 (9)
C14	0.5907 (5)	0.7645 (7)	0.66783 (15)	0.0613 (10)
H14	0.6116	0.7055	0.6987	0.074*
C15	0.5034 (5)	0.6414 (7)	0.63453 (17)	0.0694 (12)
H15	0.4671	0.4953	0.6424	0.083*
C16	0.7492 (5)	1.1144 (7)	0.69393 (15)	0.0614 (10)
H16	0.7841	1.2548	0.6786	0.074*
C17	0.6624 (6)	1.1896 (8)	0.73356 (16)	0.0706 (12)
H17A	0.5750	1.2834	0.7209	0.106*
H17B	0.7299	1.2794	0.7563	0.106*
H17C	0.6265	1.0545	0.7490	0.106*
C18	0.8911 (5)	0.9757 (10)	0.7134 (2)	0.0834 (14)

H18A	0.8599	0.8406	0.7299	0.125*
H18B	0.9583	1.0709	0.7351	0.125*
H18C	0.9454	0.9273	0.6875	0.125*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0684 (8)	0.0788 (8)	0.1002 (10)	−0.0040 (5)	0.0173 (6)	0.0148 (6)
O1	0.0799 (19)	0.0492 (16)	0.0811 (19)	−0.0082 (14)	0.0161 (15)	−0.0158 (14)
C1	0.064 (2)	0.053 (2)	0.061 (2)	−0.0177 (17)	0.0283 (18)	−0.0123 (17)
C2	0.059 (2)	0.0465 (19)	0.065 (2)	−0.0154 (16)	0.0298 (18)	−0.0121 (16)
C3	0.073 (3)	0.047 (2)	0.071 (2)	−0.0081 (18)	0.022 (2)	−0.0150 (18)
C4	0.081 (3)	0.055 (2)	0.068 (2)	−0.010 (2)	0.016 (2)	−0.0130 (19)
C5	0.059 (2)	0.058 (2)	0.079 (3)	−0.0136 (18)	0.029 (2)	−0.001 (2)
C6	0.066 (3)	0.053 (2)	0.092 (3)	−0.0027 (19)	0.027 (2)	−0.011 (2)
C7	0.077 (3)	0.053 (2)	0.078 (3)	−0.013 (2)	0.028 (2)	−0.022 (2)
C8	0.071 (3)	0.061 (2)	0.073 (3)	−0.011 (2)	0.022 (2)	−0.010 (2)
C9	0.067 (2)	0.059 (2)	0.081 (3)	−0.009 (2)	0.027 (2)	−0.013 (2)
C10	0.065 (2)	0.057 (2)	0.076 (3)	−0.0096 (19)	0.016 (2)	−0.0076 (19)
C11	0.076 (3)	0.064 (3)	0.065 (2)	−0.021 (2)	0.015 (2)	0.0024 (19)
C12	0.066 (2)	0.055 (2)	0.065 (2)	−0.0173 (18)	0.0215 (19)	0.0001 (18)
C13	0.0468 (18)	0.0501 (19)	0.066 (2)	−0.0054 (15)	0.0211 (16)	−0.0031 (16)
C14	0.060 (2)	0.053 (2)	0.071 (2)	−0.0070 (18)	0.0101 (18)	0.0066 (18)
C15	0.072 (3)	0.050 (2)	0.086 (3)	−0.0152 (19)	0.006 (2)	0.013 (2)
C16	0.060 (2)	0.057 (2)	0.069 (2)	−0.0092 (18)	0.0171 (19)	−0.0049 (18)
C17	0.079 (3)	0.066 (3)	0.069 (3)	0.002 (2)	0.020 (2)	−0.008 (2)
C18	0.053 (2)	0.092 (4)	0.105 (4)	−0.004 (2)	0.010 (2)	−0.021 (3)

Geometric parameters (Å, °)

C11—C5	1.743 (5)	C10—C15	1.364 (6)
O1—C1	1.228 (5)	C11—H11	0.9500
C1—C2	1.472 (6)	C11—C12	1.372 (6)
C1—C8	1.486 (6)	C12—H12	0.9500
C2—C3	1.393 (5)	C12—C13	1.382 (6)
C2—C7	1.400 (6)	C13—C14	1.401 (5)
C3—H3	0.9500	C13—C16	1.512 (5)
C3—C4	1.366 (6)	C14—H14	0.9500
C4—H4	0.9500	C14—C15	1.358 (6)
C4—C5	1.386 (6)	C15—H15	0.9500
C5—C6	1.359 (6)	C16—H16	1.0000
C6—H6	0.9500	C16—C17	1.521 (6)
C6—C7	1.379 (7)	C16—C18	1.538 (7)
C7—H7	0.9500	C17—H17A	0.9800
C8—H8	0.9500	C17—H17B	0.9800
C8—C9	1.321 (7)	C17—H17C	0.9800
C9—H9	0.9500	C18—H18A	0.9800
C9—C10	1.469 (6)	C18—H18B	0.9800

C10—C11	1.396 (6)	C18—H18C	0.9800
O1—C1—C2	121.0 (3)	C12—C11—H11	119.9
O1—C1—C8	121.9 (4)	C11—C12—H12	119.4
C2—C1—C8	116.9 (4)	C11—C12—C13	121.2 (4)
C3—C2—C1	120.4 (4)	C13—C12—H12	119.4
C3—C2—C7	117.0 (4)	C12—C13—C14	117.6 (4)
C7—C2—C1	122.5 (4)	C12—C13—C16	121.2 (3)
C2—C3—H3	119.4	C14—C13—C16	121.2 (4)
C4—C3—C2	121.3 (4)	C13—C14—H14	119.6
C4—C3—H3	119.4	C15—C14—C13	120.9 (4)
C3—C4—H4	120.2	C15—C14—H14	119.6
C3—C4—C5	119.6 (4)	C10—C15—H15	119.2
C5—C4—H4	120.2	C14—C15—C10	121.6 (4)
C4—C5—C11	120.0 (4)	C14—C15—H15	119.2
C6—C5—C11	118.7 (4)	C13—C16—H16	108.0
C6—C5—C4	121.3 (4)	C13—C16—C17	112.1 (3)
C5—C6—H6	120.7	C13—C16—C18	110.3 (4)
C5—C6—C7	118.6 (4)	C17—C16—H16	108.0
C7—C6—H6	120.7	C17—C16—C18	110.3 (4)
C2—C7—H7	118.9	C18—C16—H16	108.0
C6—C7—C2	122.1 (4)	C16—C17—H17A	109.5
C6—C7—H7	118.9	C16—C17—H17B	109.5
C1—C8—H8	119.9	C16—C17—H17C	109.5
C9—C8—C1	120.2 (4)	H17A—C17—H17B	109.5
C9—C8—H8	119.9	H17A—C17—H17C	109.5
C8—C9—H9	116.3	H17B—C17—H17C	109.5
C8—C9—C10	127.3 (4)	C16—C18—H18A	109.5
C10—C9—H9	116.3	C16—C18—H18B	109.5
C11—C10—C9	121.8 (4)	C16—C18—H18C	109.5
C15—C10—C9	119.7 (4)	H18A—C18—H18B	109.5
C15—C10—C11	118.5 (4)	H18A—C18—H18C	109.5
C10—C11—H11	119.9	H18B—C18—H18C	109.5
C12—C11—C10	120.3 (4)		
C11—C5—C6—C7	-179.1 (3)	C8—C9—C10—C11	-16.7 (7)
O1—C1—C2—C3	-25.1 (5)	C8—C9—C10—C15	165.8 (5)
O1—C1—C2—C7	152.0 (4)	C9—C10—C11—C12	-178.8 (4)
O1—C1—C8—C9	-13.1 (6)	C9—C10—C15—C14	179.8 (4)
C1—C2—C3—C4	177.8 (4)	C10—C11—C12—C13	0.1 (7)
C1—C2—C7—C6	-176.0 (4)	C11—C10—C15—C14	2.3 (7)
C1—C8—C9—C10	176.1 (4)	C11—C12—C13—C14	0.1 (6)
C2—C1—C8—C9	162.2 (4)	C11—C12—C13—C16	-179.6 (4)
C2—C3—C4—C5	-1.7 (6)	C12—C13—C14—C15	0.9 (6)
C3—C2—C7—C6	1.1 (6)	C12—C13—C16—C17	115.4 (4)
C3—C4—C5—C11	-179.3 (3)	C12—C13—C16—C18	-121.2 (4)
C3—C4—C5—C6	1.1 (6)	C13—C14—C15—C10	-2.1 (7)
C4—C5—C6—C7	0.5 (6)	C14—C13—C16—C17	-64.3 (5)

supporting information

C5—C6—C7—C2	-1.7 (6)	C14—C13—C16—C18	59.0 (5)
C7—C2—C3—C4	0.6 (6)	C15—C10—C11—C12	-1.3 (7)
C8—C1—C2—C3	159.6 (4)	C16—C13—C14—C15	-179.4 (4)
C8—C1—C2—C7	-23.4 (5)		
