

RESEARCH ARTICLE

Role of acid-sensing ion channels in hypoxia- and hypercapnia-induced ventilatory responses

Neil D. Detweiler, Kenneth G. Vigil, Thomas C. Resta, Benjimen R. Walker, Nikki L. Jernigan*

Vascular Physiology Group, Department of Cell Biology and Physiology, University of New Mexico Health Sciences Center, Albuquerque, NM, United States of America

* NJernigan@salud.unm.edu



OPEN ACCESS

Citation: Detweiler ND, Vigil KG, Resta TC, Walker BR, Jernigan NL (2018) Role of acid-sensing ion channels in hypoxia- and hypercapnia-induced ventilatory responses. *PLoS ONE* 13(2): e0192724. <https://doi.org/10.1371/journal.pone.0192724>

Editor: Yu Ru Kou, National Yang-Ming University, TAIWAN

Received: December 8, 2017

Accepted: January 29, 2018

Published: February 23, 2018

Copyright: © 2018 Detweiler et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data underlying this study have been uploaded to figshare and are accessible using the following link: https://figshare.com/articles/Role_of_Acid-Sensing_Ion_Channels_in_Hypoxia_and_Hypercapnia-Induced_1_Ventilatory_Responses_xlsx/5874081.

Funding: This work was supported by the National Heart, Lung, and Blood Institute, HL111084 to Dr. Nikki L. Jernigan (PI of grant), the National Heart, Lung, and Blood Institute, HL007736 to Dr. Thomas C. Resta (PI of grant), the National Institute of General Medical Sciences, GM088021 (training

Abstract

Previous reports indicate roles for acid-sensing ion channels (ASICs) in both peripheral and central chemoreception, but the contributions of ASICs to ventilatory drive in conscious, unrestrained animals remain largely unknown. We tested the hypotheses that ASICs contribute to hypoxic- and hypercapnic-ventilatory responses. Blood samples taken from conscious, unrestrained mice chronically instrumented with femoral artery catheters were used to assess arterial O₂, CO₂, and pH levels during exposure to inspired gas mixtures designed to cause isocapnic hypoxemia or hypercapnia. Whole-body plethysmography was used to monitor ventilatory parameters in conscious, unrestrained ASIC1, ASIC2, or ASIC3 knock-out (^{-/-}) and wild-type (WT) mice at baseline, during isocapnic hypoxemia and during hypercapnia. Hypercapnia increased respiratory frequency, tidal volume, and minute ventilation in all groups of mice, but there were no differences between ASIC1^{-/-}, ASIC2^{-/-}, or ASIC3^{-/-} and WT. Isocapnic hypoxemia also increased respiratory frequency, tidal volume, and minute ventilation in all groups of mice. Minute ventilation in ASIC2^{-/-} mice during isocapnic hypoxemia was significantly lower compared to WT, but there were no differences in the responses to isocapnic hypoxemia between ASIC1^{-/-} or ASIC3^{-/-} compared to WT. Surprisingly, these findings show that loss of individual ASIC subunits does not substantially alter hypercapnic or hypoxic ventilatory responses.

Introduction

Arterial O₂, CO₂, and pH (P_aO₂, P_aCO₂, and pH_a) homeostasis is maintained by reflex control of ventilation. Alterations in P_aO₂, P_aCO₂, and pH_a are detected by peripheral chemoreceptors located in type I glomus cells within the carotid and aortic bodies. P_aCO₂ homeostasis is additionally regulated by central chemoreceptors located on the ventral surface of the medulla as well as other brain regions [1]. Activation of carotid chemoreceptors in response to hypoxemia, hypercapnia, or acidosis leads to inhibition of K⁺ channels, depolarization of the chemoreceptor cells, activation of L-type Ca²⁺ channels, and release of excitatory neurotransmitters that subsequently stimulate ventilation and sympathetic activation [2–8]. Although this model

grant that supported Neil Detweiler), and the National Institute of General Medical Sciences, GM060201 (training grant that supported Kenneth Vigil). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

of carotid body chemoreception is generally accepted, there are several O₂- and CO₂/pH-sensitive ion channels and other proteins and their individual and integrated roles in chemoreception remain incompletely understood. The precise location of *central* chemoreception also remains uncertain, and it appears that several brain regions are involved [1]. A major mechanism by which central chemoreceptors detect CO₂ levels is thought to be the detection of secondary changes in cerebrospinal fluid pH [1]. Ion channels are generally the favored candidates for central chemoreception [9], but the identity of the specific channels involved remains unclear, with several candidates having been proposed [10–13]. One family of ion channels implicated in both peripheral and central chemoreception is that of the proton-gated, acid-sensing ion channels (ASICs).

ASICs are members of the degenerin/epithelial Na⁺ channel (DEG/ENaC) superfamily that form trimeric cation channels. ASIC genes (ASIC1-4) and their splice variants (ASIC1a, -1b, -2a, -2b, -3, and -4) form homo- or hetero-multimeric channels with different pH sensitivities. ASIC1 and 3 homomers are the most sensitive to acidic conditions with a half-maximal activation (pH₅₀) of around pH 6.5 [14,15]; whereas ASIC2 is least acid-sensitive (pH₅₀ ~ 4.9) [15]. ASIC1 and 3, and to a lesser extent ASIC2, are expressed in glomus cells and the transient acid-evoked depolarization of isolated glomus cells is consistent with the biophysical and pharmacological properties of ASICs [16]. ASICs are also widely expressed in the medulla where reductions in pH trigger ASIC-like currents and stimulation of phrenic nerve activity in anesthetized animals [17–19]. Based on these reports that suggest that ASICs play an important role in chemoreception, we hypothesized that ASICs contribute to hypercapnic/acidosis-induced ventilatory drive in conscious, unrestrained mice.

Recent studies from our laboratory also demonstrate that ASIC1, expressed in pulmonary arterial smooth muscle cells (PASMC), contributes to hypoxic pulmonary vasoconstriction (HPV) [20]. Although the precise mechanism remains unclear, HPV is generally thought to be an intrinsic response of O₂-sensing PASMC, supporting ventilation-perfusion matching for optimal gas exchange in the lung by diverting blood flow away from hypoxic regions of the lung. Our data showing reduced HPV in ASIC1 null mice [20] suggest ASICs are sensitive to changes in O₂. Therefore, we further hypothesized that ASICs additionally contribute to the *hypoxic* ventilatory response. To test our hypotheses, we examined ventilatory responses to isocapnic hypoxemia and hypercapnia in conscious, unrestrained, ASIC1, 2, or 3 global knockout (ASIC1^{-/-}, 2^{-/-}, or 3^{-/-}) and wild-type (WT) mice.

Materials and methods

Animals

All protocols used in this study were reviewed and approved by the Institutional Animal Care and Use Committee of the University of New Mexico School of Medicine (Protocol number: 16-200543-HSC) and abide by the National Institutes of Health guidelines for animal use. All surgeries (described below) were performed under isoflurane anesthesia and buprenorphine was used for post-operative analgesia. ASIC1 (B6.129-Asic1^{tm1Wsh/J}), ASIC2 (B6.129-Asic2^{tm1Wsh/J}), and ASIC3 (B6.129-Asic3^{tm1Wsh/J}) knockout (^{-/-}) mice (all from Jackson Laboratory, Bar Harbor, ME) were bred on a C57BL/6 background and compared to age-matched C57BL/6 wildtype (WT) controls. Disruption of the relevant ASIC was confirmed by PCR and agarose gel electrophoresis using a three-primer (ASIC1^{-/-} and ASIC2^{-/-}) or four-primer (ASIC3^{-/-}) system to detect both WT and disrupted alleles. The following primers were used for genotyping: ASIC1: 5' -CAT GTC ACC AAG CTC GAC GAG GTG-3' (WT forward primer), 5' -TGG ATG TGG AAT GTG TGC GA-3' (knockout forward primer), 5' -CCG CCT TGA GCGCA GGT TTA AAG G-3' (reverse primer); ASIC2: 5' -AGT CCT GCA

CGG TGG GAG CTT CTA-3' (reverse primer) 5'-GAA GAG GAA GGG AGC CAT GAT GAG-3' (WT forward primer), 5'-TGG ATG TGG AAT GTG TGC GA-3' (knockout forward primer); ASIC3: 5'-GAA CCT GGA AAA CAG AGG CAG GAA GGA T-3' (knockout reverse primer), 5'-CAG GGA GTA AGA TCT TAT GTA GCC TGG C-3' (knockout forward primer), 5'-TGG ATG TGG AAT GTG TGC GA-3' (WT reverse primer), 5'-CCC TGG GCA CCA GAG TTG AAG GTG TAG C-3' (WT forward primer). Males and females (~15 wk old) were used equally. Each group of knockout mice was paired with a separate, simultaneously run set of WT mice.

Femoral artery catheterization and blood gas measurement

To confirm that alterations in inspired gases were achieving the desired effect on blood gases, and that blood gases were similar between paired groups of WT and knockouts, mice were chronically instrumented with femoral artery catheters for arterial blood sampling to determine P_aO_2 , P_aCO_2 , and pH_a . The catheters were routed out through the top of the cage through spring tethers to enable blood sampling in conscious, unrestrained mice. Catheters consisted of PE-10 tubing with a stretch-tapered proximal end for insertion into the right femoral artery, and were filled with a solution of 0.9% saline containing 100 units/ml heparin. Catheterization was performed under isoflurane anesthesia (5% isoflurane for induction of anesthesia, ~2% for maintenance) and mice were given buprenorphine (0.05–0.1 mg/kg, s.c.) and enrofloxacin (15 mg/kg, s.c.) post-operatively for analgesia and protection from infection, respectively.

Blood gas measurements were performed 2 days after the implantation surgery. For these experiments, mice were placed in a polycarbonate chamber for exposure to different inspired gas mixtures. A Columbus Instruments PEGAS 4000MF gas mixer (Columbus, OH) was used to combine N_2 , O_2 , and CO_2 for the appropriate inspired gas mixtures, the compositions of which were confirmed using an OxiGraf O_2 Cap Oxygen [and CO_2] Analyzer (Sunnyvale, CA) to test samples taken from chamber inflow line. To measure blood gases, ~100 μ L of blood was allowed to flow directly from the femoral artery catheter into Abbott iStat handheld blood gas analyzer cartridges (EG6+ or G3+, Abbott Park, IL) while the mouse was conscious and unrestrained. Each measurement was taken after 5 minutes of exposure to the respective inspired gas mixture, with at least 20 minutes between subsequent exposures. The catheter was flushed with a volume of saline (containing 100 units/mL heparin) equivalent to 1.5 times the dead space of the catheter between each measurement. For each individual mouse, all blood gas measurements were performed on a single day.

Whole-body plethysmography

Whole-body plethysmography was used to assess respiratory frequency, tidal volume, and minute ventilation in conscious, unrestrained mice. First described by Drorbaugh and Fenn [21], this method utilizes a nearly sealed chamber in which the pressure transiently increases as inspired air expands due to the humidity and warmth of the lung compared to the surrounding environment. For this study we modified a method previously used for rats [22]. The plethysmography chamber consisted of a ~50 cubic centimeter transparent polycarbonate cylinder fitted with a Validyne DP45-16 differential pressure transducer and inlets for the introduction of new inspired gas mixtures. To take a measurement, the chamber was sealed with the exception of a small controlled leak through a 50 μ L glass syringe (Gastight #1705) with plunger removed, attached to one of the Luer-lock ports on the side of the chamber, which served as a physical high-pass filter to eliminate gradual changes in chamber pressure that go beyond the narrow range of the highly sensitive differential pressure transducer.

Temperature inside the chamber was monitored using an electronic probe (BAT-10, Physitemp, Clifton, NJ) and ranged from 21 to 25° C. Calibration air injections of 40 µl were used to enable calculation of tidal volume using the equation introduced by Drorbaugh and Fenn [21]:

$$V_T = \frac{P_T}{P_K} \times V_K \times \frac{T_R(P_B - P_C)}{T_R(P_B - P_C) - T_C(P_B - P_R)}$$

Where:

V_T = tidal volume

V_K = the volume of air injected into the animal chamber for calibration

P_T = the pressure deflection associated with each tidal volume

P_K = the pressure deflection associated with injection of the calibrating volume, V_K

T_R = body temperature, assumed to be 37°C for all animals

T_C = the air temperature in the chamber, which varied from 20–23°C

P_B = barometric pressure, 630 mmHg in Albuquerque, NM

P_R = vapor pressure of water at body temperature

P_C = vapor pressure of water in the chamber, derived from T_C assuming 100% humidity which was confirmed in pilot experiments (gas mixtures were humidified by bubbling through three consecutive flasks of water prior to entry into the plethysmography chamber)

Tidal volume and minute ventilation data were normalized to body mass. The plethysmography chamber was never closed from gas flow for measurement for more than 2 consecutive minutes, and the chamber was continuously flushed with fresh gas mixture for at least 2 minutes between measurements at a flow rate of 2 L/min. Mice were allowed up to 1 hour to achieve a resting state suitable for baseline measurements (lack of movement, grooming, or sniffing as observed through the transparent walls of the chamber). Mice that remained active after 1 hour were removed from the chamber and re-tested on another day. After a baseline measurement was taken, the chamber was flushed with 21% O₂, 6% CO₂, balance N₂ to achieve hypercapnia, or 7% O₂, 3.2% CO₂, balance N₂ to achieve isocapnic hypoxemia for 5 minutes before taking the subsequent measurement.

Statistics

All data are expressed as means ± SE. *n* values correspond to the numbers of animals per group. Statistical tests are specified in the figure legends or in the results section, and were made using GraphPad Prism 7.02 software (La Jolla, CA). *P* values of <0.05 were accepted as significant for all comparisons.

Results

Examining the roles of ASIC1, 2, and 3 in the hypercapnic ventilatory response

To investigate the roles of ASIC1, 2, and 3 in the hypercapnic ventilatory response, we first assessed changes in ventilation during exposure to increasing levels of inspired CO₂ (3.7, 6.0, and 9.8%) using whole-body plethysmography in conscious, unrestrained WT mice. The traces shown in Fig 1A represent the pressure difference between the inside and the outside (room pressure) of the chamber. Upward pressure deflections indicate increasing pressure in the animal chamber (inspiration). The amplitude of the pressure deflections corresponds to tidal volume. Respiratory frequency (Fig 1B, breaths/min) was determined by counting the number of pressure deflections per minute. Exact tidal volume normalized to body mass (Fig 1C, µL/breath/g) was calculated using chamber temperature and calibration air injection (see




Fig 1. Inhalation of 6% CO₂ produces reliable hypercapnia and elevation of ventilation. WT mice were exposed to a range of inspired CO₂ levels to determine optimal conditions for testing hypercapnic ventilatory responses. (A) Representative traces of whole-body plethysmography illustrate frequency and depth of breathing in conscious, unrestrained mice exposed to normal air at baseline (21% O₂, 0% CO₂, balance N₂), and to increasing levels of hypercapnia (3.7, 6.0, and 9.8% CO₂). Summary data shows (B) respiratory frequency (breaths/min), (C) tidal volume (µL/gram), and (D) minute ventilation (mL/min/gram) at each inspired CO₂ level. Arterial blood samples taken from WT mice at baseline (normal air) or during exposure to 6% CO₂ were analyzed for (E) P_aO₂, (F) P_aCO₂, and (G) pH_a. Values are means ± SE; n = 4–8 animals/group. *P < 0.05 vs. baseline (1-way ANOVA; Dunnett's post-hoc test (B) or two-tailed, paired Student's t-test (C-E)).

<https://doi.org/10.1371/journal.pone.0192724.g001>

Methods). Minute ventilation normalized to body mass (Fig 1D, mL/minute/gram) was determined by multiplying tidal volume by respiratory frequency.

First, we measured minute ventilation in mice exposed to 5 minutes of 3 levels of inspired CO₂ (3.7, 6.0, and 9.8%) to determine an appropriate level for testing hypercapnic ventilatory responses (Fig 1A–1D). In these pilot experiments, exposure to 3.7%, 6.0%, and 9.8% inspired CO₂ significantly elevated respiratory frequency (Fig 1B), but only exposure to 6.0% and 9.8%, and not 3.7% inspired CO₂ caused statistically significant elevations in tidal volume and minute ventilation (Fig 1C and 1D, respectively). We chose to use 6% CO₂ as the hypercapnic stimulus for the remaining experiments in this study because it stimulated a robust response

that was not statistically different from the response caused by 9.8% CO₂. Using arterial blood samples obtained via chronically implanted femoral artery catheters in conscious, unrestrained WT mice, we assessed P_aO₂, P_aCO₂, and pH_a at baseline (room air) and during exposure to 6% inspired CO₂. Although inspired O₂ was unchanged (21%) during exposure to 6% inspired CO₂, P_aO₂ was increased (Fig 1E) likely due to the resultant increase in alveolar ventilation. As expected, mice exhibited hypercapnia and acidosis indicated by increased P_aCO₂ and decreased pH_a during exposure to 6% inspired CO₂ (Fig 1F and 1G).

Next, we measured P_aO₂, P_aCO₂, and pH_a in ASIC1^{-/-}, ASIC2^{-/-}, ASIC3^{-/-}, and paired groups of WT mice. The purpose of this experiment was to determine if the stimuli for ventilation (P_aO₂, P_aCO₂, and pH_a) were the same between WT and knockout mice. Our results show no significant differences in P_aO₂, P_aCO₂, and pH_a between ASIC1^{-/-}, ASIC2^{-/-}, or ASIC3^{-/-} versus corresponding WT mice at baseline or during exposure to 6% inspired CO₂ (Table 1). To investigate the putative roles of ASIC1, ASIC2, and ASIC3 in the hypercapnic/acidotic ventilatory response, we exposed separate groups of ASIC1^{-/-}, ASIC2^{-/-}, and ASIC3^{-/-} mice to 6% inspired CO₂ and measured respiratory frequency, tidal volume, and minute ventilation using whole-body plethysmography as described above, and compared the responses to those of WT mice run in parallel for each experiment. Surprisingly, there were no significant differences between ASIC1^{-/-}, ASIC2^{-/-}, or ASIC3^{-/-} mice compared to WT mice in respiratory frequency, tidal volume, or minute ventilation (Fig 2). There was a tendency for ASIC2^{-/-} mice to exhibit lower respiratory frequency during hypercapnia (Fig 2D), but this trend was not statistically significant (p = 0.070).

Assessing the roles of ASIC1, 2, and 3 in the hypoxic ventilatory response

To specifically assess the roles of ASICs in the ventilatory response to hypoxia, we measured responses to isocapnic hypoxemia, which was achieved via inhalation of a hypoxic gas mixture

Table 1. Blood gas levels were not different between paired WT and knockout groups at baseline, during hypercapnia, or during isocapnic hypoxemia.

| Baseline (21% O ₂ , 0% CO ₂ , balance N ₂) | | | | | | |
|--|--------------|----------------------|--------------|----------------------|--------------|----------------------|
| | WT | ASIC1 ^{-/-} | WT | ASIC2 ^{-/-} | WT | ASIC3 ^{-/-} |
| P _a O ₂ | 63.2 ± 0.9 | 66.4 ± 0.8 | 58.6 ± 1.0 | 62.1 ± 2.2 | 64.9 ± 0.9 | 70.9 ± 2.7 |
| P _a CO ₂ | 28.0 ± 1.1 | 26.2 ± 0.5 | 28.4 ± 0.3 | 29.3 ± 1.0 | 27.4 ± 0.8 | 26.7 ± 1.2 |
| pH _a | 7.37 ± 0.01 | 7.39 ± 0.01 | 7.38 ± 0.01 | 7.39 ± 0.01 | 7.36 ± 0.01 | 7.34 ± 0.02 |
| Hypercapnia (21% O ₂ , 6.0% CO ₂ , balance N ₂) | | | | | | |
| | WT | ASIC1 ^{-/-} | WT | ASIC2 ^{-/-} | WT | ASIC3 ^{-/-} |
| P _a O ₂ | 100.2 ± 2.1* | 99.6 ± 2.1* | 90.3 ± 1.2* | 95.9 ± 2.9* | 100.1 ± 1.4* | 103.6 ± 3.5* |
| P _a CO ₂ | 45.9 ± 1.5* | 45.8 ± 0.8* | 44.8 ± 0.7* | 45.5 ± 0.9* | 47.9 ± 0.9* | 45.8 ± 0.9* |
| pH _a | 7.23 ± 0.02* | 7.22 ± 0.01* | 7.23 ± 0.01* | 7.24 ± 0.02* | 7.21 ± 0.01* | 7.20 ± 0.02* |
| Isocapnic hypoxemia (7.0% O ₂ , 3.2% CO ₂ , balance N ₂) | | | | | | |
| | WT | ASIC1 ^{-/-} | WT | ASIC2 ^{-/-} | WT | ASIC3 ^{-/-} |
| P _a O ₂ | 40.3 ± 1.5* | 40.4 ± 0.6* | 35.6 ± 1.2* | 38.5 ± 1.4* | 39.3 ± 0.8* | 42.7 ± 1.5* |
| P _a CO ₂ | 25.5 ± 0.8* | 25.1 ± 0.5 | 25.6 ± 0.4* | 26.1 ± 0.7* | 28.0 ± 0.6 | 27.3 ± 0.6 |
| pH _a | 7.39 ± 0.02 | 7.39 ± 0.00 | 7.40 ± 0.01 | 7.40 ± 0.02 | 7.35 ± 0.02 | 7.32 ± 0.2 |

Arterial blood gas levels were assessed in samples taken from conscious, unrestrained mice chronically instrumented with femoral artery catheters at baseline (room air) or during exposure to 6% inspired CO₂ or 7% O₂, 3.2% CO₂ to induce hypercapnia or isocapnic hypoxemia, respectively. No differences in P_aO₂, P_aCO₂, or pH_a between paired WT and knockout groups were detected. Values are means ± SE; n = 5–8 animals/group.

*P < 0.05 vs. baseline (2-way, repeated measures ANOVA; Sidak's post-hoc test).

<https://doi.org/10.1371/journal.pone.0192724.t001>




Fig 2. The hypercapnic ventilatory response does not require ASIC1, 2, or 3. Whole-body plethysmography was used to determine respiratory frequency (breaths/min; A, D, G), tidal volume (μL/breath/g; B, E, H), and minute ventilation (mL/min/g body wt; C, F, I) at baseline and during exposure to hypercapnia (6% CO₂) in WT, ASIC1^{-/-} (A-C), ASIC2^{-/-} (D-F), and ASIC3^{-/-} mice (G-I). Values are means ± SE; n = 8 (A-C), 7–8 (D-F), or 11–12 (G-I) animals/group. *P < 0.05 vs. baseline (2-way, repeated measures ANOVA; Sidak’s post-hoc test). Displayed P value corresponds to comparison between WT and KO (panel D).

<https://doi.org/10.1371/journal.pone.0192724.g002>

with a level of inspired CO₂ adequate to preserve baseline P_aCO₂ levels, allowing for the assessment of the hypoxic ventilatory response without confounding changes in P_aCO₂/pH_a-dependent ventilatory drive. To determine the level of inspired CO₂ required to achieve isocapnic hypoxemia, we measured baseline P_aCO₂ in WT animals, and then experimentally determined the level of inspired CO₂ necessary to maintain this P_aCO₂ in mice concurrently exposed to 7% inspired O₂. Baseline P_aCO₂ in WT mice was 29.8 ± 0.9 mmHg. Mice chronically instrumented with femoral artery catheters were exposed to 7% O₂ at a range of inspired CO₂ levels (2.5–3.5%) and the resulting P_aCO₂ was assessed (Fig 3A). Linear regression analysis of these data indicate that an inspired CO₂ of 3.2% produces isocapnic hypoxemia. Testing this exposure in a group of WT mice (n = 7) confirmed significant hypoxemia (Fig 3B) with no accompanying change in P_aCO₂ or pH_a (Fig 3C and 3D). Whole-body plethysmography measurements show that this exposure robustly increased respiratory frequency and tidal volume (Fig 3E). We used this stimulus (7% O₂, 3.2% CO₂, balance N₂) to compare hypoxic ventilatory responses in the following experiments.

Using ASIC1^{-/-}, ASIC2^{-/-}, and ASIC3^{-/-}, and WT mice chronically instrumented with femoral artery catheters, we measured P_aO₂, P_aCO₂, and pH_a to confirm that there were no differences in blood gases between respective knockout and WT mice at baseline or during




Fig 3. An inspired CO₂ of 3.2% causes isocapnic hypoxemia in mice concurrently exposed to 7.0% O₂. A scatter plot (A) showing arterial CO₂ tensions (P_aCO₂; y-axis) in mice exposed to 7.0% O₂ and a range of inspired CO₂ levels (% CO₂; x-axis) was fit using linear regression (solid line, equation displayed on the graph). The dotted lines on the graph indicate the average baseline P_aCO₂ (horizontal) measured during exposure to room air, and the estimated inspired CO₂ required to maintain this P_aCO₂ level (vertical) during exposure to 7% O₂ based on the linear regression. Arterial blood samples from WT mice exposed to 7.0% O₂, 3.2% CO₂, balance N₂, were analyzed to assess P_aO₂ (B), P_aCO₂ (C), and pH_a (D). (E) Representative whole-body plethysmography traces from a WT mouse show the ventilatory response to 7.0% O₂, 3.2% CO₂, balance N₂. Values are individual measurements (A) or means ± SE (B-D); n = 5–6 animals per group (B-D). *P < 0.05 vs. baseline (paired, two-tailed Student's t-test).

<https://doi.org/10.1371/journal.pone.0192724.g003>

exposure to 7% O₂, 3.2% CO₂, balance N₂ (Table 1), demonstrating that a similar hypoxic stimulus was achieved between groups. Although a slight but significant decrease in P_aCO₂ occurred in response to this inspired gas mixture compared to baseline in some groups (marked with asterisks, Table 1), in all cases there were no significant differences between paired WT and knockout groups. To test the roles of ASIC1, 2, and 3 in the hypoxic ventilatory response, separate groups of conscious, unrestrained ASIC1^{-/-}, ASIC2^{-/-}, ASIC3^{-/-}, and WT mice were exposed to 7% O₂, 3.2% CO₂, balance N₂ to induce isocapnic hypoxemia and their respiratory frequencies, tidal volumes, and minute ventilations were assessed using whole-body plethysmography (Fig 4). There were no differences in ventilatory parameters between




Fig 4. ASIC2, but not ASIC1 or ASIC3, contributes to the hypoxic ventilatory response. Ventilatory responses to isocapnic hypoxemia (7.0% O₂, 3.2% CO₂, bal N₂) were assessed using whole-body plethysmography and compared between (A-C) ASIC1^{-/-}, (D-F) ASIC2^{-/-}, and (G-I) ASIC3^{-/-} and WT mice. Values are means ± SE; n = 8 (A-F) or 11–12 (G-I) animals/group. *P < 0.05 vs. baseline; #P < 0.05 vs. WT (2-way, repeated measures ANOVA; Sidak's post-hoc test). Displayed P value corresponds to the comparison between WT and KO (panel D).

<https://doi.org/10.1371/journal.pone.0192724.g004>

ASIC1^{-/-} or ASIC3^{-/-} and WT mice (Fig 4A–4C and 4G–4I). ASIC2^{-/-} mice tended to have lower respiratory frequency and tidal volume than WT mice during exposure to isocapnic hypoxemia (Fig 4D and 4E). While these trends in respiratory frequency and tidal volume were not statistically significant, minute ventilation was significantly lower in ASIC2^{-/-} mice compared to WT (Fig 4F), suggesting a role for ASIC2 in the hypoxic ventilatory response.

Discussion

Previous evidence indicates that ASICs are involved in chemoreception and the control of ventilation [16–19,23,24], but most of this evidence comes from *in vitro* studies and little is known about the role of ASICs in hypercapnic and hypoxic ventilatory responses in conscious animals. Here we directly measured hypercapnic and hypoxic ventilatory responses in conscious, unrestrained mice. In contrast to our hypothesis, we found that genetic deletion of ASIC1, 2, or 3 does not alter the hypercapnic ventilatory response in mice; whereas ASIC2 appears to play a minor role in the hypoxic ventilatory response. Together, these data suggest the contribution of ASICs to ventilatory control is modest.

ASICs have been implicated in both peripheral and central CO₂ chemoreception. Isolated carotid body glomus type I cells exhibit acid (pH 6.0–6.8)-induced cation currents and

depolarization which have been attributed to ASICs [16]. Further investigation revealed that pH-sensitivity in isolated glomus cells is increased in both transgenic ASIC3 overexpressing mice and spontaneously hypertensive rats with increased expression of ASIC3, and decreased in ASIC3 null mice [23,24]. These studies suggest that ASIC3 would contribute to hypercapnic ventilatory drive. However, the pH levels used to stimulate a response in these *in vitro* experiments ranged from pH 6.0 to 6.8. In the present study, exposure to 6% inspired CO₂ caused a 70% increase in P_aCO₂, resulting in a pH_a of 7.2. This stimulus caused a robust ventilatory response (Figs 1 and 2), yet a pH_a of 7.2 may be insufficient to activate ASICs.

Other reports implicate ASICs in *central* chemoreception. The response of central chemoreceptors to changes in pH is known to be very sensitive, in that changes in cerebrospinal fluid pH from 7.30 to 7.25 cause a doubling in alveolar ventilation [25]. In neurons of the nucleus tractus solitarius in brainstem slices, acidification to pH 7.0 causes transient inward currents that are blocked by amiloride [18]. Although amiloride is a non-selective blocker of ASICs, Song et al., found that the selective ASIC1a blocker, psalmotoxin-1 (PcTX1) inhibited the increased phrenic nerve discharge in response to injection of acidified (pH 6.5) artificial cerebrospinal fluid in the lateral hypothalamus [17]. Furthermore, these authors report ASIC-like acid-induced currents in the ventrolateral medulla were absent in ASIC1^{-/-} mice [19]. These data provide strong evidence that ASIC1 contributes to central chemoreception in response to direct changes in pH (pH 6.0–7.0), however whether the observed effect is relevant to the physiological ventilatory response to hypercapnia is less clear. The threshold for activation of ASIC1 and ASIC3 is ~ pH 7.0–7.2 [26,27] and the half-maximal activation is 6.2–6.8 for ASIC1a, ~4.9 for ASIC2a, and 6.5–6.7 for ASIC3 [26,28]. Interestingly, however, Ziemann et al. reported that inhalation of higher CO₂ concentrations (10 and 20%) in anesthetized mice leads to a drop in pH ranging from 6.7–6.9 in the amygdala and lateral ventricle [29]. Consistent with our findings, the authors demonstrated an intact ventilatory response to CO₂ inhalation (5 and 10%) in ASIC1^{-/-} mice despite the important chemosensory role ASIC1 plays in the amygdala and fear-related behavior [29]. Acid-evoked ASIC responses in isolated cells tend to be transient, on the order of seconds, while the ventilatory response to hypercapnia is sustained indefinitely. This might explain why the previously reported contribution of ASIC3 to transient acid-induced currents in isolated glomus cells [16,24] does not translate to a contribution to the hypercapnic ventilatory response. However, the studies implicating ASICs in central chemoreception reported sustained changes to phrenic nerve discharge [17–19], suggesting that ASICs can mediate sustained responses to extracellular acidification in some contexts.

Thus, despite strong evidence for ASICs in chemoreception, our current findings do not support a role for ASICs in the hypercapnic ventilatory response in conscious, unrestrained mice. Although differences in ventilatory regulation may exist between species, mice are widely used in ventilatory function studies as they show dependency both on carotid bodies [30] as well as central chemoreception [1] for control of ventilation. Furthermore, mice were used to study the involvement of ASICs in carotid body glomus cells [24] and central chemoreception [19]. Therefore, it is unlikely that our current findings are due to species differences in control of ventilation. Inhalation of 6% CO₂ was chosen as the stimulus for hypercapnic ventilatory responses based on data presented in Fig 1 which indicates that 6% CO₂ causes a near-maximal ventilatory response. Although we cannot exclude the potential for ASICs to participate in hypercapnic ventilatory responses to higher levels of CO₂, this seems unlikely since the response of WT mice to 9.7% CO₂ was not different compared to the response to 6% CO₂. Additionally, our results agree with the findings of Ziemann et al. [29] which indicated that ASIC1 was not involved in ventilatory responses to 5% or 10% inspired CO₂.

Although much of the evidence supporting ASIC involvement in chemoreception focuses on the detection and ventilatory response to CO₂ or acidosis, evidence suggests ASICs may

also be hypoxia-sensitive. For example, we have previously reported that ASIC1 contributes to HPV in isolated-perfused lungs [20]. Additionally, activation of the Ca^{2+} -permeable ASIC1a contributes to neuronal cell death during ischemic brain injury [31,32]. Currently however, it is unclear whether the activation of ASICs responsible for neuronal cell death in this context results from the associated acidosis, or a direct effect of hypoxia. Liu et al reported that the hypoxia-induced chemoreceptor response and ASIC expression are both increased following exposure to *chronic* hypobaric hypoxia (380 Torr ambient air pressure for 1–7 days) in the rat petrosal ganglion, which contains chemoafferent neurons that innervate O_2 -sensitive carotid body glomus type I cells [33]. Pharmacological inhibition of ASICs using A-317576 (novel ASIC blocker [34]) or ibuprofen (non-specific ASIC inhibitor [35]) prevented the enhanced hypoxia-evoked chemoreceptor response following exposure of rats to chronic hypoxia, but did not alter the hypoxic chemoreceptor response in normoxia-exposed control animals [33]. In contrast, Lu et al. demonstrated that transgenic overexpression of ASIC3 leads to decreased glomus cell sensitivity to sodium cyanide (used to mimic hypoxia) [24]. Together, these data are consistent with the hypothesis that ASICs modulates hypoxic chemotransmission between type I cells and chemoafferent neurons. In the present study, we have differentiated between hypoxia and acidosis by utilizing *isocapnic* hypoxemia as a ventilatory stimulus that maintains baseline CO_2 and pH levels. Ventilatory responses to isocapnic hypoxemia were normal in ASIC1^{-/-} and ASIC3^{-/-} mice, but reduced in ASIC2^{-/-} mice (Fig 4F) suggesting a minor role for ASIC2 in this response. ASIC2a was reported to be expressed in carotid body glomus cells [16], and may contribute to the hypoxic ventilatory response by contributing to the depolarization of these cells during hypoxemia.

In summary, our data bring into question the physiological relevance of ASICs in chemoreception and the control of ventilation. We cannot rule out the possibility that compensatory upregulation of parallel chemoreception and ventilatory control pathways masks any effect of ASIC deletion in our global knockout mice. Furthermore, cerebral ischemia or other pathophysiological conditions might unmask a contribution of ASICs to hypercapnic and/or hypoxic ventilatory drive. Because ASICs are potential therapeutic targets for the treatment of pulmonary hypertension [20] and several neuronal diseases [36], a complete understanding of the roles ASICs play in O_2 and CO_2 homeostasis is important to the development of safe therapeutics.

Acknowledgments

We thank Anthony Gravagne in the Department of Physics and Astronomy at the University of New Mexico for his assistance with the design and fabrication of the plethysmography chamber. We also thank Lindsay Herbert for her assistance with knockout mouse colony maintenance.

Author Contributions

Conceptualization: Neil D. Detweiler, Thomas C. Resta, Benjimen R. Walker, Nikki L. Jernigan.

Data curation: Neil D. Detweiler, Kenneth G. Vigil.

Formal analysis: Neil D. Detweiler, Kenneth G. Vigil.

Funding acquisition: Thomas C. Resta, Nikki L. Jernigan.

Investigation: Neil D. Detweiler, Kenneth G. Vigil.

Methodology: Neil D. Detweiler.

Project administration: Neil D. Detweiler, Nikki L. Jernigan.

Resources: Thomas C. Resta, Benjimen R. Walker, Nikki L. Jernigan.

Software: Neil D. Detweiler.

Supervision: Neil D. Detweiler, Thomas C. Resta, Benjimen R. Walker, Nikki L. Jernigan.

Validation: Neil D. Detweiler.

Visualization: Neil D. Detweiler.

Writing – original draft: Neil D. Detweiler.

Writing – review & editing: Neil D. Detweiler, Kenneth G. Vigil, Thomas C. Resta, Benjimen R. Walker, Nikki L. Jernigan.

References

1. Nattie E, Li A. Central chemoreceptors: Locations and functions. *Compr Physiol*. 2012; 2(1):221–54. <https://doi.org/10.1002/cphy.c100083> PMID: 23728974
2. Buckler KJ. A novel oxygen-sensitive potassium current in rat carotid body type I cells. *J Physiol*. 1997 Feb 1; 498(3):649–62.
3. Delpiano MA, Hescheler J. Evidence for a PO₂-sensitive K⁺ channel in the type-I cell of the rabbit carotid body. *FEBS Lett*. 1989 Jun 5; 249(2):195–8. PMID: 2737279
4. Kim D, Cavanaugh EJ, Kim I, Carroll JL. Heteromeric TASK-1/TASK-3 is the major oxygen-sensitive background K⁺ channel in rat carotid body glomus cells. *J Physiol*. 2009 Jun 15; 587(12):2963–75.
5. López-López JR, Pérez-García MT. Oxygen sensitive Kv channels in the carotid body. *Respir Physiol Neurobiol*. 2007 Jul; 157(1):65–74. <https://doi.org/10.1016/j.resp.2007.01.022> PMID: 17442633
6. Urefña J, Fernández-Chacón R, Benot AR, Alvarez de Toledo GA, López-Barneo J. Hypoxia induces voltage-dependent Ca²⁺ entry and quantal dopamine secretion in carotid body glomus cells. *Proc Natl Acad Sci U S A*. 1994 Oct 11; 91(21):10208–11. PMID: 7937863
7. Ortega-Sáenz P, Levitsky KL, Marcos-Almaraz MT, Bonilla-Henao V, Pascual A, López-Barneo J. Carotid body chemosensory responses in mice deficient of TASK channels. *J Gen Physiol*. 2010 Apr; 135(4):379–92. <https://doi.org/10.1085/jgp.200910302> PMID: 20351062
8. Peers C. Hypoxic suppression of K⁺ currents in type I carotid body cells: Selective effect on the Ca²⁺-activated K⁺ current. *Neurosci Lett*. 1990 Nov; 119(2):253–6. PMID: 1704113
9. Guyenet PG, Stornetta RL, Bayliss DA. Central respiratory chemoreception. *J Comp Neurol*. 2010 May 20; 518(19):3883–906. <https://doi.org/10.1002/cne.22435> PMID: 20737591
10. Bayliss DA, Talley EM, Sirois JE, Lei Q. TASK-1 is a highly modulated pH-sensitive “leak” K⁺ channel expressed in brainstem respiratory neurons. *Respir Physiol*. 2001 Dec; 129(1–2):159–74. PMID: 11738652
11. Mulkey DK, Talley EM, Stornetta RL, Siegel AR, West GH, Chen X, et al. TASK Channels Determine pH Sensitivity in Select Respiratory Neurons But Do Not Contribute to Central Respiratory Chemosensitivity. *J Neurosci*. 2007 Dec 19; 27(51):14049–58. <https://doi.org/10.1523/JNEUROSCI.4254-07.2007> PMID: 18094244
12. Wang S, Benamer N, Zanella S, Kumar NN, Shi Y, Bevençut M, et al. TASK-2 Channels Contribute to pH Sensitivity of Retrotrapezoid Nucleus Chemoreceptor Neurons. *J Neurosci*. 2013 Oct 9; 33(41):16033–44. <https://doi.org/10.1523/JNEUROSCI.2451-13.2013> PMID: 24107938
13. Gestreau C, Heitzmann D, Thomas J, Dubreuil V, Bandulik S, Reichold M, et al. Task2 potassium channels set central respiratory CO₂ and O₂ sensitivity. *Proc Natl Acad Sci*. 2010 Feb 2; 107(5):2325–30. <https://doi.org/10.1073/pnas.0910059107> PMID: 20133877
14. Price MP, Lewin GR, McIlwrath SL, Cheng C, Xie J, Heppenstall PA, et al. The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature*. 2000 Oct 26; 407(6807):1007–11. <https://doi.org/10.1038/35039512> PMID: 11069180
15. Bartoi T, Augustinowski K, Polleichtner G, Grunder S, Ulbrich MH. Acid-sensing ion channel (ASIC) 1a/2a heteromers have a flexible 2:1/1:2 stoichiometry. *Proc Natl Acad Sci*. 2014 Jun 3; 111(22):8281–6. <https://doi.org/10.1073/pnas.1324060111> PMID: 24847067
16. Tan ZY, Lu Y, Whiteis CA, Benson CJ, Chappleau MW, Abboud FM. Acid-sensing ion channels contribute to transduction of extracellular acidosis in rat carotid body glomus cells. *Circ Res*. 2007; 101(10):1009–19. <https://doi.org/10.1161/CIRCRESAHA.107.154377> PMID: 17872465

17. Song N, Zhang G, Geng W, Liu Z, Jin W, Li L, et al. Acid sensing ion channel 1 in lateral hypothalamus contributes to breathing control. *PLoS One*. 2012; 7(7):1–10.
18. Huda R, Pollema-Mays SL, Chang Z, Alheid GF, McCrimmon DR, Martina M. Acid-sensing ion channels contribute to chemosensitivity of breathing-related neurons of the nucleus of the solitary tract. *J Physiol*. 2012; 590(Pt 19):4761–75.
19. Song N, Guan R, Jiang Q, Hassanzadeh CJ, Chu Y, Zhao X, et al. Acid-sensing ion channels are expressed in the ventrolateral medulla and contribute to central chemoreception. *Sci Rep*. 2016; 6 (December):38777. <https://doi.org/10.1038/srep38777> PMID: 27934921
20. Nitta CH, Osmond DA, Herbert LM, Beasley BF, Resta TC, Walker BR, et al. Role of ASIC1 in the development of chronic hypoxia-induced pulmonary hypertension. *AJP Hear Circ Physiol*. 2014 Jan 1; 306 (1):H41–52.
21. Drorbaugh JE, Fenn WO. A barometric method for measuring ventilation in newborn infants. *Pediatrics*. 1955 Jul; 16(1):81–7. PMID: 14394741
22. Walker BR, Adams EM, Voelkel NF. Ventilatory responses of hamsters and rats to hypoxia and hypercapnia. *J Appl Physiol*. 1985 Dec; 59(6):1955–60. <https://doi.org/10.1152/jappl.1985.59.6.1955> PMID: 4077803
23. Tan ZY, Lu Y, Whiteis CA, Simms AE, Paton JFR, Chappleau MW, et al. Chemoreceptor hypersensitivity, sympathetic excitation, and overexpression of ASIC and TASK channels before the onset of hypertension in SHR. *Circ Res*. 2010; 106(3):536–45. <https://doi.org/10.1161/CIRCRESAHA.109.206946> PMID: 20019330
24. Lu Y, Whiteis C a, Sluka K a, Chappleau MW, Abboud FM. Responses of glomus cells to hypoxia and acidosis are uncoupled, reciprocal and linked to ASIC3 expression: selectivity of chemosensory transduction. *J Physiol*. 2013; 591(Pt 4):919–32.
25. FencI V, Miller TB, Pappenheimer JR. Studies on the respiratory response to disturbances of acid-base balance, with deductions concerning the ionic composition of cerebral interstitial fluid. *Am J Physiol*. 1966; 210(3):459–72. <https://doi.org/10.1152/ajplegacy.1966.210.3.459> PMID: 5933194
26. Kress M, Waldmann R. Chapter 8 Acid Sensing Ionic Channels. In 2006. p. 241–76.
27. Wemmie JA, Price MP, Welsh MJ. Acid-sensing ion channels: advances, questions and therapeutic opportunities. *Trends Neurosci*. 2006 Oct; 29(10):578–86. <https://doi.org/10.1016/j.tins.2006.06.014> PMID: 16891000
28. Benson CJ, Xie J, Wemmie JA, Price MP, Henss JM, Welsh MJ, et al. Heteromultimers of DEG/ENaC subunits form H⁺-gated channels in mouse sensory neurons. *Proc Natl Acad Sci U S A*. 2002 Feb 19; 99(4):2338–43. <https://doi.org/10.1073/pnas.032678399> PMID: 11854527
29. Ziemann AE, Allen JE, Dahdaleh NS, Drebot II, Coryell MW, Wunsch AM, et al. The Amygdala Is a Chemosensor that Detects Carbon Dioxide and Acidosis to Elicit Fear Behavior. *Cell*. 2009; 139(5):1012–21. <https://doi.org/10.1016/j.cell.2009.10.029> PMID: 19945383
30. Izumizaki M. Role of the carotid bodies in chemosensory ventilatory responses in the anesthetized mouse. *J Appl Physiol*. 2004 May 14; 97(4):1401–7. <https://doi.org/10.1152/jappphysiol.00025.2004> PMID: 15194670
31. Xiong ZG, Xu T Le. The role of ASICs in cerebral ischemia. *Wiley Interdiscip Rev Membr Transp Signal*. 2012; 1(5):655–62. <https://doi.org/10.1002/wmts.57> PMID: 23181201
32. Xiong ZG, Zhu XM, Chu XP, Minami M, Hey J, Wei WL, et al. Neuroprotection in ischemia: Blocking calcium-permeable acid-sensing ion channels. *Cell*. 2004; 118(6):687–98. <https://doi.org/10.1016/j.cell.2004.08.026> PMID: 15369669
33. Liu X, He L, Dinger B, Fidone SJ. Chronic hypoxia-induced acid-sensitive ion channel expression in chemoafferent neurons contributes to chemoreceptor hypersensitivity. *AJP Lung Cell Mol Physiol*. 2011; 301(6):L985–92.
34. Dubé GR, Lehto SG, Breese NM, Baker SJ, Wang X, Matulenko MA, et al. Electrophysiological and in vivo characterization of A-317567, a novel blocker of acid sensing ion channels. *Pain*. 2005 Sep; 117 (1):88–96.
35. Voilley N, de Weille J, Mamet J, Lazdunski M. Nonsteroid anti-inflammatory drugs inhibit both the activity and the inflammation-induced expression of acid-sensing ion channels in nociceptors. *J Neurosci*. 2001 Oct 15; 21(20):8026–33. PMID: 11588175
36. Wemmie JA, Taugher RJ, Kreple CJ. Acid-sensing ion channels in pain and disease. *Nat Rev Neurosci*. 2013 Jun 20; 14(7):461–71. <https://doi.org/10.1038/nrn3529> PMID: 23783197