

Childhood maltreatment and suicide attempts in prisoners: a systematic meta-analytic review

Ioannis Angelakis¹ , Jennifer L. Austin¹ and Patricia Gooding^{2,3}

Review Article

Cite this article: Angelakis I, Austin JL, Gooding P (2020). Childhood maltreatment and suicide attempts in prisoners: a systematic meta-analytic review. *Psychological Medicine* 50, 1–10. <https://doi.org/10.1017/S0033291719002848>

Received: 31 May 2019
Revised: 11 September 2019
Accepted: 13 September 2019
First published online: 30 October 2019

Key words:

Abuse; childhood maltreatment; meta-analysis; neglect; prisoners; suicide attempts; systematic review

Author for correspondence:

Ioannis Angelakis,
E-mail: ioannis.angelakis@southwales.ac.uk

¹University of South Wales, School of Psychology, Pontypridd, Wales, UK; ²Division of Psychology and Mental Health, School of Health Sciences, Faculty of Biological, Medical and Health Sciences, University of Manchester, Manchester, UK and ³Manchester Academic Health Science Centre, MAHSC, Manchester, UK

Abstract

In the past decade, the links between core types of childhood maltreatment and suicidal acts have become an increasingly important area of investigation. However, no meta-analytic review has examined this relationship in prisoners. We undertook the first systematic meta-analytic review examining the link between childhood maltreatment and suicide attempts in prisoners to redress this important gap. We searched Medline, PsychINFO, Embase, Web of Science and CINAHL from inception until August 2019. Meta-analyses using random effect models were applied, and heterogeneity was quantified using the I^2 statistic. Publication bias and risk of bias across studies were assessed. We identified 24 studies comprising 16 586 prisoners. The rates of different types of childhood maltreatment ranged between 29% and 68% [95% confidence interval (CI) 18–81%]. The rate of suicide attempts in prisoners was 23% (95% CI 18–27%). Main results demonstrated that sexual abuse [odds ratio (OR) 2.68, 95% CI 1.86–3.86], physical abuse (OR 2.16, 95% CI 1.60–2.91), emotional abuse (OR 2.70, 95% CI 1.92–3.79), emotional neglect (OR 2.29, 95% CI 1.69–3.10), physical neglect (OR 1.57, 95% CI 1.27–1.94) and combined abuse (OR 3.09, 95% CI 2.14–4.45) were strongly associated with suicide attempts in prisoners. There was an indication of publication bias. Duval and Tweedie's trim-and-fill method was applied, which increased the odds for suicide attempts. Given the high rates of prison suicide deaths and suicide attempts, our findings suggest an urgent need for targeted suicide prevention priorities for prisoners, with a particular focus on ameliorating the effects of childhood traumatic experiences on suicidal prisoners.

Suicide fatalities are a major problem facing prisons, with male and female prisoners dying by suicide at a rate that is three- to nine-times higher than that found in the general population (Fazel *et al.*, 2011, 2017). Furthermore, Fazel *et al.* (2017) reported that such fatalities have been elevated for prisoners in England and Wales, where suicide death rates were reported to be between 6 and 20 times higher than those recorded in the community. With regard to the geographical distribution of suicide deaths, Nordic and Western European countries have reported the highest suicide death rates, with over 100 suicide fatalities per 100 000 prisoners being recorded. In Australasian and North American countries, prison suicide death rates ranged between 23 and 67 per 100 000.

Given the high rates of prison suicide deaths globally, it is imperative to identify key factors that may contribute to suicidal acts in prisoners. Engaging in suicide attempts is a strong predictor of suicide deaths (Joiner, 2005; Johnson *et al.*, 2010). Estimates of the prevalence of suicide attempts while in prison and those of the prevalence of attempts of current prisoners prior to being incarcerated have been reported as ranging between 10% and 31%, and between 10% and 54%, respectively (Godet-Mardirossian *et al.*, 2011; Bhatta *et al.*, 2014; Moore *et al.*, 2015; Power *et al.*, 2016; Sánchez *et al.*, 2019). Estimates of lifetime suicide attempts within the general population range between 0.3% and 3% (Nock *et al.*, 2008; Borges *et al.*, 2010). Thus, an in-depth examination of the key variables that are linked with suicide attempts are essential in reducing the elevated rates of prison suicide deaths.

Adverse childhood experiences have been robustly associated with an increased risk for suicide attempts (Brezo *et al.*, 2008; Kim *et al.*, 2013). For example, a meta-analytic review which examined this association in clinical and non-clinical populations demonstrated a strong link between key types of childhood maltreatment (i.e. sexual, physical and emotional abuse, and emotional neglect) and both suicide attempts and ideation (Angelakis *et al.*, 2019). Notably, these results were unaffected by key methodological variations (e.g. screening tools and/or research designs employed) and individual-level characteristics (e.g. age and gender) across studies. This demonstrates the robustness of this relationship.

In the last decade, links between myriad forms of childhood abuse and suicidal experiences have begun to be explored in incarcerated populations (Bhatta *et al.*, 2014; Boonmann *et al.*, 2016). Previous research has established that prisoners reliably report more childhood traumatic events than those in the general community (Jordan *et al.*, 1996) and that these experiences are strongly linked to suicidal acts (Angelakis *et al.*, 2019). Furthermore, with prison

© The Author(s) 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

suicide death rates being substantially higher than those in the general population, a systematic meta-analytic review examining the link between childhood maltreatment and suicide behaviors in prisoners is critical for guiding future research priorities and associated policy changes. Currently, there is no critical synthesis of evidence pertaining to the relationship between childhood abuse and suicidal acts within individuals in custody. Hence, we undertook the first systematic meta-analytic review evaluating this relationship. There were two key objectives:

- (i) to quantitatively analyze the associations between core forms of childhood maltreatment and suicide attempts in prisoners;
- (ii) to examine the degree to which any links between childhood abuse and suicide attempts also apply to suicidal ideation.

Method

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA; Moher *et al.*, 2009) and the Meta-Analysis of Observational Studies in Epidemiology (MOOSE; Stroup *et al.*, 2000) statements were employed to conduct and report the findings of this systematic review and meta-analysis. The MOOSE checklist is available in the online supplemental material (Appendix A).

Eligibility criteria

Studies were included that:

- (a) reported a quantitative outcome of the association between any type of childhood maltreatment (e.g. sexual, physical, emotional abuse; physical and/or emotional neglect) experienced before the age of 18 years and suicide attempts or any other mode of suicidal experiences, such as thoughts, plans, intentions or deaths;
- (b) focused on both juvenile (under 18) and adult (18 and over) prisoners;
- (c) employed a quantitative research design;
- (d) were written in English and published in peer-reviewed academic journals.

Studies were excluded that:

- (a) focused on adverse childhood experiences other than abuse and neglect (e.g. death of a significant other, experiences of being bullied);
- (b) reported data on the link between abuse and neglect experienced in adulthood (e.g. domestic abuse) and suicidal behaviors;
- (c) were reviews, position papers, theses, and/or reports.

Search strategy and data sources

Five electronic bibliographic databases, including Medline, PsychInfo, Embase, Web of Science, and CINAHL, were searched with an end date of August 2019. The reference lists of the identified studies were also checked to locate eligible studies, and authors were contacted if additional information was needed. The search strategy included both text words and Medical Subjective Headings (MeSH) terms and combined three main blocks of key-terms: suicide (suicid*, self* harm*, suicid* correl*), child/sexual/physical/emotional abuse or neglect or maltreatment

or adversities (child*, sex*, phys*, emoti* abuse, negl*, maltreat*, advers*) and prisoners (prison*, inmat*, offend*, jail*, custod*, imprison*, correction*, correctional facilit*, institut*, felon*, delinquen*, disciplin*, criminal*).

Study selection

Study selection was conducted in two stages. In stage 1, the first two authors (IA and JA) independently examined the titles and abstracts of the papers. In stage 2, the same raters evaluated whether all the inclusion and exclusion criteria were met. Inter-rater reliability for the title/abstract and full-text screening process was excellent ($\kappa = 0.91$ and 0.93 , respectively).

Data extraction

An electronic data extraction sheet was devised and piloted in five randomly selected papers to extract descriptive data on key study characteristics, including number and age of participants, research design, method of recruitment, type of childhood maltreatment, mode of suicidal experiences and behaviors, and quantitative data [e.g. odds ratios (ORs) or equivalent measures] for the meta-analysis. Once the data sheet was deemed appropriate, data extraction on the remaining articles was completed by the first two authors (IA and JA). Inter-rater agreement was excellent ($\kappa = 0.93$).

Risk of bias assessment

The methodological quality of the studies was critically evaluated using the criteria as adopted by the Centre for Reviews and Dissemination (CRD, 2010) guidance for undertaking reviews in health care. These assessed the research design (cross sectional = 0, prospective/experimental = 1), response rate ($\leq 70\%$ or not reported = 0, $\geq 70\%$ = 1), screening tools for childhood maltreatment and for suicide experiences (self-report scale/not reported = 0, structured/semi-structured clinical interview = 1), and control for confounding/other factors in the reported analyses (no control/not reported = 0, controlled = 1). Those studies that were scored with at least four criteria out of five were given a high-quality rating; those with less than or equal to three criteria were given a low-quality rating. A binary critical appraisal item was also created based on these ratings (1 = low quality appraisal score; 2 = high quality appraisal score) which was entered as a moderator in the meta-regression analyses.

Data analyses

The primary outcome was the association between different forms of childhood maltreatment and suicide attempts. More than half of the studies ($k = 16$) reported suicide attempts as dichotomous outcomes (number/proportions of prisoners with or without experiences of childhood maltreatment who engaged in suicide behavior in adulthood), whereas the remaining eight studies reported such outcomes in different formats (e.g. mean score of suicide attempts in participants with and without history of childhood maltreatment). ORs were selected as the preferred effect size. Continuous data were converted to ORs by utilizing a widely used formula (Borenstein *et al.*, 2009). Most of the included studies ($n = 18$) contributed more than one relevant effect size. Therefore, we ensured that all the different types of childhood




Fig. 1. PRISMA 2009 flow diagram for the entire review.

maltreatment were pooled separately to avoid double counting of studies in the same analysis.

All data were analyzed in Stata 15°. We used the *metan* command to calculate the pooled effect size of the link between the different types of childhood maltreatment and suicide attempts. A random effects model was utilized because substantial heterogeneity was anticipated, which was assessed with the I^2 statistic (Higgins *et al.*, 2003). Traditionally, values of 25%, 50% and 75% indicate low, moderate and high heterogeneity, respectively. Publication bias was examined with funnel plots and by applying the Egger *et al.*'s (1997) test in cases where the analyses were based on at least nine studies (Saveleva and Selinski, 2008). Duval and Tweedie's (2000) trim-and-fill method, which yields an estimate of the number of the missing studies, was used to provide an adjusted effect size estimate in case of publication bias. The prevalence rates of the various forms of childhood maltreatment and suicide attempts in prisoners were calculated by using the *metaprop* command (random effects model due to anticipated heterogeneity; Nyaga *et al.*, 2014), by excluding controls for those studies which recruited a mixed sample of participants (Rivlin *et al.*, 2013; Brewer-Smyth *et al.*, 2014). Finally, the extent to which methodological variations across the studies affected the link between childhood maltreatment and suicide attempts in prisoners was examined by applying univariate meta-regression models using the *metareg* command (Harbord and Higgins, 2008). We conducted such analyses only for comparisons comprising eight or more studies (Thompson and Higgins, 2002). Overall, two continuous and four categorical moderators were examined including: participants' age; percentage of males;

screening method for childhood maltreatment and suicide attempts (1 = self-report survey, 2 = interview); timeframe of suicide attempts (1 = lifetime, 2 = prison, 3 = both) and quality appraisal score (1 = low, 2 = high).

Results

We retrieved 2414 articles; 322 were removed as they were duplicates and a further 1974 were excluded because they (a) focused on other forms of adverse childhood experiences (e.g. parental death or divorce), (b) non-suicidal self-harm experiences (e.g. self-mutilation), (c) were non-empirical studies, or (d) were focused on misconduct, inappropriate behaviors, rule violation in prisons among others. The full texts of 118 articles were screened for inclusion. An additional 90 studies were excluded as they did not report data relevant to the link between childhood maltreatment and suicidal thoughts and behaviors. Of the remaining 28 studies, only four focused exclusively on either suicidal ideation (Zhang *et al.*, 2010; Boonmann *et al.*, 2016), suicide risk (Blaauw *et al.*, 2002), or suicidality (a term that may incorporate attempts, ideation, plans, urges and intentions; Sergeantanis *et al.*, 2014), resulting in 24 independent studies which were focused exclusively on the link between any form of childhood maltreatment and suicide attempts in prisoners (see Fig. 1).

The overall sample was 16 586 prisoners across the 24 studies (see Table 1). The mean age was 32.39 (s.d. = 4.64, range: 17–41). Fifty-eight percent of participants were male. Most of the studies were conducted in the United States ($k = 9$; 37.5%), Italy ($k = 4$; 16.67%), or Australia ($k = 3$; 12.5%), whereas a single study

Table 1. Study characteristics

Study	Country	Study design	Screening tool for abuse	Mode of childhood abuse	Screening tool for suicidality	Mode of suicidality	Timeframe of suicidality	Nature of offence	Sample size <i>N</i>	Men (%)	Age (years)	Q.A.
Altinas and Bilici (2018)	Turkey	Cross-sectional	CTQ	All forms of abuse and neglect	CI	SA	Lifetime	n/r	216, response rate = 92.6%	50	Range: 18 and above	3/5
Bhatta <i>et al.</i> (2014)	United States	Cross-sectional	SRQ	Sexual abuse	SRQ	SA	Lifetime	Personal, property, drug, public order and unruly offences	3156, response rate = 100%	78.0	Range: 12–17	4/5
Brewer-Smyth <i>et al.</i> (2014)	United States	Case-control	MS	Physical and sexual abuse	n/r	SA	Current	Property and drug offences	113, response rate = 91.87%	0	n/r	2/3
Butler <i>et al.</i> (2001)	Australia	Cross-sectional	HSI	Sexual Abuse	HSI	SA	Lifetime	Violent and non-violent offenses	789, response rate = 90%	83.3	Median = 30, range: 18–77	4/5
Chapman <i>et al.</i> (2014)	United States	Cross-sectional	CTQ	All forms of abuse and neglect	LPC-II	SA	Lifetime/current	Personal, property, drug offenses	104, response rate = 88.89%	0	$M_{age} = 31.94$	3/5
Chen and Gueta (2017)	Israel	Cross-sectional	CTQ	All forms of abuse and neglect	ASI-I	SA	Lifetime	Drug and property offenses, prostitution and assault	46, response rate = 100%	0	$M_{age} = 34.82$	3/5
Clements-Nolle <i>et al.</i> (2009)	United States	Cross-sectional	CTQ	All forms of abuse and neglect	SBQ	SA	Lifetime	n/r	247, response rate = 54%	0	Median = 43, range: 20–72	3/5
DeCou <i>et al.</i> (2016)	United States	Cross-sectional	LHC	Sexual abuse	LHC	SA	Lifetime	n/r	491, response rate = 23.42%	0	$M_{age} = 34$, range: 17–55	2/5
Friestad <i>et al.</i> (2014)	Norway	Cross-sectional	CI	All forms of abuse and neglect	CI	SA	Lifetime	n/r	141, response rate = 75.4%	0	$M_{age} = 33$	4/5
Godet-Mardirossian <i>et al.</i> (2011)	France	Cross-sectional	CI	Combined forms of childhood maltreatment	MNI	SA	Current	n/r	899, response rate = 100%	100	Median = 37, range: 19–84	3/5
Gorodetsky <i>et al.</i> (2016)	Italy	Cross-sectional	CTQ	All forms of abuse and neglect	MNI	SA	Lifetime	n/r	702, response rate = 100%	100	n/r	3/5
Hakansson <i>et al.</i> (2011)	Sweden	Cross-sectional	ASI	Sexual, physical and emotional abuse	ASI	SA	Lifetime/current	n/r	1561, response rate = 84.93%	80.62	n/r	4/5

Jeglic <i>et al.</i> (2013)	United States	Cross-sectional	Third party information	Combined forms of childhood maltreatment	Third party information	SA	Lifetime/current	Sexual offenses	3030, response rate = 100%	100	$M_{age} = 39.2$, range: 13–85	3/5
Kenny <i>et al.</i> (2008)	Australia	Cross-sectional	CTQ	All forms of abuse and neglect	CI	SA	Lifetime/current	Violent crimes, property, drug, public order and other offenses	319, response rate = 76%	92	$M_{age} = 16.55$, range: 16–22	3/5
Mandelli <i>et al.</i> (2011)	Italy	Cross-sectional	CTQ	All forms of abuse and neglect	MNI	SA	Lifetime	Violent and non-violent offenses	2000, response rate = 77.6%	100	$M_{age} = 40$	4/5
Moore <i>et al.</i> (2015)	Australia	Cross-sectional	CTQ	All forms of abuse and neglect	SBQ	SA	Lifetime/current	n/r	361, response rate = 86.70%	87.54	n/r	3/5
Power <i>et al.</i> (2016)	Canada	Cross-sectional	CTQ	All forms of abuse and neglect	CI	SA	Lifetime	n/r	415, response rate = 100%	64.58	n/r	4/5
Rivlin <i>et al.</i> (2013)	England	Matched-case control	CTQ	All forms of abuse and neglect	Third party referrals from prison officers	SA	Current	Violent, sexual and property offenses	120, response rate = 100%	100	Range: 18 years and above	4/5
Roy <i>et al.</i> (2014)	Italy	Cross-sectional	CTQ	All forms of abuse and neglect	MNI	SA	Lifetime	Violent and non-violent offenses	1537, response rate = 100%	100	$M_{age} = 39.6$	3/5
Sánchez <i>et al.</i> (2019)	Spain	Cross-sectional	CTQ	Emotional physical abuse, and sexual abuse	A single question	SA	Current	Property offenses	1146, response rate = 82.29%	100	$M_{age} = 36.9$	1/5
Sarchiapone <i>et al.</i> (2009)	Italy	Cross-sectional	CTQ	All forms of abuse and neglect	MNI	SA	Lifetime	Violent and non-violent offenses	903, response rate = 100%	100	$M_{age} = 40.74$	3/5
Swogger <i>et al.</i> (2011)	United States	Cross-sectional	A single question	Physical abuse	A single question	SA	Lifetime	n/r	266, response rate = 100%	74.81	$M_{age} = 33.7$, range: 18–62	4/5
Tripodi <i>et al.</i> (2014)	United States	Cross-sectional	CTQ	All forms of abuse and neglect	ASI	SA	Lifetime	n/r	150, response rate = 83%	0	$M_{age} = 34.3$	4/5
Verona <i>et al.</i> (2005)	United States	Cross-sectional	CI	Physical and sexual abuse	CI	SA	Lifetime	n/r	226, response rate = 100%	0	$M_{age} = 31.9$	3/5

ASI, addiction severity index; ASI-I, Addiction Severity Index Interview; CI, Clinical Interview; CTQ, Child Trauma Questionnaire; HSI, Health Survey Interview; LHC, Life History Calendars; LPC-II, Lifetime Para-suicide Count-II; M, mean; MNI, Mini Neuropsychiatric Interview; MS, Muenzenmaier's Scale; n/r, not reported; Q.A, Quality appraisal of the methodology of the included studies; SA, suicide attempts; SBQ, suicidal behaviours; s.d., standard deviation; SRQ, Self-Report Questions

Table 2. Rates of childhood maltreatment and suicide attempts in prisoners

	Prevalence			Heterogeneity
	<i>k</i>	%	95% CI	<i>I</i> ²
Childhood maltreatment				
Sexual abuse	11	29	18–41	98.76%
Physical abuse	8	43	29–57	98.08%
Emotional abuse	7	49	32–65	98.40%
Emotional neglect	3	68	53–81	86.16%
Suicidality				
Suicide attempts	21	23	18–27	97.24%

k, number of independent prevalence estimates.

(4.17%) was conducted in each of England, France, Spain, Sweden, Norway, Israel, Canada, and Turkey.

As detailed in Table 2, within the current sample, 29% of the prisoners reported being victims of sexual abuse [$k = 11$, 95% confidence interval (CI) 18–41%, $I^2 = 98.76%$]; 43% had experienced physical abuse ($k = 8$, 95% CI 29–57%, $I^2 = 98.08%$); 49% had experienced emotional abuse ($k = 7$, 95% CI 32–65%, $I^2 = 98.40%$); and 68% had experienced emotional neglect ($k = 3$, 95% CI 53–81%, $I^2 = 86.16%$).

With regard to suicide attempts, 58.33% ($k = 14$) of the studies reported data on lifetime suicide attempts, which incorporated suicide attempts of prisoners while in prison and attempts prior to or after leaving the prison, 16.66% ($k = 4$) reported data on suicide attempts while in prison, and the remaining 20.83% ($k = 5$) focused on a combination of attempts that occurred either during or after imprisonment. Hence, provided that lifetime suicide attempts also incorporated those attempts that occurred exclusively in prisons, we pooled and reported the rates of suicide attempts into a single variable. The overall prevalence of the suicide attempts of prisoners was 23% (95% CI 18–27%, $I^2 = 97.24%$) based on 21 studies (see Table 2; online Supplementary Appendix E).

Main meta-analyses: associations between types of childhood maltreatment and suicide attempts in prisoners

All forms of childhood maltreatment were associated with significantly increased odds for suicide attempts in prisoners (see online Supplementary Appendix B for forest plots). Sexual and emotional abuse were associated with a three-fold increased likelihood for suicide attempts ($k = 16$, OR 2.68, 95% CI 1.86–3.86, $I^2 = 92.4%$ and $k = 9$, OR 2.70, 95% CI 1.92–3.79, $I^2 = 74.0%$, respectively) whereas physical abuse and emotional neglect were associated with a two-fold increased likelihood for suicide attempts ($k = 11$, OR 2.16, 95% CI 1.60–2.91, $I^2 = 86.6%$ and $k = 7$, OR 2.29, 95% CI 1.69–3.10, $I^2 = 41.9%$, respectively). Physical neglect was associated with 1.5-fold increased likelihood for suicide attempts ($k = 7$, OR 1.57, 95% CI 1.27–1.94, $I^2 = 89.2%$). Almost half of the included studies provided data for the link between unspecified forms of childhood abuse/neglect and suicide attempts. This category was named ‘combined abuse’ and was analyzed separately. Combined abuse was associated with a three times increased likelihood for suicide attempts ($k = 13$, OR 3.09, 95% CI 2.14–4.45, $I^2 = 91.9%$). As indicated by the I^2 statistic, heterogeneity ranged from medium to

high across all analyses, except for the association between emotional neglect and suicide attempts, where heterogeneity ranged from moderate to low.

Publication bias

We assessed publication bias across comparisons that included at least nine studies. The screening of the funnel plots (see online Supplementary Appendix C) suggested that there was a publication bias for the relationship between sexual abuse and suicide attempts. Furthermore, the Egger’s *et al.* (1997) test for publication bias was significant in all the comparisons examined, which was an indication of publication bias (see Table 3). Therefore, we ran the Duval and Tweedie’s (2000) trim-and-fill method, which considerably increased the effect sizes (ORs ranged from 3.54 to 5.28; see Table 3). These results suggest that publication bias may not be an actual concern (Murad *et al.*, 2018).

Meta-regressions examining the impact of methodological variations on the associations between distinct forms of childhood maltreatment and suicide attempts in prisoners

The number of pooled studies only allowed meta-regression analyses to be conducted for the associations between sexual, physical, emotional and combined abuse, and suicide attempts. The results of the univariate meta-regression analyses revealed that participant variables, including mean age (p values ranged between 0.61 and 0.87) and percentage of males (p values ranged between 0.34 and 0.97), and methodological variations, such as screening measures for childhood maltreatment (p values ranged between 0.44 and 0.86), screening measures for suicide attempts (p values ranged between 0.73 and 0.96), timeframe of suicide attempts (p values ranged between 0.40 and 0.79) and critical appraisal score (p values ranged between 0.31 and 0.87), across studies did not affect the strength of the associations between different forms of abuse and suicide attempts (see online Supplementary Appendix D for the results of the meta-regression analyses).

Discussion

This is the first systematic meta-analytic review to examine the relationship between core forms of childhood maltreatment and suicide attempts in prisoners. We reported results from a total of 16 586 prisoners. The rates of childhood maltreatment experiences ranged from 29% to 68%, whereas the rate of suicide attempts in prisoners was 23%. There were two key findings. First, we demonstrated that suicide attempts were strongly linked, cross-sectionally, with experiences of childhood abuse and/or neglect in prisoners, which remained unaffected by methodological variations. In particular, prisoners who reported experiences of combined abuse/neglect, sexual and emotional abuse were three times more likely to engage in suicide attempts compared to those prisoners who had not experienced such events. Physical abuse and emotional neglect were associated with twice the likelihood for suicide attempts. Finally, prisoners who reported being physically neglected during childhood were found to have 1.5 more suicide attempts in adulthood compared with prisoners who had not experienced abuse or neglect. It must be highlighted that these results reflect cross-sectional rather than longitudinal data. Consequently, neither temporal precedence nor causation can be demonstrated.

Table 3. Results of meta-analyses of the association between forms of childhood maltreatment and suicide attempts in prisoners

	Total <i>k</i>	Total <i>N</i>	Effect size analyses			Heterogeneity <i>I</i> ²	Publication bias	
			OR	95% CI	<i>p</i> value		Egger's <i>p</i>	Trim-and-fill OR (95% CI)
Suicide attempts								
Sexual abuse	16 ^a	9835	2.68	1.86–3.86	<0.001	92.4%	0.001	3.92 (1.50–10.26)
Physical abuse	11 ^b	5572	2.16	1.60–2.91	<0.001	86.6%	0.001	3.66 (1.39–9.60)
Emotional abuse	9	5258	2.70	1.92–3.79	<0.001	74.0%	0.001	5.28 (1.02–27.8)
Emotional neglect	7 ^b	3123	2.29	1.69–3.10	<0.001	41.9%	–	–
Physical neglect	7 ^b	3049	1.57	1.27–1.94	<0.001	89.2%	–	–
Combined abuse	13 ^b	10 219	3.09	2.14–4.45	<0.001	91.9%	0.001	4.48 (1.33–15.13)

k, number of independent effect sizes; OR, pooled odds ratio; *N*, number of participants.

^aTwo outliers were dropped from the analyses.

^bOne outlier was dropped from the analyses.

To date there have been limited attempts to link the strong relationship between forms of childhood maltreatment and suicidal experiences, with contemporary models of suicidal thoughts and behaviors (Johnson *et al.*, 2008; O'Connor and Nock, 2014; Klonsky and May, 2015; O'Connor and Kirtley, 2018; O'Connor and Portzky, 2018). Four key priorities would advance this literature. First, researchers should identify socio-economic and epidemiological factors which are associated with a high prevalence of suicidal thoughts, acts, and fatalities in vulnerable prisoners (Rivlin *et al.*, 2013; Fazel *et al.*, 2017). Second, a better understanding of the ways in which different forms of childhood abuse may exacerbate symptoms of mental health problems is needed. Examples of such symptoms include avoidance, hyperarousal, and command hallucinations which may trigger and/or amplify suicidal thoughts and behaviors (Panagioti *et al.*, 2017). Third, future research should examine the ways in which experiences of childhood abuse feed into more transdiagnostic approaches which seek to better understand the psychological pathways leading to suicidal thoughts, urges, plans, acts and deaths (Bolton *et al.*, 2007; Johnson *et al.*, 2008; O'Connor and Kirtley, 2018). The fourth issue follows in that there should be an endeavor to investigate how symptom-specific contributors to suicidal experiences interact with putative psychological suicide mechanisms (Gooding *et al.*, 2015). Many of the contemporary models of suicide highlight that important mediators in the relationship between childhood maltreatment and suicide attempts, which include perceptions of being socially isolated, feeling a burden, being overwhelmed by, for example, strong labile emotions, feeling defeated and/or trapped, and having no hope for the future, are central to the pathways to suicidal experiences (Johnson *et al.*, 2008; Van Orden *et al.*, 2010; Palmier-Claus *et al.*, 2012; O'Connor and Kirtley, 2018). Therefore, it is paramount that future studies explore the interactions between past and current traumatic experiences in prisoners that may amplify these types of feelings and perceptions using qualitative and quantitative longitudinal and micro-longitudinal designs. Finally, a key component to consider is the development and activation of suicide schema in people who have experienced child abuse (Lau *et al.*, 2004). Suicide schemas are under-researched but have been found to be extensive in people with non-affective psychosis and also post-traumatic stress disorder (Tarrier *et al.*, 2007; Pratt *et al.*, 2010; Panagioti *et al.*, 2015). It is important to determine how suicide schema develop and become more extensive over

time as a result of current and past childhood abuse and/or neglect, especially in those vulnerable to suicidal experiences (i.e. prisoners).

None of the identified studies examined the impact of potentially important moderators on the relationship between childhood maltreatment and suicide attempts. Examples might include such variables as length of offense, life events, such as homelessness or running away from home as an adolescent (Kenny *et al.*, 2008; Rivlin *et al.*, 2013; Bhatta *et al.*, 2014; Sánchez *et al.*, 2019), and adverse prison-related experiences, including social isolation and victimization by other prisoners (Blaauw *et al.*, 2002; Suto and Arnaut, 2010). Perhaps most importantly, we should prioritize studies examining key factors that attenuate or buffer the relationship between childhood abuse/neglect and suicide attempts. For example, Rivlin *et al.* (2011) suggested that an increased level of care from professionals or prison staff, especially when prisoners engage in precursor non-suicidal self-harm behaviors, may be effective at reversing such outcomes. However, while examination of mediators and/or moderators is important, studies have rarely been able to examine these types of mediators and moderators in prisoners who have attempted or died by suicide, because converging data on pertinent and meaningful mediators and/or moderators together with suicide attempts and fatalities are rare. Clearly, future research should further explore the significance of these factors on suicidal experiences in prisoners.

Second, we identified a paucity of studies exploring the link between childhood abuse/neglect and the different modes of suicide behavior other than attempts, including suicidal ideation, urges, and plans. We only identified four studies exploring the link between any form of childhood maltreatment and suicidal ideation or suicide risk (Blaauw *et al.*, 2002; Zhang *et al.*, 2010; Sergeantanis *et al.*, 2014; Boonmann *et al.*, 2016). This is an important gap in the literature because there is compelling evidence suggesting that among those individuals experiencing lifetime suicidal ideation with or without plans the likelihood of acting upon their thoughts ranges between 30% and 55% (Nock *et al.*, 2008). Furthermore, alleviating the immense psychological distress associated with having suicidal thoughts, urges and/or forming suicide plans is an important clinical target (Tarrier *et al.*, 2013).

Recent evidence suggests that cognitive-behavioral suicide prevention therapies can be successfully extended within prison

settings. For example, Pratt *et al.* (2015) found that this treatment approach was promising in reducing suicide risk in male prisoners considered at high risk for suicide. Given preliminary evidence suggesting that the format of cognitive-behavioral therapy (e.g. individual *v.* group) does not alter its effectiveness in reducing psychological symptoms such as depression and anxiety in prisoners (Khodayarifard *et al.*, 2010), which are strongly linked to suicidal acts, research on scaling up clinical therapies for groups of high-risk prisoners is essential.

It is also worth noting that more than half of the included studies were undertaken in the USA, followed by Italy and Australia. This is not a surprising, given that the USA has among the highest rates of incarceration globally (U.S. Department of Justice, 2016). However, the USA is among the countries with the lowest rates of prison suicide (Fazel *et al.*, 2017). Countries such as England, France, Norway and Sweden, where incarceration rates are lower, contributed a single study (per country) to the analysis. Interestingly, these countries have the highest prison suicide death rates worldwide. Differences in definitions of suicide deaths in prisoners, which was initially suspected to account for this discrepancy, did not affect overall prison suicide rates. It is, therefore, important that more research is conducted to identify key factors that contribute to these outcomes, including country specific provision of mental health care, amount of daily activity, and quality of staff-prisoner relationships, among others.

There are five main limitations of the current study which warrant discussion. First, heterogeneity across the majority of the comparisons undertaken was moderate to high, which is, perhaps, to be expected. To address this, we applied random effect models to adjust for between-study variations. Second, we found an indication of publication bias for the majority of the comparisons examined. To attenuate this bias, we employed the fill-and-trim approach which increased the effect sizes. Although these results suggest that the strong link between experiences of childhood maltreatment and suicide attempts in prisoners may remain unaffected (e.g. Murad *et al.*, 2018), they should still be interpreted with caution. Third, the number of studies included in some comparisons was small. As such, these findings may warrant further investigation. Fourth, this review focused exclusively on core types of childhood maltreatment and, as such, particular types of adverse childhood events (e.g. parental deaths and divorces) were not considered. Future research should explore these relationships in more depth. Fifth, although it would have been more informative to differentiate between the prevalence of lifetime suicide attempts and those attempts that occurred while in the prison system, the lack of such information from the included studies prevented this distinction from being made. However, it is acknowledged that the most helpful information for correctional staff would be rates of suicide attempts while in prison separated from those which preceded incarceration or followed release from prison. That said, it is important to note that many prisoners return to prison after release, making the dynamic between factors which precipitate suicide attempts in prison and those which precipitate suicide attempts after release difficult to capture. In this regard, qualitative work which probes the triggers for suicidal thoughts, plans and attempts while in prison and after release, and the interactions between the effects of such different contexts with child abuse, may be of optimal practical help going forward. With these caveats acknowledged, a clear message from this meta-analysis is that prisoners who have experienced child abuse and/or neglect may be the most vulnerable with respect to having different forms of suicidal thoughts, plans and acts.

This study has three noteworthy strengths. First, it is the first systematic review with meta-analysis which incorporated an extensive synthesis of the association between core forms of childhood maltreatment and suicide attempts in prisoners confirming their robust associations with suicide attempts. Second, the current meta-analysis was conducted in compliance with PRISMA and MOOSE guidelines. Third, the methodological quality of the included studies was critically evaluated and rated.

Conclusions

In summary, this is the first systematic meta-analytic review of the relationship between childhood maltreatment and suicide attempts that was conducted in prisoners. Although this relationship has yet to be fully explored, we demonstrated that a strong link exists between forms of childhood maltreatment and suicide attempts. No evidence exists regarding other modes of suicide behaviors, including suicidal ideation, suicide plans and deaths by suicide. We recommend that further research employ more robust, higher quality designs, such as, longitudinal, micro-longitudinal, qualitative and/or mixed designs, to corroborate and expand the findings of the current review. Taken together, these findings highlight, as a matter of urgency, the need for targeted suicide prevention priorities for prisoners with a focus on ameliorating the effects of childhood maltreatment on suicidal thoughts, behaviors and attempts (House of Commons Library, 2018). These initiatives should be a priority at the levels of policy making and institutional reform, whereby the provision of different modes of suicide focused psychological therapy targeting childhood experiences of maltreatment (Pratt *et al.*, 2015) is conducted in tandem with robust staff awareness and training programs (Joe *et al.*, 2017).

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291719002848>.

Acknowledgements. We thank Matilda Angelaki for comments that greatly improved the manuscript.

Financial support. No financial support has been received for this piece of work.

Conflict of interest. All other authors declare that they have no conflicts of interest.

References

- Altintas M and Bilici M (2018) Evaluation of childhood trauma with respect to criminal behavior, dissociative experiences, adverse family experiences and psychiatric backgrounds among prison inmates. *Comprehensive Psychiatry* **82**, 100–107.
- Angelakis I, Gillespie EL and Panagioti M (2019) Childhood maltreatment and adult suicidality: a comprehensive systematic review with meta-analysis. *Psychological Medicine* **49**, 1057–1078.
- Bhatta M, Jefferis E, Kavadas A, Alemagno S and Shaffer-King P (2014) Suicidal behaviors among adolescents in juvenile detention: role of adverse life experiences. *PLoS ONE* **9**, e89408.
- Blaauw E, Arensman E, Kraaij V, Winkel F and Bout R (2002). Traumatic life events and suicide risk among jail inmates: the influence of types of events, time period and significant others. *Journal of Traumatic Stress* **15**, 9–16.
- Bolton C, Gooding P, Kapur N, Barrowclough C and Tarrier N (2007) Developing psychological perspectives of suicidal behaviour and risk in people with a diagnosis of schizophrenia: we know they kill themselves but do we understand why? *Clinical Psychology Review* **27**, 511–536.

- Boonmann C, Grisso T, Guy L, Colins O, Mulder E and Vahl P (2016) Childhood traumatic experiences and mental health problems in sexually offending and non-sexually offending juveniles. *Child and Adolescent Psychiatry and Mental Health* **10**, 45–45.
- Borenstein M, Hedges L, Higgins J and Higgins D (2009). *Introduction to Meta-Analysis*, 2nd Edn. Hoboken: Wiley.
- Borges G, Nock M, Haro-Abad J, Hwang I, Sampson N, Alonso J, Andrade LH, Angermeyer MC, Beautrais A, Bromet E, Bruffaerts R, de Girolamo G, Florescu S, Gureje O, Hu C, Karam EG, Kovess-Masfety V, Lee S, Levison D, Medina-Mora ME, Ormel J, Posada-Villa J, Sager R, Tomoy T, Uda H, Williams DR and Kessler RC (2010) Twelve-month prevalence of and risk factors for suicide attempts in the World Health Organization World Mental Health Surveys. *The Journal of Clinical Psychiatry* **71**, 1617–1628.
- Brewer-Smyth K, Burgess A and Shults J (2014) Physical and sexual abuse, salivary cortisol, and neurologic correlates of violent criminal behavior in female prison inmates. *Biological Psychiatry* **55**, 21–31.
- Brezo J, Paris J, Vitaro F, Hébert M, Tremblay R and Turecki G (2008) Predicting suicide attempts in young adults with histories of childhood abuse. *The British Journal of Psychiatry: The Journal of Mental Science* **193**, 134–139.
- Butler T, Donovan B, Fleming J, Levy M and Kaldor J (2001) Childhood sexual abuse among Australian prisoners. *Venerology* **14**, 109–115.
- Centre for reviews and dissemination (2010) *CRD's Guidance for Undertaking Reviews in Health Care*. York: CRD, University of York.
- Chapman A, Gratz K and Turner B (2014) Risk-related and protective correlates of nonsuicidal self-injury and co-occurring suicide attempts among incarcerated women. *Suicide and Life-Threatening Behavior* **44**, 139–154.
- Chen G and Gueta K (2017) Lifetime history of suicidal ideation and attempts among incarcerated women in Israel. *Psychological Trauma: Theory, Research, Practice, and Policy* **9**, 596–604.
- Clements-Nolle K, Wolden M and Bargmann-Losche J (2009) Childhood trauma and risk for past and future suicide attempts among women in prison. *Women's Health Issues* **19**, 185–192.
- DeCou C, Lynch S, Dehart D and Belknap J (2016) Evaluating the association between childhood sexual abuse and attempted suicide across the lifespan: findings from a nationwide study of women in jail. *Psychological Services* **13**, 254–260.
- Duval S and Tweedie R (2000) Trim and fill: a simple funnel-plot-based method for testing and adjusting for publication bias in meta-analysis. *Biometrics* **56**, 455–463.
- Egger M, Smith G, Schneider M and Minder C (1997) Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629–634.
- Fazel S, Grann M, Kling B and Hawton K (2011) Prison suicide in 12 countries: an ecological study of 861 suicides during 2003–2007. *Social Psychiatry and Psychiatric Epidemiology* **46**, 191–195.
- Fazel S, Ramesh T and Hawton K (2017) Suicide in prisons: an international study of prevalence and contributory factors. *The Lancet Psychiatry* **4**, 946–952.
- Friestad C, Åse-Bente R, Kjelsberg E and Friestad C (2014) Adverse childhood experiences among women prisoners: relationships to suicide attempts and drug abuse. *The International Journal of Social Psychiatry* **60**, 40–46.
- Godet-Mardirossian H, Jehel L, Falissard B and Godet-Mardirossian H (2011) Suicidality in male prisoners: influence of childhood adversity mediated by dimensions of personality. *Journal of Forensic Sciences* **56**, 942–949.
- Gooding P, Tarrier N, Dunn G, Shaw J, Awenat Y, Ulph F and Pratt D (2015) Effect of hopelessness on the links between psychiatric symptoms and suicidality in a vulnerable population at risk of suicide. *Psychiatry Research* **230**, 464–471.
- Gorodetsky E, Carli V, Sarchiapone M, Roy A, Goldman D and Enoch M (2016) Predictors for self-directed aggression in Italian prisoners include externalizing behaviors, childhood trauma and the serotonin transporter gene polymorphism 5-HTTLPR. *Genes, Brain and Behavior* **15**, 465–473.
- Hakansson A, Bradvik L, Schlyter F and Berglund M (2011) Variables associated with repeated suicide attempt in a criminal justice population. *Suicide and Life-Threatening Behavior* **41**, 517–531.
- Harbord R and Higgins JPT (2008) Meta-regression in Stata. *Stata Journal* **8**, 493–519.
- Higgins J, Thompson S, Deeks J and Altman D (2003) Measuring inconsistency in meta-analyses. *BMJ* **327**, 557–560.
- House of Commons Library (2018) Suicide prevention: Policy and strategy. Available at <https://researchbriefings.parliament.uk/ResearchBriefing/Summary/CBP-8221> (Accessed 7 February 2019).
- Jeglic EL, Spada A and Mercado CC (2013) An examination of suicide attempts among incarcerated sex offenders. *Sexual Abuse: A Journal for Research and Treatment* **25**, 21–40.
- Joe R, Drake GB, Shaffer JS and Jackson BA (2017) *Envisioning an Alternative Future for the Corrections Sector within the U.S.* Criminal Justice System. Santa Monica, CA: RAND Corporation.
- Johnson J, Gooding P and Tarrier N (2008). Suicide risk in schizophrenia: explanatory models and clinical implications. *Psychology and Psychotherapy: Theory, Research and Practice* **81**, 55–77.
- Johnson J, Gooding PA, Wood AM, Taylor PJ, Pratt D and Tarrier N (2010) Resilience to suicidal ideation in psychosis: positive self-appraisals buffer the impact of hopelessness. *Behaviour Research and Therapy* **48**, 883–889.
- Joiner T (2005) *Why People Die by Suicide*. Cambridge, MA: Harvard University Press.
- Jordan BK, Schlenger WE, Fairbank JA and Caddell JM (1996) Prevalence of psychiatric disorders among incarcerated women: convicted felons entering prison. *Archives of General Psychiatry* **53**, 513–519.
- Kenny DT, Lennings CJ and Munn OA (2008) Risk factors for self-harm and suicide in incarcerated young offenders: implications for policy and practice. *Journal of Forensic Psychology Practice* **8**, 358–382.
- Khodayarifard M, Shokoohi-Yekta M and Hamot GE (2010) Effects of individual and group cognitive-behavioral therapy for male prisoners in Iran. *International Journal of Offender Therapy and Comparative Criminology* **54**, 743–755.
- Kim S, Kang H, Kim S, Kim J, Yoon J, Jung S, Lee M, Yim H and Jun T (2013) Impact of childhood adversity on the course and suicidality of depressive disorders: the crescent study. *Depression and Anxiety* **30**, 965–974.
- Klonsky ED and May AM (2015) The Three-Step Theory (3ST): a new theory of suicide rooted in the 'Ideation-to-Action' framework. *International Journal of Cognitive Therapy* **8**, 114–129.
- Lau MA, Segal ZV and Williams JMG (2004) Teasdale's differential activation hypothesis: implications for mechanisms of depressive relapse and suicidal behaviour. *Behaviour Research and Therapy* **42**, 1001–1017.
- Mandelli L, Carli V, Roy A, Serretti A and Sarchiapone M (2011) The influence of childhood trauma on the onset and repetition of suicidal behavior: an investigation in a high-risk sample of male prisoners. *Journal of Psychiatric Research* **45**, 742–747.
- Moher D, Liberati A, Tetzlaff J and Altman DG (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Volucella* **339**, e1000097.
- Moore E, Gaskin C and Indig D (2015) Attempted suicide, self-harm, and psychological disorder among young offenders in custody. *Journal of Correctional Health Care* **21**, 243–254.
- Murad MH, Chu H, Lin L and Wang Z (2018) The effect of publication bias magnitude and direction on the certainty in evidence. *BMJ Evidence-Based Medicine* **23**, 84–86.
- Nock M, Borges G, Bromet E, Alonso J, Angermeyer M, Beautrais A, Bruffaerts R, Chiu WT, de Girolamo G, Gluzman S, de Graaf R, Gureje O, Haro J-M, Huang Y, Karam E, Kessler RC, Lepine JP, Levinson D, Medina-Mora ME, Ono Y, Posada-Villa J and Williams D (2008) Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *The British Journal of Psychiatry* **192**, 98–105.
- Nyaga VN, Arbyn M and Aerts M (2014) Mataprop: a Stata command to perform meta-analysis of binominal data. *Archives of Public Health* **72**, 39.
- O'Connor RC and Kirtley OJ (2018) The integrated motivational-volitional model of suicidal behaviour. *Philosophical Transactions of the Royal Society B-Biological Sciences* **373**, 20170268.

- O'Connor RC and Nock MK (2014) The psychology of suicidal behaviour. *The Lancet. Psychiatry* **1**, 73–85.
- O'Connor RC and Portzky G (2018) Looking to the future: a synthesis of new developments and challenges in suicide research and prevention. *Frontiers in Psychology* **9**, 2139.
- Palmier-Claus JE, Taylor PJ, Gooding P, Dunn G and Lewis SW (2012) Affective variability predicts suicidal ideation in individuals at ultra-high risk of developing psychosis: an experience sampling study. *The British Journal of Clinical Psychology* **51**, 72–83.
- Panagioti M, Angelakis I, Tarrrier N and Gooding P (2017). A prospective investigation of the impact of distinct posttraumatic (PTSD) symptom clusters on suicidal ideation. *Cognitive Therapy and Research* **41**, 645–653.
- Panagioti M, Gooding PA, Pratt D and Tarrrier N (2015) An empirical investigation of suicide schemas in individuals with posttraumatic stress disorder. *Psychiatry Research* **227**, 302–308.
- Power J, Gobeil R, Beaudette J, Ritchie M, Brown S and Smith H (2016) Childhood abuse, nonsuicidal self-injury, and suicide attempts: an exploration of gender differences in incarcerated adults. *Suicide and Life-Threatening Behavior* **46**, 745–751.
- Pratt D, Gooding P, Johnson J, Taylor P and Tarrrier N (2010) Suicide schemas in non-affective psychosis: an empirical investigation. *Behaviour Research and Therapy* **48**, 1211–1220.
- Pratt D, Tarrrier N, Dunn G, Awenat Y, Shaw J, Ulph F and Gooding P (2015) Cognitive-behavioural suicide prevention for male prisoners: a pilot randomized controlled trial. *Psychological Medicine* **45**, 3441–3451.
- Rivlin A, Fazel S, Marzano L and Hawton K (2011) The suicidal process in male prisoners making near-lethal suicide attempts. *Psychology, Crime & Law* **19**, 305–327.
- Rivlin A, Hawton K, Marzano L and Fazel S (2013) Psychosocial characteristics and social networks of suicidal prisoners: towards a model of suicidal behaviour in detention. *PLoS ONE* **8**, e68944.
- Roy A, Carli V, Sarchiapone M and Branchey M (2014) Comparisons of prisoners who make or do not make suicide attempts and further who make one or multiple attempts. *Archives of Suicide Research* **18**, 28–38.
- Sánchez FC, Ignatyev Y and Mundt A (2019) Associations between childhood abuse, mental health problems, and suicide risk among male prison populations in Spain. *Criminal Behaviour and Mental Health, CBMH* **29**, 18–30.
- Sarchiapone M, Carli V, Di Giannantonio M and Roy A (2009) Risk factors for attempting suicide in prisoners. *Suicide and Life-Threatening Behavior* **39**, 343–350.
- Saveleva E and Selinski S (2008) Meta-analyses with binary outcomes: how many studies need to be omitted to detect a publication bias? *Journal of Toxicology and Environmental Health, Part A* **71**, 845–850.
- Sergentanis T, Sakelliadis E, Vlachodimitropoulos D, Goutas N, Sergentanis I, Spiliopoulou C and Papadodima S (2014) Does history of childhood maltreatment make a difference in prison? A hierarchical approach on early family events and personality traits. *Psychiatry Research* **220**, 1064–1070.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA and Thacker SB (2000) Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group. *JAMA* **283**, 2008–2012.
- Suto I and Arnaut G (2010) Suicide in prison: a qualitative study. *The Prison Journal* **90**, 288–312.
- Swogger M, You S, Cashman-Brown S and Conner K (2011) Childhood physical abuse, aggression, and suicide attempts among criminal offenders. *Psychiatry Research* **185**, 363–367.
- Tarrrier N, Gooding P, Gregg L, Johnson J, Drake R and Socrates TG (2007) Suicide schema in schizophrenia: the effect of emotional reactivity, negative symptoms and schema elaboration. *Behaviour Research and Therapy* **45**, 2090–2097.
- Tarrrier N, Gooding P, Pratt D, Kelly J, Awenat Y and Maxwell J (2013) *Cognitive Behavioural Prevention of Suicide in Psychosis: A Treatment Manual*. London, UK: Routledge.
- Thompson SG and Higgins JPT (2002) How should meta-regression analyses should be undertaken and interpreted? *Statistics in Medicine* **21**, 1559–1513.
- Tripodi S, Onifade E and Pettus-Davis C (2014) Nonfatal suicidal behavior among women prisoners: the predictive roles of childhood victimization, childhood neglect, and childhood positive support. *International Journal of Offender Therapy and Comparative Criminology* **58**, 394–411.
- U.S. Department of Justice, Bureau of Justice Statistics (2016) Correctional populations in the United States. Available at www.bjs.gov/content/pub/pdf/cpus16.pdf (Accessed 18 February 2019).
- Van Orden KA, Witte TK, Cukrowicz KC, Braithwaite SR, Selby EA and Joiner TE (2010) The interpersonal theory of suicide. *Psychological Review* **117**, 575–600.
- Verona E, Hicks B and Patrick C (2005) Psychopathy and suicidality in female offenders: mediating influences of personality and abuse. *Journal of Consulting and Clinical Psychology* **73**, 1065–1073.
- Zhang J, Grabiner V, Zhou Y and Li N (2010) Suicidal ideation and its correlates in prisoners: a comparative study in China. *Crisis* **31**, 335–342.