

site was developed; meetings and teleconferences were held regularly. By way of a shared contract, a CSS (Theradoc, DSS Inc.) was installed at all three VAMCs between November 2015 and March 2016. Within TheraDoc, each AS program designed alerts and interventions specific to its own medical center and collaborated to develop a group of 22 interventions that were shared by all sites. There was no attempt to alter individually determined AS practices of each VAMC.

**Results.** The table demonstrates the combined AS team interventions and aggregate cost-savings generated by the shared CSS. The number of interventions and the hours spent in CSS varied between centers; the top five most heavily conducted interventions differed between sites. Over time, new interventions developed at one VAMC were adopted at one or more of the other VAMCs at the discretion of the respective AS teams. Within a short period of time the cost savings generated exceeded the CSS start-up investment.

	CY 2016	CY 2017	Combined
AS interventions	2,946	10,016	12,962
Time spent AS PharmDs (hours)	472	2,032	2,503
Time spent AS MD (hours)	199	308	508
Cost savings	\$532,520	\$1,790,906	\$2,323,426

**Conclusion.** Working collaboratively allowed each VAMC to leverage shared resources. Each AS program was able to adapt to its VAMC's specific needs, while also demonstrating significant aggregate cost savings as a result of coordinated and defined AS activities.

**Disclosures.** M. Zuzick, Document Storage Systems: Employee, Salary. J. Chang, Document Storage Systems: Employee, Salary.

### 1868. Successful Use of Telemedicine vs. On-site Infectious Diseases Consultation After Implementation of a System-Wide Antimicrobial Stewardship-Led *Staphylococcus aureus* Bacteremia Care Bundle

Jacqueline Isip, PharmD<sup>1</sup>; Leigh Ann Medaris, MD<sup>2</sup>; Lewis McCurdy, MD<sup>2</sup>; Marc Kowalkowski, PhD<sup>3</sup>; Kristin Fischer, MM<sup>1</sup>; Jennifer Onsrud, PharmD<sup>1</sup> and Lisa Davidson, MD<sup>1,2</sup>; <sup>1</sup>Antimicrobial Support Network, Atrium Health, Charlotte, North Carolina, <sup>2</sup>Division of Infectious Diseases, Atrium Health, Charlotte, North Carolina, <sup>3</sup>Center for Outcomes Research and Evaluation, Atrium Health, Charlotte, North Carolina

**Session:** 222. Antimicrobial Stewardship: Potpourri  
Saturday, October 6, 2018: 12:30 PM

**Background.** Telemedicine (TM) programs have been effectively implemented to deliver specialty care through virtual platforms to overcome geographic and resource constraints. Yet, few data exist to describe outcomes associated with TM-based management of patients with infectious diseases (ID). The purpose of this study was to compare adherence and other outcomes associated with TM and on-site (SOC) ID consultation (IDC) implementation strategies of an antimicrobial stewardship (ASP)-led *S. aureus* bacteremia (SAB) bundle.

**Methods.** We launched an SAB bundle at 10 acute care hospitals in the metro Charlotte, NC area in September 2016 for adult patients admitted with SAB and conducted a retrospective cohort study using data collected through 2017. Bundle components included (1) mandatory IDC, (2) appropriate antibiotics within 24 hours of *S. aureus* speciation, (3) repeat blood cultures at least every 72 hours until clearance, (4) obtainment of an echocardiogram, and (5) appropriate duration of intravenous antibiotic therapy based on SAB severity. ASP facilitated bundle initiation and assisted with compliance for all patients. The primary outcome was bundle adherence. Secondary outcomes included time to culture clearance and persistent SAB (i.e., positive blood cultures for >7 days). We used Wilcoxon rank-sum and chi-squared tests to compare outcomes.

**Results.** We evaluated 872 patients with SAB during the study interval. After excluding 126 patients (prematurely discharged or died/transitoned to comfort care within 48 hours of *S. aureus* speciation), we analyzed 583 SOC and 163 TM group patients. There were no differences observed in overall SAB bundle adherence (SOC 86% vs. TM 88%,  $P = 0.52$ ), or its individual components. No differences were found in time to culture clearance (median days: SOC = 2.9 vs. TM = 2.8,  $P = 0.96$ ) and persistent SAB (SOC 11% vs. TM 11%,  $P = 0.77$ ).

**Conclusion.** Our findings provide preliminary evidence to support TM-based strategies for IDC and ASP-led care bundles in resource-limited settings. Future analyses will compare mortality and hospital readmission outcomes.




Figure 1. SAB Bundle Adherence, Overall and Separate Bundle Components

**Disclosures.** All authors: No reported disclosures.

### 1869. Telepharmacy Support of an Antimicrobial Stewardship Program in a Small Rural Canadian Acute Care Hospital

Sammu Dhaliwall, BSc. Pharm, Pharm D, ACPR<sup>1</sup> and Kevin McDonald, BSc Pharm<sup>2</sup>; <sup>1</sup>Northwest Telepharmacy Solutions, Brampton, ON, Canada, <sup>2</sup>Northwest Telepharmacy Solutions, Deep River, ON, Canada

**Session:** 222. Antimicrobial Stewardship: Potpourri  
Saturday, October 6, 2018: 12:30 PM


**Background.** In 2013 Accreditation Canada identified an Antimicrobial Stewardship Program (ASP) as a Required Organizational Practice in all acute care hospitals. A successful antimicrobial stewardship program is interdisciplinary and involves clinical pharmacists as a key member of the team. This small rural acute care hospital utilizes a telepharmacy model of care with a remote clinical pharmacist. This hospital is a 20-bed acute care hospital with no on-site clinical pharmacist and has been utilizing Telepharmacy services since 2008. To prepare for Accreditation in December 2013, the remote clinical pharmacist was requested to support and lead the antimicrobial stewardship program.

**Methods.** The remote clinical pharmacist performed a gap analysis to identify areas requiring improvement for a successful ASP which included the need for an Antimicrobial Stewardship committee, an IV to PO conversion program for antibiotics, development of guidelines and clinical pathways for common infections, hospital specific antibiogram, and prospective audit with intervention and feedback. The remote clinical pharmacist participated in meetings with nurses, physicians, and other key stakeholders using videoconference technology to develop a plan for the ASP. The remote clinical pharmacist started prospective data collection in September 2013. Starting January 2014, the remote clinical pharmacists started providing prospective audit with intervention and feedback.

**Results.** The antimicrobial stewardship program was formally accepted by Accreditation Canada in December 2013.

**Figure 1.** Days of Therapy (DOT) per 1,000 Patient-Days Over 4 Years

Figure 1: Day of Therapy per 1000 Patient Days



**Conclusion.** Small rural and remote acute care hospitals without access to an on-site clinical pharmacist can successfully implement and maintain an ASP by seeking support from experienced remote clinical pharmacists.

**Disclosures.** S. Dhaliwall, Northwest Telepharmacy Solutions: Employee, Salary. K. McDonald, Northwest Telepharmacy Solutions: Employee, Salary.

### 1870. Excess Antibiotic Duration in Patients Hospitalized for Pneumonia: A Multi-Hospital Cohort Study

Valerie Vaughn, MD, MSc<sup>1</sup>; Scott Flanders, MD<sup>1</sup>; Vineet Chopra, MD, MSc<sup>2</sup>; Anna Conlon, PhD<sup>3</sup>; Anurag Malani, MD, FIDSA<sup>4</sup>; Rama Thyagarajan, MD<sup>5</sup>; Lama Hsaiky, PharmD<sup>6</sup> and Tejal Gandhi, MD<sup>7</sup>; <sup>1</sup>Internal Medicine, University of Michigan, Ann Arbor, Michigan, <sup>2</sup>Patient Safety Enhancement Program of the Ann Arbor VA Medical Center, Ann Arbor, Michigan, <sup>3</sup>University of Michigan Health System, Ann Arbor, Michigan, <sup>4</sup>St. Joseph Mercy Health System, Ypsilanti, Michigan, <sup>5</sup>Internal Medicine/Infectious Disease, Beaumont Health- Dearborn, Dearborn, Michigan, <sup>6</sup>Beaumont Hospital, Dearborn, Michigan, <sup>7</sup>Division of Infectious Diseases, University of Michigan, Ann Arbor, Michigan

**Session:** 222. Antimicrobial Stewardship: Potpourri  
Saturday, October 6, 2018: 12:30 PM

Vaughn VM, Flanders SA, Chopra V, Conlon A, Malani A, Thyagarajan R, Hsaiky L and Gandhi T

**Background.** Despite guidelines recommending patients with pneumonia receive the shortest effective duration of antibiotic therapy, patients continue to receive antibiotics for longer than necessary. Patterns and predictors of excess antibiotic duration are not well understood.

**Methods.** Cohort study of patients hospitalized with community-acquired pneumonia (CAP) or healthcare-associated pneumonia (HCAP) between November 2015 and January 2018 at 48 hospitals participating in the Michigan Hospital Medicine Safety consortium. Adult, non-ICU, medical patients were included. Patients with concomitant infections, severe immunosuppression, or unusual pathogens were excluded. Detailed patient data, included duration of antibiotic treatment, were abstracted from the medical record. Appropriate duration was calculated for each patient based on time to clinical stability, organism, and comorbidities. Patient outcomes included mortality, readmission, length of stay, transfer to intensive care, and *Clostridium difficile* infection. Disease-, patient-, provider-, and hospital-factors associated with excess antibiotic duration and association of excess duration with patient outcomes were evaluated using multivariable generalized estimating equations models.