

again formulated (FVIII <1%, inhibitor titer 19 BU) and the patient was treated with the same anti-hemorrhagic and immunosuppressive treatment, reaching CR on day +20. In addition, the patient received anti-viral drugs (lopinavir-ritonavir 400 mg/twice daily) and non-invasive mechanical ventilation with oxygen as treatment for Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) infection, leading to the resolution of the disease on day +14.


In this case, the first reported in literature, the re-appearance of AHA was triggered by acute SARS-Cov-2 infection. This finding is particularly interesting considering the close interaction between SARS-Cov-2 and the hemostatic system.⁴ The AHA cases associated with other viral infections, including hepatitis B and C viruses, have also been described.³ The recent association between immune thrombocytopenic purpura and COVID-19 further strengthens the virus-induced immune dysregulation,⁵ which could have played a key role in the pathogenesis of AHA. We therefore recommend a daily coagulation monitoring (APTT) in patients admitted with acute COVID-19 infection and the inhibitor search (through a mixing test) in those cases with otherwise unexplained onset or worsening of the hemorrhagic picture and/or aPTT prolongation.

CONFLICT OF INTEREST

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

M.F., F.S. and M.P. designed and performed research; I.T., C.G. and B.C. analyzed data; M.F., C.D.F., C.P., G.D.D. and S.C. wrote the paper together; and all authors approved the final manuscript.

Massimo Franchini¹ , Claudia Glingani¹, Giuseppe De Donno², Salvatore Casari³, Beatrice Caruso⁴, Isabella Terenziani⁴, Cesare Perotti⁵, Claudia Del Fante⁵, Filippo Sartori⁶, Mauro Pagani⁶

¹Department of Hematology and Transfusion Medicine, Carlo Poma Hospital, ASST Mantova, Mantova, Italy

²Respiratory Unit, Carlo Poma Hospital, ASST Mantova, Mantova, Italy

³Unit of Infectious Diseases, Carlo Poma Hospital, ASST Mantova, Mantova, Italy

⁴Laboratory, Carlo Poma Hospital, ASST Mantova, Mantova, Italy

⁵Immunohaematology and Transfusion Service, Apheresis and Cell Therapy Unit, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

⁶Department of Medicine, Ospedale Destra Secchia, ASST Mantova, Mantova, Italy

Correspondence

Massimo Franchini, Department of Hematology and Transfusion Medicine, Carlo Poma Hospital, 46100 Mantova, Italy.

Email: massimo.franchini@asst-mantova.it

DOI 10.1002/ajh.25865

ORCID

Massimo Franchini  <https://orcid.org/0000-0002-8795-0580>

REFERENCES

1. Franchini M, Lippi G. Acquired factor VIII inhibitors. *Blood*. 2008;112:250-255.
2. Gibson CJ, Berliner N, Miller AL, Loscalzo J. Clinical problem-solving. A bruising loss. *N Engl J Med*. 2016;375:76-81.
3. Knoebl P, Marco P, Baudo F, et al. Demographic and clinical data in acquired hemophilia A: results from the European acquired hemophilia (EACH2) registry. *J Thromb Haemost*. 2012;10:622-631.
4. Terpos E, Ntanasis-Stathopoulos I, Elalamy I, et al. Hematological findings and complications of COVID-19. *Am J Hematol*. 2020;95(7):834-847.
5. Zulfiqar AA, Lorenzo-Villalba N, Hassler P, Andrés E. Immune thrombocytopenic Purpura in a patient with Covid-19. *N Engl J Med*. 2020;382:e43

Received: 14 April 2020 | Accepted: 15 April 2020

DOI: 10.1002/ajh.25840

SARS-CoV-2 infection in beta thalassemia: Preliminary data from the Italian experience

To the Editor:

Patients with pre-existent chronic morbidities are likely to be more severely affected by SARS-Cov2 infection, but no data are available regarding Thalassemic Syndromes (TS). Note, TS and hemoglobin variants represent, according to WHO, one of the most frequent causes of anemia, affecting more than 7% of the world population.¹ Thalassemic Syndromes are classified in either transfusion-dependent thalassemia (TDT) or non-transfusion-dependent thalassemia (NTDT). Infectious complications, mainly from bacteria, constitute a common cause of mortality and morbidity in TS. Stress erythropoiesis, iron overload, splenectomy and adrenal insufficiency among others may contribute to increase susceptibility to infection.²

To verify the impact of SARS-CoV-2 infection on TS, we set-up a specific survey by electronic Case Report Form (eCRF).³ Inclusion criteria require at least 15 days of follow-up from either the onset of symptoms or SARS-CoV2 positivity. The survey was approved by Ethics Committee and eCRF was shared with the Centers of Italian Hemoglobinopathies Network. The "Società Italiana Talassemie ed Emoglobinopatie" (SITE), has estimated the presence in Italy of approximately 5000 TDT and 1900 NTDT patients.³

As of 10 April 2020, 11 cases of TS and COVID-19 have been collected (see supplementary information). All the reported patients are in Northern Italy, where the rate of infection is higher, reflecting the national epidemiology.

The mean age is 44 ± 11 years (range 31-61 years) and 55% (6/11) are females. Ten patients are TDT, and one is NTDT. All the patients have thalassemia associated comorbidities, eight are splenectomized, and one patient (#9 in the supplementary table) has pulmonary hypertension treated with sildenafil. The likely source of

infection has been detected in 55% (6/11) of cases: two had contacts with COVID-19 positive subjects, and four had occupational exposure (three are nurses working in hospital or assisted living facilities).

Three patients were asymptomatic. One patient (#3 in supplementary information) was admitted for high fever and bone marrow hypoplasia, lymphopenia, and agranulocytosis (on treatment with defiriprone) and tested positive at the third swab. Six out of 11 were hospitalized, but no one required mechanical ventilation. The patient with more severe symptoms who required more intensive ventilation support with continuous positive airway pressure (CPAP) has a history of diffuse large B-cell lymphoma, treated with chemotherapy in the previous year, currently in complete remission. Of the six people admitted to the hospital, only three received supposedly specific treatment for COVID-19: one hydroxychloroquine (HCQ), one HCQ plus ritonavir/darunavir, and one HCQ plus anakinra. Patient #3 did not receive HCQ due to concomitant therapy with amiodarone and an increased risk of *life-threatening* arrhythmia. The clinical course ranged from 10 to 29 days. Ten patients have clinically recovered and are on a daily remote phone call follow-up. Splenectomy which was present in 8/11 patients did not seem to affect the clinical course. Of note, except for the patient with myelosuppression, no increase in blood requirement was observed. When luspatercept treatment was halted in the NTD patient, hemoglobin fell from 110 to 82 g/L, a value similar to the pre-luspatercept period. Neither death nor severe SARS or signs of cytokines storm were observed in these 11 subjects, which may be surprising, taking into account the mean age and the presence of severe comorbidities.





Our data, although preliminary, do not indicate increased severity of COVID-19 in TS. A larger number of cases needs to be collected to define the impact of this new infection and its outcome in these fragile patients.

ACKNOWLEDGEMENT

We would like to thank ALT (Associazione per la Lotta alla Talassemia R.Vullo - Ferrara).

CONFLICT OF INTEREST

The authors declare no competing financial interests.

Irene Motta^{1,2}, Margherita Migone De Amicis², Valeria M. Pinto³,
Manuela Balocco³, Filomena Longo⁴, Federico Bonetti⁵,
Barbara Gianesi⁶, Giovanna Graziadei² , Maria D. Cappellini¹ ,
Lucia De Franceschi^{7†} , Antonio Piga^{4†}, Gian L. Forni^{3†} 

¹Department of Clinical Sciences and Community Health, Università Degli Studi di Milano, Milan, Italy

²Department of Internal Medicine, UOC Medicina Generale, Fondazione IRCCS Ca' Granda, Ospedale Maggiore Policlinico, Milan, Italy

³Hemoglobinopathies and Congenital Anemia Center, Ospedale Galliera, Genoa, Italy

⁴Department of Clinical and Biological Sciences, University of Turin, Turin, Italy

⁵Pediatric Haematology Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

⁶ForAnemia Foundation, Genoa, Italy

⁷Department of Medicine, Policlinico GB Rossi, Università di Verona, Verona, Italy

Correspondence

Gian Luca Forni, MD, Centro della Microcitemia e delle Anemie Congenite, Ospedale Galliera, Via Volta 6, 16128 Genoa, Italy.
Email: gianluca.forni@galliera.it

Drs. De Franceschi, Piga and Forni contributed equally to this article.
DOI 10.1002/ajh.25840

ORCID

Giovanna Graziadei  <https://orcid.org/0000-0002-6801-5730>

Maria D. Cappellini  <https://orcid.org/0000-0001-8676-6864>

Lucia De Franceschi  <https://orcid.org/0000-0001-7093-777X>

Gian L. Forni  <https://orcid.org/0000-0001-9833-1016>

REFERENCES

1. Modell B. Global epidemiology of haemoglobin disorders and derived service indicators. *Bull World Health Organ.* 2008;86(6): 480-487.
2. Taher AT, Weatherall DJ, Cappellini MD. Thalassaemia. *Lancet.* 2018; 391(10116):155-167.
3. <http://www.site-italia.org/2020/covid-19.php>. SITE communication. Accessed April 1, 2020.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

Received: 13 April 2020 | Revised: 15 April 2020 | Accepted: 21 April 2020

DOI: 10.1002/ajh.25851

Management of CLL patients early in the COVID-19 pandemic: An international survey of CLL experts

To the Editor:

The coronavirus disease 2019 (COVID-19) pandemic caused by the novel coronavirus (SARS CoV-2) has posed an unprecedented health emergency, especially to those with older age or comorbid medical conditions.¹ Patients with chronic lymphocytic leukemia (CLL) have a median age of about 70 years at diagnosis, and the majority suffer