

Effects of acetic acid on the viability of *Ascaris lumbricoides* eggs

Is vinegar reliable enough to clean the vegetables?

Yunus E. Beyhan, DVM, PhD, Hasan Yilmaz, DVM, PhD, Murat Hokelek, MD, PhD.

ABSTRACT

الأهداف: للتحقيق من تأثير حمض الخليك على بيض *Ascaris lumbricoides* (A. lumbricoides) لتحديد التركيز الفعال للخل وفترة التنفيذ لجعل استهلاك الخضار النيئة أكثر أماناً.

الطريقة: أجريت هذه الدراسة التجريبية في مايو 2015م في مختبر علم الطفيليات بكلية الطب، جامعة يوزونجويل، اسطنبول، تركيا. تم تقسيم بيض *A. lumbricoides* إلى مجموعتين. تم علاج البيض في مجموعة الدراسة بنسبة تركيز 1، 3، 5، 10% من حامض الخليك، وتم علاج البيض في المجموعة المرجعية بيوزين. وتم متابعة قابلية تحمل البيض "في النقاط التالية خلال وقت التجربة: 0، 10، 15، 20، 30، 45، و 60 دقيقة.

النتائج: نسبة حمض الخليك 1% غير كافية لمقاومة بيض الإسكارس. في الدقيقة الـ 30، أظهر حمض الخليك 3% فعاليته 95%، وعند تركيز 5%، فقدت كل البيض حيويتها. يتطلب علاج حمض الخليك عند نسبة 4.8% في 30 دقيقة، أو نسبة 4.3% في 60 دقيقة لتحقيق النجاح الكامل من العلاج.

الخاتمة: يحتوي بيض الإسكارس على 3 طبقات مقاومة جداً، ويعتقد أن تركيز حمض الخليك، والتي يمكن أن تكون فعالة على البيض قد تكون فعالة أيضاً على مقاومة العديد من العوامل الطفيلية الأخرى. لتحقيق الحماية الضرورية، يفضل غسل الخضروات وتعريضها المباشر للخل محتويًا على حامض الخليك 5% لمدة 30 دقيقة.

Objectives: To investigate the effects of acetic acid on durable *Ascaris lumbricoides* (*A. lumbricoides*) eggs to determine the effective concentration of vinegar and the implementation period to render the consumption of raw vegetables more reliable.

Methods: This experimental study was performed in May 2015 in the Parasitology Laboratory, Faculty of Medicine, Yuzuncu Yil University, Van, Turkey. The *A. lumbricoides* eggs were divided into 2 groups. Eggs in the

study group were treated with 1, 3, 5, and 10% acetic acid concentrations, and eggs in the control group were treated with Eosin. The eggs' viability was observed at the following points in time during the experiment: 0, 10, 15, 20, 30, 45, and 60 minutes.

Results: The 1% acetic acid was determined insufficient on the viability of *Ascaris* eggs. At the 30th minute, 3% acetic acid demonstrated 95% effectiveness, and at 5% concentration, all eggs lost their viability. Treatment of acetic acid at the ratio of 4.8% in 30 minutes, or a ratio of 4.3% in 60 minutes is required for full success of treatment.

Conclusion: Since *Ascaris* eggs have 3 layers and are very resistant, the acetic acid concentration, which can be effective on these eggs are thought to be effective also on many other parasitic agents. In order to attain an active protection, after washing the vegetables, direct treatment with a vinegar containing 5% acetic acid for 30 minutes is essential.

Saudi Med J 2016; Vol. 37 (3): 288-292
doi: 10.15537/smj.2016.3.13061

From the Department of Parasitology (Beyhan, Yilmaz), Faculty of Medicine, Yuzuncu Yil University, Zeve Campus, Van, the Department of Microbiology (Hokelek), Faculty of Medicine, Istanbul University Cerrahpasa, Istanbul, Turkey.

Received 19th August 2015. Accepted 23rd December 2015.

Address correspondence and reprint request to: Dr. Yunus Emre Beyhan, Department of Parasitology, Faculty of Medicine, Yuzuncu Yil University, Zeve Campus, Van, Turkey. E-mail: yebeyhan@gmail.com

Parasitic diseases are witnessed all over the world, it originally emerges in certain areas where people have the habit of defecation in random places; where human stool is used as manure and infrastructure in undeveloped areas. Approximately one billion humans are infected with *Ascaris*, 800 million humans are

infected with *Trichuris*, 200 million humans are infected with *Taenia* in the world, and each year, millions of humans lose their lives due to parasitic infections.¹⁻⁴ Food borne diseases are receiving increasing attention recently.⁵ One of the important ways that humans are infected of these diseases is the consumption of raw, or poorly washed vegetables.⁶ Vegetables growing in soil and frequently used in salads, such as lettuce, parsley, radish, cress, garden rocket, green onion, carrot, and cucumber also play an important role in the transmission of parasitic infections, in addition to many bacterial and viral agents. *Giardia*, *Entamoeba*, *Cryptosporidium*, *Hymenolepis*, *Ascaris*, *Trichuris*, *Toxocara*, *Taenia* and *Fasciola* are the parasites on top of the list, which are transmitted by such vegetables.^{3,6-10} *Ascaris lumbricoides* (*A. lumbricoides*) is the largest nematode that lives in the human small intestine. The *A. lumbricoides* eggs are of 2 types: fertilized and unfertilized. Fertilized eggs have an oval shape and a shell of 3 thick layers. The outermost layer is a hackly protein layer, beneath this, is the semipermeable layer, and at the lowermost, there is a non-permeable lipoidal vitellin membrane layer. Fertilized eggs do not have this last layer and have no infection capacity, as well. This layered structure of the egg enables its durability against different chemical and environmental factors. Thus, it could be argued that *A. lumbricoides* eggs are more resilient than many other parasitic eggs. It has been reported that the *A. lumbricoides* eggs could carry their infectious disposition when preserved in a refrigerator for 2 years; and that they keep their viability for up to 40 days at temperatures between 18 and 27°C.¹¹⁻¹⁴ The objective of the study was to investigate the effects of acetic acid (CH₃CO₂H) on the viability of durable *A. lumbricoides* eggs, to determine the effective concentrations of vinegar, and the time required to treat raw vegetables for safe consumption.

Methods. In this experimental study, several *A. lumbricoides* eggs were counted in the stool sample, which was directed to at the Parasitology Laboratory, Faculty of Medicine, Yuzuncu Yil University, Van, Turkey in 2015 May. After the stool was diluted with 0.9% physiological saline solution (PSS) in order to purify it from particulates, it was strained with a sieve

of 150 µm. Later on, the sedimentation technique using PSS was applied to the stool sample until it became clear at room temperature, and the eggs were accumulated at the bottom residue. A 1, 3, 5, and 10% concentrations of pure commercial CH₃CO₂H (Sigma-Aldrich, St. Louis, MO, USA) were prepared. The *A. lumbricoides* eggs in the control group were examined only using 0.1% Eosin, and in the study group, they were treated with 4 different CH₃CO₂H concentrations (1%, 3%, 5%, and 10%) and Eosin. Selected eggs that are viable and embryonated was the inclusion criteria, and the contrary were the exclusion criteria. Viability survey of the eggs was conducted at the following times: 0, 10, 15, 20, 30, 45, and 60 minutes under x40 magnification and approximately 19.8 (11-25) eggs were counted for each group. The eggs which were colored were identified as dead, and eggs which were not colored, were considered alive (Figure 1). This research was carried out solely on fertilized eggs, the unfertilized eggs were ignored.

Statistical assessments were conducted using the Statistical Package for Social Sciences version 15 (SPSS Inc., Chicago, IL, USA) software and Probit Regression Analysis. The level of statistical significant differences was set at $p \leq 0.05$.

Results. It has been observed that after the 45th minute, 10% of the parasitic eggs lost their viability in the control group. The effect of 1% CH₃CO₂H was found to be insufficient, even in the 90th minute, only 35.3% of the eggs (6/17) lost their viability. While 3% CH₃CO₂H showed 95% effectiveness at the 30th minute, this ratio reached 100% at the 60th minute. A 5% concentration started to demonstrate an effect at the moment of application (21/24), and all eggs lost their viability at the 30th minute (Table 1). Probit analysis demonstrates that the difference between the results was significant ($p=0.001$). The “Probit Equality value = A* + 0.78 (B) x concentration” formula has been ascertained, and A* coefficients, which change according to time are presented in Table 2 (β =regression coefficient). The CH₃CO₂H concentrations of lethal dose (LD) 10, LD25, LD50, and LD99 that statistically kills the parasitic eggs based on time are presented in Table 3. In accordance with the method, it must be treated either with 4.8% CH₃CO₂H for 30 minutes, or with 4.3% CH₃CO₂H for 60 minutes to lose viability of all existing parasitic eggs (Table 3).

Discussion. Millions of people have been infected with parasitic agents in the world, and most of them live in underdeveloped, or developing countries. In our country, the parasitosis ratio is approximately 10-85%,

Disclosure. Authors have no conflict of interest, and the work was not supported or funded by any drug company.




Figure 1 - Photograph showing the A) dead eggs (colored eggs) and B) non-colored eggs considered as alive.

Table 1 - Colored (dead) and uncolored (alive) eggs count in the control and study groups according to time.

Time, minutes	Control group	Study group Acetic acid concentration			
		1%	3%	5%	10%
0	0/25	1/20	7/23	21/24	23/24
10	0/25	2/20	17/20	20/21	21/21
15	0/20	3/20	20/24	19/20	20/20
20	0/20	4/21	21/25	21/22	18/18
30	0/20	6/24	19/20	19/19	19/19
45	2/20	4/21	17/19	18/18	14/14
60	2/20	7/19	17/17	16/16	13/13
90	3/21	6/17	14/14	15/15	11/11

Table 2 - A* coefficient values depending on time by probit equality analysis in this study.

Time, minutes	A* coefficient
0	-2.930
10	-1.839
15	-1.746
20	-1.674
30	-1.391
45	-1.393
60	-1.036
90	-1.007

Table 3 - The effective concentrations of acetic acid to *Ascaris lumbricoides* eggs obtained by statistical method.

Time, minutes	Effective concentration (g/100 ml)			
	LD10	LD25	LD50	LD99
0	2.113	2.890	3.755	6.736
10	0.715	1.493	2.357	5.338
15	0.596	1.374	2.238	5.219
20	0.503	1.281	2.145	5.126
30	0.140	0.918	1.782	4.763
45	0.143	0.921	1.785	4.766
60	-0.315	0.463	1.488	4.308
90	-0.351	0.427	1.291	4.272

LD - lethal dose

although it varies based on the region and sanitary conditions.⁴

Diseases caused by contaminated food are one of the most common health problems in the world.^{3,15} Food borne parasitic diseases are not known much, but they are becoming more and more serious for human health.⁵ Vegetables and fruits have much significance in the diet of an individual, since they are sources of various vitamins and minerals. However, they carry the risk of transmission of diseases, since they could get contaminated by many bacterial, viral, and parasitic organisms.

In recent years, the number of reported parasitic infections is increasing.^{8,9} Globalization in food supply, the increasing number of international travels, increase in the population susceptible to diseases, changes in dietary habits, and the developments in the diagnostic methods are reported as reasons for the increase in parasitic diseases caused by food originating from all over the world.⁵ This situation poses a great risk for humans susceptible to parasitic diseases, who are old, and are immunosuppressive.¹⁶ The agricultural use of water infected with parasitic eggs, unhealthy draining of human stool, and use of animal stool as manure are the main reasons for the contamination of vegetables and fruits.¹⁻³ It was reported that 19 different food-borne parasitic outbreaks occurred in the United States between the years 1993 and 1997.¹⁷ Furthermore, it was reported that 10% of *Cryptosporidium* and *Giardia* infections, and 50% of *Toxoplasma* infections were originated food ingestion.¹⁸

Studies conducted across the world demonstrated that lettuce, parsley, cabbage, cress, mint, carrot, garden rocket, cucumber, pepper, radish, green onion, red beet, tomato, and similar raw vegetables are contaminated with parasitic cysts, oocysts, and eggs. In these plants, mostly *Toxocara*, *Taenia*, *Ascaris*, *Enterobius*, *Trichuris*, hooked worm (*Ancylostoma duodenale*, *Necator Americanus*) the eggs of *Hymenolepis nana*, cysts of *Giardia*, *Entamoeba*, *Balantidium*, and oocysts of *Cryptosporidium* were found.^{3,6-10,19,20} Studies conducted in different countries showed that the rate of contamination by parasitic agents were higher: 31.2% (156) of 500 samples in Pakistan,⁶ 66.3% in Morocco,²¹ 34.7% in Poland,⁹ 25% of the vegetables bought from bazaars, and 29% of those picked from the gardens in Iran,⁸ and 23.8% bought from bazaars and 36.9% picked from the gardens in Saudi Arabia.¹⁰ In Norway,¹⁶ 2% of vegetables were found positive with *Giardia*. Similar studies were also conducted in Turkey, and 3-14% of the vegetables were found to be contaminated.^{7,19,20,22} It is of vital importance that infected vegetables should be freed

from parasitic agents. Properly washing raw vegetables and fruits may remove the parasitic cysts, oocysts and eggs, however, vegetables with leaves are usually difficult to clean.² Thus, further precautions must be taken in addition to washing the vegetables. It is more important to implement this in sensitive environments, such as hospitals, schools, restaurants, and hotels where vegetables are consumed in large amounts.

It was noted according to the standards that the total acid content of vinegar (free CH₃CO₂H in water) must be no less than 4 g/100 mL. In a study conducted in Turkey²³ on 12 grape vinegar brands, the acetic acid content was identified to be between 3.96 and 5.36 g/100mL. It was found that in 65 different wine and apple vinegar samples, total acidity varied between 5.4 and 6.6 g/100mL in Italy.²⁴ Ravdin²⁵ suggested immersing vegetables into CH₃CO₂H or vinegar for 10-15 minutes for eradication of *Entamoeba histolytica* cysts. It was observed that this period was shorter than our findings. As mentioned above, this could be due to differences between the CH₃CO₂H content of foreign vinegar brands and vinegar brands in Turkey. Since the *A. lumbricoides* eggs have 3 layers and are durable, it is conceived that an CH₃CO₂H concentration, which might be effective on these eggs, could also be effective on other parasitic agents. It was ascertained that after raw vegetables are washed, vinegar with a 5% CH₃CO₂H content must be applied for at least for 30 minutes to ascertain an effective protection, and in cases where the acid content is lower, immersing time must be prolonged (Tables 1 & 3).

This study was limited to investigate the CH₃CO₂Hs viability also on other parasitic agents, but it is an advantage that the *A. lumbricoides* eggs are one of the most durable parasitic eggs. It is a common knowledge that vegetables must be immersed into a water-vinegar mixture for health reasons. Thus, a little amount of vinegar is mixed with large amounts of water to clean the vegetation, where the CH₃CO₂H gets diluted, and its concentrations are decreased by 10s/100s of times. Contrary to popular belief, vinegar must not be mixed with water; rather it must be treated directly on the vegetation after they are prewashed.

In conclusion, the public must be informed regarding the parasitic infections that might be contaminated from vegetables and fruits. It must never be forgotten to treat raw vegetables used in salads with vinegar after washing; this would be an effective protection against parasitic infections. As a result of this study, the effectiveness of acetic acid on the viability of *Ascaris*

eggs was demonstrated. These findings are meaningful for further studies that plan to use different medications for decontamination vegetables from parasitic agents.

Acknowledgment. The authors gratefully acknowledge Prof. Dr. Siddik Keskin for his contribution in the statistical analysis.

References

1. Molins RA, Motarjemi Y, Kaferstein FK. Irradiation: a critical control point in ensuring the microbiological safety of raw foods. *Food Cont* 2001; 12: 347-356.
2. Doyle ME. Foodborne Parasites: A Review of the Scientific Literature Review. *Fri Briefings*. Madison (WI): University of Wisconsin-Madison; 2003. Available from: https://fri.wisc.edu/files/Briefs_File/parasites.pdf
3. Erdogru O, Sener H. The contamination of various fruit and vegetable with *Enterobius vermicularis*, *Ascaris* eggs, *Entamoeba histolytica* cysts and *Giardia* cysts. *Food Cont* 2005; 16: 559-562.
4. Ozel MA, Ozbel Y, Ak M. Medical Parasitic Diseases. *Publications of Turkey Parasitology Association* 2007. p 3-76.
5. Dorny P, Praet N, Deckers N, Gabriel S. Emerging food-borne parasites. *Vet Parasitol* 2009; 163: 196-206.
6. ul-Haq S, Maqbool A, Khan UJ, Yasmin G, Sultana R. Parasitic Contamination of Vegetables Eaten Raw in Lahore. *Pakistan J Zool* 2014; 46: 1303-1309.
7. Ozan Z. Prevalence of helminth eggs on vegetables for salads. Dissertation. University of Afyon Kocatepe (Turkey): University of Afyon Kocatepe; 2007.
8. Daryani A, Ettehad GH, Sharif M, Ghorbani L, Ziaei H. Prevalence of intestinal parasites in vegetables consumed in Ardabil, Iran. *Food Cont* 2008; 19: 790-794.
9. Klapek T, Borecka A. Contamination of vegetables, fruits and soil with geohelminths eggs on organic farms in Poland. *Ann Agric Environ Med* 2012; 19: 421-425.
10. Ammar AS, Omar HM. The prevalence of leafy vegetable-borne Parasites in Al-Qassim Region, Saudi Arabia. *J Agric Vet Sci* 2013; 6: 29-40.
11. World Health Organization. Integrated Guide to Sanitary Parasitology. World Health Organization Regional Office for the Eastern Mediterranean Regional Centre for Environmental Health Activities. Amman (Jordan): World Health Organization; 2004.
12. Roberts AD. Ascariasis: Introduction and Epidemiology and Transmission. In: Satoskar AR, Simon GL, Hotez PJ, Tsuji M, editors. *Medical Parasitology*. Austin (Texas): Landes Bioscience; 2009. p. 14-20.
13. Cuomo MJ, Noel, LB, White DB. Diagnosing Medical parasites: A Public Health Officers Guide to Assisting Laboratory and Medical Officers. Randolph (USA): Air Education and Training Command; 2012.
14. Ridley JW. Parasitology for Medical and Clinical Laboratory Professionals. New York (NY): Cengage Learning; 2012.
15. World Health Organization. The Role of Food Safety in Health and Development. WHO Technical Report Series No. 705. Geneva (CH): World Health Organization; 1994.
16. Robertson LJ, Gjerde B. Isolation and enumeration of *Giardia* cysts, *Cryptosporidium* oocysts, and *Ascaris* eggs from fruits and vegetables. *J Food Protect* 2000; 63: 775-778.
17. Slifko TR, Smith HV, Rose JB. Emerging parasite zoonoses associated with water and food. *Int J Parasitol* 2000; 30: 1379-1393.

18. Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, et al. Food-Related Illness and Death in the United States. *Emerg Infect Dis* 1999; 5: 607-625.
19. Ulukanligil M, Seyrek A, Aslan G, Ozbilge H, Atay S. Environmental pollution with soil-transmitted helminths in Sanliurfa, Turkey. *Mem Inst Oswaldo Cruz* 2001; 96: 903-909.
20. Kozan E, Gonenc B, Sarimehmetoglu O, Aycicek H. Prevalence of helminth eggs on raw vegetables used for salads. *Food Cont* 2005; 16: 239-242.
21. Hajjami K, Ennaji MM, Amdiouni H, Fouad S, Cohen N. Parasitic contamination on fresh vegetable consumed in Casablanca city (Morocco) and risk for consumer. *Int J Sci Technol* 2013; 2: 543-549.
22. Avcioglu H, Soykan E, Tarakci U. Control of helminth contamination of raw vegetables by washing. *Vector Borne Zoonot Dis* 2011; 11: 189-191.
23. Akbas M, Cabaroglu T. A research on the determination of compositions of grape vinegars produced in Turkey and their conformity to food legislation. *Gida* 2010; 35: 183-188.
24. Gerbi V, Zeppa G, Beltramo R, Carnacini A, Antonelli A. Characterization of white vinegars of different sources with artificial neural networks. *J Sci Food Agric* 1998; 78: 415-425.
25. Ravdin JI, Stauffer W. *Entamoeba histolytica (Amebiasis)*. In: Mandell GI, Bennett J, Dolins R, editors. Principles and Practice of Infectious Diseases. 6th ed. Philadelphia (PA): Elsevier Churchill Livingstone; 2005. p. 3107-3108.

Student Corner

We invite students from a variety of medical disciplines to submit original contributions based on their supervised research.

The Student Corner of Saudi Med J aims to help students explore research opportunities and network with other peers and mentors in the same field.

Submission Guidelines

Submitted Abstracts should include the following:

- Title should be descriptive
- Author's names and affiliation (specify college level/year, academic degree of Senior Author)
 - Abstract must be structured and not more than 300 words
 - The following are the typical headings:

Objectives (background, why the study was done, specific aims)

Methods (setting, date of study, design, subjects, intervention and analysis)

Results (findings, data and statistical tests) and

Conclusion (general interpretation of results)

General Information on Abstract Submission

Submitted Abstracts should be co-authored by a Senior Supervisor

Abstracts will be reviewed by Student's Corner Section Editor

There is no fee to submit an Abstract

Ethical Approval should be provided

Non-indexed materials