


RESEARCH ARTICLE

Open Access



# Correlation of patient characteristics with arm and finger measurements in Asian parturients: a preliminary study

Ming Jian Lim<sup>1</sup>, Chin Wen Tan<sup>1,2</sup>, Hon Sen Tan<sup>1</sup>, Rehana Sultana<sup>3</sup>, Victoria Eley<sup>4</sup> and Ban Leong Sng<sup>1,2\*</sup> 

## Abstract

**Background:** Accurate blood pressure (BP) measurement depends on appropriate cuff size and shape in relation to the arm. Arm dimensions outside the recommended range of cuff sizes or trunco-conical arms may result in inaccurate BP measurements. Measuring BP using finger cuffs is a potential solution. Arm cuff size is based on mid-arm circumference (MAC), and trunco-conicity is quantified by conicity index. We aimed to determine the correlation of MAC, body mass index (BMI), and weight with conicity index.

**Methods:** A prospective cohort study was conducted in the KK Women's and Children's Hospital where third trimester parturients scheduled for cesarean delivery were recruited after obtaining informed consent. Parturients were asked to rate their experience with time taken to obtain BP readings, cuff popping off during measurement, need to move the cuff from the upper arm to lower arm or leg, and need to change to a different cuff. Our primary outcome was the correlation between MAC and conicity index, calculated using Pearson's correlation. The correlation between BMI and weight with conicity index was also determined.

**Results:** We enrolled 300 parturients. Moderate correlation was found between left MAC and left conicity index ( $r = 0.41$ , 95% CI 0.32 to 0.51), and right MAC and right conicity index ( $r = 0.39$ , 95% CI 0.29 to 0.48). Weight ( $r = 0.35$  to 0.39) and BMI ( $r = 0.41$  to 0.43) correlated with conicity index in this study. MAC of 1 parturient fell outside the recommended range for arm cuffs, but all parturients fit into available finger cuffs. Obese parturients had increased problems with arm cuffs popping off and needing a change of cuff.

**Conclusions:** BMI better correlated with conicity index compared to MAC or weight. Standard finger cuffs were suitable for all parturients studied and may be a suitable alternative.

**Trial registration:** Clinicaltrials.gov [NCT04012151](https://clinicaltrials.gov/ct2/show/study/NCT04012151). Registered 9 Jul 2019

**Keywords:** Body weight, Blood pressure, Obstetrics

\* Correspondence: [sng.ban.leong@singhealth.com.sg](mailto:sng.ban.leong@singhealth.com.sg)

<sup>1</sup>Department of Women's Anesthesia, KK Women's and Children's Hospital, 100 Bukit Timah Road, Singapore 229899, Singapore

<sup>2</sup>Duke-NUS Medical School, 100 Bukit Timah Road, Singapore 229899, Singapore

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Accurate measurement of blood pressure (BP) is important in peripartum care [1]. Mercury sphygmomanometers or hybrid devices with cylindrical arm cuffs remain the gold standard for BP measurement [2], however, poorly fitting arm cuffs can yield inaccurate BP readings, potentially leading to inappropriate clinical management [3].

Accurate BP measurements require the cuff to fit the parturient's arm in two respects; appropriate cuff size in relation to the parturient's arm circumference, and a cuff shape that matches that of the parturient's arm [4]. Cuff size is often based on the mid-arm circumference (MAC), defined as the circumference at the mid-point of the upper arm, with appropriately-sized cuff bladder dimensions being 40% width and 80% length of the MAC, respectively [5–7]. However, sizing cuffs using these criteria overestimate BP in patients with small MAC, while underestimating BP in patients with higher MAC [8]. This issue may be magnified in obese parturients, given that appropriately sized cuffs are not always readily available. A study conducted on 450 Australian parturients reported that 12.9 and 1.3% of parturients requiring “large” and “thigh” cuffs, respectively, while none of the available cuff sizes produced an acceptable fit in 3.1% of parturients and required cuff placement on the forearm or leg [3]. The erroneous use of inappropriately small cuffs in obese parturients overestimates BP [9, 10], with potential risk for significant adverse outcomes [11].

Additionally, the shape of the cuff should match that of the arm. In most individuals, the shape of the upper arm has been described as trunco-conical (truncated cone) instead of cylindrical [4]. The degree of trunco-conicity varies depending on gender and obesity [4], and can be mathematically described using a conicity index derived from arm length, proximal and distal arm circumferences [12]. In parturients with higher conicity indices, the large gap between the BP cuff and the surface of the distal arm [9, 12] results in irregular expansion of the cylindrical BP cuff during inflation and overestimation of systolic BP by up to 9.7 mmHg and diastolic BP by up to 7.8 mmHg [13]. Again, this issue may be exacerbated in obese patients, as BMI is moderately correlated with arm conicity index ( $r = 0.51$ ,  $r^2 = 0.26$ ) [3].

A potential solution to inappropriately sized cuffs or trunco-conical arms may lie with the use of finger cuff devices such as the Nexfin™ monitor (BMEYE B.V., Holland), CNAP™ monitor (CNSystems Medizintechnik AG, Graz, Austria), and ClearSight™ (Edwards Lifesciences, Irvine, California, USA). In comparison to arm cuffs, finger cuffs from ClearSight™ accommodated up to finger circumference of 6.8 cm and failed to fit only 0.7% of Australian parturients with maximum finger circumference of 7 cm [3]. In contrast, CNAP™ finger cuff device accommodates

finger circumferences up to 8.8 cm and would fit all parturients in the study cohort.

Although the issue of inappropriate cuff size or shape for BP measurement have been established in the Caucasian population, it is yet unclear if this is applicable to Asian parturients. The MAC in Asian parturients have not been determined; a study by Wang et al. found no differences in MAC of non-pregnant Caucasians and Asians [14], but it is uncertain if the available range of arm cuffs will fit Asian parturients. Second, because arm conicity index measurements and its association with BMI, MAC, and weight have not been established in Asian parturients, we are uncertain if the issue of trunco-conicity applies to Asian parturients and how it may vary according to parturient characteristics. Finally, the suitability of available finger cuff sizes in Asian parturients has not been established. Hence, this prospective cohort study aimed to determine MAC within our Asian parturient population and its correlation with conicity index. Secondly, we investigated the association between BMI and weight with arm conicity index, as well as the ability of finger cuffs from Nexfin™ and CNAP™ to fit the finger circumferences of Asian parturients.

## Methods

This study adheres to the applicable Strengthening the Reporting of Observational studies in Epidemiology (STROBE) guidelines [15]. Informed consent was obtained from all parturients, and ethics approval was obtained from the SingHealth Centralized Institutional Review Board (Reference number: CIRB 2019/2290) and registered on Clinicaltrial.gov (ID: NCT04012151).

This was a prospective cohort study in which parturients scheduled for elective cesarean delivery were enrolled at the pre-anesthetic clinic in KK Women's and Children's Hospital (KKH) from July to December 2019. Consecutive eligible parturients were enrolled. Eligible parturients were between 21 to 50 years old, at 32 or more weeks of gestation, and American Society of Anesthesiology (ASA) Physical Status I to III. Parturients undergoing unscheduled cesarean delivery were excluded.

Information on baseline demographic and obstetric characteristics were obtained from the patients or their medical records on admission for surgery. BMI was calculated from the parturient's height and weight at enrollment. Measurements from both upper limbs were taken according to standard anthropometry protocols when available [16]. Arm length was measured from the tip of the acromion process to the tip of the olecranon process on the posterior aspect of the arm with the elbow flexed [16]. With the arm hanging by the side, MAC was measured at the mid-point of the arm length

[16], the proximal arm circumference was measured at the axilla, and the distal arm circumference was measured at the elbow above the elbow crease. Finger circumference was measured at the mid-point of the middle phalanx of the middle finger, with the hand placed flat on a table. As described by Bonso et al. [12], arm conicity index was calculated as:

$$\frac{\text{Proximal arm diameter} - \text{Distal arm diameter}}{\text{Arm length}} \times 100$$

Arm diameter was calculated by dividing the relevant arm measurement by  $\pi$ . The appropriate arm or finger cuff size was determined based on MAC and finger circumference respectively, in accordance with manufacturers' recommendations. Patients also responded to a three-point rating scale (never/sometimes/always) about their experience of the procedure for BP measurements during the current pregnancy, with questions asking about extended waiting period to obtain a reading, whether the cuff pops off during measurement, the need to take BP on the lower arm or the leg, and whether they needed to change to a different cuff.

#### Sample size calculation and statistical analysis

A sample of 300 parturients was adequate to estimate a Pearson's correlation coefficient of at least 0.4 with a 95% CI width of 0.2, based on the correlation results reported by a previous study methodology on a Caucasian population [3].

Our primary outcome was the correlation between MAC and conicity index, which was reported using Pearson's correlation with a 95% confidence interval (CI) for each arm. Univariate linear regression was used to analyze the quantitative association between conicity index and MAC, BMI, and weight. Association between conicity index and other variables were expressed as  $\beta$  estimate with its 95% CI. Parturient characteristics, arm anthropometric data and patient's experience on BP measurement was summarized according to obesity status (BMI < 30 kg/m<sup>2</sup> as non-obese; BMI  $\geq$ 30 kg/m<sup>2</sup> as obese). Categorical and continuous variables were summarized as frequency (proportions) and as mean (standard deviation (SD), minimum - maximum). Univariate logistic regression model was fitted to determine associations between each of patient's experience survey questions with obesity status. Association from logistic regression analysis was expressed as odds ratio (OR) with 95% CI. A statistical significance was set at  $p$ -value < 0.05. All the analysis was done using SAS 9.4 software.

#### Results

We screened 377 parturients, of whom 77 refused consent. Hence, 300 parturients were enrolled, with their

**Table 1** Parturient and gestational characteristics

	Non-obese <sup>a</sup> N = 194	Obese <sup>a</sup> N = 106	Total N = 300
Age (years)	33.9 ± 4.4	33.7 ± 3.7	33.8 ± 4.2
Gestational age (weeks)	38.2 ± 1.1	38.1 ± 1.0	38.2 ± 1.1
Height (m)	1.6 ± 0.1	1.6 ± 0.1	1.6 ± 0.1
Weight (kg)	66.1 ± 8.0	86.5 ± 11.4	73.3 ± 13.5
BMI (kg/m <sup>2</sup> )	26.2 ± 2.5	34.5 ± 4.0	29.1 ± 5.1
Race / ethnicity			
Chinese	119 (61.3)	48 (45.3)	167 (55.7)
Malay	36 (18.6)	34 (32.1)	70 (23.3)
Indian	15 (7.7)	14 (13.2)	29 (9.7)
Others	24 (12.4)	10 (9.4)	34 (11.3)
ASA physical status			
I	98 (50.5)	32 (30.2)	130 (43.3)
II	95 (49.0)	62 (58.5)	157 (52.3)
III	1 (0.5)	12 (11.3)	13 (4.3)
Chronic hypertension	3 (1.5)	2 (1.9)	5 (1.7)
Gestational hypertension <sup>b</sup>	6 (3.1)	5 (4.7)	11 (3.7)
Gestational diabetes	18 (9.3)	14 (13.2)	32 (10.7)

Values are represented as mean ± standard deviation (SD) or number (%)

ASA American Society of Anesthesiologists, BMI body mass index

<sup>a</sup> Obese and non-obese groups were defined as BMI  $\geq$  30 kg/m<sup>2</sup> and

BMI < 30 kg/m<sup>2</sup> respectively

<sup>b</sup> Gestational hypertension defined as having a blood pressure  $\geq$  140/90 on two separate occasions more than 6 h apart, without proteinuria and diagnosed after 20 weeks of gestation

characteristics and gestational information shown in Table 1. Of the 300 parturients, 194 (64.7%) were non-obese while 106 (35.3%) were obese. The mean (SD) BMI in non-obese and obese group parturients was 26.2 (2.5) kg/m<sup>2</sup> and 34.5 (4.0) kg/m<sup>2</sup> respectively. Arm and finger measurements, along with conicity indices are summarized in Table 2.

#### Correlation between MAC and arm conicity indices

Our primary objective was to determine the correlation between MAC and arm conicity indices. The correlation between the left MAC and left arm conicity index ( $r = 0.41$ , 95% CI 0.32–0.51) was similar to the right MAC and right arm conicity index ( $r = 0.39$ , 95% CI 0.29–0.48).

#### Association between BMI, weight, and MAC with conicity indices

We found a moderate correlation between MAC, weight, and BMI with their respective conicity indices (Table 3). Of the three variables, BMI correlated the best with conicity index for both arms (left:  $r = 0.43$ ; right:  $r = 0.41$ ). Due to significant collinearity between MAC, weight, and BMI (collinearity indices > 0.8), three separate univariate linear regression models were performed, which showed

**Table 2** Arm and finger measurements

	Non-obese <sup>a</sup> N = 194	Obese <sup>a</sup> N = 106	Total N = 300
Right arm measurements (cm)			
Length	33.9 ± 2.2 [28, 40]	34.4 ± 2.5 [26, 41]	34.1 ± 2.3 [26, 41]
Proximal circumference	29.1 ± 3.3 [20, 38]	35.0 ± 5.0 [14, 49]	31.4 ± 4.9 [14, 49]
MAC	25.5 ± 2.6 [20, 36]	31.4 ± 4.0 [15, 43]	27.6 ± 4.2 [15, 43]
Distal circumference	23.7 ± 2.3 [19, 34]	28.7 ± 3.6 [13, 40]	25.4 ± 3.7 [13, 40]
Left arm measurements (cm)			
Length	33.9 ± 2.2 [27, 41]	34.4 ± 2.4 [26, 40]	34.1 ± 2.3 [26, 41]
Proximal circumference	29.1 ± 3.3 [19, 38]	35.5 ± 4.5 [14, 49]	31.2 ± 4.9 [14, 49]
MAC	25.5 ± 2.7 [19, 37]	31.4 ± 4.0 [15, 43]	27.6 ± 4.3 [15, 43]
Distal circumference	23.7 ± 2.4 [19, 35]	28.7 ± 3.6 [13, 41]	25.5 ± 3.7 [13, 41]
Right arm conicity index (%)	5.1 ± 2.0 [1, 12]	6.4 ± 2.3 [1, 14]	5.6 ± 2.2 [1, 14]
Left arm conicity index (%)	4.8 ± 1.9 [0, 10]	6.2 ± 2.0 [1, 12]	5.3 ± 2.1 [0, 12]
Right finger circumference (cm)	5.2 ± 0.3 [5, 6]	5.5 ± 0.4 [5, 7]	5.3 ± 0.4 [4, 7]
Left finger circumference (cm)	5.1 ± 0.3 [4, 6]	5.5 ± 0.3 [5, 7]	5.2 ± 0.4 [4, 7]

Values are represented as mean ± standard deviation (SD) [min, max]

BMI body mass index; MAC mid-arm circumference

<sup>a</sup> Obese and non-obese groups were defined as BMI ≥ 30 kg/m<sup>2</sup> and BMI < 30 kg/m<sup>2</sup> respectively

significant associations between BMI, weight, and MAC with the respective conicity indices. Each unit increase in BMI increased both arm conicity indices by 0.18.

**Distribution of recommended arm and finger cuff sizes according to MAC**

Based on MAC, the frequency distribution of arm cuff sizes according to American Heart Association (AHA) recommendations are illustrated in Fig. 1. One

**Table 3** Univariate linear regression analysis describing the associations between BMI, weight, and MAC with conicity indices

	β (95% CI)	r (95% CI) <sup>a</sup>	p-value
Right conicity index			
MAC	0.20 (0.25–0.26)	0.39 (0.29–0.48)	< 0.01
Weight	0.06 (0.04–0.07)	0.35 (0.25–0.45)	< 0.01
BMI	0.18 (0.13–0.22)	0.41 (0.31–0.50)	< 0.01
Left conicity index			
MAC	0.20 (0.15–0.25)	0.41 (0.32–0.51)	< 0.01
Weight	0.06 (0.04–0.07)	0.39 (0.29–0.48)	< 0.01
BMI	0.18 (0.13–0.22)	0.43 (0.34–0.52)	< 0.01


BMI body mass index, CI confidence interval, MAC mid-arm circumference

<sup>a</sup> Pearson correlation coefficient

parturient had left and right arm circumferences of less than 17 cm, which was below the recommended range of the smallest arm cuff. None of the parturients had MAC above the range of the largest arm cuff. Similarly, fitting of the Nexfin™ or CNAP™ finger cuffs were based on finger circumferences as recommended by the manufacturer (Fig. 2). One parturient had a left finger circumference that was less than 43 mm, which was below lower range of the smallest Nexfin finger cuff. However, this parturient’s right finger circumference was 45 mm, which fell within the recommended finger cuff range. In contrast, none of the parturients’ finger circumferences fell outside the recommended measurements for CNAP™.

**Parturient experience survey**

Obese parturients were more likely to experience the arm cuff popping off (OR 10.18, 95% CI 4.04–25.69) or needing a cuff change (OR 9.72, 95% CI 3.19–29.56), compared to non-obese parturients. In addition, the parturient experience survey found that 28 (9.3%) parturients experienced an extended waiting period while taking a BP measurement, and 4 (1.3%) parturients required the cuff to be put on the lower arm or leg, of whom 3 (2.8%) were obese (Table 4).




**Discussion**

Our results showed moderate correlation between MAC and arm conicity index in third trimester parturients. Furthermore, although MAC, weight, and BMI showed moderate correlation with arm conicity, BMI had the highest correlation coefficient with arm conicity, with every unit increase in BMI increasing the conicity index by 0.18. Out of 300 parturients, one had MAC below the recommended range for arm cuffs. However, all parturients were able to fit into one of the standard Nexfin™ and CNAP™ finger cuffs. Compared to non-obese parturients, obese parturients experienced significantly more problems with BP measurement, including arm cuffs popping off, or requiring a change in BP cuffs.

Our finding that MAC is moderately correlated with conicity index is consistent with other studies of Caucasian parturients, suggesting that this correlation applies

to our multi-ethnic Asian population. Bonso et al. were the first to mathematically quantify arm conicity using a conicity index [12], and despite using non-standard anthropometry techniques and arm length measurements, Bonso et al. and Palatini et al. demonstrated that MAC was correlated with arm conicity index [12, 13]. Subsequently, Eley et al. increased the robustness and reproducibility of arm length measurements by utilizing bony landmarks based on the Anthropometry Procedures Manual of the Center for Disease Control [3, 16], and reported a similar correlation between MAC and conicity index [3].

In addition to MAC, BMI and weight were also correlated with conicity index, and of the three variables, BMI was found to correlate the best with arm conicity. These results are similar to that of Eley et al., who also reported that BMI correlated best with arm conicity,



**Table 4** Parturient satisfaction survey regarding blood pressure measurements

Question	Non-obese <sup>a</sup> (N = 194)	Obese <sup>a</sup> (N = 106)	Unadjusted odds ratio (95% CI)	p-value
Long time taken for blood pressure reading	19 (9.8)	9 (8.5)	0.86 (0.37–1.96)	0.71
Cuff pops off	6 (3.1)	26 (24.5)	10.18 (4.04–25.69)	< 0.01
Need to put cuff on lower arm or leg	1 (0.5)	3 (2.8)	5.62 (0.58–54.70)	0.14
Need to change to different cuff	4 (2.1)	18 (17.0)	9.72 (3.19–29.56)	< 0.01

Values are represented as number (%)

BMI body mass index, CI confidence interval

<sup>a</sup> Obese and non-obese groups were defined as BMI  $\geq 30$  kg/m<sup>2</sup> and BMI < 30 kg/m<sup>2</sup> respectively

accounting for 26% of the variation in conicity index [3]. Furthermore, our finding that BMI, weight, and MAC were collinear was also reported by Eley et al. [3]. This may be explained by the association between increasing BMI and higher MAC in both pregnant [3, 17, 18] and non-pregnant populations [19], and is likely due to an increase in arm fat mass as measured using bioelectrical impedance analysis [17]. Additionally, obese parturients were more likely to experience challenges during BP measurement, with an increased incidence of the arm cuffs popping off and the need to change to a different cuff, compared to non-obese parturients. These findings were echoed by Eley et al. [3].

Our findings that obesity is associated with increasing MAC and arm conicity is concerning. Although the highest MAC measured in our study did not exceed the recommended MAC range of the largest cuff, these non-standard cuff sizes may not be readily available, and shifting the point of measurement from the upper arm to lower arm is unacceptable as it has been shown to overestimate BP [20–22]. Furthermore, the increase in conicity index may cause irregular expansion of the cylindrical cuff during inflation, leading to overestimation of both systolic and diastolic BP [13]. Although the use of cone-shaped BP cuffs may improve the accuracy of BP measurements compared to cylindrical cuffs [23], the former is not available at our institution. Nonetheless, the increased risk of overestimating BP in obese parturients is clinically significant given that obese parturients are already at elevated baseline risk for hypertensive disorders [2, 13], with the prevalence of obesity in Singapore likely to increase further [24].

The potential solution to the increased MAC and conicity index of obese parturients may lie with the use of finger cuff devices like the Nexfin™ and CNAP™. Our results demonstrated that all parturients were able to fit into the standard finger cuffs provided by Nexfin™ and CNAP™, which may provide an alternative means of measuring BP in parturients who do not fit the available arm cuffs. Nexfin™ has been validated against sphygmomanometry in pregnant

patients, and passed both Association for the Advancement of Medical Instrumentation (AAMI) standards and European Society of Hypertension International Protocol Phases 1 and 2.1 [25]. Similarly, CNAP™ has been validated against intra-arterial BP in general surgery patients under anesthesia and met AAMI standards [26]. In contrast, the accuracy of finger cuff-based BP measurements were unacceptable by AAMI standards in critically ill patients receiving vasopressor infusions or with finger edema [27], and should not be used as an alternative to arm cuffs in this patient population. However, our standard practice for such patients is intra-arterial BP monitoring.

We acknowledge several limitations to this study. Since BP measurements using arm or finger cuff-based systems were not recorded and compared against another standard such as intra-arterial BP, we are unable to assess the implications of increasing MAC or conicity index on BP measurement accuracy, and this should be determined in future studies. This information will help determine if finger cuff-based BP measurements should be used in parturients with high arm conicity or who do not fit standard arm cuffs. In addition, we enrolled parturients who were scheduled for elective cesarean delivery, which may not be representative of the general obstetric population since parturients with higher BMI are more likely to undergo cesarean delivery [3, 28]. Finally, we enrolled parturients in their third trimester, raising concerns that gestational weight gain during the interval prior to cesarean delivery will lead to increased BMI and MAC. However, Hogan et al. studied MAC in parturients across all three trimesters and found that the mean MAC did not vary significantly with different trimesters of pregnancy [17].

## Conclusions

In summary, our study reported that BMI is better correlated with arm conicity index, compared to MAC or weight. Obese parturients are at increased risk of having issues during BP measurement, such as cuffs popping off

or requiring a change of cuff. Validated finger cuff devices such as the Nexfin™ and CNAP™ may be an alternative means for BP measurement in patients who are unsuited to traditional arm cuffs. Further studies are required to assess the implications of increasing MAC and arm conicity on BP measurement accuracy.

#### Abbreviations

AAMI: Association for the Advancement of Medical Instrumentation; AHA: American Heart Association; ASA: American Society of Anesthesiologists; BMI: Body mass index; BP: Blood pressure; CI: Confidence interval; MAC: Mid-arm circumference; OR: Odds ratio; SD: Standard deviation; STROBE: Strengthening the Reporting of Observational studies in Epidemiology

#### Acknowledgments

The authors would like to thank Ms. Agnes Teo (Senior Clinical Research Coordinator) and Michelle Ren (Clinical Research Coordinator) for her administrative support during this study.

#### Authors' contributions

MJL reviewed the literature, acquired funding, planned the study, oversaw patient recruitment, data analysis and interpretation, and wrote the manuscript. CWT reviewed the literature, helped in funding, oversaw data management, data analysis and interpretation. HST reviewed the literature, helped in the study design, performed data analysis and interpretation. SR reviewed the literature, helped in the study design, performed data analysis and interpretation. VE reviewed the literature, planned the study, reviewed the data analysis and interpretation and oversaw the study. SBL reviewed the literature, planned the study, and oversaw the study including the design, data analysis and interpretation. All authors approved the final version of the manuscript, and agree to be accountable for all aspects of this work.

#### Funding

This work was supported by the funding from the SingHealth Duke-NUS Anesthesiology and Perioperative Sciences Academic Clinical Program Pilot Research Grant 2019 (Grant No. ANAESPRG/02). The aforementioned sponsor was not involved in the study activities.

#### Availability of data and materials

The datasets generated and analyzed in this work are available for anyone who wishes to access the data by contacting the corresponding author.

#### Ethics approval and consent to participate

The study was approved by the SingHealth Centralized Institutional Review Board, Singapore (SingHealth CIRB Ref: 2019/2290), and registered on Clinicaltrials.gov (NCT04012151). Informed consent was obtained from all participants in written format. The authors declare that all the recruited patients provided informed consent, and that this work was conducted in accordance with the Declaration of Helsinki.

#### Consent for publication

Not applicable.

#### Competing interests

Ban Leong Sng is an associate editor of *BMC Anesthesiology*. All other authors report no conflicts of interest in this work.

#### Author details

<sup>1</sup>Department of Women's Anesthesia, KK Women's and Children's Hospital, 100 Bukit Timah Road, Singapore 229899, Singapore. <sup>2</sup>Duke-NUS Medical School, 100 Bukit Timah Road, Singapore 229899, Singapore. <sup>3</sup>Centre for Quantitative Medicine, Duke-NUS Medical School, Singapore, Singapore. <sup>4</sup>Department of Anesthesia and Perioperative Medicine, The Royal Brisbane and Women's Hospital, Brisbane, Australia.

Received: 29 June 2020 Accepted: 20 August 2020

Published online: 31 August 2020

#### References

1. Knight M, Clarke B, James R, et al. on behalf of MBRRACE-UK. Saving Lives, Improving Mothers' Care - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2015-17. Oxford: National Perinatal Epidemiology Unit, University of Oxford 2019.
2. Lowe SA, Bowyer L, Lust K, et al. The SOMANZ guidelines for the Management of Hypertensive Disorders of pregnancy 2014. *Aust N Z J Obstet Gynaecol.* 2015;55:11-6.
3. Eley VA, Roberts L, Rickards L, et al. Arm and finger measurements in the third trimester: implications for blood pressure measurement. *Pregnancy Hypertension.* 2018;14:105-9.
4. Palatini P, Asmar R. Cuff challenges in blood pressure measurement. *J Clin Hypertens (Greenwich).* 2018;20:1100-3.
5. O'Brien E, Asmar R, Beilin L, et al. European Society of Hypertension recommendations for conventional, ambulatory and home blood pressure measurement. *J Hypertens.* 2003;21:821-48.
6. Pickering TG, Hall JE, Appel LJ, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on high blood pressure research. *Circulation.* 2005;111:697-716.
7. Perloff D, Grim C, Flack J, et al. Human blood pressure determination by sphygmomanometry. *Circulation.* 1993;88:2460-70.
8. Marks LA, Groch A. Optimizing cuff width for noninvasive measurement of blood pressure. *Blood Press Monit.* 2000;5:153-8.
9. Palatini P, Parati G. Blood pressure measurement in very obese patients: a challenging problem. *J Hypertens.* 2011;29(3):425-9.
10. Fonseca-Reyes S, de Alba-García JG, Parra-Carrillo JZ, Paczka-Zapata JA. Effect of standard cuff on blood pressure readings in patients with obese arms. How frequent are arms of a 'large circumference'? *Blood Press Monit.* 2003;8:101-6.
11. Kho CL, Brown MA, Ong SL, Mangos GJ. Blood pressure measurement in pregnancy: the effect of arm circumference and sphygmomanometer cuff size. *Obstet Med.* 2009;2:116-20.
12. Bonso E, Saladini F, Zanier A, Benetti E, Dorigatti F, Palatini P. Accuracy of a single rigid conical cuff with standard-size bladder coupled to an automatic oscillometric device over a wide range of arm circumferences. *Hypertens Res.* 2010;33:1186-91.
13. Palatini P, Benetti E, Fania C, Malipiero G, Saladini F. Rectangular cuffs may overestimate blood pressure in individuals with large conical arms. *J Hypertens.* 2012;30:530-6.
14. Wang J, Thornton JC, Russell M, Burastero S, Heymsfield S, Pierson RN Jr. Asians have lower body mass index (BMI) but higher percent body fat than do whites: comparisons of anthropometric measurements. *Am J Clin Nutr.* 1994;60:23-8.
15. Von Elm E, Altman DG, Egger M, Pocock SJ, Gtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med.* 2007;147(8):573-7.
16. National Health and Nutrition Examination Survey: Anthropometry Procedures Manual 2007-08. USA: Centers for Disease Control and Prevention. 2007.
17. Hogan JL, Maguire P, Farah N, Kennelly MM, Stuart B, Turner MJ. Body mass index and blood pressure measurement during pregnancy. *Hypertens Pregnancy.* 2011;30:396-400.
18. Cooley SM, Donnelly JC, Walsh T, et al. The relationship between body mass index and mid-arm circumference in a pregnant population. *J Obstet Gynaecol.* 2011;31:594-6.
19. Wu LW, Lin YY, Kao TW, et al. Mid-arm circumference and all-cause, cardiovascular, and Cancer mortality among obese and non-obese US adults: the National Health and nutrition examination survey III. *Sci Rep.* 2017;7:2302.
20. Forsberg SA, de Guzman M, Berlind S. Validity of blood pressure measurement with cuff in the arm and forearm. *Acta Med Scand.* 1970;188:389-96.

21. Pierin AM, Alavarce DC, Gusmão JL, Halpern A, Mion D Jr. Blood pressure measurement in obese patients: comparison between upper arm and forearm measurements. *Blood Press Monit.* 2004;9:101–5.
22. Schell K, Bradley E, Bucher L, et al. Clinical comparison of automatic, noninvasive measurements of blood pressure in the forearm and upper arm. *Am J Crit Care.* 2005;14:232–41.
23. Maxwell GF, Pruijt JF, Arntzenius AC. Comparison of the conical cuff and the standard rectangular cuffs. *Int J Epidemiol.* 1985;14:468–72.
24. Executive Summary on National Population Health Survey 2016/17. Singapore: Ministry of Health Singapore and Health Promotion Board. 2017.
25. Akkermans J, Diepeveen M, Ganzevoort W, van Montfrans GA, Westerhof BE, Wolf H. Continuous non-invasive blood pressure monitoring, a validation study of Nexfin in a pregnant population. *Hypertens Pregnancy.* 2009;28: 230–42.
26. Ilies C, Bauer M, Berg P, et al. Investigation of the agreement of a continuous non-invasive arterial pressure device in comparison with invasive radial artery measurement. *Br J Anaesth.* 2012;108:202–10.
27. Hohn A, Defosse JM, Becker S, Steffen C, Wappler F, Sakka SG. Non-invasive continuous arterial pressure monitoring with Nexfin does not sufficiently replace invasive measurements in critically ill patients. *Br J Anaesth.* 2013; 111:178–84.
28. Heslehurst N, Simpson H, Ells LJ, et al. The impact of maternal BMI status on pregnancy outcomes with immediate short-term obstetric resource implications: a meta-analysis. *Obes Rev.* 2008;9:635–83.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

