

GOPEN ACCESS

Citation: Fakhour S, Renoz F, Ambroise J, Pons I, Noël C, Gala J-L, et al. (2021) Insight into the bacterial communities of the subterranean aphid *Anoecia corni*. PLoS ONE 16(8): e0256019. https:// doi.org/10.1371/journal.pone.0256019

Editor: Clara F. Rodrigues, Universidade de Aveiro, PORTUGAL

Received: December 22, 2020

Accepted: July 28, 2021

Published: August 11, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: https://doi.org/10.1371/journal.pone.0256019

Copyright: © 2021 Fakhour et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data accessibility: European Nucleotide Archive (ENA) accession number of NGS sequence generated for the 16 aphid samples reported in this paper is PRJEB35700. **RESEARCH ARTICLE**

Insight into the bacterial communities of the subterranean aphid *Anoecia corni*

Samir Fakhour^{1,2}*, François Renoz², Jérôme Ambroise³, Inès Pons², Christine Noël², Jean-Luc Gala³, Thierry Hance²

Department of Plant Protection, National Institute for Agricultural Research (INRA), Béni-Mellal, Morocco,
Earth and Life Institute, UC Louvain, Louvain-la-Neuve, Belgium, 3 Center for Applied Molecular
Technologies (CTMA), Institut de Recherche Expérimentale et Clinique (IREC), UC Louvain, Woluwe-Saint-Lambert, Belgium

• These authors contributed equally to this work.

* samir.fakhour@inra.ma

Abstract

Many insect species are associated with bacterial partners that can significantly influence their evolutionary ecology. Compared to other insect groups, aphids harbor a bacterial microbiota that has the reputation of being poorly diversified, generally limited to the presence of the obligate nutritional symbiont Buchnera aphidicola and some facultative symbionts. In this study, we analyzed the bacterial diversity associated with the dogwood-grass aphid Anoecia corni, an aphid species that spends much of its life cycle in a subterranean environment. Little is known about the bacterial diversity associated with aphids displaying such a lifestyle, and one hypothesis is that close contact with the vast microbial community of the rhizosphere could promote the acquisition of a richer bacterial diversity compared to other aphid species. Using 16S rRNA amplicon Illumina sequencing on specimens collected on wheat roots in Morocco, we identified 10 bacterial operational taxonomic units (OTUs) corresponding to five bacterial genera. In addition to the obligate symbiont Buchnera, we identified the facultative symbionts Serratia symbiotica and Wolbachia in certain aphid colonies. The detection of Wolbachia is unexpected as it is considered rare in aphids. Moreover, its biological significance remains unknown in these insects. Besides, we also detected Arsenophonus and Dactylopiibacterium carminicum. These results suggest that, despite its subterranean lifestyle, A. corni shelter a bacterial diversity mainly limited to bacterial endosymbionts.

Introduction

Insects maintain a variety of symbiotic relationships with heritable bacteria that can deeply influence their evolutionary ecology [1–3]. Thanks to their well-studied associations with a wide range of heritable symbiotic bacteria, aphids (Hemiptera: Aphididae) are valuable model systems for studying the evolution of bacterial mutualism in insects [4–6]. Like many insect species that feed on nutrient-deficient diets, aphids typically harbor an ancient nutritional obligate endosymbiont, *Buchnera aphidicola*, confined in specialized cells called bacteriocytes and

Funding: This work was supported by the Merit Scholarship Program for High Technology from the Islamic Development Bank [IBD File No. 51/MOR/ P33-600029718] and by the Fonds de la Recherche Scientifique (FNRS, FRIA Grant No. 1. E014.17F). The funders had no role in study design and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

stably maintained in host populations by vertical transmission [5,7]. In addition to their obligate partner, aphids can also host various facultative endosymbionts that can positively or negatively affect a variety of host phenotypes, depending on the ecological context [4,6].

The functional diversity of facultative symbionts associated with aphids includes γ -proteobacteria [e.g. *Arsenophonus* sp. [8,9], *Regiella insecticola* [10,11], *Serratia symbiotica* [12,13], *Hamiltonella defensa* [14,15], *Rickettsiella viridis* [16,17] and *Candidatus Fukatsuia symbiotica* [18,19]], α -proteobacteria [e.g. the genus *Rickettsia* [20,21] and *Wolbachia* [22,23]] and Mollicutes of *Spiroplasma* genus [24,25]. Ecological effects associated with these bacterial partners include defense against parasites [26–30], body color modification [31,32], heat stress tolerance [33], host plant use and nutrition [34,35] and host reproductive manipulation [36].

In addition to intracellular endosymbiotic bacteria, the aphid microbiota may also include bacterial partners involved in less lasting interactions, which may be transient or even antagonistic, and include gut bacteria, plant associates, pathogens, and environmental contaminants [1,2,37–44]. In aphids, these microorganisms have received limited consideration, notably because of the virtual absence of bacteria in the plant-phloem [45]. However, high-throughput sequencing approaches provide the opportunity to get more complete pictures of the bacterial communities associated with these insects. In this regard, recent studies suggest that the microbiota of some aphid species may be more diverse than previously thought, involving a wider a bacterial diversity that includes members of the genera *Pseudomonas, Erwinia, Acinetobacter, Staphylococcus* and *Pantoea* among others [40,42,46]. Tackling this diversity is crucial to understanding how heritable bacterial endosymbioses are established from free-living lineages. Indeed, recent studies suggest that heritable endosymbionts derive from free-living bacteria, which can sometimes reside in the host plant and in the digestive tract of aphids [39,47–51].

The diversity of the microbiota associated with aphids is linked to their living environment, and the evolutionary acquisition of certain symbionts is probably due to particular habitats. In this context, the soil is an extraordinary reserve of microbiological diversity whose functions are essential to the functioning ecosystems [52]. We therefore hypothesize that aphid species living in close association with the soil, such as the dogwood-grass aphid *Anoecia corni*, are likely to harbor a more diverse and original microbiome than the aphid species strictly present in the areal parts of plants. This particular species notably has access to the xylem tissues of the roots, the first gateway through which many soil-borne bacteria, sometimes pathogenic, transit [53–56].

A. corni is a holocylic dioecious species belonging to a genus that includes about twenty aphid species widely distributed in the Holarctic, many whose ecological and taxonomic position remain largely unknown [57,58]. In temperate areas, overwintering eggs hatch on dog-wood (*Cornus sanguinea*) during spring, giving rise to a generation of fundatrices. In summer, the alates leave dogwood and migrate onto the roots of grasses and sedges (Poaceae, Cyperaceae) where they are often attended by ants [59]. The microbiota associated with *A. corni* is unknown, and its subterranean lifestyle makes it an ideal candidate to test the hypothesis that close contact with the vast microbial community of the rhizosphere could promote the acquisition of a richer bacterial diversity compared to other aphid species. In this study, we sampled *A. corni* colonies on wheat roots in two regions of Morocco. The 16S rRNA amplicon Illumina sequencing approach was used to examine the composition of the microbiota associated with aphid samples and to clarify the relationship between these two organisms. For this, the evolutionary history was inferred using the Neighbor Joining (NJ). Our results are discussed in light of previous studies on the microbiota associated with other aphid species.

Materials and methods

Sample collection and DNA extraction

Apterous adults of *A. corni* were collected during August 2014 on roots of wheat plants (*Triticum* sp.) in Morocco with the kind permission of the landowners. A total of 16 colonies were sampled in two important regions in terms of cereal crops: eight colonies were collected in the locality of Béni Mellal-Khénifra and eight colonies were collected in the locality of Casablanca-Settat (Fig 1; S1 Table in the Supporting Information). Aphid collection consisted of three wingless parthenogenetic adult females per colony, which were immediately immersed in 95% ethanol during collection and preserved at 4°C until use.

DNA extraction, PCR amplification, library preparation and sequencing

Prior to DNA extractions, aphid samples, each comprising three adult aphids of the same colony, were surface-sterilized with 99% ethanol, 10% bleach and rinsed with sterile water to remove surface contaminants. The genomic DNA was extracted using the DNeasy Blood & Tissue kit (QIAGEN) following the instruction of the manufacturer. The quantity and quality of the DNA extractions were measured with a NanoDrop spectrophotometer (Thermo Scientific, USA). Extractions were then stored at -20°C. After extraction, the genomic 16S rRNA has



Fig 1. Geographical location of collection sites of *A. corni* colonies analyzed in this study (for details, see <u>S1 Table</u> in the supporting information). https://doi.org/10.1371/journal.pone.0256019.g001

been diluted using sterilized ultrapure water in equal concentration (5 $ng/\mu l$) from each sample for further steps.

PCR amplification, library preparation and sequencing

Sequencing libraries were prepared according to the Illumina MiSeq system instructions 16S workflow and as described previously [40]. First, DNA was amplified using universal primers with overhang adapters attached (F: 5' TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTA CGGGNGGCWGCAG and R: 5'GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVG GGTATCTAATCC) and which targets the V3-V4 variable region of the 16S bacterial rRNA gene. This first PCR step (PCR1) was carried out using a kit 2x KAPA HiFi HotStart ReadyMix in a mixture total volume of 25 $\mu l.$ The conditions of this first PCR were 95 $^\circ C$ for 3 min (1 cycle); 95°C for 30 s, 55°C for 30 s and 72°C for 30 s (25 cycles), followed by 72°C for 5 min. PCR products were cleaned with AMPure XP beads. A second-stage PCR (PCR2) was carried out using 5 µl PCR1 purified to attach dual indices and Illumina sequencing adapters using the Nextera XT Index Kit. Different combinations of index (i5 and i7) were used for each sample. The PCR2 was executed under the same conditions of PCR1 but with eight PCR cycles. A clean-up of the PCR2 products with AMPure XP beads was performed before quantification. PCR2 products were quantified and normalized at 7 ng/µl using PicoGreen dsDNA Quantitation Assay and were generated an equimolar pool (7 ng/µl). Before proceeding to highthroughput sequencing (HTS), the final pool was quantified by qPCR (kit KAPA SYBR FAST qPCR ABI Prism readymix KK4604) and 7 pM of denatured final pool was loaded on MiSeq reagent kits v3 (600 cycles). All the processes were carried out by the GIGA-Research Center of the University of Liège (ULiège, Belgium) using Illumina MiSeq Technology for paired-end sequencing $(2 \times 300 \text{ bp reads})$.

Data analysis

The data analysis was achieved as previously described [40]. The 16S rRNA Illumina Miseq sequencing reads were analyzed using UPARSE [60] bioinformatics pipeline. For each aphid sample, forward and reverse sequences from paired-end reads were assembled and the resulting consensus sequences were filtered based on their respective quality (expected number of errors <1.0 and a length >450 nt). Sequences with $\ge 97\%$ similarity were assigned to the same operational taxonomic units (OTUs). Chimera were removed using a referencebased filtering with UCHIME and the gold database of the corresponding software. A second level of quality-filtering was carried out in order to discard OTUs with a number of sequences <0.005% of the total number of sequences, as recommended previously [61]. As previously recommended [62,63], an additional filtering was performed by analyzing the negative controls in order to remove OTUs corresponding to potential contaminations. Bacterial taxonomic assignments of each OTU were obtained using the dada2 R package [64] and the Greengenes (v.13.8) database [65]. Finally, to improve the taxonomic assignment, the representative OTUs were compared to the sequences in the GenBank using BLASTn. To compare samples, the number of sequences was standardized or rarefied to 50,000 per sample. After rarefaction, the OTU table was analyzed (See S2 Table. Commands and options used to build the OUT table). All bacterial sequences found in A. corni are given in S3 and S4 Tables (Supporting information). Raw data were deposited European Nucleotide Archive (ENA) as a file under accession number PRJEB35700.

Phylogenetic relationships of bacteria associated with *A. corni* and representative endosymbionts of other aphids were established using SeaView v4.6.1 to align 16S rRNA sequences [66] and GBlocks v0.91b [67] to remove poorly aligned positions and divergent regions of DNA

alignments. We selected the best fit evolutionary models using PartitionFinder v1.1.0 [68]. The phylogenetic tree was reconstructed using the neighbor joining method with SeaView v4.6.1 and bootstrap values were computed for each branch node (N = 1000).

Results

Library basic statistics

On average, NGS sequencing produced 187,592 bacterial 16S rRNA reads per sample (Table 1).

The assembly of paired sequences resulted in consensus sequences with an average length of 466 bp. The quality-based filtering of the consensus sequences resulted in an average number of high-quality sequences per sample of 150,605.

OTU clustering and taxonomic assignment

Initially, high-quality reads were clustered using >97% sequence similarity into 23 OTUs. Based on the analysis of the negative controls, 13 OTUs that count for 0.23% of the total number of reads were identified as contaminants and removed (S4 Table). Reads were therefore clustered into 10 biologically relevant OTUs (Table 2 and S3 Table).

All OTUs correspond to the group of Proteobacteria and include three bacterial orders: *Enterobacteriales, Rhodocyclales* and *Rickettsiales.* Our results indicate that the microbial profile of *A. corni* is dominated by the order of *Enterobacteriales*, which includes the obligate symbiont *B. aphidicola* and the facultative symbionts *S. symbiotica* and *Arsenophonus*.

B. aphidicola was detected in all the samples (100%) and was represented by 5 OTUs (OTUs 1, 2, 3, 7 and 10) that account for 96.23% of all reads. OTUs 1, 7 and 10 differed by only 3 to 4 bp whereas OTUs 2 and 3 differed from these OTUs by 20 to 38 bp. Different *B. aphidicola* haplotypes are present in an aphid colony with the dominance of a single haplotype (Fig 2). The vast majority of the reads clustered into a single OTU for most aphid species (OTUs 1, 2 or 3) and many minor OTUs (OTUs 7 and 10) were detected in all samples. In aphid colonies from the Casablanca-Settat region, The OTU1 (with related minor OTUs 2, 3, 7 and 10) was detected and matched a sequence of *B. aphidicola* previously reported on *Geoica urticularia*. OTUs 2 and 3 were common to all samples from Béni Mellal-Kénifra region and OTU3 was detected on the *Anoecia* genus.

The next most abundant OTUs were presented by the facultative symbiont *S. symbiotica* (OTUs 4 and 5) that account for 3.52% of all reads. Taxonomic identification of bacterial OTUs resulted in three additional taxa including *Wolbachia* (OTU 6), *Arsenophonus* (OTU 8) and *D. carminicum* (OTU 9). Phylogenetic analyses including the symbionts associated with

Raw data			
Average size (Mb) per sample	3		
Raw number of sequence per sample	187,592		
After assembly of paired sequences			
Average size (Mb) per sample	169.4		
Average number of sequence per sample	173,117		
Average of the median sequence length	466		
After quality filtering			
Average size (Mb) per sample	72.7		
Average number of sequence per sample	150,605		
Sequence length (min; median; max)	(450; 465; 584)		

Table 1. Summary of sequencing data.

https://doi.org/10.1371/journal.pone.0256019.t001

OTU no.	PC reads.	Greengenes identification	Id%	GenBank identification	Accession	Id%
		Taxon		Three closest GenBank matches		
OTU_01	58.77	Buchnera	98.28	Buchnera aphidicola/Geoica urticularia	AJ296751.1	98.28
				Buchnera aphidicola/Myzus persicae	CP002703.1	96.34
				Buchnera aphidicola/Myzus persicae	CP002701.1	96.34
OTU_02	20.08	Buchnera	96.15	Buchnera aphidicola/Pemphigus matsumurai	KF311221.1	96.15
				Buchnera aphidicola/Pemphigus sinobursarius	KF311219.1	96.15
				Buchnera aphidicola/Pemphigus yunnanensis	HQ792326.1	96.15
OTU_03	17.34	Buchnera	91.24	Buchnera aphidicola/Anoecia oenotherae	CP033012.1	97.85
				Buchnera aphidicola/Anoecia fulviabdominalis	JX998094.1	97.2
				Buchnera aphidicola/Eulachnus mediterraneus	LT600356.1	94.22
OTU_04	2.62	Serratia	99.14	Serratia symbiotica/Aphis fabae	KT176010.1	99.35
				Serratia symbiotica/soil	MG287131.1	99.14
				Serratia symbiotica/soil	KX900450.1	99.14
OTU_05	0.89	-	90.15	Serratia symbiotica/Prociphilus longianus	MG831336.1	99.35
				Serratia symbiotica/Prociphilus longianus	MG835393.1	98.92
				Serratia symbiotica/Prociphilus longianus	MG835392.1	98.92
OTU_06	0.04	Wolbachia	96.64	Wolbachia pipientis/Pentalonia nigronervosa	KJ786950.1	96.64
				Wolbachia pipientis/Pentalonia nigronervosa	KJ786949.1	96.64
				Wolbachia pipientis/Pentalonia nigronervosa	KC522606.1	96.64
OTU_7	0.01	Buchnera	98.21	Buchnera aphidicola/Geoica urticularia	AJ296751.1	98.21
				Buchnera aphidicola/Myzus persicae	CP002703.1	95.92
				Buchnera aphidicola/Myzus persicae	CP002701.1	95.92
OTU_8	0.01	Candidatus Phlomobacter	95.05	Arsenophonus/Aleurodicus dispersus	AY264664.1	95.91
				Arsenophonus/Macrosteles sexnotatus	AB795344.1	95.27
				Arsenophonus/Stomaphis takahashii	FJ655541.1	95.27
OTU_9	0.01	Uliginosibacterium	98.92	Dactylopiibacterium carminicum/Dactylopius opuntiae	GQ853370.1	98.92
				Dactylopiibacterium carminicum/Dactylopius opuntiae	GQ853369.1	98.92
				Sphingomonas/soil	JX944513.2	96.34
OTU_10	0.01	Buchnera	98.06	Buchnera aphidicola/Geoica urticularia	AJ296751.1	98.06
				Buchnera aphidicola/Myzus persicae	CP002703.1	96.13
				Buchnera aphidicola/Myzus persicae	CP002701.1	96.13
OTUs 11-23	0.23	Contaminants identified from negative control analysis				

Table 2. Taxonomic assignment of OTUs by Greengenes and GenBank, including the three top BLAST hits, GenBank accession numbers and % identity.

PC reads, cluster size in percent; Id, identity %.

https://doi.org/10.1371/journal.pone.0256019.t002

A. corni (i.e. *B. aphidicola*, *S. symbiotica*, *Arsenophonus* and *Wolbachia*) and representative symbionts of other species of aphids and insects are shown in <u>S1-S4</u> Figs (Supporting information).

Diversity of bacterial communities in the samples

The bacterial communities of the samples were mainly composed of the obligate symbiont *B. aphidicola* and mostly complemented by the facultative symbionts *S. symbiotica* (with a high abundance of reads from three samples) and *Wolbachia* (with reads detected from four samples) (Fig 2).

The facultative symbiont *Arsenophonus* was detected in most of the samples (13/16), but with an extremely low number of reads. *D. carminicum* was also found in most of the samples with an extremely low number of reads.





https://doi.org/10.1371/journal.pone.0256019.g002

Discussion

Our approach based on 16S rRNA enabled us to identify five bacterial genera in the dogwood aphid *A. corni*, a species that spends most of its life cycle on Poaceae roots. Due to its subterranean lifestyle, we expected to find in this species a bacterial diversity that includes environmental bacteria such as gut bacteria, plant associates, pathogens and environmental contaminants. Xylem vessels are the primary entry routes for many soil-borne bacteria infecting plants [56]. Although primarily phloem-feeders, aphids are also capable of ingestion from the xylem vessels, a strategy displayed for maintaining water balance [69,70]. OTUs that were identified correspond to five bacterial genera, mostly related to symbionts already found associated with aphids: *Buchnera, Serratia, Wolbachia, Arsenophonus* and *Dactylopiibacterium*. All these genera have been previously described as symbiotic partners of insects. We did not find any bacterial partners that can be considered as environmental-related (e.g. *Pseudomonas* spp., *Erwinia* spp., etc.) as in the case of other aphid species, including those that feed on cereal crops [40,42].

The nutritional obligate symbiont *Buchnera* was found in all samples with distinctive 16S haplotypes in a single aphid colony. These results might be a consequence of *Buchnera* polyploidy, as evidenced by the 16S rRNA copy-number variation. Alternatively, a clone from a single colony may contain *Buchnera* strains with different haplotypes. Co-infection with multiple *B. aphidicola* strains was reported in several aphid genera [71,72].

The secondary endosymbiont *S. symbiotica* is one of the most common symbiont species in aphid populations [73] and was identified in three of the eight colonies surveyed. This symbiont includes a wide variety of strains ranging from co-obligate nutritional partners, that are mainly found in the Lachninae and the Chaitophorinae subfamilies [12,74], to facultative strains whose reported associated effects in the pea aphid *Acyrthosiphon pisum* are heat stress resistance and protection against parasitoids [75–77]. Strains detected in this study are probably of facultative nature, as *S. symbiotica* was not found in all colonies.

Interestingly, *Wolbachia* was detected in four colonies. This symbiont, an α -proteobacterium, is commonly found in insects and studies suggest that *Wolbachia* is present in at least 65% of arthropod species [78]. It is known to manipulate the reproduction of their host [79], promote the oogenesis in certain wasp species [80], display a nutritional function in certain bedbug and whiteflies species by producing B vitamins [81,82], and is associated with antiviral protection by influencing the vector competence of several species of mosquitoes for viruses [83,84]. Although some studies have reported the presence of *Wolbachia* in aphid populations, it is considered rare in these insects [85–90]. The biological significance of *Wolbachia* in aphids is still unknown. It has been hypothesized that the symbiont play a role in the proliferation of asexual lineages [87], and its role in the production of B vitamins in the banana aphid *Pentalonia nigronervosa* is currently debated [86,91,92]. *Wolbachia* infections in aphids could also be acquired by horizontal transmission from other insects such as parasitoid wasps, known to be infected by this symbiont [93,94]. To our knowledge, the stability of *Wolbachia* infections in aphids has never be tested and *A. corni* could be a suitable candidate to elucidate the biological significance of this symbiont in aphids.

The genera *Arsenophonus* and *Dactylopiibacterium* were also detected in most of the sampled aphids, but with a much lower read abundance than for *S. symbiotica* and *Wolbachia*. *Arsenophonus* is a bacteria found in many insect species including aphids, scale insects, leafhoppers, whiteflies and wasps [8,82,95–97]. Despite the fact that the prevalence of *Arsenophonus* can reach up to 70% in species of the *Aphis* genus [98], the phenotypes associated with this symbiont remain unclear in aphids. Bacteriophages required for protective symbiosis were found in various strains of the symbiont [99], but no defensive properties were found in *Aphis glycines* infected by *Arsenophonus* [98]. Recent studies suggest that *Arsenophonus* may be involved in host nutrition, probably by mediating host plant range [35,100–102]. In whiteflies, comparative genomics suggests that *Arsenophonus* is a source of B vitamins [82]. However, no genome of strains associated with aphids has yet been sequenced. While the presence of a symbiont in specialized host cells such as bacteriocytes and sheath cells are important clues for determining the mutualistic and heritable nature of a symbiont [4], no such information are available for this particular symbiont.

One OTU was assigned to *D. carminicum* (β -proteobacteria, family Rhodocyclaceae). So far, this bacterial species has only been reported in the scale insect species *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae), where it has been described as a nitrogen-fixing symbiont [103,104]. *Dactylopius coccus* is now well established in Morocco where it ravages the plants of *Opuntia ficus-indica*. Although *D. carminicum* is considered a symbiont capable of passing through the reproductive organs in scale insects, this species remains largely undocumented. It cannot be excluded at present that this newly discovered species resides in the soil, in the host plant or lives in other insects.

In recent years, several studies have characterized overall all the bacteria present in aphids by deep sequencing of 16S rRNA [40,42,45]. A common point throughout these studies is the reduced abundance of environmental bacteria relative to the primary and secondary endosymbionts. Besides theses heritable symbionts and in contrast to our results, many environmental bacteria have been reported in *R. padi*, e.g. the phytopathogenic members of the Pseudomonas genus and some saprophytes of plant and soil (*Acinetobacter* and *Staphyloccocus* genera). Moreover, gut symbiotic bacteria of aphids were also found, i.e. *Pantoea* and *Erwinia* genera [40,42].

In conclusion, although *A. corni* lives in the rhizosphere, an environment that is very rich in bacteria and other microorganisms, the number of bacterial taxa detected in this species is surprisingly low [105]. Despite its subterranean lifestyle, *A. corni* shelter a bacterial diversity mainly limited to known bacterial endosymbionts. Few species of facultative endosymbionts

have been detected in the context of this study. It is now recognized that some species of aphids are more likely to harbor facultative symbionts than others [4,73,106]. It should be noted that our sampling covers only a small part of Morocco, and the diversity of symbionts of aphid populations can change dramatically in response to various environmental conditions [4]. However, our study provides a snapshot of the bacterial community associated with a poorly studied aphid species, and identified bacterial taxa that may play a role in the biology of A. corni, in particular the facultative symbionts Arsenophonus and Wolbachia whose associated phenotypes in aphids are still elusive. A. corni could represent a suitable species to investigate the role of these symbiotic bacteria in aphids. Finally, insect-associated bacterial communities, and in particular heritable symbionts, have received much attention in recent decades, somewhat overshadowing the diversity of other types of microorganisms that can associate with insects, such as fungi. For example, in some species of cicadas, grasshoppers, and aphids, certain fungi species have become obligate symbionts by replacing ancestral bacterial symbionts [107–111], suggesting that fungi may establish more or less long-lasting relationships with insects and become an established part of their microbiota. Although fungal diversity was not explored in our study, it would likely deserve more attention in future work, and insects with a subterranean lifestyle are likely interesting candidates for such an investigation.

Supporting information

S1 Fig. Phylogenetic analysis of *Buchnera* associated with *A. corni* and its placement compared to strains associated with other aphid species, based on a sequence from the V3-V4 region of bacterial 16S rRNA. The evolutionary history was inferred using the Neighbor Joining (NJ) methods, with a J-C model. The percentage of replicate trees was verified with bootstrap of 1000 replicates. Designations in bold are strains sequenced in this study. Host names are followed by the GenBank accession number of each bacterial sequence. Geneious version 6.1 created by Biomatters. (TIF)

S2 Fig. Phylogenetic analysis of *Serratia symbiotica* associated with *Anoecia corni* and their placement compared to strains associated with other aphid species, based on a sequence from the V3-V4 region of bacterial 16S rRNA. The evolutionary history was inferred using the Neighbor Joining (NJ) methods, with a J-C model. The percentage of replicate trees was verified with bootstrap of 1000 replicates. Designation in bold are is strains sequenced in this study. Host names are followed by the GenBank accession number of each bacterial sequence. Geneious version 6.1 created by Biomatters. (TIF)

S3 Fig. Phylogenetic analysis of *Wolbachia* associated with *Anoecia corni* and its placement compared to strains associated with other insect species, based on a sequence from the V3-V4 region of bacterial 16S rRNA. The evolutionary history was inferred using the Neighbor Joining (NJ) methods, with a HKY model. The percentage of replicate trees was verified with bootstrap of 1000 replicates. Designation in bold is strain sequenced in this study. Host names are followed by the GenBank accession number of each bacterial sequence. Geneious version 6.1 created by Biomatters.

(TIF)

S4 Fig. Phylogenetic analysis of *Arsenophonus* associated with *Anoecia corni* and its placement compared to strains associated with other insect species, based on a sequence from the V3-V4 region of bacterial 16S rRNA. The evolutionary history was inferred using the Neighbor Joining (NJ) methods, with a HKY model. The percentage of replicate trees was verified with bootstrap of 1000 replicates. Designation in bold is strain sequenced in this study. Host names are followed by the GenBank accession number of each bacterial sequence. Geneious version 6.1 created by Biomatters.

(TIF)

S1 Table. Summary of collection details and 16S rRNA gene sequencing results for aphid samples.

(DOCX)

S2 Table. Data analysis: Commands and options used to build the OUT table. (DOCX)

S3 Table. Bacterial representative sequences (operational taxonomic units) found in *A. corni*.

(DOCX)

S4 Table. Contaminant OTUs identified in this study. (DOCX)

Acknowledgments

The authors thank Linda Dhondt, Karim Andich and Slimane Khayi for technical assistance. This paper is publication BRC359 of the Biodiversity Research center (UC Louvain).

Author Contributions

Conceptualization: Samir Fakhour, François Renoz, Thierry Hance.

Data curation: Samir Fakhour, François Renoz, Jérôme Ambroise.

Formal analysis: Samir Fakhour, François Renoz, Jérôme Ambroise, Inès Pons, Christine Noël, Jean-Luc Gala.

Funding acquisition: Samir Fakhour, Inès Pons, Thierry Hance.

Investigation: Samir Fakhour, Thierry Hance.

Methodology: Samir Fakhour, François Renoz, Thierry Hance.

Project administration: Thierry Hance.

Software: Jérôme Ambroise, Jean-Luc Gala.

Supervision: Thierry Hance.

Validation: Samir Fakhour, François Renoz, Thierry Hance.

Visualization: Samir Fakhour, François Renoz.

Writing - original draft: Samir Fakhour, François Renoz, Inès Pons.

Writing – review & editing: Samir Fakhour, François Renoz, Jérôme Ambroise, Inès Pons, Jean-Luc Gala, Thierry Hance.

References

- Engel P, Moran NA. The gut microbiota of insects—diversity in structure and function. FEMS Microbiol Rev. 2013; 37: 699–735. https://doi.org/10.1111/1574-6976.12025 PMID: 23692388
- 2. Douglas AE. Multiorganismal insects: Diversity and function of resident microorganisms. Annu Rev Entomol. 2015; 60: 17–34. https://doi.org/10.1146/annurev-ento-010814-020822 PMID: 25341109

- Gil R, Latorre A. Unity makes strength: A review on mutualistic symbiosis in representative insect clades. Life. 2019; 9: 21. https://doi.org/10.3390/life9010021 PMID: 30823538
- Oliver KM, Degnan PH, Burke GR, Moran NA. Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. Annu Rev Entomol. 2010; 55: 247–266. https://doi.org/10. 1146/annurev-ento-112408-085305 PMID: 19728837
- Moran NA, McCutcheon JP, Nakabachi A. Genomics and evolution of heritable bacterial symbionts. Annu Rev Genet. 2008; 42: 165–190. <u>https://doi.org/10.1146/annurev.genet.41.110306.130119</u> PMID: 18983256
- Feldhaar H. Bacterial symbionts as mediators of ecologically important traits of insect hosts. Ecol Entomol. 2011; 36: 533–543. https://doi.org/10.1111/j.1365-2311.2011.01318.x
- Douglas AE. Nutritional interactions in insect-microbial symbioses: Aphids and their symbiotic bacteria Buchnera. Annu Rev Entomol. 1998; 43: 17–37. <u>https://doi.org/10.1146/annurev.ento.43.1.17</u> PMID: 15012383
- Jousselin E, Cœur d'Acier A, Vanlerberghe-Masutti F, Duron O. Evolution and diversity of Arsenophonus endosymbionts in aphids. Mol Ecol. 2013; 22: 260–270. <u>https://doi.org/10.1111/mec.12092</u> PMID: 23106652
- Tian P-P, Chang C-Y, Miao N-H, Li M-Y, Liu X-D. Infections with Arsenophonus facultative endosymbionts alter performance of aphids (*Aphis gossypii*) on an amino-acid-deficient diet. Master ER, editor. Appl Environ Microbiol. 2019; 85. https://doi.org/10.1128/AEM.01407-19 PMID: 31540983
- Jamin AR, Vorburger C. Estimating costs of aphid resistance to parasitoids conferred by a protective strain of the bacterial endosymbiont *Regiella insecticola*. Entomol Exp Appl. 2019; 167: 252–260. https://doi.org/10.1111/eea.12749
- Parker BJ, Hrček J, McLean AHC, Brisson JA, Godfray HCJ. Intraspecific variation in symbiont density in an insect–microbe symbiosis. Mol Ecol. 2021; 30: 1559–1569. <u>https://doi.org/10.1111/mec.15821</u> PMID: 33512733
- Manzano-Marín A, Latorre A. Snapshots of a shrinking partner: Genome reduction in Serratia symbiotica. Sci Rep. 2016; 6. https://doi.org/10.1038/srep32590 PMID: 27599759
- Skaljac M, Kirfel P, Grotmann J, Vilcinskas A. Fitness costs of infection with Serratia symbiotica are associated with greater susceptibility to insecticides in the pea aphid Acyrthosiphon pisum: Serratia symbiotica correlates with susceptibility to insecticides in pea aphid. Pest Manag Sci. 2018; 74: 1829– 1836. https://doi.org/10.1002/ps.4881 PMID: 29443436
- Degnan PH, Yu Y, Sisneros N, Wing RA, Moran NA. *Hamiltonella defensa*, genome evolution of protective bacterial endosymbiont from pathogenic ancestors. Proc Natl Acad Sci. 2009; 106: 9063–9068. https://doi.org/10.1073/pnas.0900194106 PMID: 19451630
- Kaech H, Vorburger C. Horizontal transmission of the heritable protective endosymbiont Hamiltonella defensa depends on titre and haplotype. Front Microbiol. 2021; 11: 628755. <u>https://doi.org/10.3389/</u> fmicb.2020.628755 PMID: 33519791
- Tsuchida T, Koga R, Horikawa M, Tsunoda T, Maoka T, Matsumoto S, et al. Symbiotic bacterium modifies aphid body color. Science. 2010; 330: 1102–1104. <u>https://doi.org/10.1126/science.1195463</u> PMID: 21097935
- Polin S, Le Gallic J-F, Simon J-C, Tsuchida T, Outreman Y. Conditional reduction of predation risk associated with a facultative symbiont in an insect. Edwards OR, editor. PLOS ONE. 2015; 10: e0143728. https://doi.org/10.1371/journal.pone.0143728 PMID: 26618776
- Guay J-F, Boudreault S, Michaud D, Cloutier C. Impact of environmental stress on aphid clonal resistance to parasitoids: Role of *Hamiltonella defensa* bacterial symbiosis in association with a new facultative symbiont of the pea aphid. J Insect Physiol. 2009; 55: 919–926. <u>https://doi.org/10.1016/j.jinsphys.2009.06.006</u> PMID: 19545573
- Manzano-Marín A, Szabó G, Simon J-C, Horn M, Latorre A. Happens in the best of subfamilies: establishment and repeated replacements of co-obligate secondary endosymbionts within Lachninae aphids. Environ Microbiol. 2017; 19: 393–408. https://doi.org/10.1111/1462-2920.13633 PMID: 27902872
- Sakurai M, Koga R, Tsuchida T, Meng X-Y, Fukatsu T. *Rickettsia* symbiont in the pea aphid *Acyrthosi-phon pisum*: Novel cellular tropism, effect on host fitness, and interaction with the essential symbiont *Buchnera*. Appl Environ Microbiol. 2005; 71: 4069–4075. https://doi.org/10.1128/AEM.71.7.4069-4075.2005 PMID: 16000822
- 21. Romanov DA, Zakharov IA, Shaikevich EV. *Wolbachia, Spiroplasma*, and *Rickettsia* symbiotic bacteria in aphids (Aphidoidea). Vavilov J Genet Breed. 2020; 24: 673–682. PMID: 33659853
- 22. Ren W, Wei H, Yang Y, Shao S, Wu H, Chen X, et al. Molecular detection and phylogenetic analyses of *Wolbachia* in natural populations of nine galling aphid species. Sci Rep. 2020; 10: 12025. https://doi. org/10.1038/s41598-020-68925-z PMID: 32694524

- Moreira M, Aguiar AMF, Bourtzis K, Latorre A, Khadem M. Wolbachia (Alphaproteobacteria: Rickettsiales) infections in isolated aphid populations from oceanic Islands of the Azores archipelago: Revisiting the supergroups M and N. Environ Entomol. 2019; 48: 326–334. <u>https://doi.org/10.1093/ee/nvy189</u> PMID: 30668658
- Mathé-Hubert H, Kaech H, Ganesanandamoorthy P, Vorburger C. Evolutionary costs and benefits of infection with diverse strains of *Spiroplasma* in pea aphids*. Evolution. 2019; 73: 1466–1481. <u>https:// doi.org/10.1111/evo.13740 PMID: 30990223</u>
- Guidolin AS, Cataldi TR, Labate CA, Francis F, Cônsoli FL. Spiroplasma affects host aphid proteomics feeding on two nutritional resources. Sci Rep. 2018; 8: 2466. <u>https://doi.org/10.1038/s41598-018-</u> 20497-9 PMID: 29410456
- Leclair M, Buchard C, Mahéo F, Simon J-C, Outreman Y. A link between communities of protective endosymbionts and parasitoids of the pea aphid revealed in unmanipulated agricultural systems. Front Ecol Evol. 2021; 9: 618331. https://doi.org/10.3389/fevo.2021.618331
- Chevignon G, Boyd BM, Brandt JW, Oliver KM, Strand MR. Culture-facilitated comparative genomics of the facultative symbiont *Hamiltonella defensa*. Genome Biol Evol. 2018; 10: 786–802. https://doi. org/10.1093/gbe/evy036 PMID: 29452355
- Oliver KM, Higashi CH. Variations on a protective theme: *Hamiltonella defensa* infections in aphids variably impact parasitoid success. Curr Opin Insect Sci. 2019; 32: 1–7. https://doi.org/10.1016/j.cois. 2018.08.009 PMID: 31113620
- Heyworth ER, Ferrari J. A facultative endosymbiont in aphids can provide diverse ecological benefits. J Evol Biol. 2015; 28: 1753–1760. https://doi.org/10.1111/jeb.12705 PMID: 26206380
- Frago E, Mala M, Weldegergis BT, Yang C, McLean A, Godfray HCJ, et al. Symbionts protect aphids from parasitic wasps by attenuating herbivore-induced plant volatiles. Nat Commun. 2017; 8: 1860. https://doi.org/10.1038/s41467-017-01935-0 PMID: 29192219
- Tsuchida T. Molecular basis and ecological relevance of aphid body colors. Curr Opin Insect Sci. 2016; 17: 74–80. https://doi.org/10.1016/j.cois.2016.07.005 PMID: 27720077
- Nikoh N, Tsuchida T, Maeda T, Yamaguchi K, Shigenobu S, Koga R, et al. Genomic insight into symbiosis-induced insect color change by a facultative bacterial endosymbiont, "*Candidatus Rickettsiella viridis*". Ruby EG, editor. mBio. 2018; 9. https://doi.org/10.1128/mBio.00890-18 PMID: 29895637
- Heyworth ER, Smee MR, Ferrari J. Aphid facultative symbionts aid recovery of their obligate symbiont and their host after heat stress. Front Ecol Evol. 2020; 8: 56. https://doi.org/10.3389/fevo.2020.00056
- Lenhart PA, White JA. Endosymbionts facilitate rapid evolution in a polyphagous herbivore. J Evol Biol. 2020; 33: 1507–1511. https://doi.org/10.1111/jeb.13697 PMID: 32894786
- Wagner SM, Martinez AJ, Ruan Y, Kim KL, Lenhart PA, Dehnel AC, et al. Facultative endosymbionts mediate dietary breadth in a polyphagous herbivore. Carroll S, editor. Funct Ecol. 2015; 29: 1402– 1410. https://doi.org/10.1111/1365-2435.12459
- Simon J-C, Boutin S, Tsuchida T, Koga R, Gallic J-FL, Frantz A, et al. Facultative symbiont infections affect aphid reproduction. PLOS ONE. 2011; 6: e21831. https://doi.org/10.1371/journal.pone.0021831 PMID: 21818272
- Grenier A-M, Nardon C, Rahbé Y. Observations on the micro-organisms occurring in the gut of the pea aphid Acyrthosiphon pisum. Entomol Exp Appl. 1994; 70: 91–96. https://doi.org/10.1111/j.1570-7458. 1994.tb01762.x
- Dillon RJ, Dillon VM. The gut bacteria of insects: Nonpathogenic interactions. Annu Rev Entomol. 2004; 49: 71–92. https://doi.org/10.1146/annurev.ento.49.061802.123416 PMID: 14651457
- Grenier A-M, Duport G, Pagès S, Condemine G, Rahbé Y. The phytopathogen *Dickeya dadantii* (*Erwinia chrysanthemi* 3937) is a pathogen of the pea aphid. Appl Environ Microbiol. 2006; 72: 1956–1965. https://doi.org/10.1128/AEM.72.3.1956-1965.2006 PMID: 16517643
- Fakhour S, Ambroise J, Renoz F, Foray V, Gala J-L, Hance T. A large-scale field study of bacterial communities in cereal aphid populations across Morocco. FEMS Microbiol Ecol. 2018; 94. <u>https://doi.org/10.1093/femsec/fiy003 PMID: 29346623</u>
- Campillo T, Luna E, Portier P, Fischer-Le Saux M, Lapitan N, Tisserat NA, et al. *Erwinia iniecta* sp. nov., isolated from Russian wheat aphid (*Diuraphis noxia*). Int J Syst Evol Microbiol. 2015; 65: 3625– 3633. https://doi.org/10.1099/ijsem.0.000466 PMID: 26198254
- **42.** Gauthier J-P, Outreman Y, Mieuzet L, Simon J-C. Bacterial communities associated with host-adapted populations of pea aphids revealed by deep sequencing of 16S ribosomal DNA. PLOS ONE. 2015; 10: e0120664. https://doi.org/10.1371/journal.pone.0120664 PMID: 25807173
- **43.** Harada H, Ishikawa H. Experimental pathogenicity of *Erwinia aphidicola* to pea aphid, *Acyrthosiphon pisum*. J Gen Appl Microbiol. 1997; 43: 363–367. PMID: 12501308

- Grigorescu AS, Renoz F, Sabri A, Foray V, Hance T, Thonart P. Accessing the hidden microbial diversity of aphids: an illustration of how culture-dependent methods can be used to decipher the insect microbiota. Microb Ecol. 2018; 75: 1035–1048. https://doi.org/10.1007/s00248-017-1092-x PMID: 29119316
- Jing X, Wong AC-N, Chaston JM, Colvin J, McKenzie CL, Douglas AE. The bacterial communities in plant phloem-sap-feeding insects. Mol Ecol. 2014; 23: 1433–1444. <u>https://doi.org/10.1111/mec.12637</u> PMID: 24350573
- Zepeda-Paulo F, Ortiz-Martínez S, Silva AX, Lavandero B. Low bacterial community diversity in two introduced aphid pests revealed with 16S rRNA amplicon sequencing. PeerJ. 2018; 6: e4725. <u>https:// doi.org/10.7717/peerj.4725 PMID: 29761046</u>
- Harada H, Oyaizu H, Ishikawa H. A consideration about the origin of aphid intracellular symbiont in connection with gut bacterial flora. J Gen Appl Microbiol. 1996; 42: 17–26. <u>https://doi.org/10.2323/jgam.42.17</u>
- Harada H, Oyaizu H, Kosako Y, Ishikawa H. *Erwinia aphidicola*, a new species isolated from pea aphid, *Acyrthosiphon pisum*. J Gen Appl Microbiol. 1997; 43: 349–354. <u>https://doi.org/10.2323/jgam.</u> 43.349 PMID: 12501306
- 49. Manzano-Marín A, D'acier AC, Clamens A-L, Orvain C, Cruaud C, Barbe V, et al. Serial horizontal transfer of vitamin-biosynthetic genes enables the establishment of new nutritional symbionts in aphids' di-symbiotic systems. ISME J. 2020; 14: 259–273. <u>https://doi.org/10.1038/s41396-019-0533-6</u> PMID: 31624345
- Harada H, Ishikawa H. Gut microbe of aphid closely related to its intracellular symbiont. Biosystems. 1993; 31: 185–191. https://doi.org/10.1016/0303-2647(93)90048-h PMID: 7908835
- Renoz F, Pons I, Vanderpoorten A, Bataille G, Noël C, Foray V, et al. Evidence for gut-associated Serratia symbiotica in wild aphids and ants provides new perspectives on the evolution of bacterial mutualism in insects. Microb Ecol. 2018 [cited 19 Dec 2018]. <u>https://doi.org/10.1007/s00248-018-1265-2</u> PMID: 30276419
- 52. Wagg C, Schlaeppi K, Banerjee S, Kuramae EE, van der Heijden MGA. Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning. Nat Commun. 2019; 10: 1–10.
- Block CC, Hill JH, McGee DC. Seed Transmission of *Pantoea stewartii* in field and sweet corn. Plant Dis. 1998; 82: 775–780. https://doi.org/10.1094/PDIS.1998.82.7.775 PMID: 30856949
- 54. Salanoubat M, Genin S, Artiguenave F, Gouzy J, Mangenot S, Arlat M, et al. Genome sequence of the plant pathogen *Ralstonia solanacearum*. Nature. 2002; 415: 497–502. <u>https://doi.org/10.1038/415497a</u> PMID: <u>11823852</u>
- 55. Hildebrand M, Tebbe CC, Geider K. Survival studies with the fire blight pathogen *Erwinia amylovora* in soil and in a soil-inhabiting insect. J Phytopathol. 2001; 149: 635–639. https://doi.org/10.1046/j.1439-0434.2001.00685.x
- 56. Yadeta K, Thomma B. The xylem as battleground for plant hosts and vascular wilt pathogens. Front Plant Sci. 2013; 4. https://doi.org/10.3389/fpls.2013.00097 PMID: 23630534
- 57. Wieczorek K. Structure of the male reproductive system of Anoecia (Anoecia) corni Fabricius, 1775 (Hemiptera, Aphidoidea), a representative of the family Anoeciidae. Acta Zool. 2007; 89: 163–167. https://doi.org/10.1111/j.1463-6395.2007.00305.x
- Blackman RL, Eastop VF. Aphids on the world's herbaceous plants and shrubs. John Wiley & Sons; 2008.
- 59. Ighil ET-A, Dedryver C-A, Chaubet B, Hullé M. Les pucerons des grandes cultures: Cycles biologiques et activités de vol. 1st ed. Editions QUAE GIE; 2011.
- Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat Methods. 2013; 10: 996–998. https://doi.org/10.1038/nmeth.2604 PMID: 23955772
- Navas-Molina JA, Peralta-Sánchez JM, González A, McMurdie PJ, Vázquez-Baeza Y, Xu Z, et al. Chapter nineteen—Advancing our understanding of the human microbiome using QIIME. In: DeLong EF, editor. Methods in enzymology. Academic Press; 2013. pp. 371–444.
- Edmonds K, Williams L. The role of the negative control in microbiome analyses. FASEB J. 2017; 31: 940.3–940.3.
- Hornung BVH, Zwittink RD, Kuijper EJ. Issues and current standards of controls in microbiome research. FEMS Microbiol Ecol. 2019; 95. https://doi.org/10.1093/femsec/fiz045 PMID: 30997495
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016; 13: 581–583. <u>https://doi.org/10. 1038/nmeth.3869 PMID: 27214047</u>
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, et al. Greengenes, a chimerachecked 16S rRNA gene database and workbench compatible with ARB. Appl Env Microbiol. 2006; 72: 5069–5072. https://doi.org/10.1128/AEM.03006-05 PMID: 16820507

- Gouy M, Guindon S, Gascuel O. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. Mol Biol Evol. 2010; 27: 221–224. https://doi.org/ 10.1093/molbev/msp259 PMID: 19854763
- Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. Mol Biol Evol. 2000; 17: 540–552. <u>https://doi.org/10.1093/oxfordjournals.molbev.a026334</u> PMID: 10742046
- Lanfear R, Calcott B, Ho SYW, Guindon S. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. Mol Biol Evol. 2012; 29: 1695–1701. https://doi.org/10.1093/molbev/mss020 PMID: 22319168
- **69.** Powell G, Hardie J. Xylem ingestion by winged aphids. Entomol Exp Appl. 2002; 104: 103–108. https://doi.org/10.1046/j.1570-7458.2002.00996.x
- 70. Spiller NJ, Koenders L, Tjallingii WF. Xylem ingestion by aphids—a strategy for maintaining water balance. Entomol Exp Appl. 1990; 55: 101–104. https://doi.org/10.1111/j.1570-7458.1990.tb01352.x
- Jones RT, Bressan A, Greenwell AM, Fierer N. Bacterial communities of two parthenogenetic aphid species cocolonizing two host plants across the Hawaiian Islands. Appl Environ Microbiol. 2011; 77: 8345–8349. https://doi.org/10.1128/AEM.05974-11 PMID: 21965398
- 72. Jousselin E, Clamens A-L, Galan M, Bernard M, Maman S, Gschloessl B, et al. Assessment of a 16S rRNA amplicon Illumina sequencing procedure for studying the microbiome of a symbiont-rich aphid genus. Mol Ecol Resour. 2016; 16: 628–640. <u>https://doi.org/10.1111/1755-0998.12478</u> PMID: 26458227
- Henry LM, Maiden MCJ, Ferrari J, Godfray HCJ. Insect life history and the evolution of bacterial mutualism. Ecol Lett. 2015; 18: 516–525. https://doi.org/10.1111/ele.12425 PMID: 25868533
- 74. Monnin D, Jackson R, Kiers ET, Bunker M, Ellers J, Henry LM. Parallel evolution in the integration of a co-obligate aphid symbiosis. bioRxiv. 2020; 2020.01.14.905752. <u>https://doi.org/10.1016/j.cub.2020</u>. 03.011 PMID: 32243856
- Oliver KM, Russell JA, Moran NA, Hunter MS. Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. Proc Natl Acad Sci. 2003; 100: 1803–1807. <u>https://doi.org/10.1073/pnas.0335320100</u> PMID: 12563031
- 76. Oliver KM, Moran NA, Hunter MS. Costs and benefits of a superinfection of facultative symbionts in aphids. Proc R Soc Lond B Biol Sci. 2006; 273: 1273–1280. <u>https://doi.org/10.1098/rspb.2005.3436</u> PMID: 16720402
- 77. Pons I, Renoz F, Noël C, Hance T. New insights into the nature of symbiotic associations in aphids: infection process, biological effects and transmission mode of cultivable *Serratia symbiotica* bacteria. Appl Environ Microbiol. 2019; AEM.02445-18, aem; AEM.02445-18v1. <u>https://doi.org/10.1128/AEM.02445-18</u> PMID: 30850430
- Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species are infected with *Wolbachia*?-a statistical analysis of current data. FEMS Microbiol Lett. 2008; 281: 215– 220. https://doi.org/10.1111/j.1574-6968.2008.01110.x PMID: 18312577
- Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of invertebrate biology. Nat Rev Microbiol. 2008; 6: 741–751. https://doi.org/10.1038/nrmicro1969 PMID: 18794912
- Dedeine F, Vavre F, Fleury F, Loppin B, Hochberg ME, Boulétreau M. Removing symbiotic Wolbachia bacteria specifically inhibits oogenesis in a parasitic wasp. Proc Natl Acad Sci. 2001; 98: 6247–6252. https://doi.org/10.1073/pnas.101304298 PMID: 11353833
- Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. Wolbachia as a bacteriocyte-associated nutritional mutualist. Proc Natl Acad Sci. 2010; 107: 769–774. <u>https://doi.org/10.1073/pnas.0911476107</u> PMID: 20080750
- Santos-Garcia D, Juravel K, Freilich S, Zchori-Fein E, Latorre A, Moya A, et al. To B or not to B: Comparative genomics suggests *Arsenophonus* as a source of B vitamins in whiteflies. Front Microbiol. 2018; 9. https://doi.org/10.3389/fmicb.2018.02254 PMID: 30319574
- Dodson BL, Hughes GL, Paul O, Matacchiero AC, Kramer LD, Rasgon JL. Wolbachia enhances West Nile Virus (WNV) infection in the mosquito Culex tarsalis. PLoS Negl Trop Dis. 2014; 8: e2965. <u>https://</u> doi.org/10.1371/journal.pntd.0002965 PMID: 25010200
- Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, et al. A Wolbachia symbiont in Aedes aegypti limits infection with Dengue, Chikungunya, and Plasmodium. Cell. 2009; 139: 1268– 1278. https://doi.org/10.1016/j.cell.2009.11.042 PMID: 20064373
- Chen R, Su X, Chen J, Jiang L, Qiao G-X. *Wolbachia* infection in two species: Novel views on the colonization ability of *Wolbachia* in aphids. Environ Entomol. [cited 4 Nov 2019]. https://doi.org/10.1093/ ee/nvz122 PMID: 31644794

- 86. Gómez-Valero L, Soriano-Navarro M, Pérez-Brocal V, Heddi A, Moya A, García-Verdugo JM, et al. Coexistence of *Wolbachia* with *Buchnera aphidicola* and a secondary symbiont in the aphid *Cinara cedri*. J Bacteriol. 2004; 186: 6626–6633. https://doi.org/10.1128/JB.186.19.6626-6633.2004 PMID: 15375144
- Augustinos AA, Santos-Garcia D, Dionyssopoulou E, Moreira M, Papapanagiotou A, Scarvelakis M, et al. Detection and characterization of *Wolbachia* infections in natural populations of aphids: Is the hidden diversity fully unraveled? PLOS ONE. 2011; 6: e28695. <u>https://doi.org/10.1371/journal.pone.</u> 0028695 PMID: 22174869
- Wang Z, Shen Z-R, Song Y, Liu H-Y, Li Z-X. Distribution and diversity of *Wolbachia* in different populations of the wheat aphid *Sitobion miscanthi* (Hemiptera: Aphididae) in China. EJE. 2013; 106: 49–55. https://doi.org/10.14411/eje.2009.007
- De Clerck C, Tsuchida T, Massart S, Lepoivre P, Francis F, Jijakli MH. Combination of genomic and proteomic approaches to characterize the symbiotic population of the banana aphid (Hemiptera: Aphididae). Environ Entomol. 2014; 43: 29–36. https://doi.org/10.1603/EN13107 PMID: 24472200
- ShaoJian L, Meng G, Na W, XiaoWei C, SuLing S, WanWan F, et al. Detection and phylogenetic analysis of *Wolbachia* in *Aphis gossypii* (Hemiptera: Aphididae) populations from different areas of Henan province, central China. Acta Entomol Sin. 2019; 62: 215–221.
- De Clerck C, Fujiwara A, Joncour P, Léonard S, Félix M-L, Francis F, et al. A metagenomic approach from aphid's hemolymph sheds light on the potential roles of co-existing endosymbionts. Microbiome. 2015; 3: 63. https://doi.org/10.1186/s40168-015-0130-5 PMID: 26667400
- 92. Manzano-Marín A. No evidence for *Wolbachia* as a nutritional co-obligate endosymbiont in the aphid *Pentalonia nigronervosa.* bioRxiv. 2019; 609511. https://doi.org/10.1101/609511
- West SA, Cook JM, Werren JH, Godfray HCJ. Wolbachia in two insect host–parasitoid communities. Mol Ecol. 1998; 7: 1457–1465. https://doi.org/10.1046/j.1365-294x.1998.00467.x PMID: 9819901
- Vavre F, Fleury F, Lepetit D, Fouillet P, Boulétreau M. Phylogenetic evidence for horizontal transmission of *Wolbachia* in host-parasitoid associations. Mol Biol Evol. 1999; 16: 1711–1723. https://doi.org/ 10.1093/oxfordjournals.molbev.a026084 PMID: 10605113
- Darby AC, Choi J-H, Wilkes T, Hughes MA, Werren JH, Hurst GDD, et al. Characteristics of the genome of Arsenophonus nasoniae, son-killer bacterium of the wasp Nasonia. Insect Mol Biol. 2010; 19: 75–89. https://doi.org/10.1111/j.1365-2583.2009.00950.x PMID: 20167019
- Michalik A, Schulz F, Michalik K, Wascher F, Horn M, Szklarzewicz T. Coexistence of novel gammaproteobacterial and Arsenophonus symbionts in the scale insect Greenisca brachypodii (Hemiptera, Coccomorpha: Eriococcidae). Environ Microbiol. 2018; 20: 1148–1157. <u>https://doi.org/10.1111/1462-2920.14057</u> PMID: 29393559
- Karimi S, Askari Seyahooei M, Izadi H, Bagheri A, Khodaygan P. Effect of Arsenophonus endosymbiont elimination on fitness of the date palm hopper, *Ommatissus lybicus* (Hemiptera: Tropiduchidae). Environ Entomol. 2019; 48: 614–622. https://doi.org/10.1093/ee/nvz047 PMID: 31095275
- Wulff JA, Buckman KA, Wu K, Heimpel GE, White JA. The endosymbiont Arsenophonus is widespread in soybean aphid, Aphis glycines, but does not provide protection from parasitoids or a fungal pathogen. PLOS ONE. 2013; 8: e62145. <u>https://doi.org/10.1371/journal.pone.0062145</u> PMID: 23614027
- 99. Duron O. Arsenophonus insect symbionts are commonly infected with APSE, a bacteriophage involved in protective symbiosis. FEMS Microbiol Ecol. 2014; 90: 184–194. https://doi.org/10.1111/ 1574-6941.12381 PMID: 25041857
- Tian P-P, Chang C-Y, Miao N-H, Li M-Y, Liu X-D. The facultative endosymbiont Arsenophonus infections alter aphid (*Aphis gossypii*) performance on the amino acid-deficient diet. Appl Environ Microbiol. 2019 [cited 8 Nov 2019]. https://doi.org/10.1128/AEM.01407-19 PMID: 31540983
- Ayoubi A, Talebi AA, Fathipour Y, Mehrabadi M. Coinfection of the secondary symbionts, *Hamiltonella defensa* and *Arsenophonus* sp. contribute to the performance of the major aphid pest, *Aphis gossypii* (Hemiptera: Aphididae). Insect Sci. 2020; 27: 86–98. https://doi.org/10.1111/1744-7917.12603 PMID: 29749703
- 102. Zhao Y, Zhang S, Luo J-Y, Wang C-Y, Lv L-M, Cui J-J. Bacterial communities of the cotton aphid Aphis gossypii associated with Bt cotton in northern China. Sci Rep. 2016; 6: 22958. <u>https://doi.org/10. 1038/srep22958</u> PMID: 27079679
- 103. Vera-Ponce de León A, Ormeño-Orrillo E, Ramírez-Puebla ST, Rosenblueth M, Degli Esposti M, Martínez-Romero J, et al. *Candidatus Dactylopiibacterium carminicum*, a nitrogen-fixing symbiont of *Dactylopius* cochineal insects (Hemiptera: Coccoidea: Dactylopiidae). Genome Biol Evol. 2017; 9: 2237– 2250. https://doi.org/10.1093/gbe/evx156 PMID: 30605507
- 104. Bustamante-Brito R, Vera-Ponce de León A, Rosenblueth M, Martínez-Romero JC, Martínez-Romero E. Metatranscriptomic analysis of the bacterial symbiont *Dactylopiibacterium carminicum* from the

carmine cochineal *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae). Life. 2019; 9: 4. https://doi.org/10.3390/life9010004 PMID: 30609847

- 105. Marschner P, Yang C-H, Lieberei R, Crowley DE. Soil and plant specific effects on bacterial community composition in the rhizosphere. Soil Biol Biochem. 2001; 33: 1437–1445. <u>https://doi.org/10.1016/ S0038-0717(01)00052-9</u>
- Russell JA, Weldon S, Smith AH, Kim KL, Hu Y, Łukasik P, et al. Uncovering symbiont-driven genetic diversity across North American pea aphids. Mol Ecol. 2013; 22: 2045–2059. https://doi.org/10.1111/ mec.12211 PMID: 23379399
- 107. Suh S-O, Noda H, Blackwell M. Insect Symbiosis: Derivation of yeast-like endosymbionts within an entomopathogenic filamentous lineage. Mol Biol Evol. 2001; 18: 995–1000. <u>https://doi.org/10.1093/oxfordjournals.molbev.a003901</u> PMID: 11371588
- 108. Vogel KJ, Moran NA. Functional and evolutionary analysis of the genome of an obligate fungal symbiont. Genome Biol Evol. 2013; 5: 891–904. https://doi.org/10.1093/gbe/evt054 PMID: 23563967
- 109. Xue J, Zhou X, Zhang C-X, Yu L-L, Fan H-W, Wang Z, et al. Genomes of the rice pest brown planthopper and its endosymbionts reveal complex complementary contributions for host adaptation. Genome Biol. 2014; 15: 521. https://doi.org/10.1186/s13059-014-0521-0 PMID: 25609551
- Blackwell M. Made for each other: Ascomycete yeasts and insects. Microbiol Spectr. 2017; 5. <u>https://</u> doi.org/10.1128/microbiolspec.FUNK-0081-2016 PMID: 28597823
- 111. Matsuura Y, Moriyama M, Łukasik P, Vanderpool D, Tanahashi M, Meng X-Y, et al. Recurrent symbiont recruitment from fungal parasites in cicadas. Proc Natl Acad Sci. 2018; 115: E5970–E5979. https://doi.org/10.1073/pnas.1803245115 PMID: 29891654