



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Correspondence

Potential multicollinearity among NLR and other variables in the prediction model for the COVID-19 mortality


To the Editor,

I read with great interest the article by Vafadar Moradi et al. reporting the predictive role of neutrophil-to-lymphocyte ratio (NLR) on the one-month mortality of COVID-19 patients [1]. Based on the multivariate Cox regression analysis, authors concluded that older age, increased NLR and white blood cells (WBC) count were associated with the one-month mortality. I agree with their interpretation that increased NLR reflects an imbalance of the inflammatory response in COVID-19 patients, leading to the fatal outcome. However, I found some methodological issue to be considered in this study. When constructing the multivariate model, the authors employed both NLR and WBC as explanatory variables. Since the majority of leukocytes are neutrophils and the increase in WBC count in infectious disease is mainly due to the increase in neutrophils, there should be a significant correlation between NLR and WBC. However, the authors did not discuss this potential multicollinearity problem. In the recent study by Caillon et al., there was a strong correlation between NLR and WBC count in the COVID-19 patients, and neither of them was used in their prediction model for the in-hospital mortality [2]. Another concern in this study is the cut-off value they used in the survival analysis. The authors employed NLR value ≥ 3.3 as the threshold of predicting severe disease according to the previous study [3]. The first strange point is seen in the demographic data; the median values of NLR in the two groups, divided by oxygen saturation levels, are both higher than 3.3 (5.0 and 4.1 respectively). It is also unreasonable that the cut-off value for “disease severity” was used for the “mortality” analysis in the Kaplan-Meier curve. According to the recent meta-analysis by Li et al., half of the included studies used cut-off value greater than 6.5 for the mortality [4]. In view of this result, the cut-off value used by the authors seems to be a little too low. As a matter of fact, contributing factors to COVID-19

fatality depend on the characteristics of the cohort in which the study was conducted [5]. Therefore, the optimal cut-off value for the one-month mortality should be determined based on their own dataset. Taking into account the association between cardiovascular disease and COVID-19 severity [6], as well as the association between immune disease and cardiovascular disease [7], the NLR cut-off value can be largely influenced by the proportion of patients with cardiovascular comorbidities. Although I am not questioning the predictive role of NLR in COVID-19, above all issues should be cleared before the results of their research will be applied to the clinical practice.

Declaration of Competing Interest

None declared.

References

- [1] Vafadar Moradi E, et al. Increased age, neutrophil-to-lymphocyte ratio (NLR) and white blood cells count are associated with higher COVID-19 mortality. *Am J Emerg Med.* 2021;40:11–4.
- [2] Caillon A, et al. High systolic blood pressure at hospital admission is an important risk factor in models predicting outcome of COVID-19 patients. *Am J Hypertens.* 2021;34(3):282–90. <https://doi.org/10.1093/ajh/hpaa225>.
- [3] Yang AP, Liu J Ping, Tao W Qiang, Li H Ming. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *Int Immunopharmacol.* 2020;84.
- [4] Li X, et al. Predictive values of neutrophil-to-lymphocyte ratio on disease severity and mortality in COVID-19 patients: a systematic review and metaanalysis. *Crit Care.* 2020;24.
- [5] Macedo A, Gonçalves N, Audia Febra C. COVID-19 fatality rates in hospitalized patients: systematic review and meta-analysis. *Ann Epidemiol.* 2021. <https://doi.org/10.1016/j.annepidem.2021.02.012>.
- [6] Bae SA, Kim SR, Kim MN, Shim WJ, Park SM. Impact of cardiovascular disease and risk factors on fatal outcomes in patients with COVID-19 according to age: a systematic review and meta-analysis. *Heart.* 2020. <https://doi.org/10.1136/heartjnl-2020-317901>.
- [7] Higaki A, Caillon A, Paradis P, Schiffrin EL. Innate and innate-like immune system in hypertension and vascular injury. *Curr Hypertens Rep.* 2019;21.

Akinori Higaki

Department of Cardiology, Ehime Prefectural Central Hospital, Japan

E-mail address: keroplant83@gmail.com

5 March 2021