



# Crystal structure of 1-[2-(diethylazan-iumyl)ethyl]-3-methylimidazolium tetrachloridocuprate(II)

Gerhard Laus,<sup>a\*</sup> Volker Kahlenberg<sup>b</sup> and Herwig Schottenberger<sup>a</sup>

<sup>a</sup>University of Innsbruck, Faculty of Chemistry and Pharmacy, Innrain 80, 6020 Innsbruck, Austria, and <sup>b</sup>University of Innsbruck, Institute of Mineralogy and Petrography, Innrain 52, 6020 Innsbruck, Austria. \*Correspondence e-mail: gerhard.laus@uibk.ac.at

Received 26 March 2015; accepted 4 April 2015

Edited by K. Fejfarova, Institute of Macromolecular Chemistry, AS CR, v.v.i, Czech Republic


The title compound, (C<sub>10</sub>H<sub>21</sub>N<sub>3</sub>)[CuCl<sub>4</sub>], is composed of one 1-[2-(diethylazan-iumyl)ethyl]-3-methylimidazolium dication and a tetrachloridocuprate(II) dianion. The anion adopts a distorted tetrahedral geometry. Bifurcated interionic N—H...Cl hydrogen bonds and several C—H...Cl contacts are observed, leading to a layer-like arrangement of the components parallel to (100).

**Keywords:** crystal structure; copper(II) complex; tetrachloridocuprate; 1-[2-(diethylazan-iumyl)ethyl]-3-methylimidazolium dication; hydrogen bonding.

**CCDC reference:** 1057934

## 1. Related literature

For structures of related tetrachloridocuprates(II), see: Russell & Wallwork (1969); Głowska & Gilli (1989); Choi *et al.* (2002); Sun & Qu (2005); Elangovan *et al.* (2007*a,b*); Strasser *et al.* (2007). For details of the synthesis, see: Laus *et al.* (2012); Håkansson & Jagner (1990).



## 2. Experimental

### 2.1. Crystal data

(C<sub>10</sub>H<sub>21</sub>N<sub>3</sub>)[CuCl<sub>4</sub>]  
 $M_r = 388.64$   
 Monoclinic,  $P2_1/c$   
 $a = 17.0041$  (8) Å  
 $b = 7.1161$  (6) Å  
 $c = 14.4143$  (7) Å  
 $\beta = 112.956$  (6)<sup>o</sup>  
 $V = 1606.04$  (17) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 2.01$  mm<sup>-1</sup>  
 $T = 173$  K  
 $0.20 \times 0.16 \times 0.12$  mm

### 2.2. Data collection

Oxford Diffraction Gemini-R Ultra diffractometer  
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)  
 $T_{\min} = 0.875$ ,  $T_{\max} = 1$   
 10390 measured reflections  
 2991 independent reflections  
 2361 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.056$   
 $S = 0.98$   
 2991 reflections  
 166 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.34$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.34$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cu1—Cl4	2.2267 (7)	Cu1—Cl2	2.2456 (8)
Cu1—Cl3	2.2447 (6)	Cu1—Cl1	2.2644 (7)

**Table 2**

Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3N...Cl1	0.93	2.29	3.134 (2)	150
N3—H3N...Cl3	0.93	2.79	3.399 (2)	124
C2—H2...Cl4 <sup>i</sup>	0.95	2.66	3.480 (3)	145
C3—H3...Cl2 <sup>ii</sup>	0.95	2.75	3.423 (3)	128
C3—H3...Cl3 <sup>ii</sup>	0.95	2.77	3.537 (3)	138
C4—H4...Cl2 <sup>iii</sup>	0.95	2.84	3.608 (3)	139
C9—H9A...Cl1 <sup>iii</sup>	0.99	2.78	3.617 (3)	143

Symmetry codes: (i)  $x, y - 1, z$ ; (ii)  $-x, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iii)  $x, -y + \frac{1}{2}, z + \frac{1}{2}$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR2002* (Burla *et al.*, 2003); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97*.

## Acknowledgements

The authors are grateful to A. Schwärzler for technical assistance.

Supporting information for this paper is available from the IUCr electronic archives (Reference: FF2135).

## References

- Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
- Choi, S.-N., Lee, Y.-M., Lee, H.-W., Kang, S. K. & Kim, Y.-I. (2002). *Acta Cryst. E* **58**, m583–m585.
- Elangovan, A., Thamarachelvan, A., Ramu, A., Athimoolam, S. & Natarajan, S. (2007a). *Acta Cryst. E* **63**, m201–m203.
- Elangovan, A., Thamarachelvan, A., Ramu, A., Athimoolam, S. & Natarajan, S. (2007b). *Acta Cryst. E* **63**, m224–m226.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Główka, M. L. & Gilli, G. (1989). *Acta Cryst. C* **45**, 408–410.
- Håkansson, M. & Jagner, S. (1990). *Inorg. Chem.* **29**, 5241–5244.
- Laus, G., Wurst, K. & Schottenberger, H. (2012). *Z. Kristallogr. New Cryst. Struct.* **227**, 413–415.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Abingdon, England.
- Russell, J. H. & Wallwork, S. C. (1969). *Acta Cryst. B* **25**, 1691–1695.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Strasser, C. E., Cronje, S. & Raubenheimer, H. G. (2007). *Acta Cryst. E* **63**, m2915–m2916.
- Sun, X.-M. & Qu, Y. (2005). *Acta Cryst. E* **61**, m1360–m1362.

## supporting information

*Acta Cryst.* (2015). E71, m110–m111 [doi:10.1107/S2056989015006799]

## Crystal structure of 1-[2-(diethylazaniumyl)ethyl]-3-methylimidazolium tetrachloridocuprate(II)

Gerhard Laus, Volker Kahlenberg and Herwig Schottenberger

### S1. Comment

1-[2-(Diethylazaniumyl)ethyl]-3-methylimidazolium ions in deprotonated form can act as bidentate ligands (Laus *et al.*, 2012). In this work, it is shown that they are also capable of coordinating in protonated form. The structure of the ion pair of 3-(2-(diethylammonio)ethyl)-1-methylimidazolium tetrachlorocuprate (II) is shown in Figure 1. The anion adopts a distorted tetrahedral geometry. The Cl—Cu—Cl angles range from 97° to 134°. The heterocyclic rings of the cation are oriented parallel to the (3 7 2) and (3  $\bar{7}$  2) planes with an interplanar angle of 32.5°. The side chain is twisted out of the heterocyclic ring plane. In the crystal structure, the cations and anions are linked by N—H···Cl and C—H···Cl hydrogen bonds (Figure 2).


### S2. Experimental

The title compound, (C<sub>10</sub>H<sub>21</sub>N<sub>3</sub>)CuCl<sub>4</sub>, was obtained from the reaction of bis-(1-(2-(diethylamino)ethyl)-3-methylimidazolin-2-ylidene) di-silver(I) bis(bis(triflimide)) (Laus *et al.*, 2012) and carbonyl chlorocopper(I) (Håkansson & Jagner, 1990) in methanol under ambient conditions. This unconventional synthesis involves redox, protonation and complexation steps. Yellow-green single crystals were obtained from MeOH in modest yield. Melting point 118–120 °C.

### S3. Refinement

All hydrogen atoms were positioned geometrically and constrained to ride on their parent atoms with  $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{parent atom})$ .



**Figure 2**

Interionic contacts in the crystal structure of the title compound. Symmetry codes: (i)  $x, -1 + y, z$ ; (ii)  $-x, -1/2 + y, 3/2 - z$ ; (iii)  $x, 1/2 - y, 1/2 + z$ .

### 1-[2-(Diethylazaniumyl)ethyl]-3-methylimidazolium tetrachloridocuprate(II)

#### Crystal data

$(C_{10}H_{21}N_3)[CuCl_4]$   
 $M_r = 388.64$   
 Monoclinic,  $P2_1/c$   
 $a = 17.0041 (8) \text{ \AA}$   
 $b = 7.1161 (6) \text{ \AA}$   
 $c = 14.4143 (7) \text{ \AA}$   
 $\beta = 112.956 (6)^\circ$   
 $V = 1606.04 (17) \text{ \AA}^3$   
 $Z = 4$

$F(000) = 796$   
 $D_x = 1.607 \text{ Mg m}^{-3}$   
 Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 4805 reflections  
 $\theta = 3.1\text{--}28.5^\circ$   
 $\mu = 2.01 \text{ mm}^{-1}$   
 $T = 173 \text{ K}$   
 Fragment, yellow  
 $0.20 \times 0.16 \times 0.12 \text{ mm}$

#### Data collection

Oxford Diffraction Gemini-R Ultra  
 diffractometer  
 Graphite monochromator  
 Detector resolution:  $10.3822 \text{ pixels mm}^{-1}$   
 $\omega$  ( $1^\circ$  width) scans  
 Absorption correction: multi-scan  
 (*CrysAlis PRO*; Oxford Diffraction, 2010)  
 $T_{\min} = 0.875, T_{\max} = 1$

10390 measured reflections  
 2991 independent reflections  
 2361 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.045$   
 $\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 3.1^\circ$   
 $h = -20 \rightarrow 20$   
 $k = -6 \rightarrow 8$   
 $l = -17 \rightarrow 17$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.032$   
 $wR(F^2) = 0.056$   
 $S = 0.98$   
 2991 reflections  
 166 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0234P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

Special details

**Experimental.** Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.24089 (2)	0.54432 (5)	0.77447 (2)	0.01040 (9)
Cl1	0.33356 (4)	0.30011 (9)	0.81388 (5)	0.01280 (16)
Cl2	0.12793 (4)	0.62361 (10)	0.63485 (5)	0.01575 (16)
Cl3	0.17620 (4)	0.47770 (9)	0.87973 (4)	0.01224 (15)
Cl4	0.33226 (4)	0.76469 (10)	0.76985 (6)	0.02164 (18)
N3	0.34210 (13)	0.2088 (3)	1.03033 (15)	0.0089 (5)
H3N	0.3191	0.2461	0.9632	0.011*
N2	0.16575 (13)	-0.0096 (3)	0.88310 (15)	0.0089 (5)
N1	0.08640 (14)	0.0672 (3)	0.73070 (15)	0.0118 (5)
C9	0.35838 (17)	0.3852 (4)	1.09319 (19)	0.0130 (6)
H9A	0.3812	0.3503	1.1654	0.016*
H9B	0.3036	0.4519	1.0777	0.016*
C2	0.16480 (17)	0.0163 (4)	0.79170 (19)	0.0125 (6)
H2	0.2121	0.0012	0.7729	0.015*
C8	0.46893 (17)	0.0322 (4)	1.15266 (19)	0.0164 (6)
H8A	0.4892	0.1404	1.1978	0.025*
H8B	0.5178	-0.0452	1.1564	0.025*
H8C	0.4301	-0.0429	1.1731	0.025*
C6	0.27576 (16)	0.0917 (4)	1.04753 (19)	0.0105 (6)
H6A	0.2283	0.1748	1.0451	0.013*
H6B	0.3012	0.0373	1.1161	0.013*
C5	0.23917 (16)	-0.0678 (4)	0.97205 (19)	0.0124 (6)
H5A	0.2842	-0.1156	0.9508	0.015*

H5B	0.2219	-0.1721	1.0055	0.015*
C7	0.42237 (16)	0.1000 (4)	1.04603 (19)	0.0132 (6)
H7A	0.4613	0.1809	1.0273	0.016*
H7B	0.4074	-0.0098	1.0003	0.016*
C10	0.42071 (19)	0.5163 (4)	1.0745 (2)	0.0193 (7)
H10A	0.4779	0.4604	1.1017	0.029*
H10B	0.4218	0.6369	1.1079	0.029*
H10C	0.4028	0.5367	1.0019	0.029*
C1	0.05856 (19)	0.1108 (4)	0.62341 (19)	0.0215 (7)
H1A	0.0668	0.2451	0.6151	0.032*
H1B	-0.002	0.0793	0.5887	0.032*
H1C	0.0922	0.0373	0.5946	0.032*
C3	0.03637 (17)	0.0745 (4)	0.78593 (19)	0.0138 (6)
H3	-0.0224	0.1072	0.7612	0.017*
C4	0.08539 (16)	0.0273 (4)	0.8808 (2)	0.0134 (6)
H4	0.0682	0.0206	0.9359	0.016*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.01055 (17)	0.00944 (18)	0.01216 (18)	-0.00016 (15)	0.00545 (14)	0.00046 (14)
C11	0.0154 (4)	0.0136 (4)	0.0110 (3)	0.0043 (3)	0.0068 (3)	0.0018 (3)
C12	0.0162 (4)	0.0190 (4)	0.0114 (3)	0.0037 (3)	0.0048 (3)	0.0018 (3)
C13	0.0113 (3)	0.0140 (4)	0.0119 (3)	0.0003 (3)	0.0051 (3)	0.0012 (3)
C14	0.0185 (4)	0.0115 (4)	0.0420 (5)	-0.0021 (3)	0.0196 (4)	-0.0007 (3)
N3	0.0081 (12)	0.0116 (12)	0.0062 (11)	0.0000 (10)	0.0020 (10)	0.0009 (9)
N2	0.0094 (12)	0.0063 (13)	0.0093 (11)	-0.0013 (9)	0.0017 (10)	-0.0026 (9)
N1	0.0130 (12)	0.0092 (13)	0.0098 (12)	-0.0028 (10)	0.0007 (10)	-0.0026 (9)
C9	0.0158 (15)	0.0119 (15)	0.0110 (14)	-0.0007 (12)	0.0049 (12)	-0.0017 (11)
C2	0.0153 (15)	0.0090 (15)	0.0138 (15)	-0.0033 (12)	0.0063 (13)	-0.0046 (11)
C8	0.0107 (14)	0.0169 (16)	0.0182 (15)	0.0005 (13)	0.0020 (12)	0.0005 (13)
C6	0.0070 (14)	0.0142 (16)	0.0101 (14)	-0.0004 (11)	0.0029 (12)	0.0007 (11)
C5	0.0095 (14)	0.0109 (16)	0.0163 (15)	0.0002 (12)	0.0045 (12)	0.0023 (12)
C7	0.0092 (14)	0.0142 (16)	0.0184 (15)	0.0006 (12)	0.0079 (12)	0.0011 (12)
C10	0.0273 (17)	0.0147 (17)	0.0151 (15)	-0.0083 (13)	0.0075 (13)	-0.0039 (12)
C1	0.0277 (18)	0.0217 (18)	0.0119 (15)	0.0021 (14)	0.0044 (14)	0.0009 (13)
C3	0.0098 (14)	0.0136 (17)	0.0179 (15)	-0.0019 (12)	0.0055 (13)	-0.0039 (12)
C4	0.0106 (14)	0.0146 (15)	0.0165 (15)	-0.0034 (12)	0.0070 (13)	-0.0037 (12)

*Geometric parameters (Å, °)*

Cu1—C14	2.2267 (7)	C8—H8A	0.98
Cu1—C13	2.2447 (6)	C8—H8B	0.98
Cu1—C12	2.2456 (8)	C8—H8C	0.98
Cu1—C11	2.2644 (7)	C6—C5	1.527 (4)
N3—C6	1.499 (3)	C6—H6A	0.99
N3—C7	1.507 (3)	C6—H6B	0.99
N3—C9	1.509 (3)	C5—H5A	0.99

N3—H3N	0.93	C5—H5B	0.99
N2—C2	1.324 (3)	C7—H7A	0.99
N2—C4	1.379 (3)	C7—H7B	0.99
N2—C5	1.458 (3)	C10—H10A	0.98
N1—C2	1.329 (3)	C10—H10B	0.98
N1—C3	1.374 (3)	C10—H10C	0.98
N1—C1	1.463 (3)	C1—H1A	0.98
C9—C10	1.513 (4)	C1—H1B	0.98
C9—H9A	0.99	C1—H1C	0.98
C9—H9B	0.99	C3—C4	1.337 (4)
C2—H2	0.95	C3—H3	0.95
C8—C7	1.508 (4)	C4—H4	0.95
C14—Cu1—C13	134.47 (3)	C5—C6—H6A	108.6
C14—Cu1—C12	99.16 (3)	N3—C6—H6B	108.6
C13—Cu1—C12	100.69 (3)	C5—C6—H6B	108.6
C14—Cu1—C11	97.04 (3)	H6A—C6—H6B	107.5
C13—Cu1—C11	98.33 (3)	N2—C5—C6	112.7 (2)
C12—Cu1—C11	133.31 (3)	N2—C5—H5A	109
C6—N3—C7	112.7 (2)	C6—C5—H5A	109
C6—N3—C9	109.68 (18)	N2—C5—H5B	109
C7—N3—C9	113.2 (2)	C6—C5—H5B	109
C6—N3—H3N	107	H5A—C5—H5B	107.8
C7—N3—H3N	107	N3—C7—C8	113.7 (2)
C9—N3—H3N	107	N3—C7—H7A	108.8
C2—N2—C4	108.8 (2)	C8—C7—H7A	108.8
C2—N2—C5	126.0 (2)	N3—C7—H7B	108.8
C4—N2—C5	125.19 (19)	C8—C7—H7B	108.8
C2—N1—C3	108.4 (2)	H7A—C7—H7B	107.7
C2—N1—C1	125.8 (2)	C9—C10—H10A	109.5
C3—N1—C1	125.9 (2)	C9—C10—H10B	109.5
N3—C9—C10	112.5 (2)	H10A—C10—H10B	109.5
N3—C9—H9A	109.1	C9—C10—H10C	109.5
C10—C9—H9A	109.1	H10A—C10—H10C	109.5
N3—C9—H9B	109.1	H10B—C10—H10C	109.5
C10—C9—H9B	109.1	N1—C1—H1A	109.5
H9A—C9—H9B	107.8	N1—C1—H1B	109.5
N2—C2—N1	108.4 (2)	H1A—C1—H1B	109.5
N2—C2—H2	125.8	N1—C1—H1C	109.5
N1—C2—H2	125.8	H1A—C1—H1C	109.5
C7—C8—H8A	109.5	H1B—C1—H1C	109.5
C7—C8—H8B	109.5	C4—C3—N1	107.6 (2)
H8A—C8—H8B	109.5	C4—C3—H3	126.2
C7—C8—H8C	109.5	N1—C3—H3	126.2
H8A—C8—H8C	109.5	C3—C4—N2	106.8 (2)
H8B—C8—H8C	109.5	C3—C4—H4	126.6
N3—C6—C5	114.8 (2)	N2—C4—H4	126.6
N3—C6—H6A	108.6		



C6—N3—C9—C10	-176.3 (2)	C4—N2—C5—C6	-72.8 (3)
C7—N3—C9—C10	56.9 (3)	N3—C6—C5—N2	-88.4 (3)
C4—N2—C2—N1	-0.5 (3)	C6—N3—C7—C8	-63.9 (3)
C5—N2—C2—N1	179.5 (2)	C9—N3—C7—C8	61.3 (3)
C3—N1—C2—N2	0.4 (3)	C2—N1—C3—C4	-0.1 (3)
C1—N1—C2—N2	179.5 (2)	C1—N1—C3—C4	-179.2 (2)
C7—N3—C6—C5	-65.8 (3)	N1—C3—C4—N2	-0.2 (3)
C9—N3—C6—C5	167.1 (2)	C2—N2—C4—C3	0.4 (3)
C2—N2—C5—C6	107.2 (3)	C5—N2—C4—C3	-179.5 (2)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3 <i>N</i> ...C11	0.93	2.29	3.134 (2)	150
N3—H3 <i>N</i> ...C13	0.93	2.79	3.399 (2)	124
C2—H2...C14 <sup>i</sup>	0.95	2.66	3.480 (3)	145
C3—H3...C12 <sup>ii</sup>	0.95	2.75	3.423 (3)	128
C3—H3...C13 <sup>ii</sup>	0.95	2.77	3.537 (3)	138
C4—H4...C12 <sup>iii</sup>	0.95	2.84	3.608 (3)	139
C9—H9 <i>A</i> ...C11 <sup>iii</sup>	0.99	2.78	3.617 (3)	143

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x, y-1/2, -z+3/2$ ; (iii)  $x, -y+1/2, z+1/2$ .