






CASE REPORT

Pulmonary tuberculosis in an adult presenting with severe hyponatremia: A case report and review of literature

Shailes Paudel^{1,2}  | Maulik Dhanani^{2,3} | Krish Hitesh Patel^{2,4}  |
Naga Praneeth Vakkalagadda^{2,5}  | Vivek Sanker^{2,6}  | Abhiram Rao Damera^{2,7}  |
Umang Gupta^{2,8}  | Prakriti Bhandari^{1,2} 

¹Patan Academy of Health Sciences, Lalitpur, Nepal

²Team Erevnites, Trivandrum, India

³Southwestern University School of Medicine, Cebu City, Philippines

⁴Government Medical College, Surat, India

⁵Guntur Medical College, Guntur, Andhra Pradesh, India

⁶Noorul Islam Institute of Medical Sciences, Trivandrum, Kerala, India

⁷Mediciti Institute of Medical Sciences, Ghanpur, Telangana, India

⁸Nepalgunj medical college, Nepalgunj Banke, Nepal

Correspondence

Shailes Paudel, Patan Academy of Health Sciences, Lalitpur, Nepal.
Email: shailespaudel@gmail.com

Key Clinical Message

Identifying pulmonary pathology while evaluating electrolyte disorders is crucial for optimal patient management. Physicians working in endemic regions of tuberculosis should consider this pathology as a differential for electrolyte imbalances.

Abstract

Hyponatremia, a common electrolyte imbalance, can arise from various underlying etiologies such as diuretics, diarrhea, vomiting, congestive heart failure, and liver and renal disease. We present a case report of a 74-year-old man highlighting the association between pulmonary tuberculosis (TB) and the development of hyponatremia. GeneXpert assay of the patient's sputum sample led to the identification of underlying active pulmonary TB as the cause of hyponatremia. The patient was started on anti-TB therapy, and concurrent fluid restriction and sodium supplementation were initiated to correct the electrolyte imbalance. Over the next 3 days, the patient demonstrated clinical improvement with the resolution of hyponatremia. This case also highlights the importance of considering TB as a potential etiology in patients presenting with hyponatremia, especially in endemic areas. Further research is warranted to explore the mechanistic pathways linking pulmonary TB and hyponatremia, aiding in the development of targeted therapeutic interventions.

KEYWORDS

dyselectrolytemia, hyponatremia, mycobacterium, pulmonary tuberculosis

1 | INTRODUCTION

Hyponatremia is defined as the depletion of sodium levels in the blood below 135 mEq/L. Sodium is essential for proper nerve and muscle function and therefore hyponatremia

can cause weakness, nausea, headache, confusion, and in severe cases seizures and coma. Common causes include diuretics, vomiting, diarrhea, congestive heart failure, and renal and liver disease.¹ One of the uncommon causes of hyponatremia is pulmonary tuberculosis (TB) which is

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

caused by the bacteria *Mycobacterium tuberculosis* that commonly affects the respiratory system but is also known to cause systemic complications.

Tuberculosis can cause hyponatremia by a variety of mechanisms, including local invasion of the adrenal glands (adrenal insufficiency), local invasion of the hypothalamus or pituitary gland, Tubercular meningitis, and incorrect ADH secretion via pulmonary infection.² Timely diagnosis and early management of hyponatremia in individuals with pulmonary TB is essential to prevent complications and improve patient outcomes. We present an unusual case of a 74-year-old man who presented in the emergency department with decreased consciousness and weakness of the bilateral lower limb with retention of urine.

2 | CASE PRESENTATION

A 74-year-old man presented to the emergency department with decreased consciousness and weakness in both lower limbs for about 6 h for which he was brought to the emergency department. The patient had a history of recurrent urinary retention, but no fever or weight loss. He had a smoking history of 25 pack-years and was a non-alcoholic. The patient had a past medical history of pulmonary TB (left lung), which was successfully treated with antitubercular drugs under Directly Observed Treatment Short-course (DOTS) 15 years ago.

On examination, the patient was thin build and oriented to place but not to time and person. Chest examination revealed bilateral scattered wheeze over most of the lung fields, with bronchial breath sounds heard over the right upper lung region. Neurological examination showed bilateral lower limb weakness (power of 3/5) with intact sensations. Abdominal examination revealed a palpable bladder that was slightly tender. Other examinations were unremarkable. Bedside glucometer measurements indicated normal blood glucose levels.

Emergency blood and urine investigations (Table 1) revealed severe hyponatremia, an elevated erythrocyte sedimentation rate (ESR), and raised urine sodium concentration. A computed tomography (CT) head scan was performed to rule out stroke, which showed age-related cortical atrophy. Magnetic resonance (MR) spine imaging showed normal findings. Chest x-ray revealed areas of fibrosis (Figure 1). Medical records showed normal CT chest imaging 8 months ago and the patient refused for second CT scan of the chest (Figure 2). Sputum samples were collected for Ziehl–Neelsen (ZN) stain and Gram stain examinations, which yielded normal results. GeneXpert testing on the sputum samples detected a low amount of *Mycobacterial Tuberculosis*. Furthermore, line probe assay was conducted for drug susceptibility testing which did not show resistance to isoniazid and rifampicin.

The patient's hyponatremia was gradually corrected over 3 days, and he was discharged with antitubercular drugs. At the 1-month follow-up, the patient demonstrated significant improvement in his symptoms and electrolytes (follow-up Sodium was 142 mmol/L) and tolerated the medications without experiencing any complications.

3 | DISCUSSION

The most typical presentation of TB, an ancient human disease caused by the bacteria *Mycobacterium tuberculosis*, is respiratory symptoms.³ But TB presents itself in a variety of ways since it affects multiple systems. The respiratory system, gastrointestinal (GI) system, lymphoreticular system, skin, central nervous system, musculoskeletal system, reproductive system, and liver are the organ systems most frequently impacted.^{4,5} In this case report, we describe a patient presenting with severe hyponatremia in the context of pulmonary TB.

A serum sodium content of less than 135 mEq/L is considered hyponatremia, albeit this definition may vary

TABLE 1 Routine laboratory investigations.

Variables	Sample	Values	Laboratory reference values
Random glucose	Blood	81 mg/dL	70–100 mg/dL
Blood urea nitrogen	Blood	11 mg/dL	6–16 mg/dL
Creatinine	Blood	0.6 mg/dL	0.6–1.3 mg/dL
Calcium	Blood	9.4 mg/dL	8.6–10.3 mg/dL
Sodium	Blood	114 mmol/L	135–145 mmol/L
Potassium	Blood	4.4 mmol/L	3.5–5.0 mmol/L
Sodium	Urine	89 mmol/L	20 mmol/L
Osmolality	Urine	392 mOsm/kg	50–1200 mOsm/kg
Erythrocyte sedimentation rate (ESR)		52 mm in 1st hour	0–15 mm in 1st hour (Men)



FIGURE 1 Chest x-ray showing diffuse areas of fibrosis in the right upper lobe with deviation of trachea to the right.




FIGURE 2 HRCT Chest axial image just distal to the carina shows a focal patchy area of fibrotic change in the apico-posterior segment of the upper lobe of the left lung, as denoted by the black arrow.

slightly depending on the standards used by different laboratories. A frequent electrolyte disorder known as hyponatremia is brought on by an excess of total body water in comparison to total body sodium concentration. A ratio imbalance known as hyponatremia occurs when the body's total water content exceeds its total solute content. The two primary compartments of total body water (TBW) are extracellular fluid (ECF), which makes up one-third, and intracellular fluid (ICF), which makes up the other two-thirds. ECF's main solute is sodium, while ICF's main solute is potassium.⁶

TABLE 2 List of similar cases published in literature.

Study; Country; year	Demographic features	Past History	Duration of illness	Investigations	Treatment given
Miliary Tuberculosis Presenting with Hyponatremia and ARDS in an 82-Year-Old Immunocompetent Female; Spain; 2018 ⁹	82 years Female	Type 2 diabetes mellitus and atrial fibrillation	6 days	Normal physical examination, initial blood sodium = 123 mmol/dL, Normal chest x-ray, Chest CT with ground glass areas, and sputum culture showed M. TB.	Ciprofloxacin followed by meropenem (The patient expired before the diagnosis was made)
Miliary tuberculosis presenting with hyponatremia and thrombocytopenia; Canada; 1976 ⁸	74 years Female	None	14 days	Chest x-ray suggestive of pulmonary congestion	Isoniazid, rifampin, ethambutol, prednisone, vincristine and fluid restriction
A case of mild pulmonary tuberculosis complicated with the syndrome of inappropriate antidiuretic hormone secretion which caused impaired consciousness; Japan; 2011 ¹⁰	81 years Female	None	-	Chest x-ray suggestive of unilateral infiltration, mycobacterium detected on sputum smear	Sodium loading with fluid restriction and antitubercular therapy
A case of pulmonary tuberculosis initially presented with the syndrome of inappropriate secretion of antidiuretic hormone (SIADH); Japan; 2003 ¹¹	90 years Male	None	-	Chest x-ray showed pulmonary infiltration, mycobacterium detected on sputum smear	Sodium loading with fluid restriction and antitubercular therapy

A Study in 1969 by Chung et al. demonstrated that 11% of patients with active TB had Hyponatremia and the main cause was SIADH.⁷ Cockcroft et al. reported a 74-year-old woman with miliary TB which had been complicated by severe hyponatremia due to SIADH.⁸ Lee et al. reported an unusual case of PTB presenting as Hyponatremia with biochemical evidence of ectopic antidiuretic hormone production as a possible mechanism causing hyponatremia.²

A few similar cases in the literature are described below (Table 2):

The underlying pathophysiology of hyponatremia in pulmonary TB is likely multifactorial. Firstly, the pulmonary infection can lead to an excessive release of inflammatory cytokines, such as interleukin-6 and tumor necrosis factor-alpha, which may stimulate the secretion of antidiuretic hormone (ADH) from the posterior pituitary gland. The increased ADH levels result in impaired water excretion, leading to dilutional hyponatremia. Secondly, pulmonary TB can cause systemic inflammation and oxidative stress, which may affect the renal tubules' ability to regulate sodium and water balance. Additionally, tuberculous involvement of the central nervous system can disrupt the normal regulation of ADH release, further contributing to hyponatremia.^{12–14} Patients with PTB who are older and have an increased CRP level are likely to have hyponatremia.¹⁵

4 | CONCLUSION

In conclusion, we presented a case of a 74-year-old man with a history of pulmonary TB who showed signs of impaired consciousness, lower limb weakness, and severe hyponatremia. Mycobacterium TB was detected through GeneXpert. Our case indicates the significance of considering pulmonary TB as a possible cause of electrolyte imbalance. Particularly in regions where TB is endemic, healthcare professionals should not undervalue the link between pulmonary TB and electrolyte abnormalities. To avoid potential problems and improve patient outcomes, it is essential to identify and treat dyselectrolytemia in patients with respiratory symptoms. When dealing with electrolyte imbalances in cases presenting with these symptoms, healthcare providers should include TB in the differential diagnosis.

AUTHOR CONTRIBUTIONS

Shailes Paudel: Conceptualization; data curation; formal analysis; investigation; methodology; writing – original draft; writing – review and editing. **Maulik Dhanani:** Funding acquisition; investigation; methodology; writing – original draft; writing – review and editing. **Krish Hitesh Patel:** Funding acquisition; investigation; methodology; writing – original draft; writing – review and

editing. **Naga Praneeth Vakkalagadda:** Funding acquisition; investigation; methodology; writing – original draft; writing – review and editing. **Vivek Sanker:** Formal analysis; methodology; project administration; supervision; validation; visualization; writing – original draft; writing – review and editing. **Abhiram Rao Damera:** Investigation; methodology; visualization; writing – original draft; writing – review and editing. **Umang Gupta:** Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology. **Prakriti Bhandari:** Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology.

ACKNOWLEDGMENTS

We would like to thank the Department of Internal Medicine, Patan Academy of Health Sciences for their collaboration in this study.

CONFLICT OF INTEREST STATEMENT

None declared.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.




ETHICS STATEMENT

Ethics approval was not required for the case report as per the country's guidelines.

CONSENT

Written informed consent was obtained from the patient to publish this report.

ORCID

Shailes Paudel  <https://orcid.org/0000-0002-2405-6952>
 Krish Hitesh Patel  <https://orcid.org/0000-0001-6719-8996>
 Naga Praneeth Vakkalagadda  <https://orcid.org/0000-0002-3879-1739>
 Vivek Sanker  <https://orcid.org/0000-0003-0615-8397>
 Abhiram Rao Damera  <https://orcid.org/0009-0009-1693-5648>
 Umang Gupta  <https://orcid.org/0000-0002-7911-8663>
 Prakriti Bhandari  <https://orcid.org/0009-0002-9689-8742>

REFERENCES

- Rondon H, Badireddy M. Hyponatremia. *StatPearls [Internet]*. StatPearls Publishing; 2023.
- Jonaidi Jafari N, Izadi M, Sarrafzadeh F, Heidari A, Ranjbar R, Saburi A. Hyponatremia due to pulmonary tuberculosis: review of 200 cases. *Nephrourol Mon.* 2013;5(1):687-691. doi:10.5812/numonthly.7091

3. Patologia sempre attuale e difficile da prevenire [tuberculosis: an ever present disease but difficult to prevent]. *Ig Sanita Pubbl.* 2020;76(1):59-66.
4. Mbuh TP, Ane-Anyangwe I, Adeline W, Thumamo Pokam BD, Meriki HD, Mbacham WF. Bacteriologically confirmed extra pulmonary tuberculosis and treatment outcome of patients consulted and treated under program conditions in the littoral region of Cameroon. *BMC Pulm Med.* 2019;19(1):17. doi:[10.1186/s12890-018-0770-x](https://doi.org/10.1186/s12890-018-0770-x)
5. Mathiasen VD, Andersen PH, Johansen IS, Lillebaek T, Wejse C. Clinical features of tuberculous lymphadenitis in a low-incidence country. *Int J Infect Dis.* 2020;98:366-371. doi:[10.1016/j.ijid.2020.07.011](https://doi.org/10.1016/j.ijid.2020.07.011)
6. Overgaard-Steensen C. Initial approach to the hyponatremic patient. *Acta Anaesthesiol Scand.* 2011;55(2):139-148. doi:[10.1111/j.1399-6576.2010.02311.x](https://doi.org/10.1111/j.1399-6576.2010.02311.x)
7. Chung DK, Hubbard WW. Hyponatremia in untreated active pulmonary tuberculosis. *Am Rev Respir Dis.* 1969;99(4):595-597. doi:[10.1164/arrd.1969.99.4P1.595](https://doi.org/10.1164/arrd.1969.99.4P1.595)
8. Cockcroft DW, Donevan RE, Copland GM, Ibbott JW. Miliary tuberculosis presenting with hyponatremia and thrombocytopenia. *Can Med Assoc J.* 1976;115(9):871-873.
9. Herreros B, Plaza I, García R, Chichón M, Guerrero C, Pintor E. Miliary tuberculosis presenting with hyponatremia and ARDS in an 82-year-old immunocompetent female. *Pathogens.* 2018;7(3):72. doi:[10.3390/pathogens7030072](https://doi.org/10.3390/pathogens7030072)
10. Minami T, Wakamatsu K, Kumazoe H, Nagata N, Kajiki A, Kitahara Y. A case of mild pulmonary tuberculosis complicated with the syndrome of inappropriate antidiuretic hormone secretion which caused impaired consciousness. *Nihon Kokyuki Gakkai Zasshi.* 2011;49(8):607-611. Japanese.
11. Nishizawa Y, Yamamori C, Nishimura Y, et al. A case of pulmonary tuberculosis initially presented with syndrome of inappropriate secretion of antidiuretic hormone (SIADH). *Kekkaku.* 2003;78(1):27-31. Japanese.
12. Luies L, du Preez I. The Echo of pulmonary tuberculosis: mechanisms of clinical symptoms and other disease-induced systemic complications. *Clin Microbiol Rev.* 2020;33(4):e00036-20. doi:[10.1128/CMR.00036-20](https://doi.org/10.1128/CMR.00036-20)
13. Lee P, Ho KK. Hyponatremia in pulmonary TB: evidence of ectopic antidiuretic hormone production. *Chest.* 2010;137(1):207-208. doi:[10.1378/chest.09-0405](https://doi.org/10.1378/chest.09-0405)
14. Swart RM, Hoorn EJ, Betjes MG, Zietse R. Hyponatremia and inflammation: the emerging role of interleukin-6 in osmoregulation. *Nephron Physiol.* 2011;118(2):45-51. doi:[10.1159/000322238](https://doi.org/10.1159/000322238)
15. Yoshida T, Masuyama H, amagata H, et al. The incidence and risk factors of hyponatremia in pulmonary tuberculosis. *J Endocr Soc.* 2022;6(11):bvac130. doi:[10.1210/jendso/bvac130](https://doi.org/10.1210/jendso/bvac130)

How to cite this article: Paudel S, Dhanani M, Patel KH, et al. Pulmonary tuberculosis in an adult presenting with severe hyponatremia: A case report and review of literature. *Clin Case Rep.* 2024;12:e8469. doi:[10.1002/ccr3.8469](https://doi.org/10.1002/ccr3.8469)