

Stenting as a palliative method in the management of advanced squamous cell carcinoma of the oesophagus and gastro-oesophageal junction

Janusz Włodarczyk^{1,2}, Jarosław Kuźdżał^{1,2}

¹Department of Thoracic and Surgical Oncology, John Paul II Hospital, Krakow, Poland

²Department of Thoracic and Surgical Oncology, Jagiellonian University Medical College, Krakow, Poland

Videosurgery Miniinv 2016; 11 (1): 1–8

DOI: 10.5114/wiitm.2016.58979

Abstract

Advanced squamous cell carcinoma of the oesophagus and gastroesophageal junction usually requires palliative treatment, and the method of choice is stenting. There are several types of stents currently available, including: self-expandable metallic stents (fully or partially covered); self-expandable plastic stents; biodegradable stents. Each of the mentioned stents has its advantages and limitations, and requires a proper, patient-tailored selection. Due to the close anatomical relationship between the oesophagus and bronchial tree, some patients may require bilateral stenting. Oesophageal stenting may not only be considered as a palliative procedure, but can also be implemented to alleviate dysphagia during preoperative chemotherapy and/or radiotherapy.

Key words: dysphagia, oesophageal cancer, stenting, oesophageal fistula.

Introduction

Squamous cell carcinoma of the oesophagus is the fourth cause of death in males and seventeenth in females. There has been no change or a slight decrease in incidence over the last three decades, unlike gastro-oesophageal junction carcinoma, which exhibits a constant increase in incidence [1–5]. The development of squamous cell carcinoma of the oesophagus and gastro-oesophageal junction leads to dysphagia. More than 50% of patients present with an unresectable tumour, and progressive weight loss and dysphagia require palliative treatment. Among the many available methods of palliation, stenting, laser therapy, chemoradiation, and photodynamic therapy should be considered. However, stenting is the method of choice [6–8]. This is because of its technical simplicity, wide availability and immediate alleviation of dysphagia.

For the first time, in 1885, the prototype of the stent was used for intubation of the strictured oesophagus, and rapid development of stenting was observed with the development of modern endoscopy [9]. In the 1970s and 1980s, rigid, semi-rigid and plastic stents were used (Photos 1 and 2), associated with a high percentage of complications, such as perforation, stent migration, and mortality [10]. The introduction of self-expanding metallic stents enabled their safe and widespread use. Domschke *et al.* used such a stent for the first time in 1990 [11]. In 1991, the modified, silicone-covered Gianturco expandable metallic stent was introduced by Song *et al.* [12].

The aim of the present study is to analyse the benefits of stents, as well as their disadvantages, availability, and novel options of palliative management of oesophageal and gastro-oesophageal junction carcinoma.

Address for correspondence

Janusz Włodarczyk MD, PhD, Department of Thoracic and Surgical Oncology, John Paul II Hospital, 80 Prądnicka St, 31-202 Krakow, Poland, phone: +48 691 893 853, e-mail: jr.wlodarczyk@gmail.com




Photo 1. Uncovered and partially covered self-expanding metal stents (PCSEMS) from left to right: Ultraflex (uncovered) (Boston Scientific) Ultraflex (partial covered) (Boston Scientific), Evolution (Cook)

Stents

Currently, various stents are commercially available in Europe, America and Asia, which differ in their structure, deployment system, required pre-dilatation, coverage, and preparation for introduction. In general, stents should be characterised by a thin wall, elasticity, easy application, and low complica-

tion rate. Oesophageal stents (Photos 3 and 4) may be categorised as follows:

- A. Depending on the coverage (Photos 1, 2):
 - 1) partially covered self-expandable stents – e.g. Ultraflex, Esopha-Coil, Flamingo, Gianturco Z-stent, Evolution;
 - 2) fully covered self-expandable stents – e.g. Alimaxx-E, Niti-S, SX-Ella, Wallflex, Polyflex.
- B. Depending on the material used:
 - 1) nitinol – e.g. Ultraflex, Esopha-Coil, Wallflex, Niti-S, Choo, Hanaroo, Evolution, Alimaxx-E;
 - 2) steel – e.g. Gianturco Z-stent, Flamingo, Wallstent II;
 - 3) plastic – e.g. Polyflex.
 - 4) polydioxanone – e.g. Ella (Photo 3).
- C. Depending on function:
 - 1) with an anti-reflux valve – e.g. Ella, Dostent;
 - 2) with an anti-migration mechanisms – e.g. Alimaxx.
- D. Depending on release:
 - 1) distal – e.g. Ultraflex;
 - 2) proximal – e.g. Ultraflex.

The introduction of self-expanding metallic stents (SEMS) resulted in the elimination of rigid prostheses, which were associated with high incidence of severe complications. Covered stents constitute the largest and most commonly used




Photo 2. Fully covered self-expanding metal stents (FCSEMS) and partial covered self-expanding metal stents (PCSEMS) from left to right: Wallflex (Boston Scientific), Evolution (Cook), Niti-S double oesophageal stent (Taewoong), Niti-S Beta 2 (Taewoong) Niti-S S oesophageal stent (Taewoong), Flexella Plus (Ella), Evo – PCSEMS (Cook), Alimaxx ES (Endotek)

group. While uncovered SEMS were characterised by neoplastic tissue ingrowth, very difficult or impossible removal and difficult repositioning after implantation, the partially covered SEMS are associated with a much lower rate of these complications. They are, however, not free of problems, such as hypertrophic granulation of the proximal or distal end, leading to occlusion, migration, and fistula development [13].

The introduction of fully covered self-expanding stents in 2001 did not solve all problems. These stents are characterised by difficult implantation, due to their plastic construction. They have to be introduced through a guide, which is rigid, approximately 10 mm in diameter, and not always arranged in a manner suited to the intended site of implantation. These stents are also characterised by a tendency for migration.

Absorbable stents are a relatively new alternative for patients with oesophageal carcinoma. These stents may be used in the preoperative period, since their functioning is limited by the absorption time of ~12 weeks. They are made of an absorbable material – PDS. Their potential advantage is based on the possibility of absorption, although implantation requires wide oesophageal patency restoration, which limits the use of absorbable stents.



Photo 3. Fully covered plastic prosthesis on the left side: Polyflex (Boston Scientific) and SX Ella degradable (Ella) prosthesis on the right side for treatment of benign and malignant oesophageal stricture

An oesophageal stent may be applied after neoplastic stricture dilatation (up to the diameter of 10 mm). Commercially available oesophageal stents may be introduced under radiological or endoscopic con-



Photo 4. Types of prosthesis for oesophago-gastric junction cancer stenting, from left to right: Flamingo stent – PCSEMS (Boston Scientific), Cardia-Umbrella – FCSEMS (Micro-Tech), Cardia-Valve-stent – FCSEMS (Micro-Tech), Dua antireflux “wind sock” – PCSEMS (Cook), Boubella-E antireflux valve (FCSEMS) (Ella), Niti-S double anti-reflux valve (FCSEMS) (Taewoong)

trol, and therefore are equipped with markers, easily identified on the fluoroscopy image. Endoscopic clips can also be used to mark the area of dilatation.

Clinical evaluation

Oesophageal stenting is the method of choice in palliative management, but specific types of stents may also be used in cases of benign strictures and iatrogenic oesophageal perforations with mediastinitis [14].

The main indications for palliative oesophageal stenting include:

- dysphagia due to the inoperable oesophageal cancer;
- oesophageal compression due to mediastinal diseases (Hodgkin's disease, non-Hodgkin's lymphoma, inoperable lung cancer);
- small cell lung cancer infiltrating the oesophagus;
- thyroid cancer.

Patients requiring stenting are usually diagnosed with grade III and IV dysphagia and significant weight loss. It should also be remembered that these patients may require bronchial tree stenting, due to the close anatomical relationship between the oesophagus and bronchial tree (trachea and main bronchus). Therefore, during the endoscopic examination, the bronchial tree should also be assessed, especially when the neoplastic infiltration is located ~25 cm from the incision. In patients diagnosed with locally advanced cancer, compression of the left main bronchus and/or tracheal bifurcation may be observed, as well as oesophago-respiratory fistula. Deployment of the stent within the oesophagus may exacerbate compression of the airway and lead to acute asphyxia. Therefore, oesophageal stenting in such patients requires the preparation of the team to perform, if necessary, simultaneous bronchial tree stenting [15]. If this is anticipated on the basis of CT scan, airway stenting should be performed first.

Oesophageal dilatation

Patients with grade III and IV dysphagia require stricture dilatation before the stenting procedure. The dilatation technique in cases of oesophageal strictures varies among centres.

Generally, three methods of dilatation can be distinguished [16]:

- 1) mercury-filled rubber dilators – e.g. Maloney or Hurst;

- 2) guided polyvinyl dilators – e.g. Savary-Gilliard, Eder-Puestow, Celestin, Beuss, Biomed System American;

- 3) hydrostatic polyurethane balloons – e.g. TTS (through-the-scope), Rigiflex, Medi-Tech.

Mercury-filled rubber dilators are nowadays rarely used and have a mainly historical importance. They are most commonly employed in cases of short and medium-sized stricture dilatations. Wire-guided dilators are most commonly used. This is associated with their relative safety, 'tactile sensitivity' during the procedure, and low rate of complications.

Pneumatic balloon dilators are also very popular. The method is characterised by a low perforation rate. When using balloon dilators, there is no tactile control over stricture dilatation.

Oesophageal stent characteristics

Oesophageal stenting has been developing since the 1970s, when rigid stents were introduced by Wilson-Cook. Due to implantation difficulties and material quality, their use was associated with a high percentage of complications, especially oesophageal perforation, bronchial tree fistula formation, and a mortality rate of up to 10%. They were subsequently replaced by steel wire prostheses, which had fewer complications, although tumour ingrowth was an important drawback. Nowadays, fully or partially covered prostheses are used. These are characterised by a low complication rate [17].

The ideal prosthesis should have the following characteristics:

- appropriate diameter enabling full oral nutrition without obstruction after meals;
- low risk of migration, granulation tissue stent overgrowth, bleeding, and fistula formation;
- flexibility;
- technical simplicity of implantation, reposition, and removal;
- low risk of regurgitation.

The Ultraflex stent (Boston Scientific, Natick, USA) is made of nitinol, and fully or partially covered. It is highly flexible. These stents are also available without covering. Depending on the model, they have either proximal or distal release systems and facilitate precise implantation without fluoroscopic guidance, under simple endoscopic control. After implantation, they shorten by up to 30%. The new generation of these fully or partially covered stents are silicone-covered nitinol prostheses. In contrast to

those mentioned above, they can be refolded if dilatation amounts to 75%. This stent is equipped with 'progressive step flared ends', preventing migration.

Gianturco Z-stents (Cook, Winston Salem, USA) are partially covered steel-wire prostheses, available with an anti-reflux valve.

The Evolution stent (Cook, Winston Salem, USA) is a nitinol, partially covered prosthesis. The stent is distally dilated with recapture possibility, prior to full dilatation (in case of 50% dilatation). The stent contains an inner and outer membrane, which is intended to prevent granulation tissue ingrowth and to facilitate food passage. It has an anti-migration mechanism in the form of uncoated flanges on both ends.

Alimaxx-E (Alveolus, Charlotte, NC, USA) is a fully silicone covered, nitinol stent. The prostheses are laser-cut from a nitinol tube, and have an anti-reflux mechanism in the form of 'fish scales'. Fluoroscopy is not required. An interesting solution was proposed in cases of gastro-oesophageal junction cancer stents – the cardia umbrella stent. Due to its umbrella design, this nitinol, fully covered prosthesis has the ability to adjust to the cardia area, preventing displacement.

Niti-S (Taewong Medical, Seoul, Korea) is a 'double stent', with a non-covered external layer intended to prevent migration and a silicone-covered internal layer preventing granulation tissue ingrowth.

Choo-stent (M.I. Tech Medical, Seoul, Korea) is a nitinol, fully covered prosthesis, polyurethane-coated, distally released. An anti-reflux valve option is possible.

SX-Ella (Ella-CS, Hradec Kralove, Czech Republic) is a fully covered nitinol or steel prosthesis, distally released, with an anti-reflux valve and proximal pole ring preventing migration.

Biodegradable stents (Ella-CS, Hradec Kralove, Czech Republic) are non-covered prostheses made of PDS with a 12-week period of absorption. They are recommended during chemo-radiotherapy, prior to planned surgery, and in cases of benign lesions. The mechanism of implantation allows for multiple use. Prior to implantation, the stricture should be dilated to 12–14 Fr values; the rigidity of the guide and its location do not always allow for convenient stent dilatation.

Oesophageal stenting results and complications

The aim of stenting in patients with inoperable or unresectable oesophageal cancer is to obtain

oesophageal patency and complete oral nutrition. The development of endoscopy and technological progress currently enable the use of a number of oesophageal stents. The main characteristics include: safety and simplicity of implantation, bioelasticity, shape memory deployment, matching the shape of the stricture, resistance to deformation, and biodegradability [18]. Currently, steel, nitinol, cobalt, and an absorbable PDS polymer are used.

The procedure lasts a relatively short time, approximately 30 min, with a technical success rate reaching 100%. It is recommended to cover with the stent ~4 cm of the oesophagus above and below the stricture, although this is not always possible, especially in cases of proximal strictures [19]. Currently, with the fully covered plastic prostheses, stenting of the cervical oesophagus is also possible. Stent implantation is associated with low perioperative mortality ranging from 0 to 5%, while 30-day mortality ranges between 7% and 18% [20–25].

The stents that are currently used, despite relative good tolerance, are not free from side-effects and complications. One of the most common complications associated with stenting is granulation tissue overgrowth. Coverage with a polyurethane or silicone membrane protects from tumour ingrowth, but overgrowth beyond the ends of the stent and granulation tissue formation remain an issue. Granulation tissue is observed within two months after stent implantation in 47% of patients [26–30]. This problem, although rarely, is encountered also with fully covered plastic stents [31, 32]. The cause of granulation tissue overgrowth remains unknown. Stents of a diameter smaller than 20 mm overgrow much more slowly (SX Ella stent, Polyflex, Niti-S and Gianturco Z-stent) [33].

Another important complication associated with oesophageal stents is migration. It is usually observed in 20% of cases and it more often concerns fully covered plastic stents, as compared to those partially covered. Verschuur *et al.* demonstrated that Ultraflex stent migration was observed in 17% of patients, while Polyflex stent migration in 29%. Additionally, stents with an anti-migration mechanism (SX Ella, Niti-S, Alimaxx) do not always effectively prevent this complication [24, 28].

Fistulas to the bronchial tree are late, life-threatening complications. They occur in 10% of patients after stenting. Ferreira *et al.* observed their occurrence in 7 of 126 patients, while Uitdehaag *et al.*

observed them in 2 out of 4, after the use of SX Ella stents [29, 34, 35].

Bleeding after stenting is another life-threatening complication, and it reportedly occurs in 2–28% of patients [34, 36, 37].

Other, less severe complications include chest pain, foreign body sensation, and regurgitation. When considering the above-mentioned stents, no evidence of superiority of any of the prostheses was observed [23].

Gastro-oesophageal junction stenting

Gastro-oesophageal junction cancer stenting, as compared to proximal oesophageal stenting, is characterised by certain peculiarities and a higher complication rate. Stent migration is the most common complication. It is associated with fixation of the proximal end of the stent only – its distal segment being located in the stomach. Such a location also favours bleeding after stent placement. Siersma *et al.* connected this with two factors. First, the free segment of the stent assumes a forced position, pressing against the posterior wall of the oesophagus, which leads to its damage, ulceration, and bleeding. Secondly, the forced position of the stent, determined by the anatomy of the angle between the oesophagus and cardia, leads to exacerbation of reflux and worse swallowing quality [38, 39]. Vakil *et al.*, in a randomised study, observed migration of the stent placed across the cardia in 12% of cases [40].

It appears that the type of stent, especially its proximal diameter (flange diameter: Flamingo Wallstent – 30 mm, Wallstent II – 28 mm and Ultraflex – 28 mm), may play an important role in preventing migration. However, other studies showed that a larger stent diameter led to an increased rate of perforations and bleeding [33, 41, 42].

Vakil *et al.* demonstrated migration in 12% of cases, while the overall complication rate was 30%. Other most common complications include: fever, not necessarily associated with aspiration pneumonia, chest pain, granulation tissue overgrowth, food bolus obstruction, and perforation. No evidence of superiority of any of the stents was observed, when considering complications after stenting of the cardia [38, 43].

It is believed that adjuvant chemo- and/or radiotherapy may increase the proportion of complications. It should be remembered that dysphagia may

persist in a subset of patients following proper stent implantation, due to peritoneal metastases causing mechanical obstruction.

Quality of life

Quality of life is one of the most important factors of effective palliative treatment. Among the many questionnaires, the EORTC QLQ-C30 protocol is most often used. With 30 questions, it assesses the physical and emotional functioning, grade of dysphagia, nausea, vomiting, gastric content regurgitation, pain, degree of anxiety, and depression. The assessment of dysphagia is the key element considering the quality of life in patients with oesophageal carcinoma [44, 45].

Conclusions

Oesophageal metallic stents remain the gold standard in the palliative treatment in advanced squamous cell carcinoma of the oesophagus. Partially and fully covered stents are highly useful and well accepted. New stent designs and evolution of materials have reduced complication rates after stenting and increased the effectiveness and safety of palliative therapy. It should be remembered that some patients require bilateral stenting of the oesophagus and bronchial tree.

Conflict of interest

The authors declare no conflict of interest.

References

1. Parkin DM, Bray F, Ferlay J, et al. Global cancer statistics 2002. *Ca Cancer J Clin* 2005; 55: 74-108.
2. Brown LM, Devesa SS, Chow WH. Incidence of adenocarcinoma of the esophagus among white Americans by sex, stage, and age. *J Natl Cancer Inst* 2008; 100: 1184-7.
3. Trivers KF, Sabatino SA, Stewart SL. Trends in esophageal cancer incidence by histology, United States, 1998-2003. *Int J Cancer* 2008; 123: 1422-8.
4. Bosetti C, Levi F, Ferlay J, et al. Trends in oesophageal cancer incidence and mortality in Europe. *Int J Cancer* 2008; 122: 1118-29.
5. Siegel R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. *CA Cancer J Clin* 2014; 64: 9-29.
6. Lambert R. Palliation of carcinoma of the esophagus: is there a hope for cure? *Am J Gastroenterol* 1994; 89: S27-40.
7. Maier A, Anegg U, Fell B, et al. The effect of photodynamic therapy in a multimodale. *Lasers Surg Med* 2000; 26: 461-6.
8. Ludwig D, Dehne A, Burmester E, et al. Treatment of unresectable carcinoma of the esophagus or the gastroesophageal

- junction by mesh stents or without radiochemotherapy. *Int J Oncol* 1998; 13: 583-8.
9. Beynon J, Winston T, Thompson MH. Endoscopic insertion of Celestin tubes in carcinoma of the esophagus. *Gut* 1991; 84: 479-80.
 10. Fugger R, Niederle B, Jantsch H, et al. Endoscopic tube implantation for the palliation of malignant esophageal stenosis. *Endoscopy* 1990; 22: 100-4.
 11. Domschke W, Foerster EC, Matek W, et al. Self-expanding mesh stent for esophageal cancer stenosis. *Endoscopy* 1990; 22: 134-6.
 12. Song HY, Choi KC, Kwon HC, et al. Esophageal strictures: treatment with a new design of modified Gianturco stent. *Work in progress. Radiology* 1992; 184: 729-34.
 13. Raijman I, Siddique I, Ajani J, Lynch P. Palliation of malignant dysphagia and fistulae with coated expandable metal stents: experience with 101 patients. *Gastrointest Endosc* 1998; 48: 179-82.
 14. Safranek J, Geiger J, Vesely V, et al. Esophageal stents for less invasive treatment of mediastinitis. *Videosurgery Miniinvas* 2014; 9: 1-5.
 15. Sihoe AD, Wan IP, Yim AP. Airway stenting for unresectable esophageal cancer. *Surg Oncol* 2014; 13: 17-25.
 16. Hunter JG, Pellegrini CA. Surgery of the esophagus. *Surg Clin N Am* 1997; 77: 1202-3.
 17. Schembre D. Advances in esophageal stenting: the evolution of fully covered stents for malignant and benign disease *Adv Ther* 2010; 27: 413-25.
 18. Mijazawa T, Yamakido M, Ikeda S, et al. Implantation of ultraflex nitinol stents in malignant tracheobronchial stenoses. *Chest* 2000; 39: 1088-9.
 19. National Comprehensive Cancer Network Guidelines for the treatment of esophageal cancer 2014. Available at: www.nccn.org/professionals/physician_glsf_guidelines.asp#site 2014.
 20. McManus K, Khan I, McGuigan J. Self expanding oesophageal stents: strategies for re-intervention. *Endoscopy* 2001; 33: 601-4.
 21. Wilkes EA, Jackson IM, Cole AT, et al. Insertion of expandable metallic stents in esophageal cancer without fluoroscopy in safe and effective: a 5-year experience. *Gastrointest Endosc* 2007; 65: 923-9.
 22. Verschuur EM, Homs MY, Steyerberg EW, et al. A new esophageal stent design (Niti-S stent) for the prevention of migration: a prospective study in 42 patients. *Gastrointest Endosc* 2006; 63: 134-40.
 23. Burstow M, Kelly T, Panchani S, et al. Outcome of palliative esophageal stenting for malignant dysphagia: a retrospective analysis. *Dis Esophagus* 2009; 22: 519-25.
 24. Conigliaro R, Battaglia G, Repici A, et al. Polyflex stents for malignant oesophageal and oesophago-gastric stricture: a prospective, multicenter study. *Eur J Gastroenterol Hepatol* 2007; 19: 195-203.
 25. Ross WA, Alkassab F, Lynch PM, et al. Evolving role of self-expanding metal stents in the treatment of malignant dysphagia and fistulas. *Gastrointest Endosc* 2007; 65: 70-6.
 26. Mayoral W, Fleischer D Salcedo J, et al. Nonmalignant obstruction is a common problem with metal stents in the treatment of esophageal cancer. *Gastrointest Endosc* 2000; 51: 556-9.
 27. Homs MY, Steyerberg EW, Eijkenboom WM, et al. Single dose brachytherapy versus metal stent placement for the palliation of dysphagia from esophageal cancer: multicenter randomized trial. *Lancet* 2004; 364: 1497-504.
 28. Conio M, Repici A, Battaglia G, et al. A randomized prospective comparison of self expandable plastic stents and partially covered self expandable metal stents in the palliation of malignant esophageal dysphagia. *Am J Gastroenterol* 2007; 102: 2667-77.
 29. Uitdehaag MJ, Siersema PD, Spaander MC, et al. A new fully covered stent with antimigration properties for the palliation of malignant dysphagia: a prospective cohort study. *Gastrointest Endosc* 2010; 71: 600-5.
 30. Verschuur EM, Homs MY, Steyerberg EW, et al. A new esophageal stent design (Niti-S stent) for the prevention of migration: a prospective study in 42 patients. *Gastrointest Endosc* 2006; 63: 134-40.
 31. Sharma P, Kozarek R. Role of esophageal stents in benign and malignant disease. *Am J Gastroenterol* 2010; 105: 258-73.
 32. Bethge N, Somer A, Gross U, et al. Human tissue responses to metal stents implanted in vivo for the palliation of malignant stenoses. *Gastrointest Endosc* 1996; 43: 596-602.
 33. Verschuur EM, Steyerberg EW, Kuipers EJ, et al. Effect of stent size on complications and recurrent dysphagia in patients with esophageal or gastric cardia cancer. *Gastrointestinal Endosc* 2007; 65: 592-601.
 34. Ferreira F, Bastos P, Ribeiro A, et al. A comparative study between fluoroscopic and endoscopic guidance in palliative esophageal stent placement *Dis Esophagus* 2012; 25: 608-13.
 35. Lopes CV, Pesenti C, Bories E, et al. Self expandable metallic stents for palliative treatment of digestive cancer. *Clin Gastroenterol* 2008; 42: 991-6.
 36. Qiu G, Tao Y, Du X, et al. The impact of prior radiotherapy on fatal complications after self-expandable metallic stents (SEMS) for malignant dysphagia due to esophageal carcinoma. *Dis Esophagus* 2013; 26: 175-81.
 37. Choi SJ, Kim JH, Choi JW, et al. Fully covered, retrievable self-expanding metal stents (Niti-S) in palliation of malignant dysphagia: long-term results prospective study. *Scand J Gastroenterol* 2011; 46: 875-80.
 38. Siersma PD, Marcon M, Vakil N. Metal stents of the distal esophagus and gastric cardia. *Endoscopy* 2003; 35: 79-85.
 39. Spinelli P, Cerrai FG, Ciuffi M, et al. Endoscopic stent placement for cancer of the lower esophagus and gastric cardia. *Gastrointest Endosc* 1994; 40: 455-7.
 40. Vakil N, Morris AI, Marcon N, et al. A prospective, randomized, controlled trial of covered expandable metal stents in the palliation of malignant esophageal obstruction at the gastroesophageal junction. *Am J Gastroenterol* 2001; 96: 1791-6.
 41. Siersema PD, Hop WCJ, van Blankenstein M, et al. A new design metal stent (Flamingo stent) for palliation of malignant dysphagia: a prospective study. *Gastrointest Endosc* 2000; 51: 139-45.
 42. Siersema PD, Hop WCJ, van Blankenstein M, et al. A comparison of 3 types of covered metal stents for the palliation of patients with caused by esophagogastric carcinoma: a prospective, randomized study. *Gastrointest Endosc* 2001; 54: 145-53.

43. Sgourakis G, Gockel I, Radtke A, et al. The use of self-expanding stents in esophageal and gastroesophageal junction cancer palliation: a meta-analysis and meta-regression analysis of outcomes. *Dig Dis Sci* 2010; 55: 3018-30.
44. Blazeby JM. Measurement of outcome. *Surg Oncol* 2001; 10: 127-33.
45. Aaronson NK, Ahmedzai S, Bergman B, et al. The European Organization for Research and Treatment of Cancer QLQ-30: a quality-of life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85: 365-76.

Received: 29.04.2015, **accepted:** 7.03.2016.