

Kinetics of IL-7 and IL-15 Levels after Allogeneic Peripheral Blood Stem Cell Transplantation following Nonmyeloablative Conditioning

Muriel De Bock¹, Marianne Fillet¹, Muriel Hannon¹, Laurence Seidel², Marie-Paule Merville¹, André Gothot¹, Yves Beguin^{1,3}, Frédéric Baron^{1,3*}

1 Groupe Interdisciplinaire de Génoprotéomique Appliquée (GIGA)-I³, University of Liège, Liège, Belgium, **2** Department of Statistics, University of Liège, Liège, Belgium, **3** Department of Medicine, Division of Hematology, University and Centre Hospitalier Universitaire (CHU) of Liège, Liège, Belgium

Abstract

Background: We analysed kinetics of IL-7 and IL-15 levels in 70 patients given peripheral blood stem cells after nonmyeloablative conditioning.

Methods: EDTA-anticoagulated plasma and serum samples were obtained before conditioning and about once per week after transplantation until day 100. Samples were aliquoted and stored at -80°C within 3 hours after collection until measurement of cytokines. IL-7 and IL-15 levels were measured by ELISAs.

Results: Median IL-7 plasma levels remained below 6 pg/L throughout the first 100 days, although IL-7 plasma levels were significantly higher on days 7 (5.1 pg/mL, $P=0.002$), 14 (5.2 pg/mL, $P<0.001$), and 28 (5.1 pg/mL, $P=0.03$) (but not thereafter) than before transplantation (median value of 3.8 pg/mL). Median IL-15 serum levels were significantly higher on days 7 (12.5 pg/mL, $P<0.001$), 14 (10.5 pg/mL, $P<0.001$), and 28 (6.2 pg/mL, $P<0.001$) than before transplantation (median value of 2.4 pg/mL). Importantly, IL-7 and IL-15 levels on days 7 or 14 after transplantation did not predict grade II–IV acute GVHD.

Conclusions: These data suggest that IL-7 and IL-15 levels remain relatively low after nonmyeloablative transplantation, and that IL-7 and IL-15 levels early after nonmyeloablative transplantation do not predict for acute GVHD.

Citation: De Bock M, Fillet M, Hannon M, Seidel L, Merville M-P, et al. (2013) Kinetics of IL-7 and IL-15 Levels after Allogeneic Peripheral Blood Stem Cell Transplantation following Nonmyeloablative Conditioning. PLoS ONE 8(2): e55876. doi:10.1371/journal.pone.0055876

Editor: Maria Leite de Moraes, CNRS, France

Received: October 6, 2012; **Accepted:** January 3, 2013; **Published:** February 21, 2013

Copyright: © 2013 De Bock et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was in part supported by funds from the FNRS (<http://www2.frs-fnrs.be/>), the Belgian Foundation against Cancer (FBC) (<http://www.cancer.be/>), the anti-cancer (<http://www.cac.ulg.ac.be/>) and the Leon Fredericq foundations (<http://www.fondsleonfredericq.be/>) from the ULg, the CHU of Liège, and by the Terry Fox foundation (<http://www.terryfox.org/>). MdB and MH are Télévie Research Fellows, and MPM and FB are Senior Research Associates at the National Fund for Scientific Research (FNRS) Belgium. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: f.baron@ulg.ac.be

Introduction

Allogeneic hematopoietic stem cell transplantation (allo-HSCT) following a high dose conditioning regimen has been the best treatment option for many young patients with hematological disorders. The antitumor activity of this approach is based not only on high dose chemo-radiotherapy given in the conditioning regimen but also on immune-mediated graft-versus-tumor effects [1,2]. These observations are the basis of the development of allo-HSCT following nonmyeloablative conditioning, in which eradication of malignant cells depends on graft-versus-tumor effects [3–6].

T-cell recovery after allo-HSCT following high-dose conditioning depends on both homeostatic peripheral expansion (HPE) of donor T cells contained in the graft, and T cell neo-production from donor hematopoietic stem cells (thymo-dependent pathway) [7–15]. In young patients given myeloablative allo-HSCT, most circulating T cells during the first months following HSCT are the

progeny of T cells infused with the grafts [16], while neogenesis of T cells by the thymus plays an increasing role in reconstituting the T cell pool beyond day 100 after allo-HSCT [17–22]. Since HPE allow the expansion of both NK cells and non-tolerant T cells, it is generally accepted that HPE is one of the driving force of graft-versus-tumor effects.

Several studies have demonstrated that IL-7 and IL-15 are the main driving forces of HPE after allo-HSCT following high-dose conditioning [7,23]. IL-7 is a γ -common chain cytokine that is secreted by stromal cells from multiple organs including thymus, bone marrow, and lymphoid organs. IL-7 is required for human T cell development since mutations in the IL-7 receptor alpha can lead to severe combined immunodeficiency [24]. Administration of IL-7 has been shown to dramatically increase peripheral T cell numbers, primarily through augmentation of HPE [25–31]. IL-15 is another γ -common chain cytokine secreted by antigen-presenting cells, bone marrow stroma, thymic epithelium, and epithelial cells in the kidney, skin, and intestines [32]. IL-15 plays

an important role in the development and function of NK cells, and of NK/T cells, and is required for optimal proliferation of CD8⁺ T cells and for homeostatic proliferation of CD8⁺ memory T cells [33–39].

While high-dose conditioning regimens typically induce a profound lymphodepletion, progressive replacement of host-derived T cells by donor-derived T cells is the rule after nonmyeloablative conditioning [40,41]. This prompted us to analyze the kinetics of IL-7 and IL-15 blood levels after allo-HSCT following a nonmyeloablative conditioning with the aim of determining whether there is a rationale for boosting HPE and perhaps graft-versus-tumor effects in patients with high risk disease given grafts after nonmyeloablative conditioning by administering IL-7 and/or IL-15.

Patients and Methods

Patients and Donors

Data from 70 patients transplanted between March 2007 and April 2011 at the University of Liège were included in the study (Table 1). All patients were given G-CSF-mobilized peripheral blood stem cells (PBSC) after low-dose [2 Gy (n = 60), or 4 Gy (n = 10)] total body irradiation (TBI)-based nonmyeloablative regimen. Twenty-three nonmyeloablative recipients who were given PBSC from HLA-mismatched unrelated donors were co-transplanted with third party mesenchymal stromal cells (MSCs) as a potential way to prevent severe GVHD [42]. Further, 3 nonmyeloablative recipients were included in a double blind randomized study assessing the impact of MSC co-transplantation on transplantation outcomes. No patient was given in-vivo T cell depletion.

Ethics

Written informed consent was obtained from each patient to undergo allo-HSCT and to collect, store and analyze blood samples for research purposes. The Ethics Committee of the University of Liège (“Comité d’Ethique Hospitalo-Facultaire Universitaire de Liège”) approved the consent form as well as the current research study protocol (protocol #B707201112193).

Clinical Management

The clinical management has been performed as previously reported [43,44]. Chimerism levels among peripheral T-cells were generally measured with PCR-based analysis of polymorphic microsatellite regions (AmpFISTR® Identifiler®, Applied Biosystems, Lennik, Belgium) [43]. CD3 (T-cell) selection was carried out with the RosetteSep^K human T-cell enrichment kit (StemCell Technologies, Vancouver, Canada) [43,44].

Cytokines Levels

EDTA-anticoagulated plasma and serum samples were obtained before conditioning and about once time per week after transplantation until day 100. Samples were aliquoted and stored at –80°C within 3 hours after collection until measurement of cytokines. Kinetic courses of IL-7 production in plasma samples were evaluated before conditioning and approximately at days 7, 14, 28, 40, 60, 80 and 100 after allo-HSCT. IL-15 serum sample levels were assessed before conditioning and approximately at days 7, 14 and 28 after allo-HSCT. IL-7 and IL-15 levels were measured by ELISAs following the manufacturer’s protocol (High sensitivity IL-7 and IL-15 quantikine, R&D Systems, Minneapolis, MN, USA). The standard curve ranges for IL7 were 0.25 to 16 pg/mL, and the minimal detectable dose was <0.1 pg/mL. No patient had IL-7 levels below this threshold in the current study.

The standard curve ranges for IL15 were 3.9 to 250 pg/mL, and the minimal detectable dose was <2 pg/mL. IL-15 levels were between 0 and 2 pg/mL in our study in 15 patients before transplantation, in no patient on days 7 and 14, and in 1 patient on day 28. No sample dilution was performed for IL-15 assay. For IL-7 analysis, samples were diluted twice. Patient samples whose cytokine level were out of standard curve range, were re-assessed after dilution.

Immune Recovery

Immune recovery was prospectively assessed as previously described [43,44]. Briefly, patients’ peripheral white blood cells were phenotyped using 4 color flow cytometry after treatment with a red blood cell lysing solution. The following antibodies were used: CD3-ECD (Beckman Coulter, Iotest #A07748); CD4-V450 (Becton Dickinson Horizon #560345); CD8-FITC (Beckman Coulter Iotest #A07756); CD56-PC7 (Beckman Coulter Iotest #A21692); CD45RA-PE (Dako #R7086). The percentage of positive cells was calculated relative to total nucleated cells, after subtraction of non-specific staining. Absolute counts were obtained by multiplying the percentages of positive cells by the white blood cell counts (XE-5000 hematology analyzer, Sysmex, Kobe, Japan). Absolute lymphocytes counts (ALC) were measured directly by the XE-5000 analyzer or after microscopic review of the blood smears when the automated differential was flagged. Absolute white blood cell counts were used instead of ALC when white blood cell counts were below 150 cells ×10⁹/L.

Statistical Analyses

The Mann Whitney test was used to compare counts of lymphocyte subset and cytokine levels in patients given grafts after 2 Gy or 4 Gy TBI. The Wilcoxon matched pair test was used to compare cytokines levels before and at various time points after transplantation. Generalized linear mixed models were used to analyze factors affecting immune recovery and cytokine levels after transplantation. Factors included in the models included : (1) dose of TBI (2 Gy vs 4 Gy), MSC infusion or not, number of days after allo-HSCT, number of CD3⁺ cells transplanted, donor type (related vs unrelated), patient age, and donor age for analyses examining lymphocyte counts; (2) dose of TBI (2 Gy vs 4 Gy), MSC infusion or not, grade II–IV acute GVHD the first 100 days after transplantation, number of CD3⁺ cells transplanted, donor type (related vs unrelated), patient age, and donor age, and either IL-7 or IL-15 levels on days 7–14 (median) for analyses examining lymphocyte count increments from days 14–28 (median) to days 80–100 (median); and (3) number of days after allo-HSCT, number of CD3⁺ cells transplanted, donor type (related vs unrelated), dose of TBI (2 Gy vs 4 Gy), ALC, CRP levels, donor and patient ages, and MSC infusion or not, for analyses of cytokine levels. Incidences of acute GVHD according to the cytokines levels were assessed using cumulative incidence methods. A Cox model was constructed for determining potential factors associated with the occurrence of grade II–IV acute GVHD the first 200 days after transplantation. Factors included in the model included median day 7 and day 14 IL-7 levels, median day 7 and day 14 IL-15 levels, dose of TBI (2 Gy vs 4 Gy), donor type (related vs unrelated), female donor to male recipient versus other gender combination, MSC infusion or not, patient age, and donor age. Spearman’s correlation was used to examine the relationship between parameters. Statistical analyses were carried out with Graphpad Prism (Graphpad Software, San Diego, CA) and SAS version 9.2 for Windows (SAS Institute, Cary, NC, USA).

Table 1. Patients' characteristics.

	Nonmyeloablative conditioning (n = 70)
Median age (range)	50 (16–73)
Gender (male/female)	48/22
Diagnostic (# of patients)	
Acute myeloid leukemia in CR	21
Acute lymphoblastic leukemia in CR	4
Chronic myeloid leukemia	1
Chronic lymphocytic leukemia	6
Lymphoma	16
Myelodysplastic syndrome/myeloproliferative disorder	9
Multiple myeloma	13
Donor (# of patients)	
Sibling	13
Unrelated	57
Conditioning regimen (# of patients)	
TBI 2 Gy	1
Fludarabine 90 mg/m ² +TBI 2 Gy	59
Fludarabine 90 mg/m ² +TBI 4 Gy	10
Immunosuppressive regimen (# of patients)	
Tacrolimus+MMF	70
Co-transplantation with MSC	
Yes	23
No	44
Unknown*	3
Graft composition; median (range) x 10 ⁶ /kg	
CD34	5.4 (1.1–14.5)
CD3	314 (92–1216)

*double blind randomized study: The information of which of these 3 patients (if any) have been given MSC has been given by the director of the Cell Laboratory only to LS (the statistician); TBI, total body irradiation; MMF, mycophenolate mofetil.
doi:10.1371/journal.pone.0055876.t001

Results

Immune Recovery

Median ALC count on day 0 was 110 (range, 10–5440) cells/ μ L, demonstrating the persistence of recipient T cells at the time of transplantation. While median CD8⁺ T cell levels reached the lower limit of normal values from day 60 after transplantation, median CD4⁺ T cell (including naive CD4⁺ T cells) remained below the lower limit of normal values the first 6 months after transplantation (Figure 1). No significant difference of T cell subset counts were observed between 2 Gy and 4 Gy TBI regimen. Using generalized linear mixed models taking into consideration data from day 14, 28, 40, 60, 80 and 100 for each patient, counts of CD3⁺ T cells ($P < 0.001$), CD8⁺ T cells ($P < 0.001$), CD4⁺ T cells ($P = 0.024$), NK cells ($P < 0.001$) and NK/T cells ($P < 0.001$) increased over time but not those of naive CD4⁺ T cells ($P = 0.13$). Further, high numbers of transplanted CD3⁺ T cells were associated with higher counts CD3⁺ T cells ($P = 0.009$), CD8⁺ T cells ($P = 0.003$), and CD4⁺ T cells ($P = 0.0099$), while high donor age was associated with lower counts of CD3⁺ T cells ($P = 0.04$), CD4⁺ T cells ($P = 0.05$), and naive CD4⁺ T cells ($P = 0.021$). There was no significant association between MSC administration and lymphocyte subset counts after transplantation.

IL-7 Plasma Levels

Median IL-7 plasma levels remained below 6 pg/L throughout the first 100 days (the upper limit of normal range being 9.2 pg/mL (Quantikine© HS catalog number HS750)), although IL-7 plasma levels were significantly higher on days 7 (5.1 pg/mL, $P = 0.002$), 14 (5.2 pg/mL, $P < 0.0001$) and 28 (5.1 pg/mL, $P = 0.03$) (but not thereafter) than before transplantation (median value of 3.8 pg/mL) (Figure 1G). Using generalized linear mixed models, low number of transplanted CD3⁺ T cells ($P = 0.001$), low ALC level the day of IL-7 assessment ($P < 0.0001$), high donor age ($P = 0.003$), having received PBSC from unrelated donors ($p = 0.006$), and high level of CRP the day of IL-7 assessment ($P = 0.033$) were associated with high levels of IL-7 (Table 2).

IL-15 Serum Levels

Median IL-15 serum levels were significantly higher on days 7 (12.5 pg/mL, $P < 0.001$), 14 (10.5 pg/mL, $P < 0.001$) and 28 (6.2 pg/mL, $P < 0.001$) than before transplantation (median value of 2.4 pg/mL) (Figure 1H). IL-15 levels on day 7 and 14 were significantly higher in 4 Gy than 2 Gy TBI. Using generalized linear mixed models, conditioning with 4 versus 2 Gy TBI ($P = 0.002$), having received PBSC from unrelated donors ($P = 0.001$), low ALC level the day of IL-15 assessment ($P < 0.001$), and high level of CRP the day of IL-15 assessment




Figure 1. Median ALC (A), median MNC-subset cell counts (B–F), and median IL-7 (G) and IL-15 (H) after allogeneic hematopoietic cell transplantation following 2 Gy (continuous line) or 4 Gy (broken line) total body irradiation. The error bars shows the 25th and 75th percentiles. For ALC and MNC-subset, horizontal lines show the medians and the grey square the limit of normal value (if not truncated) in 47 healthy volunteer donors; for IL-7, horizontal line shows the medians and the grey square the limit of normal value according to the manufacturer brochure. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. doi:10.1371/journal.pone.0055876.g001

($P = 0.006$) were each associated with high IL-15 levels on days 7 and 14 after allo-HSCT (Table 2).

Correlation between IL-7 and IL-15 Levels and Lymphocyte Subset Counts on Days 14 or 28 after allo-HSCT

Day 14 IL-7 levels inversely correlated with day 14 counts of CD3⁺ T cells ($R = -0.46$, $P = 0.002$; Figure 2A), CD8⁺ T cells ($R = -0.41$, $P = 0.006$), CD4⁺ T cells ($R = -0.44$, $P = 0.004$), and memory CD4⁺ T cells ($R = -0.45$, $P = 0.003$), but not with counts of naïve CD4⁺ T cells ($R = -0.28$, $P = 0.07$), NK/T cells ($R = -0.04$, $P = 0.8$) nor NK cells ($R = -0.14$, $P = 0.4$). There was a weak association between day 14 IL-7 and IL-15 levels ($R = 0.27$, $P = 0.049$). Further, day 14 IL-15 levels correlated with day 14 counts of NK cells ($R = -0.32$, $P = 0.039$; Figure 2B) and of NK/T cells ($R = -0.32$, $P = 0.037$), but not with those of other T cell subsets.

Day 28 IL-7 levels inversely correlated with day 28 counts of CD3⁺ T cells ($R = -0.47$, $P < 0.001$; Figure 2A), CD8⁺ T cells ($R = -0.41$, $P = 0.002$), CD4⁺ T cells ($R = -0.39$, $P = 0.002$), naïve CD4⁺ T cells ($R = -0.40$, $P = 0.002$), and memory CD4⁺ T cells ($R = -0.38$, $P = 0.004$), but not with counts of NK/T cells ($R = -0.17$, $P = 0.2$), nor NK cells ($R = -0.02$, $P = 0.9$), nor with day-28 donor T cell chimerism levels ($R = 0.0$, $P = 0.95$). There was no significant association either between day 28 IL-7 and IL-15 levels ($R = 0.07$, $P = 0.6$). Further, day 28 IL-15 levels correlated with day 28 counts of NK cells ($R = -0.32$, $P = 0.015$; Figure 2B) but not with those of T cell subsets, nor with day-28 donor T cell chimerism levels ($R = 0.14$, $P = 0.29$).

To further assess the potential association between early IL-7 or IL-15 levels on immune recovery, we analysed whether there was a relationship between median cytokine levels on days 7 and 14 and the difference of lymphocyte subset counts between days 80–100 (median) and days 14–28 (median). Interestingly, in multivariate analyses, early IL-7 levels did not correlate with any




Figure 2. Correlation between CD3⁺ T cell counts and IL-7 levels on day 14 (black circles and continuous line) and on day 28 (open triangles and broken lines) after transplantation (A). Correlation between NK cell counts and IL-15 levels on day 14 (black circles and continuous line) and on day 28 (open triangles and broken lines) after transplantation (B). doi:10.1371/journal.pone.0055876.g002

lymphocyte subset increment from days 14–28 to day 80–100 after transplantation, while high IL-15 levels early after transplantation correlated with a lower increment of NK cells over time ($P = 0.04$).

IL-7 and IL-15 Levels did not Predict for Subsequent Acute GVHD

The 180-day cumulative incidence of grade II–IV acute GVHD was 30%, a rate similar to what has been observed by other group of investigators using similar conditioning regimen [45]. As shown in the Figure 3, no statistically significant association between cytokines levels on days 7 or 14 after transplantation and occurrence of grade II–IV acute GVHD were observed.

Specifically, the 180-day cumulative incidence of grade II–IV acute GVHD was 29% in patients with day 7 IL-7 levels > median (5.1 pg/mL) versus 20% in patients with day 7 IL-7 levels ≤ median ($P = 0.38$) (Figure 3A). Similarly, the 180-day cumulative

Table 2. Multivariable analyses of factors affecting cytokines levels on days 7 and 14 after allo-HSCT.

Factor(s) associated with higher levels**†	
IL-7	- Low ALC on day 7 or 14 ($P < 0.001$).
	- Low # of transplanted T cells (CD3 ⁺) ($P = 0.001$).
	- High CRP levels on day 7 or 14 ($P = 0.033$).
	- Unrelated donors ($P = 0.006$).
	- High donor age ($P = 0.003$).
IL-15	- 4 vs 2 Gy TBI ($P = 0.002$).
	- Unrelated donors ($P = 0.001$).
	- High CRP levels on day 7 or 14 ($P = 0.006$).
	- Low ALC on day 7 or 14 ($P < 0.001$).

*Other factors assessed were number of days after allo-HSCT, patient age, and mesenchymal stromal cells infusion or not; †P values were determined according to generalized linear mixed models; TBI, total body irradiation.

doi:10.1371/journal.pone.0055876.t002




Figure 3. Cumulative incidence of grade II–IV acute GVHD according to day 7 IL-7 plasma levels among nonmyeloablative recipients ($P = 0.4$) (A). Cumulative incidence of grade II–IV acute GVHD according to day 14 IL-7 plasma levels among nonmyeloablative recipients ($P = 0.18$) (B). Cumulative incidence of grade II–IV acute GVHD according to day 7 IL-15 serum levels among nonmyeloablative recipients ($P = 0.8$) (C). Cumulative incidence of grade II–IV acute GVHD according to day 14 IL-15 serum levels among nonmyeloablative recipients ($P = 0.6$) (D). doi:10.1371/journal.pone.0055876.g003

incidence of grade II–IV acute GVHD was 19% in patients with day 14 IL-7 levels > median (5.2 pg/mL) versus 37% in patients with day 14 IL-7 levels \leq median ($P = 0.18$) (Figure 3B).

The 180-day cumulative incidence of grade II–IV acute GVHD was 24% in patients with day 7 IL-15 levels > median (12.5 pg/mL) versus 28% in patients with day 7 IL-15 levels \leq median ($P = 0.8$) (Figure 3C). Similarly, the 180-day cumulative incidence of grade II–IV acute GVHD was 25% in patients with day 14 IL-15 levels > median (10.5 pg/mL) versus 33% in patients with day 14 IL-15 levels \leq median ($P = 0.8$) (Figure 3D).

Finally, in a multivariate Cox model, neither median IL-7 levels ($P = 0.17$ with a trend for an inverse correlation) on days 7–14 nor median IL-15 levels ($P = 0.21$ with a trend for a positive correlation) on days 7–14 correlated with occurrence of grade II–IV acute GVHD the first 200 days after transplantation. Similarly, the use of MSC was not associated with decreased incidence of grade II–IV acute GVHD. This could be explained by the fact that all 23 MSC recipients versus 9 of the remaining 49 patients (18%) received PBSC from HLA-mismatched donors. None of the other factors tested (dose of TBI, donor type, female donor to male recipient versus other gender combination, patient age, and donor age) were significantly associated with the incidence of grade II–IV acute GVHD in the current study.

IL-15 Levels did not Predict for Subsequent Relapse/Progression

Given that a previous publication showed an association between high IL-15 levels and low risk of relapse/progression [46], we compared the cumulative incidence of relapse/progression according to IL-15 levels 14 days after transplantation in our cohort of patients. The 6-month and 1-year cumulative incidences of relapse/progression were 29% and 32%, respectively, in patients with day 14 IL-15 levels > median (10.5 pg/mL) versus 37% and 46%, respectively, in patients with day 14 IL-15 levels \leq median ($P = 0.57$).

Discussion

Following allo-HSCT, eradication of residual tumor cells depends in part (in case of high-dose conditioning) or mainly (in case of nonmyeloablative conditioning) on immune-mediated graft-versus-tumor effects [1,2,4]. Prior studies have demonstrated a close relationship between T cell reconstitution and graft-versus-tumor effects after allo-HSCT [4,14,47–49]. Given that HPE allows the expansion of potentially alloreactive T cell clones, it has been generally accepted that HPE plays a major role in graft-versus-tumor effects, but could also cause or favor acute GVHD.

This prompted us to investigate the kinetics of IL-7 and IL-15 levels in a cohort of 70 patients given grafts after truly nonmyeloablative conditioning.

First, patients given grafts after nonmyeloablative conditioning had only a modest (<2 fold) increase of IL-7 levels after transplantation (contrarily to what we observed in another cohort of patients given grafts after myeloablative conditioning [50]), that persisted up to day 21. This is probably due to the fact that nonmyeloablative patients experienced relatively mild lymphopenia (and thus continue to consume the IL-7 produced by stromal cells) as demonstrated by the persistence of median ALC counts of 110 cells/ μ L at the time of transplantation. Although the first T cell chimerism assessment in current patient was usually around day 28 after HSCT, a prior study analyzing data from patients given similar conditioning regimen demonstrated that a median of 50 CD3⁺ T cells of recipient origin/ μ L persisted on day 14 after HSCT [40]. Further, as observed by other groups of investigators [46,51,52], there was a strong inverse correlation between IL-7 levels and absolute lymphocyte counts [46,52], as well as a strong inverse correlation between IL-7 levels and T cell subsets on days 14 and 28 after transplantation. Other factors associated with IL-7 levels included high CRP levels, and low numbers of transplanted T cells. Levels of IL-7 in current nonmyeloablative recipients were lower to what was observed by Thiant *et al.* in a cohort of 45 patients given grafts after fludarabine +2 Gy TBI (n = 18) or more intense but still reduced-intensity conditioning (n = 27) [52], and where much lower than what was observed by Dean *et al.* in patients given grafts after sequential chemotherapy followed by a chemotherapy/fludarabine-based reduced-intensity conditioning [53]. This apparent discrepancy is probably explained the fact that median ALC counts on day 0 were 110 (range, 10–5440) cells/ μ L in current patient versus 0 (range, 0–322) cells/ μ L in the Dean *et al.* study, while median counts of CD3⁺ T cells were 0 (range, 0–1900) cells/ μ L at the time of transplantation in Thiant *et al.* study.

IL-15 levels were lower in nonmyeloablative patients conditioned with 2 Gy TBI than in those conditioned with 4 Gy TBI, demonstrating that the release of IL-15 was proportional to the intensity of the conditioning regimen. As observed by Thiant *et al.* [46,52], there was a correlation between IL-7 and IL-15 levels on day 14 (but not on day 28) after transplantation, and an inverse correlation between IL-15 levels and NK cell counts. Other factors affecting IL-15 levels included high CRP levels.

Several observations demonstrate that immune recovery depended mainly on HPE the first year after nonmyeloablative conditioning regimen in current patients. Firstly, there was a strong correlation between the number of infused T cells and high counts of CD4⁺ and CD8⁺ T cells, as previously observed [43,54]. Secondly, thymic function was minimal during the first 100 days after allo-HSCT given that levels of naïve CD4⁺ T cells did not

significantly increase the first 100 days after transplantation despite that some naïve T cells can undergo HPE and keep their naïve phenotype. Third, there was a correlation between high donor age and low counts of CD3⁺ T cells (P = 0.04), CD4⁺ T cells (P = 0.05), and naïve CD4⁺ T cells (P = 0.021), as previously observed in patients given grafts after nonmyeloablative conditioning [55]. Despite that, we failed to find any significant association between IL-7 and/or IL-15 levels early after transplantation and increment of T cell subset counts from days 14–28 to day 80–100, even after adjusting for potentially confounding cofactors.

A number of previous studies have demonstrated that high levels of IL-7 [46,52,53] and/or IL-15 [46,52] early after transplantation correlated with subsequent occurrence of grade II–IV acute GVHD, while others study failed to find such an association [51,56]. The largest study including data from 153 consecutive allogeneic transplant recipients given grafts after high-dose conditioning and ATG observed no correlation between IL-7 levels early after transplantation and acute GVHD, while, interestingly, there was an inverse correlation between IL-15 levels early after transplantation and grade II–IV acute GVHD [57]. Further, a recent study demonstrated that administration of IL-7 after allogeneic T cell-depleted transplantation in humans did not increase acute GVHD [58]. In the current study, we did not observe any association between levels of IL-7 or IL-15 early after allo-HSCT and grade II–IV acute GVHD. The same was true after adjusting the analyses for potentially confounding cofactors. Differences in postgrafting immunosuppression might be the cause for these apparent discrepancies between studies. As example, it has been shown that tacrolimus (given in patients included in the current study) decreased T cell proliferation induced by IL-7 [59], and tacrolimus levels were kept high in our patients the first weeks after transplantation (median 18.6, 16.4, 14.9 and 14.3 μ g/L on days 0, 7, 14 and 21 after transplantation, respectively) probably explaining the low relatively incidence of acute GVHD observed [60].

In summary, these data suggest that IL-7 and IL-15 levels remain relatively low after nonmyeloablative transplantation, and that IL-7 and IL-15 levels early after nonmyeloablative transplantation do not predict for acute GVHD.

Acknowledgments

The authors are grateful to O. Dengis for excellent technical support.

Author Contributions

Taking care of the patients: YB FB. Conceived and designed the experiments: MF YB FB. Performed the experiments: MdB MF MH MPM AG. Analyzed the data: MdB MF LS FB. Contributed reagents/materials/analysis tools: MdB MF YB FB. Wrote the paper: MdB FB.

References

- Weiden PL, Flourmoy N, Thomas ED, Prentice R, Fefer A, et al. (1979) Antileukemic effect of graft-versus-host disease in human recipients of allogeneic-marrow grafts. *N Engl J Med* 300: 1068–1073.
- Miller JS, Warren EH, van den Brink MR, Ritz J, Shlomchik WD, et al. (2010) NCI First International Workshop on The Biology, Prevention, and Treatment of Relapse After Allogeneic Hematopoietic Stem Cell Transplantation: Report from the Committee on the Biology Underlying Recurrence of Malignant Disease following Allogeneic HSCT: Graft-versus-Tumor/Leukemia Reaction. *Biol Blood Marrow Transplant* 16: 565–586.
- Sandmaier BM, Mackinnon S, Childs RW (2007) Reduced intensity conditioning for allogeneic hematopoietic cell transplantation: current perspectives. *Biol Blood Marrow Transplant* 13: 87–97.
- Baron F, Maris MB, Sandmaier BM, Storer BE, Sorror M, et al. (2005) Graft-versus-tumor effects after allogeneic hematopoietic cell transplantation with nonmyeloablative conditioning. *J Clin Oncol* 23: 1993–2003.
- Baron F, Petersdorf EW, Gooley T, Sandmaier BM, Malkki M, et al. (2009) What is the role for donor natural killer cells after nonmyeloablative conditioning? *Biol Blood Marrow Transplant* 15: 580–588.
- Baron F, Labopin M, Niederwieser D, Vigouroux S, Cornelissen JJ, et al. (2012) Impact of graft-versus-host disease after reduced-intensity conditioning allogeneic stem cell transplantation for acute myeloid leukemia: a report from the Acute Leukemia Working Party of the European group for blood and marrow transplantation. *Leukemia*. 26: 2462–2468.
- Crooks GM, Weinberg K, Mackall C (2006) Immune reconstitution: from stem cells to lymphocytes. *Biol Blood Marrow Transplant* 12: 42–46.
- Peggs KS (2006) Reconstitution of adaptive and innate immunity following allogeneic hematopoietic stem cell transplantation in humans. *Cytotherapy* 8: 427–436.

9. Gress RE, Emerson SG, Drobyski WR (2010) Immune reconstitution: how it should work, what's broken, and why it matters. *Biol Blood Marrow Transplant* 16: S133–S137.
10. Clave E, Busson M, Douay C, Peffault de Latour R, Berrou J, et al. (2009) Acute graft-versus-host disease transiently impairs thymic output in young patients after allogeneic hematopoietic stem cell transplantation. *Blood* 113: 6477–6484.
11. Bahecci E, Epperson D, Douek DC, Melenhorst JJ, Childs RC, et al. (2003) Early reconstitution of the T-cell repertoire after non-myeloablative peripheral blood stem cell transplantation is from post-thymic T-cell expansion and is unaffected by graft-versus-host disease or mixed chimerism. *Br J Haematol* 122: 934–943.
12. Fallen PR, McGreavey L, Madrigal JA, Potter M, Ethell M, et al. (2003) Factors affecting reconstitution of the T cell compartment in allogeneic hematopoietic cell transplant recipients. *Bone Marrow Transplant* 32: 1001–1014.
13. Toubert A, Glauzy S, Douay C, Clave E (2012) Thymus and immune reconstitution after allogeneic hematopoietic stem cell transplantation in humans: never say never again. *Tissue Antigens* 79: 83–89. doi:10.1111/j.1399-0039.2011.01820.x.
14. Bosch M, Khan FM, Storek J (2012) Immune reconstitution after hematopoietic cell transplantation. *Curr Opin Hematol* 19: 324–335. doi:10.1097/MOH.0b013e328353bc7d.
15. Federmann B, Hagele M, Pfeiffer M, Wirths S, Schumm M, et al. (2011) Immune reconstitution after haploidentical hematopoietic cell transplantation: impact of reduced intensity conditioning and CD3/CD19 depleted grafts. *Leukemia* 25: 121–129. doi:10.1038/leu.2010.235.
16. Storek J, Dawson MA, Maloney DG (2003) Correlation between the numbers of naive T cells infused with blood stem cell allografts and the counts of naive T cells after transplantation. *Biol Blood Marrow Transplant* 9: 781–784.
17. Douek DC, Vescio RA, Betts MR, Brenchley JM, Hill BJ, et al. (2000) Assessment of thymic output in adults after haematopoietic stem-cell transplantation and prediction of T-cell reconstitution. *Lancet* 355: 1875–1881.
18. Weinberg K, Blazar BR, Wagner JE, Agura E, Hill BJ, et al. (2001) Factors affecting thymic function after allogeneic hematopoietic stem cell transplantation. *Blood* 97: 1458–1466.
19. Hochberg EP, Chillemi AC, Wu CJ, Neuberger D, Canning C, et al. (2001) Quantitation of T-cell neogenesis in vivo after allogeneic bone marrow transplantation in adults. *Blood* 98: 1116–1121.
20. Lewin SR, Heller G, Zhang L, Rodrigues E, Skulsky E, et al. (2002) Direct evidence for new T-cell generation by patients after either T-cell-depleted or unmodified allogeneic hematopoietic stem cell transplantations. *Blood* 100: 2235–2242.
21. Hazenberg MD, Otto SA, de Pauw ES, Roelofs H, Fibbe WE, et al. (2002) T-cell receptor excision circle and T-cell dynamics after allogeneic stem cell transplantation are related to clinical events. *Blood* 99: 3449–3453.
22. Krenger W, Blazar BR, Hollander GA (2011) Thymic T-cell development in allogeneic stem cell transplantation. *Blood* 117: 6768–6776.
23. Mackall CL, Fry TJ, Gress RE (2011) Harnessing the biology of IL-7 for therapeutic application. *Nat Rev Immunol* 11: 330–342.
24. Puel A, Ziegler SF, Buckley RH, Leonard WJ (1998) Defective IL7R expression in T(-)B(+)NK(+) severe combined immunodeficiency. *Nat Genet* 20: 394–397.
25. Mackall CL, Fry TJ, Bare C, Morgan P, Galbraith A, et al. (2001) IL-7 increases both thymic-dependent and thymic-independent T-cell regeneration after bone marrow transplantation. *Blood* 97: 1491–1497.
26. Alpdogan O, Schmaltz C, Muriglan SJ, Kappel BJ, Perales MA, et al. (2001) Administration of interleukin-7 after allogeneic bone marrow transplantation improves immune reconstitution without aggravating graft-versus-host disease. *Blood* 98: 2256–2265.
27. Fry TJ, Moniuszko M, Creekmore S, Donohue SJ, Douek DC, et al. (2003) IL-7 therapy dramatically alters peripheral T-cell homeostasis in normal and SIV-infected nonhuman primates. *Blood* 101: 2294–2299.
28. Chu YW, Memon SA, Sharrow SO, Hakim FT, Eckhaus M, et al. (2004) Exogenous IL-7 increases recent thymic emigrants in peripheral lymphoid tissue without enhanced thymic function. *Blood* 104: 1110–1119.
29. Storek J, Gillespy TI, Lu H, Joseph A, Dawson MA, et al. (2003) Interleukin-7 improves CD4 T-cell reconstitution after autologous CD34 cell transplantation in monkeys. *Blood* 101: 4209–4218.
30. Sportes C, Hakim FT, Memon SA, Zhang H, Chua KS, et al. (2008) Administration of rhIL-7 in humans increases in vivo TCR repertoire diversity by preferential expansion of naive T cell subsets. *J Exp Med* 205: 1701–1714.
31. Beq S, Nugeyre MT, Ho Tsong FR, Gautier D, Legrand R, et al. (2006) IL-7 induces immunological improvement in SIV-infected rhesus macaques under antiviral therapy. *J Immunol* 176: 914–922.
32. Fehniger TA, Caligiuri MA (2001) Interleukin 15: biology and relevance to human disease (Review). *Blood* 97: 14–32.
33. Lodolce JP, Boone DL, Chai S, Swain RE, Dassopoulos T, et al. (1998) IL-15 receptor maintains lymphoid homeostasis by supporting lymphocyte homing and proliferation. *Immunity* 9: 669–676.
34. Tan JT, Ernst B, Kieper WC, LeRoy E, Sprent J, et al. (2002) Interleukin (IL)-15 and IL-7 jointly regulate homeostatic proliferation of memory phenotype CD8+ cells but are not required for memory phenotype CD4+ cells. *J Exp Med* 195: 1523–1532.
35. Goldrath AW, Sivakumar PV, Glaccum M, Kennedy MK, Bevan MJ, et al. (2002) Cytokine requirements for acute and Basal homeostatic proliferation of naive and memory CD8+ T cells. *J Exp Med* 195: 1515–1522.
36. Berard M, Brandt K, Bulfone-Paus S, Tough DF (2003) IL-15 promotes the survival of naive and memory phenotype CD8+ T cells. *J Immunol* 170: 5018–5026.
37. Alves NL, Hooibrink B, Arosa FA, van Lier RA (2003) IL-15 induces antigen-independent expansion and differentiation of human naive CD8+ T cells in vitro. *Blood* 102: 2541–2546.
38. Alpdogan O, Eng JM, Muriglan SJ, Willis LM, Hubbard VM, et al. (2005) Interleukin-15 enhances immune reconstitution after allogeneic bone marrow transplantation. *Blood* 105: 865–873.
39. Huntington ND, Alves NL, Legrand N, Lim A, Strick-Marchand H, et al. (2011) IL-15 transpresentation promotes both human T-cell reconstitution and T-cell-dependent antibody responses in vivo. *Proc Natl Acad Sci U S A* 108: 6217–6222.
40. Baron F, Baker JE, Storb R, Gooley TA, Sandmaier BM, et al. (2004) Kinetics of engraftment in patients with hematologic malignancies given allogeneic hematopoietic cell transplantation after nonmyeloablative conditioning. *Blood* 104: 2254–2262.
41. Baron F, Sandmaier BM (2006) Chimerism and outcomes after allogeneic hematopoietic cell transplantation following nonmyeloablative conditioning. *Leukemia* 20: 1690–1700.
42. Baron F, Lechanteur C, Willems E, Bruck F, Baudoux E, et al. (2010) Cotransplantation of mesenchymal stem cells might prevent death from graft-versus-host disease (GVHD) without abrogating graft-versus-tumor effects after HLA-mismatched allogeneic transplantation following nonmyeloablative conditioning. *Biol Blood Marrow Transplant* 16: 838–847.
43. Castermans E, Baron F, Willems E, Schaaaf-Lafontaine N, Meuris N, et al. (2008) Evidence for neo-generation of T cells by the thymus after non-myeloablative conditioning. *Haematologica* 93: 240–247.
44. Castermans E, Hannon M, Dutrieux J, Humblet-Baron S, Seidel L, et al. (2011) Thymic recovery after allogeneic hematopoietic cell transplantation with non-myeloablative conditioning is limited to patients younger than 60 years of age. *Haematologica* 96: 298–306.
45. Lange T, Hubmann M, Burkhardt R, Franke GN, Cross M, et al. (2011) Monitoring of WT1 expression in PB and CD34(+) donor chimerism of BM predicts early relapse in AML and MDS patients after hematopoietic cell transplantation with reduced-intensity conditioning. *Leukemia* 25: 498–505. doi:10.1038/leu.2010.283.
46. Thiant S, Yakoub-Agha I, Magro L, Trauet J, Coiteux V, et al. (2010) Plasma levels of IL-7 and IL-15 in the first month after myeloablative BMT are predictive biomarkers of both acute GVHD and relapse. *Bone Marrow Transplant* 45: 1546–1552. doi:10.1038/bmt.2010.13.
47. Maraninchi D, Gluckman E, Blaise D, Guyotat D, Rio B, et al. (1987) Impact of T-cell depletion on outcome of allogeneic bone-marrow transplantation for standard-risk leukaemias. *Lancet* 2: 175–178.
48. Soiffer RJ, Lerademacher J, Ho V, Kan F, Artz A, et al. (2011) Impact of immune modulation with anti-T-cell antibodies on the outcome of reduced-intensity allogeneic hematopoietic stem cell transplantation for hematologic malignancies. *Blood* 117: 6963–6970. doi:10.1182/blood-2011-01-332007.
49. Mohty M, Avinens O, Faucher C, Viens P, Blaise D, et al. (2007) Predictive factors and impact of full donor T-cell chimerism after reduced intensity conditioning allogeneic stem cell transplantation. *Haematologica* 92: 1004–1006.
50. De Bock M, Fillet M, Merville MP, Gothot A, Beguin Y, et al. (2012) Kinetics of IL-7 and IL-15 Levels After Allogeneic Peripheral Blood Stem Cell Transplantation (allo-PBSC-T) Following High-Dose or Nonmyeloablative Conditioning. *Biol Blood Marrow Transplant* 18 suppl 2: s275.
51. Bolotin E, Annet G, Parkman R, Weinberg K (1999) Serum levels of IL-7 in bone marrow transplant recipients: relationship to clinical characteristics and lymphocyte count. *Bone Marrow Transplant* 23: 783–788.
52. Thiant S, Labalette M, Trauet J, Coiteux V, de Berranger E, et al. (2011) Plasma levels of IL-7 and IL-15 after reduced intensity conditioned allo-SCT and relationship to acute GVHD. *Bone Marrow Transplant* 46: 1374–1381. doi:10.1038/bmt.2010.300.
53. Dean RM, Fry T, Mackall C, Steinberg SM, Hakim F, et al. (2008) Association of serum interleukin-7 levels with the development of acute graft-versus-host disease. *J Clin Oncol* 26: 5735–5741. doi:10.1200/JCO.2008.17.1314.
54. Baron F, Schaaaf-Lafontaine N, Humblet-Baron S, Meuris N, Castermans E, et al. (2003) T-cell reconstitution after unmanipulated, CD8-depleted or CD34-selected nonmyeloablative peripheral blood stem-cell transplantation. *Transplantation* 76: 1705–1713.
55. Baron F, Storer B, Maris MB, Storek J, Piette F, et al. (2006) Unrelated donor status and high donor age independently affect immunologic recovery after nonmyeloablative conditioning. *Biol Blood Marrow Transplant* 12: 1176–1187.
56. Abu-Ghosh A, Goldman S, Slone V, van de Ven C, Suen Y, et al. (1999) Immunological reconstitution and correlation of circulating serum inflammatory mediators/cytokines with the incidence of acute graft-versus-host disease during the first 100 days following unrelated umbilical cord blood transplantation. *Bone Marrow Transplant* 24: 535–544. doi:10.1038/sj.bmt.1701921.
57. Hagel LM, Liu Y, Ugarte-Torres A, Williamson TS, Russell JA, et al. (2010) High IL-15 serum levels on day 7 after hematopoietic cell transplantation are associated with a low likelihood of graft-versus-host disease and a high likelihood of infections. *ASH Annual Meeting Abstracts* 116: 1271.
58. Perales MA, Goldberg JD, Yuan J, Koehne G, Lechner L, et al. (2012) Recombinant human interleukin-7 (CYT107) promotes T cell recovery

- following allogeneic stem cell transplantation. *Blood*. doi:10.1182/blood-2012-06-437236.
59. Almawi WY, Assi JW, Chudzik DM, Jaoude MM, Rieder MJ (2001) Inhibition of cytokine production and cytokine-stimulated T-cell activation by FK506 (tacrolimus). *Cell Transplant* 10: 615–623.
60. Ram R, Storer B, Mielcarek M, Sandmaier BM, Maloney DG, et al. (2012) Association between calcineurin inhibitor blood concentrations and outcomes after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 18: 414–422. doi:10.1016/j.bbmt.2011.08.016.