

Draft Genome Sequence of *Streptococcus gordonii* Type Strain CCUG 33482^T

Francisco Salvà-Serra,^{a,b,c} Hedvig E. Jakobsson,^b Kaisa Thorell,^d Lucia Gonzales-Siles,^b Erika T. Hallbäck,^b Daniel Jaén-Luchoro,^c Fredrik Boulund,^e Per Sikora,^{f,g} Roger Karlsson,^{b,h} Liselott Svensson,^{a,b} Antoni Bennasar,^c Lars Engstrand,^d Erik Kristiansson,^e Edward R. B. Moore^{a,b}

Culture Collection University of Gothenburg (CCUG), Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden^a; Department of Infectious Diseases, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden^b; Microbiology, Department of Biology, University of the Balearic Islands, Palma de Mallorca, Spain^c; Department of Microbiology, Tumor and Cell Biology, Karolinska Institute, Stockholm, Sweden^d; Department of Mathematical Sciences, Chalmers University of Technology, Gothenburg, Sweden^e; Integrative Clinical Genomics, Clinical Diagnostics, SciLifeLab, Stockholm, Sweden^f; Laboratory Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden^g; Nanoxis Consulting AB, Gothenburg, Sweden^h

F.S.-S. and H.E.J. contributed equally to this work.

***Streptococcus gordonii* type strain CCUG 33482^T is a member of the *Streptococcus mitis* group, isolated from a case of subacute bacterial endocarditis. Here, we report the draft genome sequence of *S. gordonii* CCUG 33482^T, composed of 41 contigs of a total size of 2.15 Mb with 2,061 annotated coding sequences.**

Received 5 February 2016 Accepted 9 February 2016 Published 24 March 2016

Citation Salvà-Serra F, Jakobsson HE, Thorell K, Gonzales-Siles L, Hallbäck ET, Jaén-Luchoro D, Boulund F, Sikora P, Karlsson R, Svensson L, Bennasar A, Engstrand L, Kristiansson E, Moore ERB. 2016. Draft genome sequence of *Streptococcus gordonii* type strain CCUG 33482^T. Genome Announc 4(2):e00175-16. doi:10.1128/genomeA.00175-16.

Copyright © 2016 Salvà-Serra et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Edward R. B. Moore, erbmoore@ccug.se.

Streptococcus gordonii is a commensal bacterium commonly found in human oral cavities and pharynges (1) and has been implicated in the formation of dental biofilms (2). *S. gordonii* also has been isolated from samples of bacterial endocarditis (3). *S. gordonii* was described as a novel species in 1989 after analyses of viridans streptococci (4). In those studies, *Streptococcus sanguis* type I-II, strain SK3^T (=CCUG 33482^T ←NCTC 7865^T ←ATCC 10558^T), isolated from a patient with subacute bacterial endocarditis (5), was reclassified as a distinct species, *S. gordonii*, with strain SK3^T designated the type strain. *S. gordonii* is one of 14 species within the *Streptococcus mitis* group (6).

S. gordonii CCUG 33482^T was cultivated with 5% CO₂, at 37°C, on Columbia agar base plus 5% horse blood. A fresh, pure-culture biomass was lysed in low-TE buffer (1 mM Tris, 0.1 mM EDTA, pH 8.0) supplemented with Triton 1.2% (wt/vol), lysozyme (20 mg/ml), and proteinase K (1 mg/ml), at 56°C overnight. Genomic DNA was isolated, using a MagNA pure compact nucleic acid isolation kit version I (Roche Diagnostics, Mannheim, Germany). DNA was sequenced with an Illumina MiSeq instrument (SciLifeLab, Stockholm, Sweden), generating 633,858 paired-end reads of 300 bp, yielding a total of 190.2 Mb. Sequence reads were trimmed and assembled *de novo* with CLC Genomic Workbench version 8 (CLC bio, Aarhus, Denmark) and default settings. Assembly quality was assessed using Quast version 3.1 (7). Trimming of the sequence reads left 632,358 paired-end reads (80.2% of initial sequence data) with an average of 241 bp, producing an assembly length of 2,166,763 bp, consisting of 41 contigs; the longest contig measures 632,335 bp. The *N*₅₀ is 233,538 bp, and the average genome coverage of the assembly is 71×. The G+C content of the genome sequence is 40.5%, which is consistent with the 41% reported when the species was described (4).

The genome sequence was annotated using the NCBI Prokary-

otic Genome Annotation Pipeline (PGAP) (8). Annotation identified 2,061 coding sequences, 46 tRNAs, and 3 repeat regions. Comparative analyses of the sequence, by average nucleotide identity based on BLAST (ANiB) (9), using JSpecies version 1.2.1 (10), with the genome sequences of the type strains of species of the *S. mitis* group ranged from 73.1% to 81.8%. The analysis of the genome sequence for the presence of CRISPR-Cas systems (11) with CRISPRFinder (12) confirmed the presence of 4 CRISPR regions. Analyses against the Comprehensive Antibiotic Resistance Database (CARD) (13) and against the virulence factor database (VFDB) (14) identified 8 putative antibiotic resistance genes and 72 putative virulence factors. At the beginning of this project, the genome sequence of the type strains of all species included within the *S. mitis* group were available, except for that of *S. gordonii*. The genome sequence of *S. gordonii* CCUG 33482^T represents essential data for the characterization of metabolic features of *Streptococcus* and phylogenomic studies of the *S. mitis* group.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [LQWV000000000](https://www.ncbi.nlm.nih.gov/nuclink/LQWV000000000). The version described in this paper is the first version, [LQWV010000000](https://www.ncbi.nlm.nih.gov/nuclink/LQWV010000000).

ACKNOWLEDGMENTS

The CCUG is supported by the Sahlgrenska University Hospital. This research was supported by the European Commission: Tailored-Treatment, project no. 602860, and by the Swedish Västra Götaland regional funding, project no. ALFGBG-437221. F.S.-S. was supported by a stipend for Basic and Advanced Research from the CCUG through the Institute of Biomedicine, Sahlgrenska Academy, University of Gothenburg. A.B. acknowledges Spanish MINECO support (project no. CGL2012-39604 and Consolider project no. CSD2009-00006).

FUNDING INFORMATION

This work, including the efforts of Francisco Salvà-Serra, Hedvig E. Jakobsson, Lucia Gonzales-Siles, Fredrik Boulund, Roger Karlsson, Erik Kristiansson, and Edward R. B. Moore, was funded by European Commission (602860). This work, including the efforts of Francisco Salvà-Serra, Hedvig E. Jakobsson, Lucia Gonzales-Siles, Fredrik Boulund, Roger Karlsson, Liselott Svensson, Erik Kristiansson, and Edward R. B. Moore, was funded by Swedish Västra Götaland (ALFGBG-437221). This work, including the efforts of Francisco Salvà-Serra, Daniel Jaén-Luchoro, and Antoni Bennasar, was funded by Spanish MINECO (CSD2009-00006).

REFERENCES

- Jenkinson HF, Lamont RJ. 1997. Streptococcal adhesion and colonization. *Crit Rev Oral Biol Med* 8:175–200. <http://dx.doi.org/10.1177/10454411970080020601>.
- Zhou P, Liu J, Li X, Takahashi Y, Qi F. 2015. The sialic acid binding protein, Hsa, in *Streptococcus gordonii* DL1 also mediates intergeneric co-aggregation with *Veillonella* species. *PLoS One* 10:e0143898. <http://dx.doi.org/10.1371/journal.pone.0143898>.
- Douglas CW, Heath J, Hampton KK, Preston FE. 1993. Identity of viridans streptococci isolated from cases of infective endocarditis. *J Med Microbiol* 39:179–182. <http://dx.doi.org/10.1099/00222615-39-3-179>.
- Kilian M, Mikkelsen L, Henrichsen J. 1989. Taxonomic study of viridans streptococci: description of *Streptococcus gordonii* sp. nov. and emended descriptions of *Streptococcus sanguis* (White and Niven 1946), *Streptococcus oralis* (Bridge and Sneath 1982), and *Streptococcus mitis* (Andrewes and Horder 1906). *Int J Syst Bacteriol* 39:471–484. <http://dx.doi.org/10.1099/00207713-39-4-471>.
- White JC, Niven CF. 1946. *Streptococcus* s.b.e.: a *Streptococcus* associated with subacute bacterial endocarditis. *J Bacteriol* 51:717–722.
- Kawamura Y, Hou XG, Sultana F, Miura H, Ezaki T. 1995. Determination of 16S rRNA sequences of *Streptococcus mitis* and *Streptococcus gordonii* and phylogenetic relationships among members of the genus *Streptococcus*. *Int J Syst Bacteriol* 45:406–408. <http://dx.doi.org/10.1099/00207713-45-2-406>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <http://dx.doi.org/10.1093/bioinformatics/btt086>.
- Tatusova T, Ciufu S, Federhen S, Fedorov B, McVeigh R, O'Neill K, Tolstoy I, Zaslavsky L. 2015. Update on RefSeq microbial genomes resources. *Nucleic Acids Res* 43:D599–D605. <http://dx.doi.org/10.1093/nar/gku1062>.
- Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM. 2007. DNA-DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57: 81–91. <http://dx.doi.org/10.1099/ijs.0.64483-0>.
- Richter M, Rosselló-Móra R. 2009. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* 106: 19126–19131. <http://dx.doi.org/10.1073/pnas.0906412106>.
- Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA, Horvath P. 2007. CRISPR provides acquired resistance against viruses in prokaryotes. *Science* 315:1709–1712. <http://dx.doi.org/10.1126/science.1138140>.
- Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <http://dx.doi.org/10.1093/nar/gkm360>.
- McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJV, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Tennyson Yu WGD. 2013. The Comprehensive Antibiotic Resistance Database. *Antimicrob Agents Chemother* 57: 3348–3357. <http://dx.doi.org/10.1128/AAC.00419-13>.
- Chen L, Zheng D, Liu B, Yang J, Jin Q. 2016. VFDB 2016: hierarchical and refined dataset for big data analysis—10 years on. *Nucleic Acids Res* 44:D694–D697. <http://dx.doi.org/10.1093/nar/gkv1239>.