



Original article

MepmiRDB: a medicinal plant microRNA database

Dongliang Yu^{1,†}, Jiangjie Lu^{1,2,†}, Weishan Shao^{1,2}, Xiaoxia Ma¹,
Tian Xie^{3,4,*}, Hidetaka Ito⁵, Tingzhang Wang⁶, Min Xu^{1,2},
Huizhong Wang^{1,2,*} and Yijun Meng^{1,2,*}

¹College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou, 310036, China, ²Zhejiang Provincial Key Laboratory for Genetic Improvement and Quality Control of Medicinal Plants, Hangzhou Normal University, Hangzhou, 310036, China, ³Department of Pharmacology, Holistic Integrative Pharmacy Institutes, College of Medicine, Hangzhou Normal University, Hangzhou, 311121, China, ⁴Key Laboratory of Elemene Class Anti-cancer Chinese Medicine of Zhejiang Province and Engineering Laboratory of Development and Application of Traditional Chinese Medicine from Zhejiang Province, Hangzhou Normal University, Hangzhou, 311121, China, ⁵Faculty of Science, Hokkaido University, Sapporo, 060-0810, Japan and ⁶Key Laboratory of Microbiological Technology and Bioinformatics Research in Zhejiang Province, Hangzhou, 310012, China

*Corresponding author: Tel.: +86-571-28860237; E-mail: xbs@hznu.edu.cn

Correspondence may also be addressed to Huizhong Wang. Tel.: +86-571-28865198; E-mail: whz62@163.com

and Yijun Meng. Tel.: +86-571-28865198; E-mail: mengyijun@zju.edu.cn

†These authors contributed equally to this work.

Citation details: Yu, D., Lu, J., Shao, W. *et al.* MepmiRDB: a medicinal plant microRNA database. *Database* (2019) Vol. 2019: article ID baz070; doi:10.1093/database/baz070

Received 14 December 2018; Revised 27 April 2019; Accepted 7 May 2019

Abstract

MicroRNAs (miRNAs) have been recognized as a key regulator in plant development and metabolism. Recent reports showed that the miRNAs of medicinal plants not only act as a critical modulator in secondary metabolism but also had a great potential of performing cross-kingdom gene regulation. Although several plant miRNA repositories have been publicly available, no miRNA database specific for medicinal plants has been reported to date. Here, we report the first version of MepmiRDB (medicinal plant microRNA database), which is freely accessible at <http://mepmirdb.cn/mepmirdb/index.html>. This database accommodates thousands of miRNA candidates belonging to 29 medicinal plant species. The miRNA information on sequences, expression patterns and regulatory networks has been included in the functional modules of the database. Specifically, the 'Sequence' module provides the sequences of the mature miRNAs and their precursors, and the structure information of the precursors. Moreover, the processing and small RNA accumulation signals on the miRNA precursors are also included in the 'Sequence' module. The organ/growth condition-specific expression information of the mature miRNAs has been stored in the 'Expression' module. The 'Interaction' module offers the information of the degradome-validated miRNA–target pairs of eight plant species. The 'Search' module enables users to search for the miRNAs

by plant species and miRNA families, or by sequences. All data in this database are available for download. Taken together, the functional modules of MepmiRDB ensure its importance and timeliness for mechanistic and functional studies on the medicinal plant miRNAs.

Database URL: <http://mepmirdb.cn/mepmirdb/index.html>

Introduction

MicroRNAs (miRNAs) play critical roles in diverse biological processes of plants, such as growth and development, stress response and metabolism (1). Currently, there are several databases accommodating information of plant miRNAs, such as miRBase (2), PMRD (3), miRNEST (4), miREX (5), PmiRKB (6) and PNRD (7). However, to date, no database has been established specifically for the miRNAs of medicinal plants. Here, we report the first version of MepmiRDB (medicinal plant microRNA database) including 29 medicinal plant species, most of which are listed in the pharmacopoeia of the People's Republic of China (8).

Increasing pieces of evidence point to the pivotal role of plant miRNAs in modulating the biosynthesis and accumulation of secondary metabolites (9, 10). It is particularly important for the medicinal plants since a dominant portion of the bioactive ingredients are produced through the secondary metabolic pathways (11–14). On the other hand, miRNAs have become a charming candidate for therapeutic management of cancer and other intractable diseases (15). Notably, several reports indicated the ability of plant miRNAs in cross-kingdom communication (16–19), and some studies showed the high stability of certain ingested miRNAs during circulation in human or animal bloodstream (20–22). These lines of evidence point to the possibility that the exogenous, plant-derived miRNAs could perform target regulation in human body. In this regard, the miRNAs emerged as the bioactive components of medicinal plants (23–25). Taken together, the above-mentioned points formed the driving force for design and construction of MepmiRDB, which has been freely available online (<http://mepmirdb.cn/mepmirdb/index.html>).

Database construction and contents

The database structure has been illustrated in Figure S1 and S2. Five major functional modules, including 'Sequence', 'Expression', 'Interaction', 'Search' and 'Download', are available in MepmiRDB (Figure 1A) and are described below.

The 'Sequence' module

Sequence information of thousands of the miRNAs belonging to 29 plant species was included in this

module. All of the miRNAs were predicted by using the recently published workflow PmiRDiscVali (26). These miRNAs were generally divided into conserved and non-conserved ones. By comparing to all of the plant miRNAs registered in miRBase (release 21), the sRNAs with identical sequences were extracted from the sRNA-seq (small RNA sequencing) data sets, and were regarded as the conserved miRNAs existed in specific medicinal plants (Figure 2A for example). Besides, 86 RNA-seq datasets of 23 plant species were employed for transcriptome assembly by Trinity (trinityrnaseq_r20140717, default parameters) (27). The assembled transcripts were served as the basic data for miRNA precursor discovery by using miRDeep-P (v1.3, default parameters) (28). We noticed that a large portion of the miRDeep-P predictions belonged to the non-conserved miRNAs (Table 1), suggesting that these miRNA candidates might be evolutionarily young or species-specific.

An overview of the sequence characteristics shows that around 50% of the miRNA candidates start with 5' U and are 21 nt in length (Figure 1B and C), which is consistent with the known sequence features of plant miRNAs (1).

Degradome-seq data is useful for tracking the processing signals of miRNA precursors (29, 30). For eight plant species with degradome-seq data, the 'Sequence' module provides the detectable processing signals on the secondary structures of the pri-miRNAs (Figure 2C and D). Additionally, sRNA expression levels on the miRNA precursors were achieved by mapping the sRNA-seq data onto the pri-miRNAs. In many cases, the highly enriched accumulation regions of the sRNAs correlated well with the mature miRNA-coding regions, indicating the high reliability of the pri-miRNA candidates (Figure 2C and D).

The 'Expression' module

The module provides the organ/growth condition-specific expression information of the mature miRNA candidates (Figure 2B).

The 'Interaction' module

For the eight species with degradome-seq data, miRNA target prediction was performed by using the psRNATarget algorithm (2017 Update, default parameters) (31), treating the assembled transcripts and the mature miRNA candidates as the input data. Then, the predicted miRNA-target

Table 1. The number of microRNAs reported by the current version of MepmiRDB.

Organism	miR_total	miR_miRBase	miR_miRDP	miR_(miRBase \cap miRDP)	Pri-miR_miRDP	Pre-miR_miRDP
<i>Actinidia chinensis</i> (ach)	141	141	0	0	0	0
<i>Ammopiptanthus mongolicus</i> (amo)	456	332	151	27	148	76
<i>Aquilaria sinensis</i> (asi)	235	202	44	11	23	22
<i>Catharanthus roseus</i> (cro)	217	166	68	17	38	37
<i>Citrus sinensis</i> (csi)	948	470	530	52	641	273
<i>Carthamus tinctorius</i> (cti)	287	250	42	5	22	21
<i>Dimocarpus longan</i> (dlo)	195	154	55	14	30	30
<i>Dendranthema morifolium</i> (dmo)	497	199	312	14	317	168
<i>Dendrobium officinale</i> (dof)	669	283	408	22	370	225
<i>Eucommia ulmoides</i> (eul)	313	313	0	0	0	0
<i>Ginkgo biloba</i> (gbi)	438	298	150	10	106	77
<i>Juglans regia</i> (jre)	329	224	148	43	176	88
<i>Litchi chinensis</i> (lch)	298	219	85	6	84	43
<i>Lonicera japonica</i> (lja)	315	161	171	17	140	117
<i>Lycium chinense</i> (lyc)	298	245	67	14	41	35
<i>Nelumbo nucifera</i> (nnu)	627	453	207	33	198	107
<i>Oenanthe javanica</i> (oja)	25	25	0	0	0	0
<i>Panax ginseng</i> (pgi)	249	117	135	3	167	77
<i>Punicat granatum</i> (pgr)	261	258	4	1	2	2
<i>Paeonia lactiflora</i> Dafugui (plaD)	192	192	0	0	0	0
<i>Paeonia lactiflora</i> Hongyanzhenghui (plaH)	154	154	0	0	0	0
<i>Paeonia lactiflora</i> Zifengyu (plaZ)	163	163	0	0	0	0
<i>Panax notoginseng</i> (pno)	339	281	66	8	45	34
<i>Papaver somniferum</i> (pso)	164	138	26	0	14	13
<i>Ricinus communis</i> (rco)	178	159	34	15	29	18
<i>Raphanus sativus</i> (rsa)	124	120	4	0	2	2
<i>Salvia miltiorrhiza</i> (smi)	345	220	143	18	122	75
<i>Taxus mairei</i> (tma)	180	132	49	1	25	25
<i>Taxus x media</i> (tme)	581	315	281	15	257	145




Figure 1. Brief summary of MepmiRDB (medicinal plant microRNA database). (A) Functional flow chart of MepmiRDB. The database provides four major functional modules, ‘Sequence’, ‘Expression’, ‘Interaction’ and ‘Search’. All of the sequence and expression data could be retrieved from ‘Download’. (B) 5’ first nucleotide composition of the miRNAs deposited in MepmiRDB. (C) Sequence length distribution of the miRNAs deposited in MepmiRDB. For (B) and (C), the 29 plant species are indicated by different colors, as shown on the bottom.

pairs were subject to degradome-seq data-based validation according to the previously reported method (32). As a result, 380 pairs involving 119 miRNAs and 47 target transcripts in *Ammopiptanthus mongolicus*, 2341 pairs involving 192 miRNAs and 208 targets in *Citrus sinensis*, 13 023 pairs involving 389 miRNAs and 4506 targets in *Dendrobium officinale*, 6933 pairs involving 216 miRNAs and 2968 targets in *Nelumbo nucifera*, 9660 pairs involving 143 miRNAs and 2009 targets in *Panax ginseng*, 1560 pairs involving 130 miRNAs and 864 targets in *Salvia miltiorrhiza*, 149 pairs involving 72 miRNAs and 55 targets in *Taxus mairei* and 262 pairs involving 22 miRNAs and 98 targets in *Taxus × media* were identified. For each pair, t-plot (target plot) showing target cleavage signal(s) and visual network showing regulatory relationship between the miRNA and its target are available for users (Figure 2E and F). Besides, the transcripts interacting with the miRNA targets at the protein level were retrieved from the STRING database (33) by using BLASTX (34) (v2.2.26,

E -value $< 1e-3$, query coverage $> 50\%$, identity $> 35\%$). As a result, the protein–protein interactions were also visible in the regulatory networks (Figure 2E).

The ‘Search’ module

The module enables users to search for the miRNAs by plant species and miRNA families, or by sequences. For sequence search, it enables users to perform either ‘exact’ or ‘fuzzy’ search.

Other functional modules

The ‘Statistics’ module provides users with statistical information in both table and figure formats. Besides, all of the sequence and expression information reported in MepmiRDB has been made freely available in the ‘Download’ module. A help document is available in the ‘Help’ module for the first-time visitors.

Funding

National Natural Science Foundation of China (81730108, 31601062 and 31771436); Major Increase Or Decrease Program In The Central Finance Level (2060302); Key Project of Zhejiang Province Ministry of Science and Technology (2015C03055); Key Project of Hangzhou Ministry of Science and Technology (20162013A07).

Conflict of interest. None declared.

References

- Jones-Rhoades,M.W., Bartel,D.P. and Bartel,B. (2006) MicroRNAs and their regulatory roles in plants. *Annu. Rev. Plant Biol.*, **57**, 19–53.
- Kozomara,A. and Griffiths-Jones,S. (2014) miRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.*, **42**, D68–D73.
- Zhang,Z., Yu,J., Li,D. *et al.* (2010) PMRD: plant microRNA database. *Nucleic Acids Res.*, **38**, D806–D813.
- Szczesniak,M.W. and Makalowska,I. (2014) miRNEST 2.0: a database of plant and animal microRNAs. *Nucleic Acids Res.*, **42**, D74–D77.
- Bielewicz,D., Dolata,J., Zielezinski,A. *et al.* (2012) mirEX: a platform for comparative exploration of plant pri-miRNA expression data. *Nucleic Acids Res.*, **40**, D191–D197.
- Meng,Y., Gou,L., Chen,D. *et al.* (2011) PmiRKB: a plant microRNA knowledge base. *Nucleic Acids Res.*, **39**, D181–D187.
- Yi,X., Zhang,Z., Ling,Y. *et al.* (2015) PNRD: a plant non-coding RNA database. *Nucleic Acids Res.*, **43**, D982–D989.
- The Pharmacopoeia Editorial Committee (2015) *Pharmacopoeia of the People's Republic of China*. Chemical Technology Press, Shanghai, China.
- Bulgakov,V.P. and Avramenko,T.V. (2015) New opportunities for the regulation of secondary metabolism in plants: focus on microRNAs. *Biotechnol. Lett.*, **37**, 1719–1727.
- Gupta,O.P., Karkute,S.G., Banerjee,S. *et al.* (2017) Contemporary understanding of miRNA-based regulation of secondary metabolites biosynthesis in plants. *Front. Plant Sci.*, **8**, 374.
- Boke,H., Ozhuner,E., Turktas,M. *et al.* (2015) Regulation of the alkaloid biosynthesis by miRNA in opium poppy. *Plant Biotechnol. J.*, **13**, 409–420.
- Vashisht,I., Mishra,P., Pal,T. *et al.* (2015) Mining NGS transcriptomes for miRNAs and dissecting their role in regulating growth, development, and secondary metabolites production in different organs of a medicinal herb, *Picrorhiza kurroa*. *Planta*, **241**, 1255–1268.
- Yu,Z.X., Wang,L.J., Zhao,B. *et al.* (2015) Progressive regulation of sesquiterpene biosynthesis in arabidopsis and patchouli (*Pogostemon cablin*) by the miR156-targeted SPL transcription factors. *Mol. Plant*, **8**, 98–110.
- Zhang,M., Dong,Y., Nie,L. *et al.* (2015) High-throughput sequencing reveals miRNA effects on the primary and secondary production properties in long-term subcultured *Taxus* cells. *Front. Plant Sci.*, **6**, 604.
- Rupaimoole,R. and Slack,F.J. (2017) MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat. Rev. Drug Discov.*, **16**, 203–222.
- Li,J., Zhang,Y., Li,D. *et al.* (2015) Small non-coding RNAs transfer through mammalian placenta and directly regulate fetal gene expression. *Protein Cell*, **6**, 391–396.
- Sala-Cirtog,M., Marian,C. and Anghel,A. (2015) New insights of medicinal plant therapeutic activity—the miRNA transfer. *Biomed. Pharmacother.*, **74**, 228–232.
- Chin,A.R., Fong,M.Y., Somlo,G. *et al.* (2016) Cross-kingdom inhibition of breast cancer growth by plant miR159. *Cell Res.*, **26**, 217–228.
- Zhao,Y., Cong,L. and Lukiw,W.J. (2018) Plant and animal microRNAs (miRNAs) and their potential for inter-kingdom communication. *Cell. Mol. Neurobiol.*, **38**, 133–140.
- Arroyo,J.D., Chevillet,J.R., Kroh,E.M. *et al.* (2011) Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc. Natl. Acad. Sci. U. S. A.*, **108**, 5003–5008.
- Liang,G., Zhu,Y., Sun,B. *et al.* (2014) Assessing the survival of exogenous plant microRNA in mice. *Food Sci. Nutr.*, **2**, 380–388.
- Yang,J., Farmer,L.M., Agyekum,A.A. *et al.* (2015) Detection of dietary plant-based small RNAs in animals. *Cell Res.*, **25**, 517–520.
- Hong,M., Wang,N., Tan,H.Y. *et al.* (2015) MicroRNAs and Chinese medicinal herbs: new possibilities in cancer therapy. *Cancers*, **7**, 1643–1657.
- Zhou,Z., Li,X., Liu,J. *et al.* (2015) Honeysuckle-encoded atypical microRNA2911 directly targets influenza A viruses. *Cell Res.*, **25**, 39–49.
- Xie,W., Weng,A. and Melzig,M.F. (2016) MicroRNAs as new bioactive components in medicinal plants. *Planta Med.*, **82**, 1153–1162.
- Yu,D., Wan,Y., Ito,H. *et al.* (2019) PmiRDiscVali: an integrated pipeline for plant microRNA discovery and validation. *BMC Genomics*, **20**, 133.
- Grabherr,M.G., Haas,B.J., Yassour,M. *et al.* (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnol.*, **29**, 644–652.
- Yang,X. and Li,L. (2011) miRDeep-P: a computational tool for analyzing the microRNA transcriptome in plants. *Bioinformatics*, **27**, 2614–2615.
- Ma,X., Tang,Z., Qin,J. *et al.* (2015) The use of high-throughput sequencing methods for plant microRNA research. *RNA Biol.*, **12**, 709–719.
- Yu,D., Ma,X., Zuo,Z. *et al.* (2017) Bioinformatics resources for deciphering the biogenesis and action pathways of plant small RNAs. *Rice*, **10**, 38.
- Dai,X. and Zhao,P.X. (2011) psRNATarget: a plant small RNA target analysis server. *Nucleic Acids Res.*, **39**, W155–W159.
- Shao,C., Chen,M. and Meng,Y. (2013) A reversed framework for the identification of microRNA-target pairs in plants. *Brief. Bioinform.*, **14**, 293–301.
- Szklarczyk,D., Morris,J.H., Cook,H. *et al.* (2017) The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res.*, **45**, D362–D368.
- Altschul,S.F., Madden,T.L., Schaffer,A.A. *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.*, **25**, 3389–3402.