

Review Article

Novel Drug Design for Treatment of COVID-19: A Systematic Review of Preclinical Studies

Sarah Mousavi ¹, Shima Zare,² Mahmoud Mirzaei ³, and Awat Feizi ⁴

¹Department of Clinical Pharmacy and Pharmacy Practice, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

²School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

³Child Growth and Development Research Center, Research Institute for Primordial Prevention of Non-Communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran

⁴Department of Epidemiology and Biostatistics, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran

Correspondence should be addressed to Sarah Mousavi; s.mousavi@pharm.mui.ac.ir

Received 19 March 2022; Revised 23 May 2022; Accepted 3 August 2022; Published 25 September 2022

Academic Editor: Siddhartha Pati

Copyright © 2022 Sarah Mousavi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Since the beginning of the novel coronavirus (SARS-CoV-2) disease outbreak, there has been an increasing interest in discovering potential therapeutic agents for this disease. In this regard, we conducted a systematic review through an overview of drug development (in silico, in vitro, and in vivo) for treating COVID-19. **Methods.** A systematic search was carried out in major databases including PubMed, Web of Science, Scopus, EMBASE, and Google Scholar from December 2019 to March 2021. A combination of the following terms was used: coronavirus, COVID-19, SARS-CoV-2, drug design, drug development, In silico, In vitro, and In vivo. A narrative synthesis was performed as a qualitative method for the data synthesis of each outcome measure. **Results.** A total of 2168 articles were identified through searching databases. Finally, 315 studies (266 in silico, 34 in vitro, and 15 in vivo) were included. In studies with in silico approach, 98 article study repurposed drug and 91 studies evaluated herbal medicine on COVID-19. Among 260 drugs repurposed by the computational method, the best results were observed with saquinavir ($n = 9$), ritonavir ($n = 8$), and lopinavir ($n = 6$). Main protease ($n = 154$) following spike glycoprotein ($n = 62$) and other nonstructural protein of virus ($n = 45$) was among the most studied targets. Doxycycline, chlorpromazine, azithromycin, heparin, bepridil, and glycyrrhizic acid showed both in silico and in vitro inhibitory effects against SARS-CoV-2. **Conclusion.** The preclinical studies of novel drug design for COVID-19 focused on main protease and spike glycoprotein as targets for antiviral development. From evaluated structures, saquinavir, ritonavir, eucalyptus, *Tinospora cordifolia*, aloe, green tea, curcumin, pyrazole, and triazole derivatives in in silico studies and doxycycline, chlorpromazine, and heparin from in vitro and human monoclonal antibodies from in vivo studies showed promised results regarding efficacy. It seems that due to the nature of COVID-19 disease, finding some drugs with multitarget antiviral actions and anti-inflammatory potential is valuable and some herbal medicines have this potential.

1. Introduction

Coronavirus disease 2019 (COVID-19), which was first identified in December 2019, and shortly after, declared a pandemic by World Health Organization (WHO) [1]. As of January 18, 2022, there have been more than 326 million confirmed cases and 5.54 million deaths globally [2]. Coronaviruses belong to the family of Coronaviridae, RNA viruses with crown-like spikes on the surface of the

coronavirus particles. According to a meta-analysis of Macedo et al. [3], the mortality rate of COVID-19 was 17.1% for patients admitted to hospitals, whereas WHO estimated a fatality rate of 6.73%, which was much lower than that calculated from published studies. Among the critical cases of COVID-19, the mortality rate reaches 40% [4].

Substantial efforts have been made in the treatment of patients with COVID-19. The WHO recommendations in the treatment of COVID-19 are as follows [5]: molnupiravir

(conditional), baricitinib (strong), ruxolitinib and tofacitinib (conditional), sotrovimab (conditional), casirivimab and imdevimab (conditional), IL-6 receptor blockers (tocilizumab and sarilumab) (strong), remdesivir (conditional), and systemic corticosteroids (strong). The WHO recommends not to use ivermectin, lopinavir/ritonavir, hydroxy-chloroquine, and convalescent plasma.

The pathogenesis of COVID-19 was explained by cytokine storm, reduction in ACE2 expression, and activation of complement pathway-induced microvascular injury and thrombosis [6]. The mechanisms of the recommended agents are focused on the mentioned pathogenesis to improve the clinical outcome of COVID-19, and antiviral therapies are missing. The anticoronaviral strategies include preventing the synthesis of viral RNA, inhibiting virus replication, blocking the virus binding to human cell receptors, or inhibiting the viruses' self-assembly process [7]. The SARS-CoV-2 contains at least four structural proteins: spike (S) protein, envelope (E) protein, membrane (M) protein, and nucleocapsid (N) protein, and 16 nonstructural proteins (NSPs). Among the translated NSPs, the main protease, also called chymotrypsin-like protease (3C-like protease), and the papain-like protease are two essential proteases for proteolytic processing of the coronavirus replicase polyprotein, therefore generating functional replication complex of the virus, whereas RNA-dependent RNA polymerase is the central enzyme for RNA synthesis. These three NSPs play crucial roles in coronavirus replication, making them attractive targets for anticoronaviral drug design [8].

The S protein, a surface-located trimeric glycoprotein of coronaviruses, promotes the attachment of viruses to host cells through binding to angiotensin-converting enzyme 2 (ACE2) and virus-cell membrane fusion during viral infection. Thus, the S protein has been considered as a major target for the development of vaccines and drug [9].

The development of a new therapeutic agent is a complex, lengthy, and expensive process, which can take 2–4 years of preclinical development and 3–6 years of clinical development and over 500 million dollar cost. There are three critical steps to develop a new drug including discovery and development, preclinical research, and clinical development [10].

Drug discovery involves screening hits, medicinal chemistry, and optimization of hits to reduce potential drug side effects. For drug discovery, two different complementary approaches can be applied: classical pharmacology, also known as phenotypic drug discovery, which is the historical basis of drug discovery, and reverse pharmacology, also known as designated target-based drug discovery. Screening methods based on phenotypic drug discovery have been used to discover new natural products mainly from the terrestrial origin [11]. These two strategies have advantages and disadvantages and promote very different screening assays. The frequent re-discovery of the same compounds, the technical difficulties associated with the isolation of compounds from extracts, and the incompatibility of natural product extracts with high-throughput screening (HTS) campaigns were the disadvantages of phenotypic drug

discovery. On the other hand, natural product structures have the characteristics of high chemical diversity, biochemical specificity, and other molecular properties that make them favorable as lead structures for drug discovery, which serve to differentiate them from libraries of synthetic and combinatorial compounds [12]. Overly simplified assays, acting of drug on more than one target, the multifactorial nature of diseases, and challenges to identify a single molecular target are some limitations of target-based drug discovery. Therefore, a comprehensive screening strategy will incorporate both targeted and phenotypic assays, with one format designated as the primary screen and the other as a secondary or follow-up assay. During the spread of COVID-19 outbreak, great efforts have been made in therapeutic drug discovery against the virus. Because COVID-19 is a new, acute, severe infectious disease, the anti-SARS-CoV-2 drug development strategies are to screen existing drugs to identify potentially effective drugs, to expand indications, and to develop a vaccine [12]. The safety of conventional drugs has been mostly verified; if effective, they can be quickly applied in clinical practice (repurposing of existing drugs). The recent rises of several high transmissible strains sounded alarms for currently used vaccines and drugs. Therefore, developing broad-spectrum antiviral drugs not only to combat COVID-19 but also to provide protective arsenals against future viral outbreaks is a requirement. Scientists continue the development of broad-spectrum antiviral drugs from natural or chemical sources, which have the potential advantages of broad-spectrum therapeutic effect and insensitivity to viral evasion. Given the urgency of the SARS-CoV-2 outbreak, here we discuss the discovery and development of new therapeutics for SARS-CoV-2 infection based on the strategies from preclinical drug discovery.

2. Methods

We report this systematic review based on the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines [13].

2.1. Data Sources and Searches. Studies published in PubMed, Scopus, Web of Science, EMBASE, Google Scholar, and DrugBank were searched from December 2019 to March 2021 using the following search terms: "Coronavirus," "Covid-19," "SARS-CoV-2," "Drug design," "Drug development," "In silico," "In vitro," and "In vivo" alone or in combination without language restrictions. The keywords were selected using expert opinions, mesh, and related article titles.

All articles with full text or in the absence of full text with abstract are included in the screening of this study. Studies were excluded if studies were comments, editorial, letters, review, and preprints.

2.2. Data Extraction. Two researchers independently extracted data from included studies using a predefined data extraction form. All disagreement was discussed and solved

after rechecking the source data with a third investigator. The data extracted, including the last name of the first author, type of the study (in silico, in vitro, in vivo), and name of the agent (chemical compound, drug, herb, etc.), studied the mechanism and efficacy of the agent according to study design that specified according to the following definition.

Computer-aided drug design can be divided into three different categories. All are based on ligands and receptors, which are briefly described [14] as follows:

- (1) *Dock Receptor-Based Approach* [15]. Once the three-dimensional structure of the ligand molecules and their receptor is known, the receptor-based method is a good candidate for identifying or optimizing drugs. Due to the presence of three-dimensional structures of compounds and receptors, the nature of the interaction between the ligand and the receptor and the type of structure that the ligand can have to interact with them in favorable conditions can be identified using this method. The compound is simulated on the active site of the dock (meaning anchoring) and on the interaction of the ligand with the receptor by molecular mechanics and molecular dynamics. In this method, due to the ligation of the ligand in the active position, the ligand changes in terms of conformity and changes its position in different conditions and shows interaction with the receptor in different types of situations. To determine the type of ligands that can be docked into the receptor site, the matching of the shape and the complementarity of the hydrophobic, hydrophilic, and charged parts must be considered. Various software packages such as AUTODOCK-Glide, LUDI and LigandFit are used to design the drug based on the structure of the receptor.
- (2) *Ligand-Based Approach* [16]. This method is used where the three-dimensional structures of the receptor are unknown and instead the structure of the ligands is known, which is one of the common methods. In this method, by indirectly studying compounds that react with biomolecules, they seek to design compounds that are pharmacologically active. In ligand-based drug design methods, in the absence of biomolecule structure, by studying specific ligands, it seeks to identify the structural and physicochemical properties of the compounds so that the desired compound can be designed based on data extracted from the study of previous compounds. This method is a kind of drug design based on pharmacophore (pharmacophore refers to the part of the drug to which the effect of the drug depends on that part of the molecule), and by studying the quantitative relationship between structure and their activity, drugs can be designed by this method. It can be said that it is a method for designing the pharmacophores of drugs.
- (3) *Denovo Design-Based Approach* [17]. This method is used when the structure of the ligand is unknown but the structure of the receptor is known. In this method, there is information about the structures of the receptor or quasi-receptors, but there is no structure of the main composition that can interact with the active site of the receptor. One of the functions of drug design based on this method is to suggest and present the main composition that is complementary to the active site. The basis of the method is that the database of existing 3D structures is used to find small molecules that can interact with the active site of the receptor in terms of size, geometry, and functional groups. Software packages such as GROW and LEGEND are used to design drugs by this method.

Drug design methods in the computer include quantitative structure-activity relationship (RASQ), docking, molecular dynamic simulation, and computational modeling. In these studies, the efficacy is evaluated based on the function of the drug or compound agent and the mechanism of action. Figure 1 shows the methods of computer-aided drug design (CADD).

2.2.1. Computational Methods in Drug Design.

Quantitative Structure-Activity Relationship (QSAR). QSAR provides studies on the relationship between chemical structure and biological activity or other biological activities that are important in selecting or removing a compound before synthesis and testing. QSAR [18] is especially important to predict the result, especially when it is not possible to experiment with a compound. Molecular descriptors, which are the most important components of QSAR, can be obtained experimentally or through mathematical formulas from various theories such as quantum mechanics, chemical graph theory, and study theories. QSAR seeks to establish a statistically significant relationship between structure and performance. It also explains the specific effect of a drug and can ultimately predict the effect of newly synthesized chemical compounds. QSAR model is also an equation that predicts a property through molecular descriptors and their coefficients. Evaluation of the effectiveness of new compounds that have been studied using this method can be reported as a percentage of enzyme inhibition if the modeling has been done and mentioned in the article.

2.2.2. Docking. In this technique, to achieve a combination with a pharmacological effect and increase the pharmacological activity of the drug, different formulations of a drug interact with the receptor, and the structure that has the best interaction with the receptor and the lowest energy level is selected for laboratory steps [19]. In this way, possible structures that have a stronger interaction with the receiver can be isolated at this stage. The same issue is considered and reported as an effectiveness measure.




FIGURE 1: Computational methods in drug design (CADD: computer-aided drug design).

2.2.3. Molecular Dynamic Simulation. In this technique, which is based on the simulation of drug-receptor interaction in the body, docking problems are solved and in fact play a complementary role in this technique [20]. Due to the time-consuming work with this technique, effective structures cannot be achieved directly through it, and the final stages of the study of drug-receptor interactions should be evaluated before starting laboratory work by having effective compounds from the previous stages. Molecular dynamic simulations produce information at the microscopic level (position and velocity of atoms). The conversion of these data into macroscopic values (pressure, energy, etc.) is done using statistical mechanics. In fact, molecular dynamics and statistical mechanics link microscopic concepts and macroscopically observable quantities. Molecular dynamic simulations are only able to predict the thermodynamic behavior and stability of the ligand binding mechanism at the active site of the target enzyme. This is reported as a criterion.

2.2.4. In Vitro. Study of drug in cell culture medium: effectiveness in these studies means inhibition of the replication of COVID-19 by the compound or drug under study [21].

The half-maximal inhibitory concentrations (IC_{50}) are a measure of the effectiveness of a compound in inhibiting biological function [22].

In vivo studies are those in which the effects of drugs are tested on whole living organisms or cells usually animals as opposed to a tissue organism or dead organism. In vivo testing is better studied for observing the overall effects of an experiment on living subjects [23].

3. Results

In this review, we reported a significant number of articles with *in silico*, *in vitro*, and *in vivo* approaches for drug development of COVID-19. We retrieved a total of 2538 articles from the initial database search. After the removal of duplication and screening, 317 studies were selected for inclusion in this review. Figure 2 shows the PRISMA diagram.

The analysis of article contents indicated that 266 studies performed *in silico* approaches against viral targets; 34 studies used *in vitro* approaches against SARS-CoV-2; and 15 studies used *in vivo* (animal) models.

3.1. Results from In Silico Drug Discovery. From 267 studies used in *in silico* approaches, 98 article studies repurposed approved drugs with a new mechanism of action and 91 studies evaluated natural products (e.g., herbal medicine) on COVID-19. The characteristics of these studies are summarized in Tables 1 and 2. Also, Table 3 shows the characteristics of the remaining studies ($N=87$).

In *in silico* studies used the following component of novel coronavirus as targets: main protease ($N=154$), spike glycoprotein ($N=62$), nonstructural protein ($N=45$), RNA-dependent RNA polymerase ($N=21$), papain-like protease ($N=19$).

About 260 drugs were repurposed by the computational methods for COVID-19 therapy such as about 120 drugs candidate against the main protease, 52 drugs against the spike glycoprotein, 14 drugs against RNA-dependent RNA polymerase, and 28 drugs against other nonstructural proteins.




FIGURE 2: PRISMA diagram of the study.

Among the studied repurposed drugs, the best results (regarding efficacy) were observed with saquinavir ($N=9$ study), ritonavir ($N=8$ study), lopinavir ($N=6$ study), remdesivir ($N=3$ study), and amikacin, danoprevir, favipiravir, and telaprevir.

Table 1 shows target-based synthesis of data for COVID-19 drug repurposing. As presented, at least two studies show the efficacy of aprepitant, cobicistat, dipyrindamole, and dihydroergotamine against the main protease and tegobuvir ($N=2$) against spike protein. The following list of drugs had multitarget action: avapritinib, famotidine, bicitegravir, ziprasidone, capmatinib, pexidartinib, amprenavir, zafirlukast, cilostazol, paromomycin, lopinavir, and remdesivir.

A total of 91 studies used in silico methods to evaluate the effects of natural products including herbal medicine against SARS-CoV-2. Among them, 54 studies used the main protease as main target, in which eucalyptus ($N=3$), *Tinospora cordifolia* ($N=3$), and flavonoids (e.g., hesperidin, rutin, and herbacetin) were the most studied and effective. Other studied targets were as follows: spike ($N=22$) and multitarget ($N=20$). From the plant metabolites, oleanolic acid, hesperidin, epigallocatechin gallate, jensenone, tinosponone, and anistone show promising results in computational methods against COVID-19. Aloe, green tea, eucalyptus, curcumin, and many Chinese, Indian, and African plants were also effective on COVID-19 in silico.

The derivatives of pyrazoles, oxadiazoles, phenyl-triazolinones, triazoles, benzoylpinostrobin, benzoic acid, benzylidenechromanones, coumarin, and selenium show efficacy in computational methods and could be considered as lead molecules for drug design and synthesis against COVID-19. Melatonin, ebselen, phenyl furoxan, thimerosal,

isatin, romidepsin, phenyl mercurin, pleconaril, and tyrosine kinase inhibitors (such as nilotinib and imatinib) are examples of other drugs that are effective in in silico methods.

3.2. Results from In Vitro and In Vivo Studies. Screening of in vitro studies leads to finding 34 studies. Different compounds were evaluated and most studies focused on the inhibition of viral replication, which was assessed by the quantification of viral RNA by PCR and IC_{50} value reported. Most of the studies use the Vero E6 cell line for the assessment of replication of SARS-CoV-2 (Table 4). In particular, main protease, papain-like protease, RNA-dependent RNA polymerase, NSP-14, NSP-15, spike protein, and TMPRSS2 were evaluated in vitro as targets. Also, three studies focused on in vitro inhibition of inflammatory markers such as interleukins and the effects of the cytopathic effect on infected cell.

Nitric oxide, ginkgolic acid, anacardic acid, troxerutin, bisindolylmaleimide derivatives, small molecules (GRL-172, and 5 h), baicalin and baicalein (phytochemicals), and bepridil were effective drugs in vitro against the main protease of COVID-19.

The following drugs could inhibit the spike/ACE2-mediated cell entry of COVID-19 in vitro: romidepsin, panobinostat, givinostat, sirtinol, saquinavir, lipopeptides, hydroxyzine, azelastine, heparin, and glycyrrhizic acid.

Alkenyl sulfonylurea derivatives [7], an IMU-8381 inhibitor of human dihydroorotate dehydrogenase, pyrazole derivatives, and phillyrin, regulated the expression of inflammatory cytokines (e.g., IL, TNF- α , and NF- κ B) induced

TABLE 1: Summary of studies with in silico method that used repurposed drugs for novel drug discovery against COVID-19.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Abosheasha and El-Gowily [24]	2020	In silico	Drug repurposing molecular docking-based virtual screening	15 antiplatelet FDA-approved drugs	Main protease (Mpro) and spike glycoprotein (S)	Cilostazol has favorable binding interaction with Mpro (PDB ID: 6LU7) cilostazol, iloprost, epoprostenol, prasugrel, and icosapent ethyl that have a higher binding affinity on spike glycoprotein (S) Drugs that shared >70% similarity to the binding sites of those targets were reversin, pentagastrin, remdesivir, norfloxacin, and nitazoxanide against COVID-19 papain-like protease whereas benzylglutathione, lopinavir, and hydroxymethylglutathione against RNA polymerase Best MM/GBSA binding energy; ChEMBL275592, montelukast, ChEMBL28834. Bromocriptine and saquinavir demonstrate stability in the active site of Mpro Isavuconazonium, a P2-P3 α -ketoamide derivative, and pentagastrin are the top three molecules (Lig13b as the benchmark) based on docking energy	Cilostazol is a promising FDA drug against COVID-19 by inhibiting both Mpro and S protein
Abdul Kadhim et al. [25]	2020	In silico	Drug repurposing, docking	Experimental and approved drugs	Papain-like protease and RNA polymerase		Antiresistance reversin showed the highest inhibitory efficacy against COVID-19 papain-like protease, and benzylglutathione is an experimental compound; however, it had the highest RNA polymerase inhibiting efficacy
Abu-Saleh et al. [26]	2020	In silico	Ligand-based/structure-based virtual screening, MD simulations, and binding energy calculations	Approved drugs and bioactive compounds listed in the DrugBank and ChEMBL databases	Main protease		
Achilonu et al. [27]	2020	In silico	High-throughput virtual screening and ligand docking	FDA-approved drugs	Main protease		
Aftab et al. [28]	2020	In silico	Repositioning/target-based virtual screening and molecular docking	Ten antiviral drugs were screened: ribavirin, remdesivir, sofosbuvir, penciclovir, nitazoxanide, nafamostat, chloroquine, galidesivir, favipiravir, and interferon	RNA-dependent RNA polymerase (RdRp)	Galidesivir and its drug-like compounds CID123624208 and CID11687749 have shown an effective attachment to the priming site of viral RdRp	CID123624208 and CID11687749 may be considered for in vitro and in vivo clinical trials
Ahmadi et al. [29]	2021	In silico	Drug repurposing study using molecular docking	Enfuvirtide, an HIV-1 fusion inhibitor peptide	SARS-CoV-2 fusion inhibitor	Enfuvirtide binding to the S2 protein of SARS-CoV-2 was remarkably stable and can act as a strong SARS-CoV-2 fusion inhibitor	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Ahmed et al. [30]	2021	In silico	Drug repurposing (high-throughput virtual screening) (HTVS) followed by re-docking with standard precision (SP) and extra-precision (XP) molecular docking	FDA-approved antiviral and anti-infection drugs	Main protease	Of 1397 potential drugs, 157 showed considerable affinity towards Mpro. High-affinity lead drugs (iodixanol, amikacin, troxerutin, and rutin) were identified. Amikacin was found to be the most potent inhibitor of main protease	Aminoglycosides may serve as a scaffold to design potent drug molecules against COVID-19
Anand et al. [31]	2021	In silico	Molecular docking	130 US FDA-approved drugs including hypertension, cardiovascular diseases, respiratory tract infections (RTI), antibiotics, and antiviral drugs	Structural and nonstructural proteins of SARS-CoV-2 (nsp3, nsp5, nsp10, nsp16)	15 potent drugs exhibiting significant inhibitory potential against SARS-CoV-2 like baloxavir marboxil, danoprevir and sofosbuvir, fosinopril, moexipril, quinapril, telmisartan, azilsartan, verapamil, and doxazosin	Azithromycin, doxycycline, clarithromycin, rifamycin, and augmentin; nsp10; virginiamycin, tunicamycin, quinupristin, fidaxomicin, digoxin, and azithromycin; main protease; caspofungin, amphotericin B, ketoconazole, and micafungin; E and N proteins; virginiamycin and amphotericin B; S protein
Ancy et al. [32]	2020	In silico	Molecular docking, molecular dynamics, and binding free energy simulation study	HIV-1 protease, namely, TMB607 and TMC-310911	Main protease	TMB607 molecule binds strongly with the SARS-CoV-2 main protease enzyme	
Ansari et al. [33]	2020	In silico	Repurposing drug molecular docking	TAT-peptide 47-57 (GRKKRRQRRRP)-conjugated repurposed drugs (i.e., lopinavir, ritonavir, favipiravir, and hydroxychloroquine) Drugs available in the super DRUG2 database	Main protease	TP-conjugated ritonavir, lopinavir, favipiravir, and hydroxychloroquine have superior and significantly enhanced interactions with main protease	
Arun et al. [34]	2020	In silico	Repurposing drug molecular docking	FDA-approved drugs	Main protease	Binifibrate and bamifylline bind strongly to the enzyme active site	
Arya et al. [35]	2020	In silico	Molecular docking	FDA-approved drugs	Papain-like protease	15 FDA-approved drugs, including chloroquine and formoterol, bind the target enzyme with significant affinity and good geometry, suggesting their potential to be utilized against the virus	
Baby et al. [36]	2020	In silico	Schrodinger's computer-aided drug discovery tools for in silico drug repurposing	FDA-approved library of drugs	RNA-dependent RNA polymerase (RdRp)	Pitavastatin, ridogrel, rosoxacin	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Baby et al. [36]	2021	In silico	Schrodinger's computer-aided drug discovery tools for in silico drug repurposing	FDA-approved library of drugs	Main protease	Tipiracil and aprepitant interacted with the main protease	
Baker et al. [37]	2021	In silico	Repurposing drug molecular docking	50 compounds with activity against main protease	Main protease	Drugs including boceprevir, ciluprevir, nartlaprevir, and telaprevir may be more potent against main protease than boceprevir and suitable for rapid repurposing	
Bharath et al. [38]	2020	In silico	Drug repurposing computer-aided drug design (CADD)	4015 known and approved small molecules	Spike glycoprotein	Glycyrrhizic acid (GA) of plant origin may be repurposed for SARS-CoV-2 intervention	
Bhowmik et al. [39]	2021	In silico	Repurposing drugs, docking, and molecular dynamic simulation	Orientin (phytochemical)	Inhibitor of SARS-CoV-2 spike and host cell receptor GRP78 binding	Binding of orientin in the overlapping residues of GRP78 binding region of SARS-CoV-2 spike model	As a promising precautionary or therapeutic measure for COVID-19
Bolelli et al. [40]	2021	In silico	Drug repurposing, virtual screening method	FDA-approved drugs	Main protease	Three compounds (dobutamine and its two derivatives) Sovaprevir, elbasvir, danoprevir, samatasvir, Candesartan, saquinavir ritonavir, indinavir, lopinavir, brilacidin, flovagatran, aplidin, desmopressin, and felypressin listed as potential inhibitors of main protease	
Cavasotto et al. [41]	2021	In silico	Drug repurposing, docking-based screening using a quantum mechanical scoring	FDA-approved drugs	Spike protein, main protease papain-like protease	Conivaptan exhibited the highest binding of the Nsp9 replicase. Tegobuvir exhibited maximum stability along with the highest binding energy at the active site of the spike proteins	
Chandel et al. [42]	2020	In silico	Drug repurposing, molecular dynamics, and docking	FDA-approved drugs	Nsp9 replicase and spike proteins		
Chen et al. [43]	2020	In silico	Drug repurposing, molecular dynamics, and docking	FDA-approved drugs	Spike (S)-mediated cell entry	Cepharanthine, abemaciclib, osimertinib, trimipramine, colforsin, and ingenol	
Chidambaram et al. [44]	2020	In silico	Molecular docking	Coumarins and their analogs	Main protease	Natural coumarin analog toddacoumarinone displayed remarkable inhibition ability. Synthetic coumarin analog (1 m) also displayed the comparable inhibition ability main protease in intricate with α -ketoamide	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Choudhary et al. [45]	2020	In silico	Drug repurposing, molecular dynamics, and docking	FDA-approved drugs	Spike glycoprotein and cellular angiotensin-converting enzyme 2 (ACE2) receptor	GR 127935 hydrochloride hydrate, GNF-5, RS504393, TNP, and eptifibatide acetate were found binding to virus binding motifs of ACE2 receptor. KT203, BMS195614, KT185, RS504393, and GSK1838705A were identified to bind at the receptor-binding site on the viral S protein. Racemic mixtures of the ibuprofen enantiomers might be a potential treatment for main protease	
Clemente et al. [46]	2021	In silico	Molecular docking, molecular dynamic (MD) simulations	Ibuprofen	Main protease	Ivermectin could interfere with activity of spike proteins	
Cosic et al. [47]	2021	In silico	Extended resonant recognition model (RRM)	Ivermectin	Spike proteins	Lopinavir, remdesivir, ritonavir, saquinavir, and raltegravir were successfully docked into the binding site of main protease	
Daoud et al. [48]	2021	In silico	Structure-based pharmacophore approach, molecular docking, and repurposing studies	FDA-approved drugs	Main protease	24 best scored ligands (14 traditional herbal isolates and 10 approved drugs) as potential candidates to inhibit the S protein	Quinupristin, nilotinib, acetyldigitoxin
de Oliveira et al. [49]	2021	In silico	Molecular modeling and virtual screening and repurposing studies	9091 FDA-approved drugs	Spike protein	Zafirlukast as the best repurposing candidate for COVID-19	
Delijewski and Haneczok [50]	2021	In silico	Supervised machine learning model and repurposing studies	FDA-approved drugs	Against SARS-CoV-2		
Dey et al. [51]	2021	In silico	Virtual database screening, molecular docking, all-atom molecular dynamic simulation, and MM-PBSA analysis	Tretinoin, mefenamic acid, ondansetron, and artemether	Envelope (E) protein	Tretinoin as a potential SARS-CoV-2 E protein ion channel blocker and virus assembly inhibitor	
Durdagi [52]	2020	In silico	Molecular modeling approach in virtual drug screening repurposing study	FDA-approved drugs	Type 2 transmembrane serine protease (TMPRSS2)	Benzquercin as strong TMPRSS2 inhibitor	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Durdagi et al. [53]	2020	In silico	Molecular docking, MM-GBSA-based predictions, and molecular dynamic repurposing study	FDA-approved drugs	Main protease and spike receptor-binding domain bound with ACE2 COVID-19 target proteins	Pinelautide, rotigaptide, telinavir, ritonavir, pinokalan, terlakiren, cefotiam, and cefpiramide as SARS-CoV-2 main protease inhibitors. Denopamine, bomelolol, naminterol, rotigaptide, and benzquercin as potential ACE2/spike protein domain inhibitors HCV protease, DPP-4, α -thrombin, and coagulation factor Xa known inhibitors Four potential inhibitors against Mpro enzyme, two available drugs (Talampicillin and Lurasidone), and two novel drug-like compounds (ZINC000000702323 and ZINC0000012481889)	
Eleftheriou et al. [54]	2020	In silico	Molecular docking	34 approved and on-trial protease inhibitors	Main protease		
Elmezayen et al. [55]	2021	In silico	Molecular modeling approach in virtual drug screening repurposing study	Commercially available drugs and ZINC15 library	Main proteases		
Encinar et al. [56]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	9000 US Food and Drug Administration (FDA)-approved investigational and experimental drugs from the DrugBank repository	S-Adenosyl-L-methionine-binding pocket of nsp16, [2] the unique "activating surface" between nsp16 and nsp10, and [3] the RNA-binding groove of nsp16	Tegobuvir, sonidegib, siramesine, antrafenine, bemcentinib, itacitinib, or phthalocyanine antagonism of SARS-CoV-2 RNAs lacking 20-O-methylation	
Farag et al. [57]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	2000 FDA-approved drugs	Main protease	Darunavir, nelfinavir, and saquinavir bound to the central site of main protease substrate-binding pocket rosvastatin, montelukast, and the anti-histaminic fexofenadine bound to the terminal site of main protease substrate-binding pocket	Starting point for further in vitro and in vivo testing
Feng et al. [58]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	FDA-approved drugs	Spike protein	Eltrombopag possesses a high binding affinity to S protein plus human ACE2	
Ferraz et al. [59]	2020	In silico	Ligand and structure-based virtual screening, repurposing study	FDA-approved drugs	Main protease	Two oral (bedaquiline and glibenclamide) and one buccal drug (miconazole)	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Fischer et al. [60]	2020	In silico	Molecular docking approach in virtual drug screening and repurposing study	Over 606 million compounds	Main protease	12 purchasable compounds, with binding affinity to the target protease the natural compounds (–)—taxifolin and rhamnetin as potential inhibitors of main protease	Initiative for in vitro testing
Gimeno et al. [61]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	FDA-approved drugs	Main protease	Perampanel, carprofen, celecoxib, alprazolam, trovafloxacin, sarafloxacin, and ethyl biscoumacetate. Carprofen and celecoxib	Including the HIV protease inhibitor lopinavir/ritonavir combination (phase 4), glucocorticoid receptor agonist dexamethasone (phase 3/4), DNA replication inhibitor niclosamide (phase 2/3), antineoplastic agent lenalidomide (phase 4), and calcineurin inhibitor tacrolimus (phase 3), ABT-737 (BCL inhibitor), brefeldin-A (protein synthesis inhibitor), indirubin (CDK inhibitor), TPCA-1 (IKK inhibitor), lopinavir (HIV protease inhibitor), GW-441756 (growth factor receptor inhibitor), treprostinil (prostaglandin analog), typhostin-AG-1478 (EGFR inhibitor) and epoxysterol (LXR agonist), fostamatinib (SYK inhibitor), VER-155008 (HSP inhibitor), KU-0063794 (MTOR inhibitor), PIK-90 (PI3K inhibitor), linsitinib (IGF-1 inhibitor), TAK-715 (p38 MAPK inhibitor), Y-27632 (Rho-associated kinase inhibitor), AZ-628 (RAF inhibitor), and lestaurtinib (FLT3 inhibitor)
Guo et al. [62]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study single-cell RNA sequencing	US FDA-approved drugs	Against SARS-CoV-2	281 FDA-approved drugs that have the potential to be effective against SARS-CoV-2 infection, 16 of which are currently undergoing clinical trials to evaluate their efficacy against COVID-19	
Gupta et al. [63]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	FDA-approved drugs	Main protease	Cobicistat is the most efficient inhibitor of Mpro both in silico and in vitro	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Huynh et al. [64]	2021	In silico	Docking and molecular dynamics and repurposing study	FDA-approved drugs	Papain-like protease	The chances of drug repurposing for PLpro might be low	
Ibrahim et al. [65]	2020	In silico	Molecular dynamic simulations, molecular docking, MM-GBSA analysis, and repurposing study	DrugBank database	Main protease	DB02388 and cobicistat (DB09065)	
Ifikhar et al. [66]	2020	In silico	Molecular modeling approach in virtual drug screening and repurposing study	4574 compounds also containing FDA-approved drugs	RdRp, main protease, and helicase	Rimantadine, bagrosin, and grazoprevir showed binding to main protease. Casopitant is a neurokinin-1 receptor that showed binding to RdRp. Meclonazepam and oxiphenisatin showed specific interactions with helicase	
Jain and Mujiwar [67]	2020	In silico	Computational drug repurposing docking simulations	2880 FDA-approved drugs	Main protease	Metocurine, dihydroergotamine, imatinib, daunorubicin, bromocriptine, irinotecan, azelastine, gestodene, adapalene, and simvastatin	Metocurine was chosen as a safe and effective drug candidate for developing therapy against the viral Mpro enzyme of SARS-CoV-2 for the treatment of COVID-19
Jarvis et al. [68]	2020	In silico	Tier-based scoring system repurposing study	Clinically developed drugs	Potential repurposing against COVID-19	Four drug classes (antimalarial amino-quinolones, selective estrogen receptor modulators (SERMs), low potency tricyclic antidepressants, and tricyclic antidepressants) as potential drug candidates for COVID-19	The tricyclic antidepressants and tricyclic antidepressants were further excluded based on a high adverse event profile
Kadioglu et al. [69]	2021	In silico	Repositioning/virtual drug screening, molecular docking, and supervised machine learning algorithm drug repositioning	FDA-approved drug natural compound dataset ZINC database	Spike protein, nucleocapsid protein, and 2'-O-ribose methyltransferase	Conivaptan, paritaprevir, simeprevir, dihydroergotamine, ZINC000027215482, ZINC00025215584, loniflavone, procyanidin	
Kandeel et al. [70]	2020	In silico	Drug repurposing molecular dynamic (MD) simulations followed by molecular mechanics/generalized born surface area (MM/GBSA) binding energy calculations	1697 clinical FDA-approved drugs	Papain-like protease	Phenformin, quercetin, and ritonavir	Phenformin was more stable than quercetin and ritonavir

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Kandwal and Fayne [71]	2020	In silico	Repurposing drug computational design pharmacophore features	In-development/approved drugs	Viral nucleocapsid and nonstructural proteins	Isepanamic and streptomycin (nsp3); coenzyme-I, rutin, epigallocatechin gallate-(-), and procyanidin-b-2 (nsp7/nsp8/nsp12); paromomycin (nsp10/nsp16); olomoucine, sapropterin, tetrahydrofolic acid, INS316, and adenosine phosphate (nsp15); varespladib, hexanoic acid, citric acid, OSI-027, MK-5108, stepronin, calcium gluceptate, CPP, pirenexine, midafotel, and maltobionic acid (nucleocapsid)	
Khan et al. [72]	2020	In silico	Drugs repurposing molecular dynamic simulation	31 FDA-approved anti-HIV drugs, and traditional Chinese medicines (TCM) database	Main protease	Saquinavir and TCM5280805	
Khan et al. [73]	2020	In silico	Drugs repurposing molecular docking	23 prospective drug candidates	Main protease	Epirubicin, vapreotide, and saquinavir exhibited better binding affinity	Synergistic interaction
Kouznetsova et al. [74]	2020	In silico	Drugs repurposing molecular docking	FDA-approved drugs	Papain-like protease	Inhibitors of HIV, hepatitis C, and cytomegalovirus (CMV) demonstrated some activity	
Krishnaprasad et al. [75]	2020	In silico	Drugs repurposing molecular docking	FDA-approved library of drugs	RNA-dependent RNA polymerase	Pitavastatin, ridoqrel, and rosoxacin displayed superior binding with the active site	
Kumar et al. [76]	2020	In silico	Drugs repurposing docking and molecular dynamic (MD) simulations combined with molecular mechanics/generalized born surface area (MM/GBSA)	12 FDA-approved drugs (darunavir, indinavir, saquinavir, tipranavir, diosmin, hesperidin, rutin, raltegravir, velpatasvir, ledipasvir, rosvastatin, and borteozomib)	Main protease	Saquinavir as a potent inhibitor of dimeric main protease	
Kumar et al. [77]	2020	In silico	Drugs repurposing molecular docking molecular dynamic simulations MM/GBSA	Withaferin A (Wi-A), withanone (Wi-N) (active withanolides of ashwagandha), and caffeic acid phenethyl ester (CAPE, bioactive ingredient of propolis)	Main protease	Wi-N and CAPE possess the potential to inhibit the functional activity of main protease	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Kumar et al. [78]	2020	In silico	Drugs repurposing molecular docking	FDA-approved drugs	Main protease	Lopinavir-ritonavir, tipranavir, and raltegravir show the best molecular interaction with the main protease	
Kumar et al. [79]	2020	In silico	Drugs repurposing molecular docking molecular dynamic simulations	FDA-approved library of drugs	Main protease	Hyaluronic acid and acarbose show strong interactions with catalytic site residues of main protease	
Li et al. [80]	2020	In silico	Drug repurposing free energy perturbation-based absolute binding free energy (FEP-ABFE) predictions	Virtual screening of existing drugs	Main protease	25 drugs were predicted, and 15 were confirmed as potent inhibitors of SARS-CoV-2 main protease. The most potent one is dipyrindamole. Hydroxychloroquine (ki = 0.36 μ M) and chloroquine (ki = 0.56 μ M) were also found to potentially inhibit main protease 29 drugs that could actively interact with two or more target proteins, with 5 drugs (avapritinib, bicittegravir, ziprasidone, capmatinib, and pexidartinib) being common candidates for all four key host proteins and 3 of them possessing the desirable molecular properties	
Liang et al. [81]	2021	In silico	Drug repurposing molecular docking	2,631 FDA-approved small molecules	Multiple main proteins		
Lokhande et al. [82]	2021	In silico	Drugs repurposing molecular docking molecular dynamic simulations	FDA-approved drugs	Main protease	Mitoxantrone, leucovorin, birinapant, and dynasore	
Mahanta et al. [83]	2020	In silico	Drugs repurposing molecular docking molecular dynamic simulations	U.S. Food and Drug Administration-approved antimicrobial drugs	Main protease	Viomycin	
Mahdian et al. [84]	2021	In silico	Drugs repurposing molecular docking molecular dynamic simulations	FDA-approved drugs	Viral entry receptors (ACE2 and CD147) and integral enzyme of the viral polymerase (RdRp)	Ledipasvir, estradiol benzoate, and vancomycin and paritaprevir	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Marak et al. [85]	2020	In silico	Repurposing drug homology modeling molecular docking	108 FDA-approved antiparasitic and anti-inflammatory drugs	10 SARS-CoV-2 targets (PLpro, 3CLpro, RdRp, spike, helicase, NSP1, NSP3, NSP4, NSP9, and NSP16-NSP10)	Ivermectin, atovaquone, posaconazole, doxycycline, moxidecin, amphotericin B, chlortetracycline, spiramycin, sulfasalazine, parecoxib, and etoricoxib exhibited good binding affinities	
Mohapatra et al. [86]	2020	In silico	Repurposing drug machine learning (ML) technology	FDA-approved drugs	Against COVID-19	10 FDA-approved commercial drugs that can be used for repurposing amprenavir would probably be the most effective drug based on the selected criteria Nilotinib, imatinib, and dihydroergotamine for 3clpro and dexasone and raltegravir for RdRp. Raltegravir, an anti-HIV drug, was observed to be the best compound against RdRp based on docking binding energy dihydroergotamine is a suitable candidate for main protease BCR-ABL tyrosine kinase inhibitor, imatinib, inhibits SARS-CoV-2	Via fusion inhibition
Molavi et al. [87]	2021	In silico	Repurposing drug molecular docking	1760 FDA-approved drugs	RNA-dependent RNA polymerase (RdRp) and main protease		
Mulgaonkar et al. [88]	2020	In silico/in vitro	Repurposing drug molecular docking	FDA-approved drugs	Spike glycoprotein		
Mycroft-West et al. [89]	2020	In silico	Repurposing drug molecular docking molecular dynamic simulations	Heparin	Spike (S1) protein receptor-binding domain	Inhibition of viral infection arises from an overlap between the binding sites of heparin/HS on S1-RBD	Repurposing heparin and its derivatives as antiviral agents against SARS-CoV-2
Nayariseri et al. [90]	2020	In silico	Shape-based machine learning assisted by molecular docking and molecular dynamic simulations. ADMET studies	31 repurposed compounds	Main protease	Remdesivir, valrubicin, aprepitant, and fulvestrant	The novel compound nCorv-EMBS herein proposed stands as a promising inhibitor to be evaluated further for COVID-19 treatment
Odhar et al. [91]	2020	In silico	Molecular docking molecular dynamic simulations	1615 FDA-approved drugs	Main protease	Convaptan azelastine	
Ortega et al. [92]	2020	In silico	Repurposing drug molecular docking	Famotidine	Against SARS-CoV2	Famotidine could interact within the catalytic site of the three proteases associated with SARS-CoV2 replication	Weak binding affinity could be reached only upon intravenous administration
Pandey et al. [93]	2021	In silico	Repurposing drug molecular docking	9 flavonoids	Spike glycoprotein	Baicalin	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Parveen and Alhoman [94]	2021	In silico	Molecular docking molecular dynamic simulation density functional theory (DFT) ADME-Tox	FDA-approved anticancer drugs (capmatinib, pemigatinib, selpercatinib, and tucatinib)	Spike glycoprotein (S1) and the main protease	Potential of selected anticancer drugs for plausible drug development to fight COVID-19	Capmatinib, pemigatinib, selpercatinib, and tucatinib
Peele et al. [95]	2020	In silico	Molecular docking molecular dynamic simulations	USFDA-approved drugs, plant-derived natural drugs	Main protease	Lopinavir, amodiaquine, theaflavin digallate 22 candidates with putative SARS-CoV-2 Mpro inhibitory activity. Enalkiren, ethylsulfonamide-D-Trp-Gln-p-aminobenzamide, delparantag ritonavir and lopinavir, saquinavir	Beneficial polypharmacological effects
Pinzi et al. [96]	2021	In silico	Drug repurposing molecular docking molecular mechanic Poisson-Boltzmann surface area (MM-PBSA)	DrugBank database	Main protease		
Pokhrel et al. [97]	2020	In silico	Drug repurposing molecular dynamic simulations	US Food and Drug Administration (FDA)-approved drugs	RNA-dependent RNA polymerase	Quinupristin is particularly interesting because it is expected to bind across the RNA tunnel, blocking access from both sides Ethacrynic acid, naproxen, allopurinol, butenafine hydrochloride, raloxifene hydrochloride, tranlycypromine hydrochloride, saquinavir mesylate	Quinupristin represents a potential anti-SARS-CoV-2 therapeutic
Ray et al. [98]	2020	In silico	Drug repurposing intramolecularly quenched fluorescence (IQF) peptide substrate	774 FDA-approved drugs	Main protease		
Sachdeva et al. [99]	2020	In silico	Drug repurposing molecular docking	Antimalarial drugs	Spike protein and main protease	Doxycycline showed the most effective binding to the spike protein, whereas halofantrine and mefloquine bound effectively with the main protease	Doxycycline could potentially be a good candidate for repurposing for COVID-19
Sang et al. [100]	2020	In silico	Drug repurposing molecular docking molecular mechanics Poisson-Boltzmann surface area (MM-PBSA)	6 approved anti-HIV drugs	Main protease	Darunavir	
Saxena et al. [101]	2021	In silico/In vitro	Drug repurposing molecular docking	FDA-approved DrugBank database	Spike protein	Ertugliflozin possesses several desired properties	Good candidate for immediate repurposing for the treatment of COVID-19
Setianingsih et al. [102]	2020	In silico	Drug repurposing molecular docking, molecular dynamic simulations	160 potential drugs from therapeutic target database	13 protein targets (12 SARS-CoV-2 proteins and 1 human protein)	Suramin, the strongest binding affinity against 3 protein targets (spike protein, nucleocapsid protein, ACE2)	Suramin is the most potential to bind nucleocapsid and spike protein of SARS-CoV-2

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Shah et al. [103]	2020	In silico	Drug repurposing molecular docking	61 molecules that are already being used in clinics or under clinical scrutiny as antiviral agents	Against the SARS-CoV-2	37 molecules were found to interact with >2 protein structures of COVID-19. HIV protease inhibitors and RNA-dependent RNA polymerase inhibitors showed promising features of binding to COVID-19 enzyme	Methisazone, an inhibitor of protein synthesis; CGP42112A, an angiotensin AT2 receptor agonist; and ABT450, an inhibitor of the nonstructural protein 3-4A, might become convenient treatment option as well against COVID-19
Sharma and Mishra [104]	2020	In silico	Drug repurposing target-based virtual ligand screening	ZINC drug database and our own database of natural products	Against the SARS-CoV-2	Antivirus drugs (ribavirin, valganciclovir, and thymidine), antibacterial drugs (cefpiramide, sulfasalazine, phenethicillin, lymecycline, demeclocycline, doxycycline, oxytetracycline, and tigecycline), anti-asthmatic drugs (montelukast, fenoterol, and reproterol), and hepatoprotective drug silybin have antiviral activity. Natural hesperidin was targeting the binding between spike RBD and human ACE2	The natural products, such as flavonoids like neohesperidin, hesperidin, baicalin, kaempferol 3-O-rutinoside, and rutin from different sources, andrographolide, neoandrographolide, and 14-deoxy-11,12-didehydroandrographolide from <i>A. paniculata</i> , and a series of xanthones from the plants of <i>Sweetia</i> genus, with antiviral, antibacterial, and anti-inflammation activity could effectively interact with these targets of SARS-CoV-2
Shekhar et al. [105]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulations	2,625 FDA-approved small molecules	Spike (S) protein fusion peptide region	Chloramphenicol succinate, imipenem, and imidurea	
Singh et al. [106]	2021	In silico	Drug repurposing molecular docking molecular dynamic simulations	1749 FDA-approved drugs	NSP12, a RNA polymerase	5 compounds which include 3a (paritaprevir), 3d (glecaprevir), 3h (velpatasvir), 3j (remdesivir), and 3l (ribavirin) had the best binding affinity	
Sinha et al. [107]	2021	In silico	Drug repurposing systematic pharmacokinetics, drug-likeness, basicity predictions, virtual screening, and molecular dynamic analysis	Hydroxychloroquine (HCQ), chloroquine (CQ)	Spike protein	1-[1-(6-Chloroquinolin-4-yl)piperidin-4-yl]piperidin-3-ol and (1r,2R)-2-N-(7-chloroquinolin-4-yl)cyclohexane-1,2-diamine interact with the active site of the spike protein similar to HCQ and CQ, respectively, with augmented safety profile	

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Soni et al. [108]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	Rifampicin	Main protease	Rifampicin docking score was $-7.24 \text{ kcal}\cdot\text{mol}^{-1}$, and it can predict as a very good inhibitor of main protease	
Tariq et al. [109]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulations	15 antimalarial drugs (including chloroquine) and 2413 US Food and Drug Administration- approved drugs	Main protease spike (S) protein	Paromomycin with activity against two targets spike protein and protease domain	
Tatar et al. [110]	2021	In silico	Drug repurposing molecular docking molecular dynamic simulations Drug repurposing quantitative structure- activity relationship (QSAR) mode molecular docking molecular dynamic simulation MM- PBSA method	34 antiviral compounds	RNA-binding domain	Rapamycin, saracatinib, camostat, trametinib, and nafamostat were the top hit compounds	
Tejera et al. [111]	2020	In silico	activity relationship (QSAR) mode molecular docking molecular dynamic simulation MM- PBSA method	DrugBank database	Main protease	Levothyroxine, amobarbital, and ABP-700	
Terahı et al. [112]	2020	In silico	Drug repurposing molecular docking	7,173 clinically approved drug	Angiotensin- converting enzyme 2 (ACE2)	Lividomycin, burixafor, quisinostat, fluprolyline, pemetrexed, spirofylline, edotecarin, diniprofylline	
Trezza et al. [113]	2020	In silico	Drug repurposing docking simulations, with molecular dynamics (MD), supervised MD (SuMD), and steered MD (SMD) simulations	FDA-approved drugs	Spike glycoprotein	Simeprevir, lumacaftor	
Ugurel et al. [114]	2020	In silico	Drug repurposing structure-based drug design genome sequences were analyzed	FDA-approved drugs	Helicase (Nsp13)	Cangrelor, fludarabine, folic acid, and polydatin inhibit both the wild-type and mutant SARS- CoV-2 helicase	
Unni et al. [115]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulations	DrugBank and PubChem library	Spike protein (S protein)	Bisoxatin (DB09219)	A laxative drug

TABLE 1: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Vaishali et al. [116]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulation ADME properties	FDA-approved compounds	Nonstructural protein 9 (Nsp9) replicase and spike proteins	Conivaptan exhibited the highest binding energy and maximum stability of the Nsp9 replicase. Tegobuvir exhibited maximum stability along with the highest binding energy at the active site of the spike proteins	
Verma et al. [117]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulation MM-GBSA-based energy	FDA-approved drugs	Main protease	Top-ranked drugs including adefovir, lumefantrine, dipyridamole, dihydroergotamine, hexoprenaline, riboflavin (vitamin B2), and pantethine (vitamin B5)	
Wei et al. [118]	2020	In silico	Drug repurposing molecular docking molecular dynamic simulations	US Food and Drug Administration (FDA)-approved drugs from DrugBank and natural compounds from traditional Chinese medicine systems pharmacology (TCMSP)	Spike protein (S protein)	Digitoxin and bisindigotin in TCMSP had the highest docking scores Forsythiae fructus and Isatidis radix are components of Lianhua Qingwen, and raltegravir had relatively high binding scores	
Xu et al. [119]	2021	In silico	Drug repurposing molecular docking molecular dynamic simulations	FDA-approved drugs	Spike protein	Thymoquinone, a phytochemical compound obtained from the plant <i>Nigella sativa</i> , is a potential drug candidate	

TABLE 2: Summary of studies with in silico method that used natural products for novel drug discovery against COVID-19.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Abdelli et al. [120]	2021	In silico	Molecular docking	Isothymol, thymol, limonene, P-cymene, and γ -terpinene derived from the essential oil of the antiviral and antimicrobial plant Ammoides verticillata (Desf.) Briq.	Inhibition of ACE2 cellular receptor	Isothymol, a major component of this plant, gives the best docking scores, as good ACE2 inhibitor	
Abouelela et al. [121]	2021	In silico	Molecular docking, dynamic simulation, and binding free energy calculation	Aloe	Main protease and spike protein	132, 134, and 159 were the best scoring compounds against main protease, while compounds 115, 120, and 131 were the best scoring ones against spike glycoprotein. Compounds 120 and 131 were able to achieve significant stability and binding free energies during molecular dynamic simulation	
Adem et al. [122]	2020	In silico	Molecular docking	Medicinal plant-based bioactive compounds (80 flavonoid compounds)	Main protease	Hesperidin, rutin, diosmin, apiin, diacetylcurcumin, (E)-1-(2-hydroxy-4-methoxyphenyl)-3-[3-[(E)-3-(2-hydroxy-4-methoxyphenyl)-3-oxoprop-1-enyl]phenyl]prop-2-en-1-one, and beta'-[4-methoxy-1,3-phenylene]bis(2'-hydroxy-4',6'-dimethoxyacrylophenone) have been found as more effective on COVID-19 than nelfinavir	
Allam et al. [123]	2020	In silico	Molecular docking, 3D shape similarity study (rapid overlay chemical similarity-ROCS) to the clinically used drugs in COVID-19 patients	3'-Hydroxy-4'-methoxy-chroman-7-O- β -d-glucopyranoside 4, ferulic acid heptyl ester 1, naringenin 2, and 4,2',4'-trihydroxy-6'-methoxychalcone-4'-O- β -d-glucopyranoside 3, which were isolated from peach (Prunus persica (L.) Batsch) fruits	Main protease, spike protein	Naringenin 2 and 4,2',4'-trihydroxy-6'-methoxychalcone-4'-O- β -d-glucopyranoside 3 have a strong binding mode to a protease receptor and spike protein and also block the inflammatory storm	Recommendation of peach fruits in controlling and managing COVID-19 cases
Al-Sehemi et al. [124]	2020	In silico	Molecular docking	31000 natural compounds of the natural product activity and species source (NPASS) library	Spike glycoprotein	Castanospermine and karuquinone B were shown to be the best-in-class derivatives in silico able to target an essential structure of the virus and to act in the early stage of infection	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Attia et al. [125]	2021	In silico	Molecular docking	10 phenolic antiviral	Against SARS-CoV-2	Hesperidin showed the highest docking score	Hesperidin and its mediated ZnO nanoparticles are willing antiviral agents
Azim et al. [126]	2020	In silico	Virtual screening methods molecular docking	27 plant metabolites	Main protease proteins (MPP), Nsp9 RNA-binding protein, spike receptor-binding domain, spike ectodomain, and HR2 domain	Asiatic acid, avicularin, guajaverin, and withaferin showed a maximum binding affinity with all key proteins in terms of lowest global binding energy	
Bhowmik et al. [39]	2021	In silico	Repurposing drugs, docking, and molecular dynamic simulation	Orientin (phytochemical)	Inhibitor of SARS-CoV-2 spike and host cell receptor GRP78 binding	Binding of orientin in the overlapping residues of GRP78 binding region of SARS-CoV-2 spike model	As a promising precautionary or therapeutic measure for COVID-19
Çakır et al. [127]	2021	In silico	Molecular docking	Peptides derived from beta-lactoglobulin	Inhibit the host cell membrane receptors PLpro (papain-like protease), RdRp (RNA-dependent RNA polymerase), Mpro or 3cl protease, and spike protein	Ala-Leu-Pro-Met-His-Ile-Arg (ALMPHIR) and Ile-Pro-Ala-Val-Phe-Lys (IPAVFK) peptides	β -Lactoglobulin (BLG) is the major whey protein of cow and sheep's milk (~3 g/l)
Chatterjee et al. [128]	2021	In silico	Molecular docking	Hesperidin, kaempferol, quercetin, epigallocatechin		Hesperidin, kaempferol, quercetin, epigallocatechin	Lead to conclusive data for the treatment of polyphenols, flavonoids, and bioflavonoids against SARS-CoV-2
Chikhale et al. [129]	2020	In silico	Molecular docking, dynamics	Plant Withania somnifera (Indian ginseng)	NSP15 endoribonuclease and receptor-binding domain of prefusion spike protein	Withanoside X and quercetin glucoside from W. somnifera have favorable interactions at the binding site of selected proteins	Immunomodulatory, antioxidant, and anti-inflammatory roles
Chikhale et al. [130]	2021	In silico	Molecular docking, dynamics, and network pharmacology analysis	Saikosaponins	Adjuvant therapy in the treatment of COVID-19	Saikosaponins interact with the proteins CAT gene CAT (catalase) and checkpoint kinase 1 (CHEK1)	Possible improvement in immune response towards COVID-19
Chikhale et al. [131]	2020	In silico	Molecular docking, dynamics	Asparagus racemosus (Willd.)	NSP15 endoribonuclease and spike receptor-binding domain	Asparoside -C and Asparoside -F have good binding with target proteins	Asparagus racemosus holds promise as SARS-CoV-2 (S) and (N) protein inhibitor
Chowdhury [132]	2020	In silico	Molecular docking, dynamics	Tinospora cordifolia (Giloy)	Main protease	Berberine can regulate main protease protein's function	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Dahab et al. [133]	2020	In silico	Molecular docking	10 phenolic compounds of different classes (phenolic acids, flavonoids, and coumarins)	Main protease and RNA polymerase	The top 7 hits are flacourtacin [3], sagerinic acid [16], hordatine A [23], hordatine B [134], N-feruloyl tyramine dimer [135], bisavenanthramides B-5 [27], and vulnibactins [38] and have better binding scores than remdesivir, the native ligand in RNA polymerase target (PDB ID: 7bV2)	Hordatines are phenolic compounds present in barley and were found to exhibit the highest binding affinity to both protease and polymerase
Das et al. [136]	2020	In silico	Molecular docking	Flavonoid-based phytochemicals of <i>Calendula officinalis</i>	Main protease	Rutin, isorhamnetin-3-O- β -D, calendoflaside, narcissin, calendulaglycoside B, calenduloside, and calendoflavoside have better binding energy than the native ligand	Rutin and calendoflaside showed better stability, compactness, and flexibility
Debnath et al. [137]	2020	In silico	Sequential E-pharmacophore and structure-based virtual screening (VS)	113687 number of commercially available natural compounds	ADP-ribose phosphatase	6 potential inhibitors having good binding affinity towards active sites	Commercially available
Dev and Kaur [138]	2020	In silico	Molecular docking	Eucalyptus essential oil	Main protease	Jensenone may represent potential treatment potential to act as main protease inhibitor	
Duru et al. [139]	2021	In silico	Molecular docking	Oil of <i>Nigella sativa</i> seed	Replicase polypeptide 1a, RNA-binding protein of NSP9, ADP ribose phosphatase of NSP3, 3-chymotrypsin-like protease 3CLpro, and RNA-dependent RNA polymerase RdRp, and ACE2-angiotensin-converting enzyme from the <i>Homo sapiens</i>	The binding affinity of caryophyllene oxide was the highest on NSP9 and RdRp targets, while α -bergamotene gave the best binding affinity on RPIA target. The binding affinity of β -bisabolene on the ACE2 was almost the same as remdesivir	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
El-Demerdash et al. [140]	2021	In silico	Molecular dynamic simulations, molecular docking	15 guanidine alkaloids	Main protease (Mpro) (PDB ID: 6lu7), spike glycoprotein (PDB ID: 6VYB), nucleocapsid phosphoprotein (PDB ID: 6VYO), membrane glycoprotein (PDB ID: 6M17), and a nonstructural protein (nsp10) (PDB ID: 6W4H)	Crambesidin 786 [5] and crambescidin 826 had the highest binding affinities. The examined 15 alkaloids especially 5 and 13 showed promising docking, ADMET, toxicity, and MD results	
Elekofehinti et al. [141]	2020	In silico	Molecular docking studies, molecular dynamics, and ADME/Tox	50,000 natural compounds retrieved from IBS database	Papain-like protease	STOCK1N-69160 [(S)-2-((R)-4-((R)-2-amino-3-methylbutanamido)-3-(4-chlorophenyl) butanamido) propanoic acid hydrochloride] has been proposed as a novel inhibitor against COVID-19 PLpro	
El-Hawary et al. [142]	2021	In silico	Molecular docking (a combination of metabolomics and in silico approaches)	A. terreus, the endophytic fungus associated with soybean roots	Main protease	Aspergillide B1 and 3 α -hydroxy-3,5-dihydromonacolin L were found to be potent anti-COVID-19 drug candidates	
Emirik [143]	2020	In silico	Molecular docking, MM-GBSA-based predictions, and molecular dynamics	Turmeric contents	SARS-CoV-2 vital proteins	Turneric spice has the potential to inhibit the SARS-CoV-2 vital proteins and can be used a therapeutic or protective agent against SARS-CoV-2 via inhibiting key protein of the SARS-CoV-2. Compounds 4, 23, and 6 are the most prominent inhibitor for the main protease, the spike glycoprotein, and RNA polymerase of virus, respectively	
Fakhar et al. [144]	2020	In silico	Structure-based pharmacophore modeling, virtual screening-based PHASE screen score, molecular modeling	Anthocyanin derivatives	Main protease	6 best anthocyanin-derived natural compounds, which could be used as promising lead compounds against main protease SARS-CoV-2	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Falade et al. [145]	2021	In silico	Molecular docking	Saponins and tannins	Main protease	Ellagic acid, arjunic acid, theasapogenol B, and euscaphic acid as potential inhibitors of SARS-CoV-2 (Mpro) with better pharmacokinetics and bioavailability compared with remdesivir	
Fitriani et al. [146]	2020	In silico	Molecular docking	Phytochemical compounds (Moringa oleifera, Allium cepa, Cocos nucifera, Psidium guajava, and Eucalyptus globulus)	Main protease	Oleanolic acid in Allium cepa, α -tocotrienol in Cocos nucifera, asiatic acid in Psidium guajava, and culinoid in Eucalyptus globulus were the most recommended compound in each medicinal plant	Oleanolic acid in Allium cepa found as a potential inhibitor of COVID-19 Mpro
Gangadevi et al. [147]	2021	In silico	Molecular dynamic simulations, molecular docking	Library of natural compounds	Host ACE2 receptor with spike RBD domain of SARS-CoV-2	Kobophenol A, identified through docking studies, is the first compound that inhibits SARS-CoV-2 binding to cells through blocking S1-RBD to the host ACE2 receptor and thus may serve as a good lead compound against COVID-19	
Gangarapu et al. [148]	2020	In silico	Molecular docking online pkCSM and SwissADME Web server	Phytoconstituents of Siddha official formulation kabasura kudineer and novel herbal preparation—JACOM	Spike protein	37 compounds were screened, and of these, 9 compounds showed high binding affinity against spike protein	SNACK-V formulations could be used for effective treatment of COVID-19
Ghosh et al. [149]	2020	In silico	Molecular dynamic simulations, molecular docking	8 polyphenols from green tea	Main protease	3 polyphenols (epigallocatechin gallate, epicatechin gallate, and galocatechin-3-gallate) interact strongly with one or both catalytic residues (His41 and Cys145) of main protease	
Ghosh et al. [150]	2021	In silico	Molecular dynamic simulations, molecular docking, MM-GBSA analysis	Justicia adhatoda alkaloids	Main protease	1 alkaloid (anisotone) had interaction with both the catalytic residues (His41 and Cys145) of Mpro and exhibited good binding affinity (-7.9 kcal/mol)	More potent Mpro inhibitor than the two previously recommended antiviral drugs (lopinavir and darunavir)
Gorla et al. [151]	2020	In silico	Molecular docking	Essential flavonoids	SARS-CoV-2 spike glycoprotein receptor-binding domain (RBD-S) and host angiotensin-converting enzyme-2 protease domain (PD-ACE2)	Biochanin A and silymarin bind significantly at the active sites of RBD-S and PD-ACE2	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Gurung et al. [152]	2020	In silico	Virtual screening, molecular docking	Antiviral compounds from plants	Main protease	Bonducellin D was identified as the best lead molecule, which shows higher binding affinity 4 nontoxic, druggable plant-derived alkaloids (10-hydroxyusambarensine and cryptoquindoline) and terpenoids (6-oxoisoguesterin and 22-hydroxyhopan-3-one)	
Gyebi et al. [153]	2021	In silico	Molecular docking, ADME/Tox, and Lipinski filter analysis	African plants derived alkaloids and terpenoids	Main protease	Octatriacontyl pentafluoropropionate is well oriented inside the enzyme pockets, in addition to an excellent binding manner with the active site of the target macromolecules	Menoufia propolis could be a promising candidate in the combat against the pandemic COVID-19
Elwakil et al. [154]	2021	In silico	Gas chromatography/mass spectrometry analysis molecular docking	Egyptian propolis	RNA-dependent RNA polymerase, spike protein S1, and main protease	13 phytochemicals were studied, eight showed very strong binding affinities to main protease, and four showed moderate to strong binding affinities	
Hasan et al. [155]	2020	In silico	Molecular docking	Compounds present in the plant <i>Solanum surattense</i>	Main protease	6 main compounds possess high binding energy with the receptor active site of the main protease	
Hashem [156]	2020	In silico	Molecular docking	Honeybee and propolis	Main protease	9 potent natural products with binding affinities (ΔG binding) >-48.0 kcal/mol four bis([1, 3] dioxolo)pyran-5-carboxamide derivatives were identified as potential drug candidates	MolPort-004-849-765, MolPort-000-708-794, MolPort-002-513-915 and MolPort-000-702-646 are bis([1,3]dioxolo)pyran-5-carboxamide derivatives
Ibrahim et al. [157]	2020	In silico	Molecular dynamic simulations, molecular docking, MM-GBSA analysis	MolPort database that contains over 100,000 natural products	Main protease	High potency of salvianolic acid A and curcumin as main protease inhibitors	Salvianolic acid A as an in silico natural product inhibitor against the SARS-CoV-2 main protease
Ibrahim et al. [158]	2020	In silico	Molecular dynamic simulations, molecular docking	Metabolites present in several common spices	Main protease	Six compounds had good binding energies. CID_9910474 and CID_10503282 had a better stability when compared to other selected phytochemicals	
Isa et al. [159]	2020	In silico	Docking and molecular dynamic (MD) simulation	Extracts of <i>Zingiber officinale</i> and <i>Anacardium occidentale</i>	Main protease		

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Istifli et al. [160]	2020	In silico	Molecular dynamics and molecular mechanic Poisson-Boltzmann surface area (MM/PBSA) methods	23 phytochemicals belonging to different flavonoid subgroups	Spike glycoprotein cellular proteases (transmembrane serine protease 2 (TMPRSS2), cathepsin B and L (CatB/L)).	(-)-Epicatechin gallate interacted strongly with all the proteins studied	Epicatechin gallate can be evaluated as a candidate molecule in drug development studies against 2019-nCoV since it was not the substrate of P-gp (P-glycoprotein), did not inhibit any of the cytochrome Ps, and did not show AMES toxicity or hepatotoxicity on eukaryotic cells
Jan et al. [161]	2021	In silico	Cell-based infection assay molecular modeling	2,855 small molecules and 190 traditional herbal medicines	Main protease RNA-dependent RNA polymerase	Mefloquine, nelfinavir, and extracts of <i>Ganoderma lucidum</i> (RF3), <i>Perilla frutescens</i> , <i>Mentha haplocalyx</i> Baicalin, herbacetin, and pectolinarin have been discovered to block the proteolytic activity. Baicalin showed an effective inhibitory activity against main protease	
Jo et al. [162]	2020	In silico	Docking	Flavonoids	Main protease		
Joshi et al. [163]	2021	In silico	Docking	7100 molecules	Main protease	Several natural molecules such as δ -viniferin, myricitrin, taiwanhomoflavone A, lactucopicrin 15-oxalate, nympholide A, afzelin, biorobin, hesperidin, and phyllaemblicin B strongly binds to main protease	These molecules also showed strong binding with other potential targets of SARS-CoV-2 infection such as viral receptor human angiotensin-converting enzyme 2 (hACE2) and RNA-dependent RNA polymerase (RdRp)
Junior et al. [164]	2021	In silico	Docking and molecular dynamic (MD) simulation	Lapachol(1,4-naphthoquinone)	SARS-CoV-2 nonstructural proteins (nsps)	Lapachol derivatives VI and IX demonstrated the strongest binding	Lapachol derivatives are potential ligands for SARS-CoV-2 Nsp9
Kar et al. [165]	2020	In silico	Molecular docking molecular dynamic simulations and analysis of MM-PBSA energy	Indian plants including <i>Justicia adhatoda</i> , <i>Ocimum sanctum</i> , and <i>Swertia chirata</i>	Spike protein, main protease enzyme Mpro, and RNA-dependent RNA polymerase (RdRp)	Anisotone against SARS-CoV-2 spike and Mpro proteins and amarogentin against RdRp as potential inhibitors	
Khalifa et al. [166]	2020	In silico	Molecular docking modeling structural-relationship activity	10 anthocyanins	Main protease	Phacelianin, gentiodelphin, cyanodelphin, tecophilin	Leading molecules for further optimization and drug development process to combat COVID-19
Khalifa et al. [167]	2020	In silico	Molecular operating environment molecular docking	19 structural different hydrolysable tannins	Main protease	Pedunculagin, tercatatin, and castalin	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Khan et al. [168]	2020	In silico	Molecular docking	Marine natural products	Main protease	C-1 (PubChem CID 11170714) exhibited good activity	
Kiran Raj et al. [169]	2020	In silico	Molecular docking	C-Phycocyanin of <i>Spirulina platensis</i>	Nonstructural proteins 12	C-Phycocyanin inhibits the active site of nsp12	
Krupanidhi et al. [170]	2020	In silico	Molecular docking molecular dynamic simulation ADME along with toxicity analysis	Phytochemical constituents of <i>Tinospora cordifolia</i>	Main protease	Tinosponone	
Kumar et al. [171]	2020	In silico	Molecular docking molecular dynamic simulations and analysis of MM-PBSA energy	Novel natural metabolites	Main protease	Ursolic acid, carvacrol, and oleanolic acid could	
Kumar et al. [172]	2021	In silico	Molecular docking, ADMET, and molecular dynamic simulations	Phytoconstituents from natural herbs	Main protease	Laurolitine	Laurolitine, an active constituent of roots of <i>Lindera aggregata</i>
Kumar et al. [173]	2021	In silico	Molecular docking molecular dynamic simulations and analysis of MM-PBSA energy	Strychnos nux-vomica	Main protease	Demethoxyguiaflavine, strychnoflavine	
Li et al. [174]	2021	In silico	Network pharmacology and in vitro experiment verification molecular docking	Maxing Shigan decoction (MXSGD)	ACE2, Mpro, and RdRp	The components with strong potential affinity (top 10) with ACE2, Mpro, and RdRp are mainly from <i>Glycyrrhiza uralensis</i> (Chinese name: Ganciao) and <i>Semen armeniacae</i> amarum (Chinese name: Kuxingren). Among them, amygdalin was selected as the optimal candidate component binds to all three key targets, and eugenone, glycyrrhizin, and glycyrol also exhibited superior affinity interactions with ACE2, Mpro, and RdRp, respectively	Multicomponent, multitarget, and multi-approach intervention
Maiti and Banerjee [175]	2021	In silico	Bioinformatic molecular docking	Tea flavonoid catechin products	Spike glycoproteins	Epigallocatechin gallate and theaflavin gallate interact better than hydroxychloroquine	
Mahmud et al. [176]	2021	In silico	Molecular docking molecular dynamic simulation MM-GBSA scores	3063 compounds from more than 200 plants from the Asian region	Main protease	Curcumin, gartanin, robinetin	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Mahmud et al. [177]	2020	In silico	Molecular docking molecular dynamic simulation MM-GBSA scores	Plant-derived natural compounds	Main protease	Carinol, albanin, myricetin	
Mesli et al. [178]	2021	In silico	Molecular docking molecular dynamic simulations	Leaves of <i>Corchorus olitorius</i> Linn.	Angiotensin-converting enzyme 2	Méthyl-1,4,5-tri-O-caféoyl quinate has a stronger bond, high affinity, and gives the best docking scores compared with the co-crystallized inhibitor (PRD_002214) of the enzyme ACE2	
Mohammadi et al. [179]	2020	In silico	Molecular docking	Marine polypeptides were isolated from the hydrolysate of Pacific oyster	Main protease	The score of Leu-Leu-Glu-Tyr-Ser-Ileu ligand was higher than remdesivir	Pacific oyster compounds may have the potency to be evolved as an anti-COVID-19 main protease
Murugan et al. [180]	2020	In silico	Molecular docking molecular dynamic simulation MM-GBSA scores	Andrographis paniculata phytochemicals	3 nonstructural proteins (3 L main protease (3CLpro), papain-like proteinase (PLpro) and RNA-directed RNA polymerase (RdRp)), and a structural protein (spike protein (S))	Neoandrographolide (AGP3) has shown promising binding affinity towards all the four targets	
Naik et al. [181]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	Natural product activity and species source (NPASS) database	Endoribonuclease exoribonuclease RNA-dependent RNA polymerase (RdRp) methyltransferase and main protease	21 compounds showed maximum docking scores NPC214620, NPC52382, and NPC270578 are targeting five, four, and three-drug targets, respectively	Multitarget-based drug design
Narkhede et al. [182]	2020	In silico	Molecular docking molecular dynamic simulations	Natural products	Main protease	Glycyrrhizin, bicyclogermacrene, tryptanthrin, β -sitosterol, indirubin, indican, indigo, hesperetin, chrysophanic acid, rhein, berberine, and β -caryophyllene	Interactions with the COVID-19 main protease were highest in the case of glycyrrhizin and rhein
Nivetha et al. [183]	2020	In silico	Molecular docking molecular dynamic simulation MM-PBSA	Seselin purified from the leaf extracts of <i>Aegle marmelos</i>	Spike protein S2, main protease, and free enzyme of the SARS-CoV-2	Seselin had inhibitory potential over multiple SARS-CoV-2 targets	
Ogunyemi et al. [184]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	226 bioactive compounds from African herbs and medicinal plants	RNA-dependent RNA polymerase	Drugable alkaloids (10' - hydroxyusambarensine, cryptospirolepine, strychnopentamine) and flavonoids (usarotenoid A and 12 α -epi-milletosin)	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Padhi et al. [185]	2021	In silico	Molecular docking ADME properties	415 natural metabolites isolated from several plants, fungi, and bacteria	Spike glycoprotein (S1) and the main protease	Fusaric acid, jasmonic acid, jasmonic acid methyl ester, putaminoxin, putaminoxins B and D, and stagonolide K were predicted to have considerable (ADME) and safety indices	Jasmonic acid and putaminoxins B and D bind best to main protease
Pandey et al. [186]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	10 potential naturally occurring compounds (flavonoids/non-flavonoids)	Spike glycoprotein	Fisetin, quercetin, and kaempferol consist of drug-likeness property	
Kumar et al. [187]	2020	In silico	Molecular docking	Kabasura kudineer and thonthasura kudineer are two Siddha formulations	Spike glycoprotein	Cucurbitacin B (−112.09), cardiofolioside (−111.5), apigenin (−98.84), and pyrethrin (−92.98) were observed as more effective with less bind energies	Kabasura kudineer could be a potential Siddha medicine for COVID-19
Rahman et al. [188]	2021	In silico	Molecular docking ADMET properties	Rutin	Main protease (Mpro), RNA-dependent RNA polymerase (RdRp), papain-like protease (PLpro), and spike (S) protein	Significant binding of rutin with Mpro, RdRp, PLpro, and S proteins. Main protease exhibited the strongest binding affinity	Optimal solubility, nontoxic, and noncarcinogenic properties
Rahman et al. [189]	2020	In silico	Molecular operating environment (MOE) ligand-based pharmacophore approach and a molecular docking-based screening	Natural product activity and species source (NPASS)	Type II transmembrane serine protease (TMPRSS2)	85 compounds with molecular docking comparable to or greater than that of the standard inhibitor (camostat mesylate) were identified. The top 12 compounds with the most favorable structural features were studied. The low-molecular-weight compound NPC306344 showed significant interaction with the active site residues of TMPRSS2	
Rakib et al. [190]	2020	In silico	Molecular docking	Bioactive phytochemicals isolated from <i>Tinospora crispa</i>	Main protease	The top nine hits might serve as potential anti-SARS-CoV-2 lead molecules, with three of them exerting biological activity	
Ramadhan et al. [191]	2020	In silico	Molecular docking	<i>Erlingera elatior</i> plant	Main protease	Ergosterol peroxide sitostenone	
Rangsinth et al. [192]	2021	In silico	Molecular docking ADMET properties	Natural products isolated from edible and medicinal mushrooms	Main protease	6 of 25 compounds are the best drug-like property candidates, including colossolactone VIII, colossolactone E, colossolactone G, ergosterol, helianthol F, and velutin	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Rivero-Segura et al. [193]	2021	In silico	Molecular docking molecular dynamic simulation ADME properties	Mexican natural products	Against the SARS-CoV-2	Quercetin, riolozatrione, and cichoriin target the key proteins of SARS-CoV-2	Cichoriin reaches higher lung levels (100 mg/kg, IV); therefore, it may be considered in developing therapeutic tools
Selvaraj et al. [194]	2020	In silico	Homology modeling and molecular dynamic (MD) simulation MM/GBSA, MD simulations, and PCA calculations	Traditional Chinese medicine (TCM) database	Nsp 14 guanine-N7 methyltransferase (N7- MTase)	TCM 57025, TCM 3495, TCM 5376, TCM 20111, and TCM 31007 are the compounds from the TCM database, which can occupy and interact nicely with the substrate-binding site of N7- MTase	
Sharma and Kaur [195]	2021	In silico	Molecular docking, protein interaction calculator ADME studies	12 bioactive molecules present in essential oils of eucalyptus plant leaves	Spike (S) protein	Toruatone	
Sharma [196]	2020	In silico	Molecular docking, protein interaction calculator	Eucalyptol (1,8 cineole), an essential oil component from eucalyptus oil	Main protease	Eucalyptol may represent potential treatment potential to act as main protease inhibitor	Effective binding of eucalyptol to COVID-19 protease
Sharma and Kaur [197]	2020	In silico	Molecular docking, protein interaction calculator	Jensenone, an essential oil component from eucalyptus oil	Main protease	Jensenone may represent potential treatment potential to act as main protease inhibitor	
Shawan et al. [198]	2021	In silico	Pharmacophore study molecular docking molecular dynamic simulation ADME properties	43 flavonoids of 7 different classes	Against the SARS-CoV-2	Luteolin and abyssinone II were found to develop successfully docked complex within the binding sites of target proteins	
Sindhu et al. [199]	2020	In silico	Molecular docking	Clerodendrum paniculatum leaves	Main protease	Clerodol	
Singh et al. [200]	2021	In silico	Molecular docking and structural dynamic studies	Tea (Camellia sinensis) polyphenols	Nonstructural protein 16 (NSP16)	Theaflavin compound demonstrated lower binding free energy in comparison with the standard molecule sinefungin EGCG, theaflavin (TF1), theaflavin-3'-O-gallate (TF2a), theaflavin-3'-gallate (TF2b), theaflavin 3,3'-digallate (TF3), hesperidin, quercetagenin, and myricetin strongly bind to the active site of RdRp	EGCG, TF2a, TF2b, and TF3 can inhibit RdRp and represent an effective therapy for COVID-19
Singh et al. [201]	2020	In silico	Molecular docking and structural dynamic studies molecular mechanics Poisson-Boltzmann surface area (MM-PBSA) ADME properties	Polyphenols	RNA-dependent RNA polymerase (RdRp)		

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Srimathi et al. [202]	2020	In silico	Molecular docking	Traditional herbal medicine: apo-quinine, catechin, cinchonidine, cinchonine, cupreidine, epicatechin, epiprocumumenol, epiquinine, procumumenol, quinidine, quinine, zedoaronediol, procumumadiol	Against the SARS-CoV-2	Epicatechin, apo-quine	
Subbaiyan et al. [203]	2020	In silico	Molecular docking	Active constituents present in common herbs	Spike (S) protein	Epigallocatechin gallate (EGCG) was found to have the highest binding affinity with the viral S protein, followed by compounds, "F" (curcumin), "D" (apigenin), and "E" (chrysophanol)	Green tea
Surti et al. [204]	2020	In silico	Molecular docking molecular dynamic simulations	Ilimaquinone (marine sponge metabolite)	Spike receptor-binding domain, RNA-dependent RNA polymerase, Nsp10, Nsp13, Nsp14, Nsp15, Nsp16, main protease, and papain-like protease	Ilimaquinone exhibited promising inhibitory potential against all the SARS-CoV-2 target proteins, as evident from the binding energies	Most promising inhibitory candidate against the SARS-CoV-2 papain-like protease
Tao et al. [205]	2020	In silico	Network pharmacology and molecular docking.	Huashi Baidu formula (HSBDF): Chinese	Against the SARS-CoV-2	Baicalcin and quercetin were the top two compounds of HSBDF, which had high affinity with ACE2	Regulate multiple signaling pathways through ACE2
Umar et al. [206]	2021	In silico	Molecular docking molecular dynamic simulation ADME properties	Azadirachta indica, Mangifera indica, and Moringa oleifera: African plants	Main protease	Most of the active phytochemicals of the study plants exhibited relative inhibitory potentials against main protease and preferred pharmacological features when compared with hydroxychloroquine	Caffeic acid, chlorogenic acid, catechin, ellagic acid, gallic acid, etc.
Umesh et al. [207]	2021	In silico	Molecular docking molecular dynamic simulation ADME properties	Chemical compounds from Indian spices	Main protease	Carnosol exhibited the highest binding affinity for arjunglucoside-I and rosmanol showed a strong and stable binding affinity with favorable ADME properties	

TABLE 2: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Yang et al. [208]	In silico	2020	High-throughput virtual screening	Natural Products Research Laboratories (NPRL)	Main protease	Curcuminoid derivatives (including NPRL334, NPRL339, NPRL342, NPRL346, NPRL407, NPRL415, NPRL420, NPRL472, and NPRL473) display strong binding affinity to COVID-19 3Lpro polypeptide	NPRL-334 revealed the strongest binding affinity
Yu et al. [209]	2020	In silico	Metascape analysis protein docking molecular docking	Mongolian medicine	SARS-CoV-2 S protein RBD domain	253 active components were predicted. Phillyrin and chlorogenic acid can effectively prevent the combination of SARS-CoV-2 S protein and ACE2 at the molecular level	
Zhang et al. [210]	2020	In silico	Molecular docking molecular dynamic simulation ADME property network pharmacology analysis	Chinese herbal medicines	Anti-2019-nCoV activity	13 compounds that exist in traditional Chinese medicines were found to have potential anti-2019-nCoV activity. 125 Chinese herbs were found to contain 2 or more of these 13 compounds. Of these 125 herbs, 26 are classically cataloged as treating viral respiratory infections	Regulating viral infection, immune/inflammation reactions, and hypoxia response

TABLE 3: Summary of other studies with in silico method.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Abu-Melha et al. [211]	2020	In silico	Molecular docking combined with molecular dynamic simulation (MDS)	Hydrazones, pyrazoles, and pyrazines bearing thiazole moiety	Main protease	The average binding affinities of the compounds 3a, 3b, and 3c (-8.1 ± 0.33 kcal/mol, -8.0 ± 0.35 kcal/mol, and -8.2 ± 0.21 kcal/mol, respectively) are better than that of the positive control nelfinavir	
Aghaee et al. [212]	2021	In silico	Pharmacophore model molecular docking combined with molecular dynamic simulation (MDS), MM/PBSA, ADME studies	Pharmit website	Main protease	ML188, nelfinavir, lopinavir, ritonavir, and α -ketoamide	
Ahmad et al. [213]	2021	In silico	Structure-based virtual screening (SBVS) of ASINEX antiviral library, molecular dynamic (MD) simulations	ASINEX antiviral library	Main protease	SCHEMBL12616233, SCHEMBL18616095, and SCHEMBL20148701 compounds conformation with main protease show good stability after initial within active cavity moves, a rich intermolecular network of chemical interactions, and reliable relative and absolute binding free energies Raltegravir, simeprevir, cobicistat, and daclatasvir have higher binding energy and strong interaction with active sites of the receptor proteins (with a precision of 85%)	BBB_26580140 lead and its similar analogs to be explored in vivo lead molecules
Ahmed et al. [214]	2020	In silico	Molecular docking, molecular dynamics, and structure-activity relationship	76 prescription antiviral drugs	RNA-dependent RNA polymerase (RdRp) and main protease (Mpro)		

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Alabboud and Javadmanesh [215]	2020	In silico	Molecular docking combined with molecular dynamic simulation	88 conventional drugs, 16 vitamins, and 63 natural (plant)	Main protease	Various vitamins (B9, A, K, and E vitamins) exhibited a significantly strong interaction with the studied receptor. Pleconaril, adefovir dipivoxil, and stavudine in addition to plant-based compounds such as curcumin (<i>Curcuma longa</i>), anolignan A (<i>Anogeissus acuminata</i>), and phyllamycin B (<i>Phyllanthus myrtifolius</i>) had strong ligand-protein interactions ADM_13083841, LMG_15521745, and SYN_15517940 showed stable conformation and interacted well with the active residues of papain-like protease	
Alamri et al. [216]	2020	In silico	Structure-based virtual screening coupled with all-atom molecular dynamic (MD) simulations	Protease inhibitors database composed of ~7,000 compounds	Papain-like protease	Elvitegravir and oxolinic acid are able to interact with the NTP entry channel and thus interfere with the RNA-directed 5'-3' polymerase activity of SARS-CoV-2 RdRp. Rilapladib is the only quinoline that can interrupt the spike-RBD-ACE2 complex	Quinoline, 1,2,3,4-tetrahydro-1-[(2-phenylcyclopropyl)sulfonyl]-trans-(8Cl), saquinavir, elvitegravir, oxolinic acid, and rilapladib are suggested for the treatment of COVID-19
Alexpandi et al. [217]	2020	In silico	Molecular docking	113 quinoline drugs	Main protease, RNA-dependent RNA polymerase (RdRp) inhibitors spike-RBD-ACE2 inhibitor		
Al-Sehemi et al. [218]	2020	In silico	Molecular docking, MD simulation, and molecular mechanics Poisson-Boltzmann surface area (MM-PBSA) results	Phenyl furoxan, an exogenous nitric oxide donor	Main protease	Spiro-isoquinolino-piperidine-furoxan moieties can be used as an effective ligand for main protease inhibition due to the presence of key isoquinolino-piperidine skeleton with additional NO effect	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Al-Shar'i [219]	2020	In silico	Molecular docking, MD simulation, and molecular mechanics Poisson-Boltzmann surface area (MM-PBSA) results	Different databases	Main protease	9 compounds with different chemotypes	
Badavath et al. [220]	2020	In silico/in vitro	Molecular docking, molecular dynamics, and structure-activity relationship studies	Screening of 118 compounds with 16 distinct heterocyclic moieties in comparison with 5 natural products and 7 repurposed drugs	Main protease	Oxidiazoles (A2 and A4) derivatives have the best docking scores. Structure-activity relationship studies showed a good comparison with a known active main protease and repurposed drug ebelsen with an IC ₅₀ value of $-0.67 \mu\text{M}$	
Barros et al. [221]	2020	In silico	Molecular docking	24 ligands	SARS-CoV-2 receptors, Nsp9 replicase, main protease (Mpro), NSP15 endoribonuclease, and spike protein (S protein) interacting with human ACE2	Antimalarial drug mefloquine and anti-HIV antiretroviral saquinavir interacted with all the studied receptors	
Basit et al. [222]	2020	In silico	Protein-protein docking and molecular dynamic simulation Pharmacophore-based screening, docking consensus approach (DCA), molecular dynamic simulations, common hit approach (CHA)	Truncated version of human ACE2 (tACE2)	S Glycoprotein	tACE2 provides a high-affinity protein inhibitor for S glycoprotein	
Battisti et al. [223]	2020	In silico		Aldrich Market Select (AMS) database from ChemNavigator/Sigma-Aldrich with over 8 million unique chemical structures	Against SARS-CoV-2	10 compounds with high coronavirus inhibition potential	Lead molecules
Benítez-Cardoza and Vique-Sánchez [224]	2020	In silico	Molecular docking	500,000 compounds	Potential inhibitors of the interaction between ACE2 and SARS-CoV-2 (RBD)	20 compounds were determined by docking focused on the region of interaction between ACE2 and RBD	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Cava et al. [225]	2020	In silico	Gene Ontology and enrichment analysis protein-protein interaction (PPI) network virtual screening method	—	Against SARS-CoV-2	A protein-protein interaction network of 193 genes, 22 interactions, and 36 potential drugs for future treatment strategies including nimesulide, fluticasone propionate, thiabendazole, photofrin, and didanosine	Only didanosine is a real antiviral drug, while the others are mostly anti-inflammatory
Chen et al. [226]	2020	In silico	Crystal structure, virtual screening	7173 purchasable drugs (drugs-lib), with 4574 unique compounds and their stereoisomers	Main protease	16 candidates for consideration, ledipasvir velpatasvir	
Choudhary et al. [227]	2020	In silico	Molecular docking, MM-GBSA predictive binding energy calculations, and molecular dynamic simulation	15,754 natural and synthetic compounds	Main protease	Compound 2 (molecular bank code AAA396) and compound 3 (molecular bank code AAD146)	
Chunduru et al. [228]	2021	In silico	Molecular docking	Novel drug-like inhibitors for COVID-19	Main protease	Structure 61 was found to be more stable and can be further assessed for their antiviral activity to combat COVID-19	
Coelho et al. [229]	2020	In silico	Biochemical high-throughput screening	Compound library containing known drugs, bioactive molecules, and natural products	Main protease	Organomeric compounds thimerosal and phenylmercuric acetate, benzophenone, Evans blue, a sulfonic acid-containing dye	
Dai et al. [230]	2020	In silico	Structure-based design	—	Main protease	Designed and synthesized two lead compounds (11a and 11b) targeting main protease. Both exhibited excellent inhibitory activity	
de Lima Menezes and da Silva [231]	2020	In silico	Molecular dynamic simulations, molecular docking	DrugBank database	Nonstructural protein 1 (nsp1)	Tirilazad, phthalocyanine, and Zk-806450 showed better energy score than control molecules that have in vitro activity against nsp1 from SARS-CoV-2	Tirilazad, phthalocyanine, and Zk-806450

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Debnath et al. [232]	2020	In silico	Pharmacophores studies, structure-based virtual screening, molecular dynamic (MD) simulation	Drug molecule information retrieved from DrugBank	Main protease	DB07456 and DB13592 displayed a similar type of binding interaction with co-ligands and remdesivir, and the predicted Ki values of 2 inhibitors were found to be superior to remdesivir	
Di Micco et al. [233]	2021	In silico	Molecular docking, MM-GBSA-based predictions, and molecular dynamics	Zonulin inhibitor larazotide acetate (also called AT1001)	Main protease	AT1001, besides its well-demonstrated effect in ameliorating mucosal permeability in ALI/ARDS Compound 11 as the best potential nsp16 inhibitor herein identified, as it displayed a better stability and average binding free energy for the ligand-enzyme complex compared to sinefungin	
El Hassab et al. [234]	2021	In silico	Structure-based virtual screening, molecular dynamic simulation, and MM-PBSA approaches	48 million drug-like compounds of the ZINC database	SARS-CoV-2 2'-O-methyltransferase (nsp16)	The prominent drug-like and potent inhibitory compounds are 2-[2-(2-aminoacetyl)amino-3-(4-hydroxyphenyl)propanamide (ZINC000004762511), 6'-fluoroaristeromycin (ZINC000001483267), and cyclo(L-histidyl-L-histidyl) (ZINC000005116916) scaffolds	
Elmessaudi-Idrissi et al. [235]	2020	In silico	In silico screening, molecular docking, and dynamic approaches	5000 compounds of the ZINC database	Main protease	Melatonin can have response potential in early stages for its possible effects on ACE2 and main protease, although it is also promising in more severe stages of the disease for its action against hyper-inflammation	Do not confirm antiviral activity, but can rather be used as a basis for further preclinical and clinical trials
Feitosa et al. [236]	2020	In silico	Molecular docking	—	Main protease		

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Gurung et al. [237]	2021	In silico	Virtual screening, molecular docking	1,36,191 molecules	Spike (S) protein receptor-binding domain (RBD) to the host cell surface receptor, angiotensin-converting enzyme 2 (ACE2)	ZINC33039472 exhibited binding free energy value lower as compared to the control (emodin) with a higher contribution by gas-phase energy and van der Waals energy to the total binding free energy ~200 compounds were identified as having strong interaction with Mpro (ZINC20291569, ZINC90403206, ZINC95480156) that showed the highest binding energy	
Haider et al. [238]	2020	In silico	Computer-aided drug design (CADD) molecular docking	ZINCPharmer	Main protease	Zanamivir, indinavir, saquinavir, and remdesivir are among the exciting hits of main protease	Flavin adenine dinucleotide (FAD) deficiency medicine, and coenzyme A, a coenzyme may also be potentially used for the treatment of SARS-CoV-2 infections
Hall and Ji [239]	2020	In silico	Homology modeling molecular docking	ZINC15 database: 3447 entries	Spike glycoprotein and main protease	Phenyltriazolinones (PubChem ID: 104161460) and allosteric HCV NS5B polymerase thumb pocket 2 (PubChem ID: 163632044) have shown antiviral activity and also have high affinity towards the main protease	
Havranek and Islam [240]	2020	In silico	Docking and molecular dynamics	2692 protease inhibitor compounds	Main protease	Promising binding affinities of TMC-310911 and ritonavir	
Ibrahim et al. [241]	2021	In silico	Molecular dynamic simulations, molecular docking, MM-GBSA	18 anti-COVID-19 drug candidates against SARS-CoV-2 main protease	Main protease	The proposed inhibitor ΔABP-D25Y	
Jaiswal and Kumar [242]	2020	In silico	Docking studies and molecular dynamic simulation	Designed a protein inhibitor	Spike (S) glycoprotein		

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Jamalan et al. [243]	2021	In silico	Docking and molecular dynamic (MD) simulation	Virtual screening based on GRL-0617	Papain-like proteinase	5-(aminomethyl)-2-methyl-N-[(1R)-1-naphthalen-1-ylethyl]benzamide outperformed GRL-0617 in terms of binding affinity (-9.7 kcal/mol). 2-(4-fluorobenzyl)-5-nitro-1H-isoindole-1,3(2H)-dione, 3-nitro-N-[(1r)-1-phenylethyl]-5-(trifluoromethyl)benzamide, 5-acetamido-2-methyl-N-[(1S)-1-naphthalen-1-ylethyl]benzamide	
Jin et al. [244]	2020	In silico	Structure-assisted drug design, virtual drug screening, and high-throughput screening	10,000 compounds	Main protease	Ebselen, disulfiram, tideglusib, carmofur, shikonin, PX-12	Ebselen also exhibited promising antiviral activity in cell-based assays
Kanhed et al. [245]	2021	In silico	Systematic virtual screening approach	ASINEX BioDesign library approved drug library	Main protease	Ritonavir, nelfinavir, and saquinavir were predicted to be the most potent Mpro inhibitors. 20 molecules (pyrazoles, cyclic amides, pyrrolidine-based compounds, and miscellaneous derivatives)	
Kavitha et al. [246]	2020	In silico	Molecular docking molecular dynamic simulations	1000 protease-inhibitor-like compounds available in the ZINC database	Main protease	1,2,4 triazolo[1,5-a]pyrimidin-7-ones	
Krishnan et al. [247]	2020	In silico	Molecular docking	3978 compounds with potential antiviral activity	Endoribonuclease (NSP15)	8 compounds with good docking score and docking energy e.g., Z595015370, Z1343129850, and Z2760938911	
Kumar et al. [248]	2021	In silico	Molecular docking molecular dynamic simulation molecular mechanic Poisson-Boltzmann surface area approaches	Million molecules and natural compound databases	Main protease	Three compounds namely ZINC14732869, ZINC19774413, and ZINC19774479 displayed better binding affinities	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Kumar et al. [249]	2020	In silico	Molecular docking molecular dynamic simulations	13 approved antiviral drugs	Main protease	Indinavir was described as a lead drug. Indinavir possesses an important pharmacophore	Novel compound 16(hydroxyethylamine derivative) suitability as a strong candidate for therapeutic discovery against COVID-19
Kwarteng et al. [250]	2020	In silico	Bioinformatic approach molecular docking molecular dynamic simulations	—	Nucleocapsid (N) protein	Zidovudine triphosphate, an anti-HIV agent, as a potential inhibitor of the N-terminal domain of SARS-CoV2 N protein	
Li et al. [251]	2021	In silico	Molecular docking	21 antiviral, antifungal, and anticancer compounds	Papain-like protease	Neobavaisoflavone	
Mathpal et al. [252]	2020	In silico	Molecular docking molecular dynamic simulation MM-PBSA	3180 FDA-approved drugs from “the ZINC database”	Main protease	ZINC03831201, ZINC08101052, ZINC01482077, and ZINC03830817	
Maurya et al. [253]	2020	In silico	Molecular docking	Antiviral, anti-infectious, and anti-protease compounds	NSP10/NSP16 methyltransferase and main protease	Cyclocytidine hydrochloride, trifluridine adonitol, and meropenem penciclovir bound with a good docking score NSP10/NSP16 methyltransferase complexed with telbivudine, oxytetracycline dihydrate, methyl gallate, 2-deoxyglucose, and daphnetin Romidepsin and its active form (RedFK)	
Mohamed et al. [254]	2020	In silico	Molecular docking	12 histone deacetylases (HDACs)	Main protease		
Monajemi and Zain [255]	2021	In silico	ONIOM (own N-layered integrated molecular orbital and molecular mechanics; QM/MM) approach	N3, ebselen, disulfiram, tideglusib, carmofur, shikonin, and PX-12	Main protease		Better inhibition from PX-12 than ebselen
Motiware et al. [256]	2020	In silico	Molecular docking molecular dynamic simulations	Previously reported SARS-3CL protease inhibitors	Main protease	N3, ebselen, and PX-12 inhibitors	
Mutlu et al. [257]	2020	In silico	Structure-based approach molecular docking molecular dynamic simulations	FDA-approved and investigational drugs	Nsp12/Nsp8	N-substituted isatin derivatives and pyrazolone	Two drugs, RX-3117 (fluorocyclopentenyl cytosine) and nebulol

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Naidoo et al. [258]	2020	In silico	Molecular docking molecular dynamic simulation MM-PBSA	Cyanobacterial metabolites	Main protease (Mpro) and the papain-like protease (PLpro)	Deoxycyclin, drospormopsin	Cylindrospermopsin, deoxycylindrospermopsin, carrageenan, cryptophycin 52, eucapsitrione, tipranazole, tolyporphin, and apratoxin A exhibited promising inhibitory potential against the SARS-CoV- 2 Mpro. The compounds cryptophycin 1, cryptophycin 52, and deoxycylindrospermopsin were observed to display encouraging binding energy scores with the PLpro of SARS- CoV-2
Olubiyi et al. [259]	2020	In silico	Molecular docking molecular dynamic simulations	Approved drugs, investigational drugs, natural products, and organic compounds	Main protease	Several tyrosine kinase inhibitors, which include a bioflavonoid and steroid hormones, bind best to main protease	Nilotinib, enasidenib, afatinib, ertapenem, phthalocyanine, hypericin, amrubicin, theacitrin A, theaflavin, amentoflavone, epigallocatechin gallate, glabrolide, cortisol, estradiol, testosterone
Özdemir et al. [260]	2020	In silico	Molecular docking density functional theory (DFT) ADME-Tox	42 coumarin derivatives	Main protease	6,7-Dihydroxy-3- phenylcoumarin derivatives gave relatively higher scores, and for all coumarins, and 4- trifluoromethylphenyl substituted coumarin had the highest score The highest score (-10.01 kcal/mol) in the coumarin group is 2- morpholinoethan-1-amine substituted coumarin	
Ozdemir et al. [261]	2020	In silico	Molecular docking molecular mechanics Poisson-Boltzmann surface area (MM-PBSA)	Coumarin derivatives	Spike S1 subunit, NSP5, NSP12, NSP15, and NSP16	Ritonavir, nelfinavir, and saquinavir were predicted to be the most potent Mpro inhibitors 20 molecules categorized into four classes viz. disubstituted pyrazoles, cyclic amides, pyrrolidine- based compounds, and miscellaneous derivatives	
Patel et al. [262]	2021	In silico	Pharmacophore studies	Drug library (having drugs and diagnostic agents, which are approved by FDA or other world authorities) and the ASINEX BioDesign library	Main protease		

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Peng et al. [263]	2020	In silico	Drug repositioning through virus-drug association prediction	—	Against SARS-CoV-2	Ribavirin was predicted to be the best small molecular drug, with a higher molecular binding energy with human angiotensin-converting enzyme 2 (ACE2), followed by remdesivir, mycophenolic acid, and chloroquine (−6.29 kcal/mol) Three 5-O-benzoylpinostrobin derivatives each with fluoro, tertiary butyl, and trifluoromethyl substituents at 4-position of benzoyl group showed the lowest free energy of binding value and the highest similarity of ligand-receptor interactions with co-crystallized ligands	
Pratama et al. [264]	2020	In silico	Molecular docking	Novel 5-O-benzoylpinostrobin derivatives	Main protease	Two compounds: PubChem3408741 and PubChem4167619 had the binding free energy of −94.02 kJ·mol ^{−1} and −122.75 kJ·mol ^{−1} , respectively, as compared to reference X77 (−76.48 kJ·mol ^{−1})	Lead molecules for targeting Mpro enzyme
Pundir et al. [265]	2020	In silico	Pharmacophore-based virtual screening molecular mechanics Poisson-Boltzmann surface area (MM-PBSA)	PubChem database	Main protease	Papain-like protease, RNA-dependent RNA polymerase, main protease, spike glycoprotein, nonstructural protein 15 (nsp15)	Quinadoline B [19] was predicted to confer favorable ADMET values, high gastrointestinal absorptive probability, and poor blood-brain barrier crossing capacities
Quimque et al. [266]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	97 antiviral secondary metabolites from fungi		Two fumiquinazoline alkaloids quinadoline B (19), scedapin C (15), and polyketide isochaetochromin D1 (8)	
Rakib et al. [267]	2021	In silico	Molecular docking molecular dynamic simulation ADME properties	Selenium-containing heterocyclic compounds	Main protease	Selection of the 16 most effective selenocompounds as potential anti-COVID-19 drug candidates. Ethaselen showed potential binding affinities	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Rane et al. [268]	2020	In silico	Molecular docking molecular dynamic simulations	Diarylpyrimidine analogs	Spike glycoprotein	AP-NP (2-(2-amino-5-(naphthalen-2-yl)pyrimidin-4-yl)phenol), AP-3-OMe-Ph (2-(2-amino-5-(3-methoxyphenyl)pyrimidin-4-yl)phenol), and AP-4-Me-Ph (2-(2-amino-5-(p-tolyl)pyrimidin-4-yl)phenol) from a group of diarylpyrimidine derivatives, which appears to bind at the interface of the hACE2-S complex with low binding free energy	
Rao et al. [269]	2020	In silico	Molecular docking molecular dynamic simulations	Various small molecules	Main protease	Pyranonigrin A, a secondary fungal metabolite	
Salman et al. [270]	2020	In silico	Molecular docking molecular dynamic simulation ADME properties	Library of immunomodulatory medicinal compounds with antiviral capability	SARS proteases, spike protein, and nonstructural proteins (NSP-9, 15)	6 compounds: arzanol, ferulic acid, genistein, resveratrol, rosmarinol, and thymohydroquinone ZINC00003118440 is a theophylline derivative.	Good pharmacokinetic properties and low acute toxicity of these compounds
Sarma et al. [271]	2020	In silico	Molecular docking GBSA binding free energy molecular dynamic simulations	56,079 compounds from ASINEX and Maybridge library	RNA-binding N-terminal domain (NTD) of the N protein	Pyrimidone derivatives as possible inhibitors of RNA binding to the N-terminal domain of N protein of coronavirus	Lead molecules
Sepay et al. [272]	2020	In silico	Molecular docking, bioinformatics, and molecular electrostatic potential ADME studies	Benzylidenechromanones, naturally occurring oxygen heterocyclic compounds	Main protease	(Z)-3-(4'-chlorobenzylidene)-thiochroman-4-	Effective pharmacological properties
Shehroz et al. [273]	2020	In silico	Pharmacophore modeling molecular docking	DrugBank, ZINC, and TIMBLE databases	Spike (S) protein	Only eight molecules fit the criteria for safe oral drugs	Lead molecules
Singh and Das [274]	2021	In silico	Molecular docking	Chloroquine (CQ) hydroxychloroquine (HCQ) azithromycin	Spike (S) protein main protease host cathepsin L (CTSL) receptor-binding domain (RBD)	Azithromycin affinity scores (ΔG) with strong interactions with ACE2, CTSL, Mpro, and RBD. CQ firm bond score with Mpro HCQ and two results (ACE2 and Mpro) were firmly bound to the receptors	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Stefaniu et al. [275]	2020	In silico	Molecular docking density functional theory (DFT) computations, drug-likeness assessment	Derivatives of benzoic acid	Main protease	2,5-dihydroxybenzoic acid (gentisic acid) and octyl	A combination of the two compounds can be considered
Tachoua et al. [276]	2020	In silico	Molecular docking and structural dynamic study molecular mechanic Poisson-Boltzmann surface area (MM-PBSA) ADMET analysis	Chloroquine, quinine, nitazoxanide, doxycycline, lymecycline, cetirizine, mizolastine, indinavir	Main protease	Lymecycline mizolastine	
Uniyal et al. [277]	2020	In silico	Structure-based virtual screening molecular docking MM-GBSA binding free energy molecular dynamic simulations	Commercially available chemical libraries	Main protease	Compound AG-690/11203374_1 and AG-690/11203374_2 emerged as the best in silico hits based on the docking, MM-GBSA, dynamics, and ADMET studies	Lead molecules
Welker et al. [278]	2020	In silico/in vitro	Structure-activity relationships molecular docking fluorescence-based enzyme-activity assay Vero E6 cells	A series of rationally designed competitive, noncovalent, nonpeptidic active site-directed SARS-CoV PLpro inhibitors	Papain-like cysteine proteases (PLpro)	<i>R</i>)-5-amino-2-methyl-N-(1-(naphthalen-1-yl)ethyl) benzamide (2b), which is known to bind into the S3 and S4 pockets of the SARS-CoV PLpro. Isoindoline as a new class of potent PLpro inhibitors	IC ₅₀ value of $2.9 \pm 0.2 \mu\text{M}$
Wen et al. [279]	2021	In silico	Structure-based screening	8,820 compounds	Main protease	Trichostatin A	A histone deacetylase inhibitor and an antifungal compound
White et al. [280]	2020	In silico	Homology modeling and molecular dynamics approach molecular docking	~970,000 chemical compounds	Helicase (Nsp13)	Nilotinib and lumacaftor have significant activity in inhibiting purified recombinant SARS-CoV-2 helicase	
Wu et al. [281]	2020	In silico	Molecular docking	11 HIV-1 protease inhibitors, 12 nucleotide-analog inhibitors, 728 approved drugs	Main protease RNA-dependent RNA polymerase	Remdesivir shows the best binding energy on RdRp and saquinavir is the best inhibitor of main protease	

TABLE 3: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Zaher et al. [282]	2020	In silico	Design, synthesis SAR study molecular docking	Newly synthesized sixteen halogenated triazole compounds	Helicase (Nsp13)	The most potent compounds were 4-(cyclopent-1-en-3-ylamino)-5-(2-(4-iodophenyl)hydrazinyl)-4H-1,2,4-triazole-3-thiol [16] and 4-(cyclopent-1-en-3-ylamino)-5-[2-(4-chlorophenyl)hydrazinyl]-4H-1,2,4-triazole-3-thiol [12] ZINC12562757 and 112,260,215 were the best potential inhibitors of the ACE2 and main protease, respectively. Evo_1 compound enjoys the highest docking energy among the designed de novo ligands	
Zarezade et al. [283]	2021	In silico	3D-QSAR pharmacophore modeling ADMET properties, molecular docking molecular dynamic simulation MM-PBSA, hybrid QM-MM method de novo ligand design	PubChem and ZINC databases	Human angiotensin-converting enzyme 2 and main protease	Evo_1 has a stronger potential for specific inhibition of main protease, as compared to the 112, 260, 215 compound. Lead molecules	

TABLE 4: Summary of studies with *in vitro* method.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Agarwal et al. [284]	2020	In vitro	IL-1 β assay using THP-1 cells, <i>in vivo</i> pharmacokinetic investigation in C57BL/6 mice	Alkenyl sulfonylurea derivatives	NLRP3 inhibition and reduction in the release of interleukin-1 β	IL-1 β inhibition IC ₅₀ of 35 nM, good oral absorption showing C _{max} of 8.49 μ g/mL with an AUC of 48.9 μ g·h/mL and terminal half-life of 2.86 h, after oral route of administration at 3 mg/kg dose. Although the viral replication was not completely abolished (at 200 μ M and 400 μ M), SNAP delayed or completely prevented the development of viral cytopathic effect in treated cells (IC ₅₀ = 440.95 μ M $\pm$ 36.15 SE)	Novel thiazolo-alkenyl sulfonylurea derivative 7 as potent, selective, and orally bioavailable NLRP3 inflammasome inhibitor
Akaberi et al. [285]	2020	In vitro	Vero E6 cells, RT-qPCR for the quantification of viral RNA, expression and purification of SARS-CoV-2 3CL protease <i>in vitro</i> enzymatic assay	NO-Donor S-nitroso-N-acetylpenicillamine (SNAP); nitric oxide	Main protease		A dose-dependent inhibition of the 3CL protease (400 μ M)
Bernstein and Zhang [286]	2020	In vitro	Vero E6 cells: virus was then added at a multiplicity of infection (MOI) of 0.01, with 1 h allowed for infection. Cytotoxicity: Cell Counting Kit-8 (CCK-8) colorimetric assay	Gallium maltolate (GaM)	Inhibition of viral replication	EC ₅₀ (concentration producing 50% inhibition of viral replication) of about 14 μ M (CI 95%: 8.9–22.8 μ M)	No cytotoxicity was observed at concentrations up to at least 200 μ M
Bocci et al. [287]	2020	In vitro	Ligand-based virtual screening/Vero E6 cells. Uninfected cells and chloroquine as control inhibition of the SARS-CoV-2-mediated CPE following infection in Vero E6 cells tissue culture infectious dose (TCID ₅₀) assay	4000 approved drugs/ glafenine, amodiaquine vorinostat, zuclopenthixol, isosuprine, ambroxol, panobinostat, and pracinostat	Inhibition of viral replication	In silico: glafenine, amodiaquine (amodiaquine), vorinostat, zuclopenthixol, isosuprine, nebiivolol, ambroxol, panobinostat, and pracinostat. <i>In vitro</i> : zuclopenthixol and amodiaquine show anti-SARS-CoV-2 activity comparable with chloroquine. EC ₅₀ values were estimated as 0.13 μ M for AQ, 1.35 μ M for ZPX, and 2.72 μ M for nebiivolol	Zuclopenthixol and nebiivolol exhibit <i>in vitro</i> antiviral activities and potencies that are comparable or better than chloroquine and hydroxychloroquine
Cao et al. [288]	2020	In vitro	African green monkey kidney Vero E6 cells were infected with virus at an MOI of 0.01 for 1 h. Immunofluorescence assay physiologically based pharmacokinetic modeling and simulations	9 artemisinin-related compounds	Inhibition of viral replication	Arteannuin B showed the highest anti-SARS-CoV-2 potential with an EC (50) of 10.28 $\pm$ 1.12 μ M. Artesunate and dihydroartemisinin showed similar EC (50) values of 12.98 $\pm$ 5.30 μ M and 13.31 $\pm$ 1.24 μ M, respectively	Lumefantrine could inhibit SARS-CoV-2 <i>in vitro</i> with an EC ₅₀ of 23.17 $\pm$ 3.22 μ M. Impair viral infection by modulating host cell metabolic pathways

TABLE 4: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Chen et al. [289]	2021	In vitro	Fluorescence-based high-throughput screen, docking/pre-treated with various concentrations of test compound for 1 h, followed by infection with SARS-CoV-2 (MOI of 0.0001) in the presence of test compounds, expression and purification of SARS-CoV-2 PLpro, 3CL protease from <i>Escherichia coli</i> enzymatic assay of SARS-CoV-2 PLpro cytotoxicity assay	1920 natural products	Main protease, papain-like protease	Ginkgolic acid and anacardic acid were also identified as inhibitors of 3CL ^{pro} , with IC ₅₀ values of 1.79 ± 0.58 and 2.07 ± 0.35 µM, respectively, both ginkgolic acid and anacardic acid dose-dependently inhibited PLpro activity, with IC ₅₀ values of 16.30 ± 0.64 and 17.08 ± 1.30 µM, respectively	Ginkgolic acid act as an irreversible inhibitor against both PLpro and 3CL ^{pro} , suggesting it is a covalent inhibitor
Gendrot et al. [290]	2020	In vitro	Vero E6 cells, replication was estimated by RT-PCR/chloroquine as control determination of the inhibition stage	Doxycycline	Inhibition of viral replication	Median effective concentration (EC ₅₀) and 90% effective concentration (EC ₉₀) for doxycycline were 4.5 ± 2.9 µM and 23.5 ± 16.5 µM, respectively, EC ₅₀ = 4.5 µM	Interacted at both entry and postentry stages of the SARS-CoV-2 infection doxycycline has anti-inflammatory effects
Gupta et al. [291]	2021	In vitro	High-throughput virtual screening, molecular dynamic simulation, prime MM-GBSA/Vero E6 cells infected with the SARS-CoV-2 fluorescence-based biochemical assay for inhibitors of 3CLpro	FDA-approved drugs	Main protease, PLpro, RNA-directed RNA polymerase, helicase, NSP-15, NSP14, and 2'-O-methyltransferase	Troxerutin (main protease and PLpro) and bisindolylmaleimide derivatives (main protease and exon) had possible dual targets. Ivermectin has nonselective toxicity to the ATCC E6 Vero cells at ≤50 µM and 16.67 µM based on the number of nuclei counted	BIM IX specifically blocked 3CLpro <i>in vitro</i> enzymatic assay inhibition was observed with an IC ₅₀ value of 113.7 ± 5.2 µM a known inhibitor of protein kinase C isoforms, bisindolylmaleimide IX (BIM IX), was found to be a potent inhibitor of SARS-CoV-2
Hahn et al. [292]	2020	In vitro	Vero B4 and 76 cells 9 Vero E6 cells/remdesivir RT-qPCR for the detection of extracellular SARS-CoV-2 in cell immunostaining for the detection of Intracellular SARS-CoV-2 Western blot analysis viral plaque and yield reduction assay neutral red assay (NRA)	(IMU-838) inhibitor of human dihydroorotate dehydrogenase (DHODH)	Inhibition of viral replication/immunomodulator	A virus-specific EC ₉₀ of 6.2 ± 1.9 µM was measured, with no cytotoxicity observed with drug concentrations up to 100 µM at 3 days p.i.	Active moiety of IMU-838, vidofludimus, possesses broad-spectrum antiviral activity. IMU-838 reduces T lymphocyte proliferation, cytokine production, and organ infiltration by leukocytes in various <i>in vivo</i> and <i>in vitro</i> models for autoimmunity

TABLE 4: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Hattori et al. [293]	2021	In vitro	VeroE6 cell-based assays with RNA-qPCR, cytopathic assays	Two small-molecule-compounds, named GRL-1720 and 5 h, containing an indoline and indole moiety	Main protease	EC(50) values of 15 ± 4 and $4.2 \pm 0.7 \mu\text{M}$ for GRL-1720 and 5 h	Combination of 5 h and remdesivir exhibits synergism against SARS-CoV-2.5 h might serve as a lead M (pro) inhibitor
Jang et al. [294]	2021	In vitro	Vero CCL-81 cells/Calu-3 human lung epithelial cells	Gemcitabine and its analog 2'-fluoro-2'-deoxycytidine (2FdC)	Inhibition of viral replication	50% effective concentration (EC(50)) of $1.2 \mu\text{M}$ compare to remdesivir 2FdC was marginally active (EC(50) = $175.2 \mu\text{M}$)	Gemcitabine has a synergistic effect when combined with remdesivir
Jeon et al. [295]	2020	In vitro	Drug repositioning/chloroquine, lopinavir, and remdesivir were used as reference drugs/VeroE6 cell/immunofluorescence analysis	48 FDA-approved drugs	Inhibition of viral replication	Niclosamide (IC_{50} $0.28 \mu\text{M}$), ciclesonide (IC_{50} $4.33 \mu\text{M}$)	24 potential antiviral drug candidates against SARS-CoV-2 infection
Liu et al. [296]	2020	In vitro	Virtual screening/spiked pseudotyped lentivirus for imitating SARS-CoV-2 cell entry/ACE2-GFP expressing HEK293T 2 cells, HeLa Cells, 293T (human, kidney) cells, and Vero-E6 (African green monkey, kidney) cells	FDA-approved drugs	Angiotensin I-converting enzyme 2 (ACE2) and a spike protein	Nine potential candidates were selected and submitted to experimental studies. Three (romidepsin, saquinavir, and nelfinavir) of nine drugs possess the ability to suppress SARS-2-S pseudotyped particles to enter the ACE2 expressing cells in a concentration-dependent manner	Five clinical HDAC inhibitors including romidepsin, panobinostat, givinostat hydrochloride monohydrate, CAY10603, and sirtinol could inhibit noticeably the spike/ACE2-mediated cell entry of SARS-CoV-2
Ma et al. [297]	2020	In vitro	Vero E6 cells using cytopathic effect and plaque reduction in assay. HCoV-229E infection in Huh-7 cells. Western blot assay	Phillyrin (KD-1) ingredient of Forsythia suspensa	Inhibition of viral replication	Inhibit SARS-CoV-2. Markedly reduce the production of pro-inflammatory cytokines (TNF- α , IL-6, IL-1 β , MCP-1, and IP-10) at the mRNA levels. p-NF- κB p65, NF- κB p65, and p-I $\kappa\text{B}\alpha$	KD-1 could significantly inhibit virus proliferation in vitro, the upregulated expression of pro-inflammatory cytokines induced by SARS-CoV-2
Masih et al. [298]	2020	In vitro	LPS-stimulated RAW267.4 cells by enzyme immunoassay Western blot assay	Pyrazole derivatives	Anti-inflammatory levels of interleukin-1 β , tumor necrosis factor- α , and interleukin-6	Compound 6c as the most potent analog among the tested series. Inhibition of inhibitor kappa B- α and NF- κB	
Omotuyi et al. [299]	2020	In vitro	Kinetic assay	Aframomum melegueta. 100 secondary metabolites	Main protease, 2'-O-ribose methyltransferase (NSP16), and surface glycoprotein/ACE2 receptor interface	Diarylheptanoid (Ietestuianin A), phenylpropanoid (4-cinnamoyl-3-hydroxy-spiro[furan-5,2'-(1'H)-indene]-1',2,3'-(2'H,5'H)-trione), flavonoids (quercetin, apigenin, and tectochrysin) have been identified as high binding compounds to SARS-CoV-2 targets in a polypharmacology manner	Di-ethyl-ether (IC_{50} = 0.03 mg/L), acetone (IC_{50} = 1.564 mg/L), ethyl-acetate (IC_{50} = 0.382 mg/L), and methanol (IC_{50} = 0.438 mg/L) fractions demonstrated the best inhibition in kinetic assay

TABLE 4: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Outlaw et al. [300]	2020	In vitro	Vero E6 cell β -Gal complementation-based fusion assay. Viral titration and plaque reduction neutralization assay quantitative RT-PCR.	A lipopeptide that is derived from the C-terminal heptad repeat (HRC) domain of SARS-CoV-2 S	Inhibition cell-cell fusion	Inhibits cell-cell fusion mediated by SARS-CoV-2 S and blocks infection by live SARS-CoV-2 in Vero E6 cell monolayers (IC ₅₀) of ~10 nM and 90% inhibitory concentration (IC ₉₀) of ~100 nM. Selection of 30 compounds.	
Pitsillou et al. [301]	2021	In vitro	Molecular docking, molecular dynamic simulation/papain-like protease (SARS-CoV-2) assay kit GRL-0617 as a control enzymatic inhibition assay	300 small compounds derived predominantly from our OliveNet™ library (222 phenolics) and supplemented with synthetic and dietary compounds	Papain-like protease	Hypericin possessed inhibition activity, and both rutin and cyanidin-3-O-glucoside resulted in a concentration-dependent inhibition of the PLpro	The IC ₅₀ values were not clarified in experiment
Pizzorno et al. [302]	2020	In vitro	Vero E6 cells/remdesivir, lopinavir, chloroquine as comparator	Favipiravir, ribavirin, umifenovir (arbidol), berberine, cyclosporine A, diltiazem	Inhibition of viral replication	Umifenovir, berberine, and cyclosporine A with estimated 50% inhibitory concentrations of 0.99, 5.2, 1.38, 3.5, 10.6, and 3 μ M, respectively (IC ₅₀) of 8.2 μ M in monkey VeroE6 cells (CC ₅₀) of 13.5 μ M, and selectivity index (SI) of 1.65. IC ₅₀ of 11.3 μ M, CC ₅₀ of 23.1 μ M, and SI of 2.04 in human A549-ACE2 cells	A strong antagonism between remdesivir and berberine, in contrast with remdesivir/diltiazem, for which we describe high levels of synergy
Plaze et al. [303]	2021	In vitro	Monkey VeroE6 cells and human alveolar basal epithelial A549-ACE2 cell RT-qPCR for the presence of SARS-CoV-2 RNA	Chlorpromazine	Inhibition of viral replication		
Reznikov et al. [304]	2021	In vitro	Repurposing study molecular docking/human lung A549 cells that were transfected with hACE2 (ACE2-A549) cells, Vero E6 cells	Electronic health records of over 219,000 patients with antihistamines	Inhibition of viral replication/spike protein	IC ₅₀ 15.3 μ g/ml for hydroxyzine, 17.4 μ g/ml for diphenhydramine, and 2.24 μ g/ml for azelastine	Hydroxyzine and possibly azelastine bind angiotensin-converting enzyme-2 (ACE2) and the sigma-1 receptor as off-target disease prevention
Sakurai et al. [305]	2021	In vitro	VeroE6 cells/human colon-derived Caco-2 cells/infection assay with immunofluorescence	5-Amino levulinic acid (5-ALA)	Inhibition of viral replication	IC ₅₀ of 570 μ M/IC ₅₀ of 63 μ M in human Caco-2 cells. No significant cytotoxicity	5-ALA is synthesized in most animals and plants and we are continuously consuming it in our food. Show anti-inflammation effects in humans
Stone et al. [306]	2021	In vitro	Vero E6 monkey kidney and Calu-3 human lung adenocarcinoma cells/RT-qPCR of RNA	Stenoparib poly(ADP ribose) polymerase (PARP) inhibitor	Inhibition of viral replication	Dose-dependent activity against SARS-CoV-2 at concentrations up to 30 μ M with negligible cytotoxicity. Synergistic effect with remdesivir	Stenoparib impedes entry and postentry processes

TABLE 4: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Su et al. [307]	2020	In vitro	Vero E6 cells, quantitative real-time RT-PCR	Shuanghuanglian preparation, a Chinese traditional patent medicine (chlorogenic acid, phillyrin, baicalin, and baicalin)	Main protease	Dose-dependent inhibition of SARS-CoV-2 3CLpro, and the resulting IC ₅₀ values were 0.090, 0.064, and 0.076 $\mu\text{g}/\text{mL}$ for three Shuanghuanglian oral liquids produced by three different pharmaceutical companies	Baicalin and baicalin, which have good drug-like properties
Touret et al. [308]	2020	In vitro	VeroE6 cells/human colon-derived Caco-2 cell real-time RT-PCR/remdesivir as control	FDA-approved chemical library	Inhibition of viral replication	23 drugs were selected to cover the 12 different groups. 11 compounds such as macrolide antibiotics, proton pump inhibitors, antiarrhythmic agents, or CNS drugs emerged showing antiviral potency with $2 < \text{EC}_{50} \leq 20 \mu\text{M}$	Two of the highest antiviral activity were obtained for azithromycin ($\text{EC}_{50} = 2.12 \mu\text{M}$) and hydroxychloroquine ($\text{EC}_{50} = 4.17 \mu\text{M}$). Voronoprazan and sulfadoxine show a moderate activity with EC_{50} above $30 \mu\text{M}$
Tree et al. [309]	2021	In vitro	Plaque inhibition assay with Vero E6 cells/differential scanning fluorimetry/ELISA assays	Unfractionated heparin (UFH), low MW heparins	Inhibition of viral replication/spike protein RBD	All the UFH preparations had potent antiviral effects, with IC ₅₀ values ranging between 25 and $41 \mu\text{g}/\text{mL}$, whereas LMWHs were less inhibitory by ~ 150 -fold (IC ₅₀ range $3.4\text{--}7.8 \text{ mg}/\text{mL}$)	Heparin directly inhibits the binding of RBD to the human ACE2 protein receptor
Vatansever et al. [310]	2021	In vitro	Docking, screening assay/Vero E6, and A549 cell culture	30 FDA/EMA-approved drugs	Main protease	Six FDA/EMA-approved drugs can potentially inhibit Mpro with an IC ₅₀ value lower than $100 \mu\text{M}$. Bepridil exhibited strong inhibition of SARS-CoV-2 from entry and replication inside Vero E6 and A549 cells (IC ₅₀ = $72 \mu\text{M}$)	Bepridil indicated that it had low micromolar EC ₅₀ values
Wang et al. [311]	2020	In vitro	FRET-based enzyme activity assay of Mpro measurement of human TMPRSS2 activity by a FRET-based enzymatic assay surface plasmon resonance (SPR) analysis molecular docking African green monkey kidney cell Vero E6 Virtual drug screening, docking/VeroE6 cells detecting the viral RNA by real-time RT-PCR or viral proteins by immunofluorescence	Catechin, kaempferol, quercetin, proanthocyanidins, resveratrol, and tannic acid	TMPPSS2 (transmembrane protease serine 2) main protease	Tannic acid inhibited the activities of the two proteases with an IC ₅₀ of 13.4 mM for Mpro and 2.31 mM for TMPRSS2	A potent dual inhibitor
Watashi [312]	2021	In vitro		FDA-approved drugs	Main protease, Plpro, RdRp, helicase, spike, ACE2, and TMPRSS2	Abiraterone, amodiaquine, anidulafungin, arbidol, astemizole, atazanavir, auranofin, azithromycin, bazedoxifene, bexarotene, camostat IC_{50} values $< 5 \mu\text{M}$	Cepharanthine, cetilistat, chloroquine, ciclesonide, cyclosporine A, digoxin, ivermectin, lopinavir, mefloquine, nelfinavir, niclosamide

TABLE 4: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Yu et al. [313]	2021	In vitro	Computer-aided drug design and biological verification surface plasmon resonance (SPR) assays and NanoBIT assays	Traditional Chinese medicine	Spike proteins	Glycyrrhizic acid, the most efficient and nontoxic broad-spectrum anticoronavirus molecule IC_{50} was 22 μM	
Zandi et al. [314]	2021	In vitro	African green monkey kidney cells (Vero CCL-81 cells) and reverse transcription-quantitative PCR (qRT-PCR). Remdesivir and β -D- N^4 -hydroxycytidine as positive drug controls	Nucleoside analogs/anti-HCV agents	Inhibition of viral replication	Sofosbuvir and favipiravir demonstrated no antiviral effect against COVID-2	
Zhang et al. [315]	2020	In vitro	Hybrid virtual screening, docking, force field-based simulation/qRT-PCR assay, indirect immunofluorescence assay (IFA) and CCK-8 assay, surface plasmon resonance (SPR) assay/Vero cells	1906 approved drugs from TargetMol-approved drug library	RNA-dependent RNA polymerase (RdRp)	Pralatrexate and azithromycin (EC_{50}) values of 0.008 and 9.453 μM	A new therapeutic agent pralatrexate against COVID-19 by targeting RdRp
Zhang et al. [316]	2020	In vitro	African green monkey kidney Vero E6 cell line, quantitative real-time PCR (RT-PCR) analysis, immunofluorescence microscopy, Western blot analysis	Lopinavir/ritonavir, rupintrivir, and AG7404	Inhibition of viral replication	(IC_{50}) and half cytotoxic concentration (CC_{50}) values of lopinavir were $\sim 12.01 \mu M/L$ and 80.82 $\mu M/L$; for ritonavir, the IC_{50} and CC_{50} values were 19.88 $\mu M/L$ and 94.71 $\mu M/L$ ($SI = 4.76$)	Rupintrivir inhibited SARS-CoV-2 infection only at high drug concentrations (Vero E6: $IC_{50} = 34.08 \mu M/L$, AG7404: Vero E6: $IC_{50} \sim 195.8 \mu M/L$). Lopinavir/ritonavir should be stopped for clinical use due to the huge gap between in vitro IC_{50} and free plasma concentration
Zhu et al. [317]	2021	In vitro	Human fibroblast lung cells (MRC-5), cytopathic effect (CPE) assay, primary human airway air-liquid interface (ALI) cultures, TEER assay, CCK-8 assays	Stimulator of interferon genes (STING): dimeric amidobenzimidazole (diABZI)	Inhibition of viral replication	Potent anticoronavirus activity against both the common cold human coronavirus 229E (HCoV-229E) and SARS-CoV-2 in cell culture systems. EC_{50} of 120 μM	

TABLE 5: Summary of studies with in vivo method.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Lei et al. [318]	2020	In vivo	BALB/c mice to determine the pharmacokinetic profiles of the fusion proteins	A recombinant protein by connecting the extracellular domain of human ACE2 to the Fc region of the human immunoglobulin IgG1	Spike protein	The IC ₅₀ values of ACE2-Ig for SARS-CoV and SARS-CoV-2 neutralization were 0.8 and 0.1 µg·ml ⁻¹ , respectively. Desirable pharmacological properties in mice	Potential applications in the diagnosis, prophylaxis, and treatment of SARS-CoV-2
Rathnayake et al. [319]	2020	In vivo	Old male hDPP4-KI mice infected with MERS _{MA} -CoV	Dipeptidyl and tripeptidyl series of compounds	Main protease	7a , 6c , 7e , 7h , and 6j : EC ₅₀ values ranging from 0.15 to 0.9 µM in Vero E6 cells, IC ₅₀ : 0.17 to 0.82 µM. 40% of mice treated with compound 6h survived, and all mice treated with compound 6j were alive at the end of the study Neutralizing hu-mAbs targeting SARS-CoV-2 promotes reduction in viral load and prevent infection in macaques and hamsters effective dose: 5.3 and 16 mg/kg inhibitory concentration (IC) ₅₀ /IC90 (half-maximal/90% IC) of 4.4/18 to 26/140 ng/ml	6j resulted in the survival of MERS _{MA} -CoV-infected hDPP4-KI mice
Schäfer et al. [320]	2020	In vivo	An immune-competent mouse model of COVID-19 by remodeling the SARS-CoV-2 S RBD at the mACE2-binding interface. Human Ace2-expressing HT1080 cells and Syrian hamster model of SARS-CoV-2	Human mAbs (hu-mAbs)	Reduction in viral load and prevent infection	Direct inhibitory activity on human PPAR-1 was detected in the low micromolar range (IC ₅₀ = 1.55 µM). MP1032 works as a ROS scavenger, reduced the secretion of TNF-α and IL-6. MP1032 pre-treatment significantly reduced plasma cytokine levels compared with vehicle-treated mice by 50% (TNF-α) and 25% (IL-6). Significant reduction in SARS-CoV-2 replication	MP1032 could further attenuate prolonged virus replication by preventing oxidative stress or by limiting ADP ribosylation of the viral nucleocapsid protein via PPAR-1 inhibition
Schumann et al. [321]	2020	In vivo	Female C57BL/6 mice, LPS stimulation	MP1032:5-amino-2,3-dihydro-1,4-phthalazinedione sodium salt	Immune-modulating		

TABLE 5: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Shang et al. [322]	2021	In vivo	African green female hACE2 transgenic mice, SARS-CoV-2 injection RNA extraction and qPCR, RNA in situ hybridization assay, histopathological examination	Endosomal acidification inhibitors, including chloroquine, bafilomycin A1, and NH ₄ Cl	Antiviral actions against SARS-CoV-2	(40 µM), bafilomycin A1 (100 nM), and NH ₄ Cl (12.5 mM) suppressed the replication of SARS-CoV-2 in all cell types chloroquine (60 mg·kg ⁻¹) and bafilomycin A1 (0.1 mg·kg ⁻¹) markedly reduced virus yields in lung tissues. Alleviated viral pneumonia in hACE2 transgenic mice	Endosomal acidification inhibitors exhibited antiviral actions against SARS-CoV-2
Sheahan et al. [323]	2020	In vivo	Male and female 20- to 29-week old SPF C57BL/6J (Stock 000664 Jackson Labs) mice, whole-body plethysmography 50, 150, or 500 mg/kg EIDD-2801 2 hr prior to intranasal infection with 5E + 04 PFU of mouse-adapted SARS-CoV Sprague Dawley rats. (i) HCQ-IV: 12 rats received a single dose of 0.590 mg HCQ sulfate per animal via IV injection; (ii) HCQ-IT: 20 rats received a single dose of 0.590 mg HCQ sulfate per animal via intratracheal (IT) administration; and (iii) liposomal HCQ-IT: 20 rats received a single dose of 0.284 mg liposomal HCQ sulfate per animal via IT administration 0.25, 1, 4, 24, and 72 hours postdose and for tissue/organ samples were 0.25, 4, 24, and 72 hours postdose	Ribonucleoside analog β-d-N ⁴ -hydroxycytidine	Antiviral efficacy against the three most recently emerged human CoV: SARS-CoV, MERS-CoV, and SARS-CoV-2	Bioavailable NHC prodrug (β-d-N ⁴ -hydroxycytidine-5'-isopropyl ester), improved pulmonary function and reduced virus titer and body weight loss Compared with unformulated HCQ administered intravenously, liposomal HCQ showed higher (~30-fold) lung exposure, longer (~2.5-fold) half-life in lungs, but lower blood exposure with ~20% of peak plasma concentration (C _{max}) and 74% of area under the curve from 0 to 72 hours (AUC 0–72) and lower heart exposure with 23% of C _{max} and 58% of AUC 0–24 (normalized for dose)	Continued development of EIDD-2801 as a potent broad-spectrum antiviral
Tai et al. [324]	2021	In vivo		Inhalable liposomal hydroxychloroquine (HCQ)	Pharmacokinetic study		Inhalable liposomal HCQ may provide clinical benefit and serve as a potential treatment of COVID-19

TABLE 5: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Tortorici et al. [325]	2020	In vivo	From two individuals recovering from severe COVID-19 disease. Surface plasmon resonance (SPR) and flow cytometry. Cell-cell fusion assay using VeroE6 cells/Syrian hamster model	Human neutralizing antibodies (S2E12 and S2M11)	Against SARS-CoV-2	IC ₅₀ values were 1.2 to 6.6 ng/ml to protect hamsters against SARS-CoV-2 challenge competitively block angiotensin-converting enzyme 2 (ACE2) attachment and that S2M11 also locks the spike in a closed conformation by recognition of a quaternary epitope spanning two adjacent receptor-binding domains	Antibody cocktails for prophylaxis or therapy
Wahl et al. [326]	2021	In vivo	Coronavirus replication in human lung-only mice (LoM) and histopathologic analysis LoM were administered EIDD-2801 starting 24 h or 48 h post-SARS-CoV-2 exposure and every 12 h thereafter	Ribonucleoside analog β -D-N ⁴ -hydroxycytidine (NHC): oral prodrug EIDD-2801 (also known as molnupiravir or MK-4482)	Inhibited SARS-CoV-2 replication	EIDD-2801 pre-exposure prophylaxis significantly reduced virus titers in the human lung tissues of LoM by over 100,000-fold in two independent experiments Ten polypeptide drugs were selected. Dalbavancin showed the strongest inhibitory ability. (EC50) was ~12 nM. Significant inhibition of SARS-CoV-2 pseudovirion entry into HEK293/hACE2 cells, with an IC ₅₀ of ~53 nM in both mouse and rhesus macaque models, viral replication, and histopathological injuries caused by SARS-CoV-2 infection are significantly inhibited by dalbavancin administration	Prophylactic administration EIDD-2801 is highly effective
Wang et al. [327]	2021	In vivo	Male hACE2 mice inoculated intranasally with SARS-CoV-2 stock virus rhesus macaque model of SARS-CoV-2 infection	Screening the FDA-approved peptide drug library with LibDock	Against SARS-CoV-2	Dalbavancin directly binds to human angiotensin-converting enzyme 2 (ACE2) with high affinity, thereby blocking its interaction with the SARS-CoV-2 spike protein	

TABLE 5: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Weston et al. [328]	2020	In vivo	Mice were inoculated intranasally with SARS-CoV	20 FDA-approved drugs	Against SARS-CoV-2	7 of these inhibit SARS-CoV-2 at non-cytotoxic concentrations. Many of the compounds have IC ₅₀ under 10 μ M, chloroquine and chlorpromazine did not inhibit viral replication in mouse lungs based on viral titers recovered at 4 dpi neither drug inhibited viral replication in the lungs, but both protected against clinical disease IC90 of 1.76 nM; Vero E6 IC90 of 0.88 nM: human cell line significantly reduced genomic RNA content reduction of nearly 2 log units in SARS-CoV-2 viral titers in the lungs of the 0.3 mg/kg plitidepsin group, reduction of 2 log units in viral lung titers at day 3, similar to two daily 50 mg/kg doses of remdesivir No mouse death and no significant alteration in mouse activity or body weight (IHC) data revealed that the expression levels of both ACE2 and TMPRSS2 were markedly diminished in the GB-2 group compared with the control group 50 μ g/mL of theaflavin could inhibit protein expression of ACE2 and TMPRSS2	Plitidepsin as a host-targeted anti-SARS-CoV-2 agent with in vivo efficacy
White et al. [329]	2021	In vivo	A human ACE2-expressing adenovirus to transduce the naturally resistant wild-type BALB/c mice and sensitize them to SARS-CoV-2 infection, K18-hACE2 mouse model	eEF1A inhibitor plitidepsin (aplidin): targeted the eukaryotic translation machinery	Inhibited SARS-CoV-2 replication		
Wu et al. [330]	2020	In vivo	10 male C57BL/6 mice related to 200 mg/kg/day GB-2 everyday by oral administration. immunohistochemistry (IHC) assessment	GB-2, the formula from Tian Shang Sheng Mu of Chiayi Puzi Peitian Temple: in Taiwan: the index compound of <i>Camellia sinensis</i> var. assamica extract	The protein and mRNA expression of ACE2 and TMPRSS2		GB-2 and theaflavin could act as potential compounds for ACE2 and TMPRSS2 inhibitors

TABLE 5: Continued.

Author	Year	Method	Detail of method	Name of compound/drug	Target	Efficacy	Comments
Yuen et al. [331]	2021	Ex vivo	Human lung tissues were treated with the indicated drugs overnight, followed by SARS-CoV-2 infection at 2×10^6 PFU/well	Small molecules targeting the ULK1/Atg1 complex involved in the induction stage of autophagy (ULK1 inhibitor SBI0206965), the ATG14/Beclin1/VPS34 complex involved in the nucleation step of autophagy (class III PI3-kinase inhibitor VPS34-IN1), and a widely used autophagy inhibitor that persistently inhibits class I and temporary inhibits class III PI3-kinase (3-MA) and a clinically approved autophagy inhibitor that suppresses autophagy by inhibiting lysosomal acidification and prevents the formation of autophagolysosome (HCQ)	Target the key cellular factors involved in key steps of the autophagy pathway	Not all the tested autophagy inhibitors suppressed SARS-CoV-2 infection inhibition of class III PI3-kinase Vps34 downstream of ULK1, in contrast to inhibition of ULK1, which caused a significant decrease in SARS-CoV-2 replication (EC50) of HCQ was $19 \mu\text{M}$, that of Vps34-IN1 was $0.82 \mu\text{M}$ Vps34-IN1 potently inhibited SARS-CoV-2 viral replication in normal ex vivo human lung tissue culture in a dose-dependent manner	Class III PI3-kinase may be a possible target for COVID-19 therapeutic development
				Mouse anti-SARS-CoV-2-neutralizing MAbs: 2H2 and 3C1	Spike (S) protein	(IC ₅₀) of 12 ng/mL effectively treat SARS-CoV-2-infected mice even when administered as late as 24 h post-infection	
Zhang et al. [332]	2021	In vitro/ in vivo	Preparation of MAbs from female BALB/c mice and wild-type Balb/c mice were intranasally inoculated with hACE2-encoded adenovirus 5 (Ad5-hACE2) to allow expression of the hACE2 receptor in the lung, followed by intranasal infection with live SARS-CoV-2 3 days later				

by SARS-CoV-2 markedly and could be considered as adjuvant treatment of COVID-19 severe disease.

Doxycycline, chlorpromazine, azithromycin, heparin, bepridil, tannic acid, and glycyrrhizic acid are well-known drugs that show both *in silico* and *in vitro* inhibitory effects against SARS-CoV-2 and should be considered for this purpose.

Also, *in vitro* studies show that lopinavir/ritonavir, sofosbuvir, and favipiravir have no antiviral effects against SARS-CoV-2 (huge gap between *in vitro* IC₅₀ and free plasma concentration).

We included 16 *in vivo* studies in our final analysis. Most of them use a combination of *in vitro* and *in vivo* methods for the evaluation of novel drugs. Table 5 shows the detail of these studies.

Human monoclonal antibodies were the most evaluated drugs ($N=4$), promote the reduction in viral load (*in vitro*), and prevent infection in animal models of SARS-CoV-2.

A prodrug of hydroxycytidine (molnupiravir) improved pulmonary function and reduced viral titer *in vivo* and was introduced as a potential broad-spectrum antiviral agent against SARS-CoV-2.

Chloroquine and chlorpromazine did not inhibit viral replication in mouse lungs, but protected them against clinical disease.

Dalbavancin shows significant inhibitory ability in both *in vitro* and *in vivo* models of COVID-19. The drug binds directly to ACE2 and blocks its interaction with the SARS-CoV-2 spike protein.

4. Discussion

Drug development is a multistep process, typically requiring more than five years to assure the safety and efficacy of the new compound. There are several strategies in antiviral drugs for coronaviruses including empirical testing of known antiviral drugs, large-scale phenotypic screening of compound libraries, and target-based drug discovery. To date, an increasing number of drugs have been shown to have anticoronavirus activities *in vitro* and *in vivo*, but only remdesivir and several neutralizing antibodies have been approved by the US FDA for treating COVID-19. However, remdesivir's clinical effects are controversial and new antiviral drugs are still urgently needed. Given the urgency of the SARS-CoV-2 outbreak, here we discuss the discovery and development of new therapeutics for SARS-CoV-2 infection, which have been conducted in basic research (*in silico*) and preclinical study (*in vitro*, *in vivo*). Our database search identified about 3000 studies, which means a global effort for drug development in the current COVID-19 pandemic. Although the chance of successful drug development is very low (less than about 10%) and till today, there are no approved drugs. Many potential candidates (at least 420 drugs) are in clinical trials.

As summarized in the results, many compounds are in the development process for COVID-19 disease. Some of these compounds are completely new and could serve as seeds (or leads) for developing antiviral drugs against COVID-19, but as we need therapeutics as soon as possible,

half of the studies focused on drug repurposing (repositioning), which is a process of investigation of existing drugs for new therapeutic purposes. With the emergence of a growing COVID-19 pandemic, the drug repurposing process was being accelerated. Clinical trials using repurposed drugs may take less time and have a lower overall cost of manufacturing and could have a wide distribution of drugs. According to our results, 260 drugs repurposed by the computational methods for COVID-19, among them saquinavir, ritonavir, and lopinavir, showed the best efficacy in *in silico* environment. These drugs can be rapidly repurposed for clinic application for treating COVID-19 patients given their proven safety.

Many trials are performed using a combination of ritonavir-lopinavir. The results of a systematic review and meta-analysis showed that this drug combination has no more treatment effects than other therapeutic agents in COVID-19 patients and is currently not used anymore [134]. We could not find any clinical trial on saquinavir, which is the most studied drug *in silico* and show high potency against COVID-19. Saquinavir could be a suggestion for further clinical research.

Given that the development of synthetic chemicals for therapeutic use is a random process that might result in serendipitous discovery, many pharmaceutical companies are now focused on the development of plant-derived drugs. Natural products and their structural analogs have historically made a major contribution to pharmacotherapy, especially for cancer and infectious diseases. Nevertheless, natural products also present challenges for drug discovery, such as technical barriers to screening, isolation, characterization, and optimization. In recent years, several technological and scientific developments—including improved analytical tools, genome mining and engineering strategies, and microbial culturing advances—are addressing such challenges and opening up new opportunities. Consequently, interest in natural products as drug leads is being revitalized. Medicinal plants have attracted significant attention to treat infectious diseases. Complex molecular structures and a wide variety of natural compounds make medicinal plants an excellent biological resource for drug discovery. Our results show that various plants have potential antiviral activities and could use or be a basis for drug development against COVID-19.

Some of the studied plants are among what used by people on a daily basis, such as green tea, aloe, curcumin, and eucalyptus. A systematic review and meta-analysis of RCTs on herbal medicine in the management of COVID-19 show the significant effects of the combined therapy of herbal medicine in treating COVID-19 without any significant side effects [135].

Our results retrieved 91 *in silico* studies of natural products including herbal medicine. The results of this *in silico* approach showed that some of these studied active ingredients have a high affinity for each of the four important viral proteins compared with the inhibitors previously reported for each of these proteins. They could possibly have an inhibitory effect on the SARS-CoV-2 and COVID-19. Most evaluated chemical compounds

had inhibitory effects against one or two proteins of SARS-CoV-2. In addition to having a great affinity to attach to the viral proteins, these herbal compounds have antioxidant, vasoprotective, anticarcinogenic, and antiviral properties. Thus, they can be applied as extremely safe therapeutic natural compounds and clinical assessments might have notable outcomes for controlling COVID-19.

Due to the nature of preclinical (in vitro and in vivo) studies, the number of drug development studies in this area was less than in silico studies. Again, repurposed drugs were the most studied drugs in vitro (both herbal and synthetic). Some of them such as saquinavir, heparin, glycyrrhizic acid, and chlorpromazine show efficacy in both in vitro and in silico environments. Chlorpromazine is the single agent that was found to have efficacy in in silico, in vitro, and in vivo areas.

In terms of mechanism of action, different targets from the structural and nonstructural proteins of COVID-19 were evaluated. Most of the studies focused on the main protease, papain-like protease, and spike glycoprotein. Apart from the specific protein that leads to viral replication, SARS-CoV-2 causes a surge of pro-inflammatory cytokines and chemokines, which cause damage to lung tissue and deterioration of lung function. Therefore, the design of a drug with multitarget of action against different proteins of COVID-19 and also anti-inflammatory potential could be valuable.

Currently, there is no highly efficacious and specific treatment for SARS-CoV-2. Herein, we provided data on novel compounds in therapeutic drug discovery and development. Due to the nature of SARS-CoV-2 and the rises of several high transmissible strains, repurposing existing drugs has demonstrated power by bringing several drugs to approval for treating COVID-19 patients, such as remdesivir. Our results confirmed that a large number of repurposed agents are currently being explored for treating SARS-CoV-2 infection. However, these drugs still suffer from suboptimal therapeutic effect or known strong side effect. To accelerate drug discovery and development, especially during the current pandemic, natural products capture attention again. Our results showed that various natural bioactive compounds are being investigated in the pre-clinical step of drug development for COVID-19. In addition to having high affinity, these herb active ingredients have antioxidant, vasoprotective, anticarcinogenic, and antiviral properties. Therefore, they can be used as extremely safe therapeutic compounds in drug design studies to control COVID-19. However, the pharmacological effects and adverse reactions of some drugs under development are still unclear, and hence, well-designed high-quality studies are needed to further study the effectiveness and safety of these potential drugs to accelerate drug development targeting SARS-CoV-2 and thus promote progress towards ending the pandemic.

Data Availability

All data generated or analyzed during this study are included in this published article.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This manuscript was financially supported by a grant from the Research and Technology Department of Isfahan University of Medical Sciences (grant no. 340068), Isfahan, Iran.

References

- [1] C. Sohrabi, Z. Alsafi, N. O'Neill et al., "World Health Organization declares global emergency: a review of the 2019 novel coronavirus (COVID-19)," *International Journal of Surgery*, vol. 76, pp. 71–76, 2020.
- [2] WHO, "WHO coronavirus (COVID-19) dashboard," 2022, <https://covid19.who.int/>.
- [3] A. Macedo, N. Gonçalves, and C. Febra, "COVID-19 fatality rates in hospitalized patients: systematic review and meta-analysis," *Annals of Epidemiology*, vol. 57, pp. 14–21, 2021.
- [4] D.-H. Kim, Y. J. Choe, and J.-Y. Jeong, "Understanding and interpretation of case fatality rate of coronavirus disease 2019," *Journal of Korean Medical Science*, vol. 35, no. 12, 2020.
- [5] WHO, *Therapeutics and COVID-19: Living Guideline*, World Health Organization, Geneva, Switzerland, 2022.
- [6] H. A. Rothan and S. N. Byrareddy, "The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak," *Journal of Autoimmunity*, vol. 109, Article ID 102433, 2020.
- [7] M. Mei and X. Tan, "Current strategies of antiviral drug discovery for COVID-19," *Frontiers in Molecular Biosciences*, vol. 8, p. 310, 2021.
- [8] M. Mohamadian, H. Chiti, A. Shoghli, S. Biglari, N. Parsamanesh, and A. Esmailzadeh, "COVID-19: virology, biology and novel laboratory diagnosis," *The Journal of Gene Medicine*, vol. 23, no. 2, Article ID e3303, 2021.
- [9] A. A. Momtazi-Borojeni, M. Banach, Ž. Reiner et al., "Interaction between coronavirus S-protein and human ACE2: hints for exploring efficient therapeutic targets to treat COVID-19," *Angiology*, vol. 72, no. 2, pp. 122–130, 2021.
- [10] C. Liu, Q. Zhou, Y. Li et al., *Research and Development on Therapeutic Agents and Vaccines for COVID-19 and Related Human Coronavirus Diseases*, ACS Publications, Washington, DC, USA, 2020.
- [11] O. M. Lage, M. C. Ramos, R. Calisto, E. Almeida, V. Vasconcelos, and F. Vicente, "Current screening methodologies in drug discovery for selected human diseases," *Marine Drugs*, vol. 16, no. 8, p. 279, 2018.
- [12] A. G. Atanasov, S. B. Zotchev, V. M. Zotchev, C. T. Dirsch, and C. T. Supuran, "Natural products in drug discovery: advances and opportunities," *Nature Reviews Drug Discovery*, vol. 20, no. 3, pp. 200–216, 2021.
- [13] D. Moher, D. G. Altman, A. Liberati, and J. Tetzlaff, "PRISMA statement," *Epidemiology*, vol. 22, no. 1, p. 128, 2011.
- [14] S. Surabhi and B. Singh, "Computer aided drug design: an overview," *Journal of Drug Delivery and Therapeutics*, vol. 8, no. 5, pp. 504–509, 2018.
- [15] R. T. Kroemer, "Structure-based drug design: docking and scoring," *Current Protein & Peptide Science*, vol. 8, no. 4, pp. 312–328, 2007.

- [16] F. L. Stahura and J. Bajorath, "New methodologies for ligand-based virtual screening," *Current Pharmaceutical Design*, vol. 11, no. 9, pp. 1189–1202, 2005.
- [17] G. Schneider and U. Fechner, "Computer-based de novo design of drug-like molecules," *Nature Reviews Drug Discovery*, vol. 4, no. 8, pp. 649–663, 2005.
- [18] X. Lin, X. Li, and X. Lin, "A review on applications of computational methods in drug screening and design," *Molecules*, vol. 25, no. 6, p. 1375, 2020.
- [19] L. Pinzi and G. Rastelli, "Molecular docking: shifting paradigms in drug discovery," *International Journal of Molecular Sciences*, vol. 20, no. 18, p. 4331, 2019.
- [20] H. Alonso, A. A. Bliznyuk, and J. E. Gready, "Combining docking and molecular dynamic simulations in drug design," *Medicinal Research Reviews*, vol. 26, no. 5, pp. 531–568, 2006.
- [21] P. J. Eddershaw and M. Dickins, "Advances in in vitro drug metabolism screening," *Pharmaceutical Science & Technology Today*, vol. 2, no. 1, pp. 13–19, 1999.
- [22] N. Georgakis, E. Ioannou, C. Varotsou, G. Premetis, E. G. Chronopoulou, and N. E. Labrou, "Determination of half-maximal inhibitory concentration of an enzyme inhibitor," *Targeting Enzymes for Pharmaceutical Development*, Springer, Berlin, Germany, 2020.
- [23] M. Pellegatti, "Preclinical in vivo ADME studies in drug development: a critical review," *Expert Opinion on Drug Metabolism & Toxicology*, vol. 8, no. 2, pp. 161–172, 2012.
- [24] M. A. Abosheasha and A. H. El-Gowily, "Superiority of ciltazolidine among antiplatelet FDA-approved drugs against COVID 19 Mpro and spike protein: drug repurposing approach," *Drug Development Research*, vol. 82, 2020.
- [25] A. Abdul Kadhim, N. Hadi, M. Abdulhussein, M. Abdulhussein, S. Zamil, and S. Zamil, "Preprocessing of the candidate antiviral drugs against COVID-19 in models of SARS cov2 targets," *Prensa Medica Argentina*, vol. 106, p. 2, 2020.
- [26] A. A. Abu-Saleh, I. E. Awad, A. Yadav, and R. A. Poirier, "Discovery of potent inhibitors for SARS-CoV-2's main protease by ligand-based/structure-based virtual screening, MD simulations, and binding energy calculations," *Physical Chemistry Chemical Physics*, vol. 22, no. 40, pp. 23099–23106, 2020.
- [27] I. Achilonu, E. A. Iwuchukwu, O. J. Achilonu, M. A. Fernandes, and Y. Sayed, "Targeting the SARS-CoV-2 main protease using FDA-approved Isavuconazonium, a P2-P3 α -ketoamide derivative and pentagastrin: an in-silico drug discovery approach," *Journal of Molecular Graphics and Modelling*, vol. 101, 2020.
- [28] S. O. Aftab, M. Z. Ghouri, M. U. Masood et al., "Analysis of SARS-CoV-2 RNA-dependent RNA polymerase as a potential therapeutic drug target using a computational approach," *Journal of Translational Medicine*, vol. 18, no. 1, p. 275, 2020.
- [29] K. Ahmadi, A. Farasat, M. Rostamian, B. Johari, and H. Madanchi, "Enfuvirtide, an HIV-1 fusion inhibitor peptide, can act as a potent SARS-CoV-2 fusion inhibitor: an in silico drug repurposing study," *Journal of Biomolecular Structure & Dynamics*, vol. 40, pp. 5566–5576, 2021.
- [30] M. Z. Ahmed, Q. Zia, A. Haque et al., "FDA-approved antiviral and anti-infection agents as potential inhibitors of SARS-CoV-2 main protease: an in silico drug repurposing study," *Journal of Infection and Public Health*, vol. 14, pp. 611–619, 2021.
- [31] J. Anand, T. Ghildiyal, A. Madhwal, R. Bhatt, D. Verma, and N. Rai, "Computational guided approach for drug repurposing against SARS-CoV-2," *Future Virology*, vol. 16, no. 3, 2021.
- [32] I. Ancy, M. Sivanandam, and P. Kumaradhas, "Possibility of HIV-1 protease inhibitors-clinical trial drugs as repurposed drugs for SARS-CoV-2 main protease: a molecular docking, molecular dynamics and binding free energy simulation study," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 5368–5375, 2020.
- [33] M. A. Ansari, Q. M. S. Jamal, S. Rehman et al., "TAT-peptide conjugated repurposing drug against SARS-CoV-2 main protease (3CLpro): potential therapeutic intervention to combat COVID-19," *Arabian Journal of Chemistry*, vol. 13, no. 11, pp. 8069–8079, 2020.
- [34] K. Arun, C. Sharanya, J. Abhithaj, D. Francis, and C. Sadasivan, "Drug repurposing against SARS-CoV-2 using E-pharmacophore based virtual screening, molecular docking and molecular dynamics with main protease as the target," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 4647–4658, 2020.
- [35] R. Arya, A. Das, V. Prashar, and M. Kumar, "Potential inhibitors against papain-like protease of novel coronavirus (COVID-19) from FDA approved drugs," 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c7483f842e655654db2b42>.
- [36] K. Baby, S. Maity, C. H. Mehta, A. Suresh, U. Y. Nayak, and Y. Nayak, "Targeting SARS-CoV-2 main protease: a computational drug repurposing study," *Archives of Medical Research*, vol. 52, no. 1, pp. 38–47, 2021.
- [37] J. D. Baker, R. L. Uhrich, G. C. Kraemer, J. E. Love, and B. C. Kraemer, "A drug repurposing screen identifies hepatitis C antivirals as inhibitors of the SARS-CoV2 main protease," *PLoS One*, vol. 16, no. 2, Article ID e0245962, 2021.
- [38] B. R. Bharath, H. Damle, S. Ganju, and L. Damle, "In silico screening of known small molecules to bind ACE2 specific RBD on spike glycoprotein of SARS-CoV-2 for repurposing against COVID-19," *F1000Research*, vol. 9, p. 663, 2020.
- [39] A. Bhowmik, S. Biswas, S. Hajra, and P. Saha, "In silico validation of potent phytochemical orientin as inhibitor of SARS-CoV-2 spike and host cell receptor GRP78 binding," *Heliyon*, vol. 7, no. 1, 2021.
- [40] K. Bolelli, T. Ertan-Bolelli, O. Unsalan, and C. Altunayar-Unsalan, "Fenoterol and dobutamine as SARS-CoV-2 main protease inhibitors: a virtual screening study," *Journal of Molecular Structure*, vol. 1228, Article ID 129449, 2021.
- [41] C. N. Cavasotto and J. I. Di Filippo, "In silico drug repurposing for COVID-19: targeting SARS-CoV-2 proteins through docking and consensus ranking," *Molecular Informatics*, vol. 40, no. 1, Article ID e2000115, 2021.
- [42] V. Chandel, P. P. Sharma, S. Raj, R. Choudhari, B. Rathi, and D. Kumar, "Structure-based drug repurposing for targeting NSP9 replicase and spike proteins of severe acute respiratory syndrome coronavirus 2," *Journal of Biomolecular Structure and Dynamics*, vol. 40, 2020.
- [43] C. Z. Chen, M. Xu, M. Pradhan et al., "Identifying SARS-CoV-2 entry inhibitors through drug repurposing screens of SARS-S and MERS-S pseudotyped particles," *ACS Pharmacology & Translational Science*, vol. 3, no. 6, pp. 1165–1175, 2020.
- [44] S. K. Chidambaram, D. Ali, S. Alarifi, S. Radhakrishnan, and I. Akbar, "In silico molecular docking: evaluation of coumarin based derivatives against SARS-CoV-2," *Journal of Infection and Public Health*, vol. 13, no. 11, pp. 1671–1677, 2020.

- [45] S. Choudhary, Y. S. Malik, and S. Tomar, "Identification of SARS-CoV-2 cell entry inhibitors by drug repurposing using in silico structure-based virtual screening approach," *Frontiers in Immunology*, vol. 11, p. 1664, 2020.
- [46] C. Clemente, M. Freiburger, S. Ravetti, D. Beltramo, and A. Garro, "An in silico analysis of Ibuprofen enantiomers in high concentrations of sodium chloride with SARS-CoV-2 main protease," *Journal of Biomolecular Structure and Dynamics*, vol. 40, pp. 5653–5664, 2021.
- [47] I. Cosic, D. Cosic, and I. Loncarevic, "Analysis of ivermectin as potential inhibitor of SARS-CoV-2 using resonant recognition model," *International Journal of Sciences*, vol. 10, no. 1, pp. 1–6, 2021.
- [48] S. Daoud, S. J. Alabed, and L. A. Dahabiyeh, "Identification of potential COVID-19 main protease inhibitors using structure-based pharmacophore approach, molecular docking and repurposing studies," *Acta Pharmaceutica*, vol. 71, no. 2, pp. 163–174, 2021.
- [49] O. V. de Oliveira, G. B. Rocha, A. S. Paluch, and L. T. Costa, "Repurposing approved drugs as inhibitors of SARS-CoV-2 S-protein from molecular modeling and virtual screening," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 11, pp. 3924–3933, 2021.
- [50] M. Delijewski and J. Haneczok, "AI drug discovery screening for COVID-19 reveals zafirlukast as a repurposing candidate," *Medicine in Drug Discovery*, vol. 9, Article ID 100077, 2021.
- [51] D. Dey, S. Borkotoky, and M. Banerjee, "In silico identification of tretinoin as a SARS-CoV-2 envelope (E) protein ion channel inhibitor," *Computers in Biology and Medicine*, vol. 127, Article ID 104063, 2020.
- [52] S. Durdagi, "Virtual drug repurposing study against SARS-CoV-2 TMPRSS2 target," *Turkish Journal of Biology*, vol. 44, no. SI-1, pp. 185–191, 2020.
- [53] S. Durdagi, B. Aksoydan, B. Dogan, K. Sahin, A. Shahraki, and N. Birgül-Iyison, "Screening of clinically approved and investigation drugs as potential inhibitors of SARS-CoV-2 main protease and spike receptor-binding domain bound with ACE2 COVID19 target proteins: a virtual drug repurposing study," 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c74aa5702a9b4c1c18b2d4>.
- [54] P. Eleftheriou, D. Amanatidou, A. Petrou, and A. Geronikaki, "In silico evaluation of the effectivity of approved protease inhibitors against the main protease of the novel SARS-CoV-2 virus," *Molecules*, vol. 25, no. 11, p. 2529, 2020.
- [55] A. D. Elmezayen, A. Al-Obaidi, A. T. Şahin, and K. Yelekçi, "Drug repurposing for coronavirus (COVID-19): in silico screening of known drugs against coronavirus 3CL hydrolase and protease enzymes," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 8, pp. 2980–2992, 2021.
- [56] J. A. Encinar and J. A. Menendez, "Potential drugs targeting early innate immune evasion of SARS-coronavirus 2 via 2'-O-methylation of viral RNA," *Viruses*, vol. 12, no. 5, 2020.
- [57] A. Farag, P. Wang, M. Ahmed, and H. Sadek, "Identification of FDA approved drugs targeting COVID-19 virus by structure-based drug repositioning," 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c74b2a567dfe0f38ec4ee7>.
- [58] S. Feng, X. Luan, Y. Wang et al., "Eltrombopag is a potential target for drug intervention in SARS-CoV-2 spike protein," *Infection, Genetics and Evolution: Journal of Molecular Epidemiology and Evolutionary Genetics in Infectious Diseases*, vol. 85, Article ID 104419, 2020.
- [59] W. R. Ferraz, R. A. Gomes, A. L. S. Novaes, and G. H. Goulart Trossini, "Ligand and structure-based virtual screening applied to the SARS-CoV-2 main protease: an in silico repurposing study," *Future Medicinal Chemistry*, vol. 12, no. 20, pp. 1815–1828, 2020.
- [60] A. Fischer, M. Sellner, S. Naranjan, M. Smieško, and M. A. Lill, "Potential inhibitors for novel coronavirus protease identified by virtual screening of 606 million compounds," *International Journal of Molecular Sciences*, vol. 21, no. 10, 2020.
- [61] A. Gimeno, J. Mestres-Truyol, M. J. Ojeda-Montes et al., "Prediction of novel inhibitors of the main protease (M-pro) of SARS-CoV-2 through consensus docking and drug repositioning," *International Journal of Molecular Sciences*, vol. 21, no. 11, 2020.
- [62] K. Guo, Z. Wang, P. Gao et al., "Identification of repurposal drugs and adverse drug reactions for various courses of coronavirus disease 2019 (COVID-19) based on single-cell RNA sequencing data," 2020, <https://arxiv.org/abs/2005.07856>.
- [63] A. Gupta, C. Rani, P. Pant et al., "Structure-based virtual screening and biochemical validation to discover a potential inhibitor of the SARS-CoV-2 main protease," *ACS Omega*, vol. 5, no. 51, pp. 33151–33161, 2020.
- [64] T. Huynh, W. Cornell, and B. Luan, "In silico exploration of inhibitors for SARS-CoV-2's papain-like protease," *Frontiers in Chemistry*, vol. 8, Article ID 624163, 2021.
- [65] M. A. Ibrahim, A. H. Abdelrahman, and M.-E. F. Hegazy, "In-silico drug repurposing and molecular dynamics puzzled out potential SARS-CoV-2 main protease inhibitors," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 5756–5767, 2020.
- [66] H. Iftikhar, H. N. Ali, S. Farooq, H. Naveed, and S. Shahzad-Ul-Hussan, "Identification of potential inhibitors of three key enzymes of SARS-CoV2 using computational approach," *Computers in Biology and Medicine*, vol. 122, Article ID 103848, 2020.
- [67] R. Jain and S. Mujwar, "Repurposing metocurine as main protease inhibitor to develop novel antiviral therapy for COVID-19," *Structural Chemistry*, vol. 31, no. 6, pp. 2487–2499, 2020.
- [68] M. A. Jarvis, F. A. Hansen, K. Rosenke et al., "Evaluation of drugs for potential repurposing against COVID-19 using a tier-based scoring system," *Antiviral Therapy*, vol. 25, no. 4, pp. 223–231, 2020.
- [69] O. Kadioglu, M. Saeed, H. J. Greten, and T. Efferth, "Identification of novel compounds against three targets of SARS CoV-2 coronavirus by combined virtual screening and supervised machine learning," *Computers in Biology and Medicine*, vol. 133, Article ID 104359, 2021.
- [70] M. Kandeel, A. H. Abdelrahman, K. Oh-Hashi et al., "Repurposing of FDA-approved antivirals, antibiotics, anthelmintics, antioxidants, and cell protectives against SARS-CoV-2 papain-like protease," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 5129–5136, 2020.
- [71] S. Kandwal and D. Fayne, "Repurposing drugs for treatment of SARS-CoV-2 infection: computational design insights into mechanisms of action," *Journal of Biomolecular Structure and Dynamics*, vol. 40, pp. 1316–1330, 2020.
- [72] A. Khan, S. S. Ali, M. T. Khan et al., "Combined drug repurposing and virtual screening strategies with molecular dynamics simulation identified potent inhibitors for SARS-CoV-2 main protease (3CLpro)," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 4659–4670, 2020.

- [73] M. A. Khan, S. Mahmud, A. Ul Alam, M. E. Rahman, F. Ahmed, and M. Rahmatullah, "Comparative molecular investigation of the potential inhibitors against SARS-CoV-2 main protease: a molecular docking study," *Journal of Biomolecular Structure & Dynamics*, vol. 39, 2020.
- [74] V. L. Kouznetsova, A. Zhang, M. Tatineni, M. A. Miller, and I. F. Tsigelny, "Potential COVID-19 papain-like protease PLpro inhibitors: repurposing FDA-approved drugs," *PeerJ*, vol. 8, Article ID e9965, 2020.
- [75] B. Krishnaprasad, S. Maity, C. H. Mehta, A. Suresh, U. Y. Nayak, and Y. Nayak, "Targeting SARS-CoV-2 RNA-dependent RNA polymerase: an in silico drug repurposing for COVID-19," *F1000 Research*, vol. 9, pp. 1–23, 2020.
- [76] D. Kumar, V. Chandel, S. Raj, and B. Rath, "In silico identification of potent FDA approved drugs against coronavirus COVID-19 main protease: a drug repurposing approach," *Chemical Biology Letters*, vol. 7, no. 3, pp. 166–175, 2020.
- [77] V. Kumar, J. K. Dhanjal, S. C. Kaul, R. Wadhwa, and D. Sundar, "Withanone and caffeic acid phenethyl ester are predicted to interact with main protease (M-pro) of SARS-CoV-2 and inhibit its activity," *Journal of Biomolecular Structure & Dynamics*, vol. 39, 2020.
- [78] Y. Kumar, H. Singh, and C. N. Patel, "In silico prediction of potential inhibitors for the main protease of SARS-CoV-2 using molecular docking and dynamics simulation based drug-repurposing," *Journal of Infection and Public Health*, vol. 13, no. 9, pp. 1210–1223, 2020.
- [79] P. Kumar, T. Bhardwaj, A. Kumar et al., "Reprofiling of approved drugs against SARS-CoV-2 main protease: an in-silico study," *Journal of Biomolecular Structure & Dynamics*, vol. 40, pp. 3170–3184, 2020.
- [80] Z. Li, X. Li, Y. Y. Huang et al., "Identify potent SARS-CoV-2 main protease inhibitors via accelerated free energy perturbation-based virtual screening of existing drugs," *Proceedings of the National Academy of Sciences*, vol. 117, no. 44, pp. 27381–27387, 2020.
- [81] H. Liang, L. Zhao, X. Gong, M. Hu, and H. Wang, "Virtual screening FDA approved drugs against multiple targets of SARS-CoV-2," *Clinical and Translational Science*, vol. 14, 2021.
- [82] K. B. Lokhande, S. Doiphode, R. Vyas, and K. V. Swamy, "Molecular docking and simulation studies on SARS-CoV-2 Mpro reveals mitoxantrone, leucovorin, birinapant, and dynasore as potent drugs against COVID-19," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 7294–7305, 2020.
- [83] S. Mahanta, P. Chowdhury, N. Gogoi et al., "Potential antiviral activity of approved repurposed drug against main protease of SARS-CoV-2: an in silico based approach," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 3802–3811, 2020.
- [84] S. Mahdian, M. Zarrabi, Y. Panahi, and S. Dabbagh, "Repurposing FDA-approved drugs to fight COVID-19 using in silico methods: targeting SARS-CoV-2 RdRp enzyme and host cell receptors (ACE2, CD147) through virtual screening and molecular dynamic simulations," *Informatics in Medicine Unlocked*, vol. 23, 2021.
- [85] B. N. Marak, J. Dowarah, L. Khiangte, and V. P. Singh, "Step toward repurposing drug discovery for COVID-19 therapeutics through in silico approach," *Drug Development Research*, vol. 82, no. 3, pp. 374–392, 2020.
- [86] S. Mohapatra, P. Nath, M. Chatterjee et al., "Repurposing therapeutics for COVID-19: rapid prediction of commercially available drugs through machine learning and docking," *PLoS One*, vol. 15, no. 11, Article ID e0241543, 2020.
- [87] Z. Molavi, S. Razi, S. A. Mirmotalebisohi et al., "Identification of FDA approved drugs against SARS-CoV-2 RNA dependent RNA polymerase (RdRp) and 3-chymotrypsin-like protease (3CLpro), drug repurposing approach," *Bio-medicine & Pharmacotherapy*, vol. 138, Article ID 111544, 2021.
- [88] N. S. Mulgaonkar, H. Wang, S. Mallawarachchi, D. Ruzek, B. Martina, and S. Fernando, "Bcr-Abl tyrosine kinase inhibitor imatinib as a potential drug for COVID-19," 2020, <https://www.biorxiv.org/content/10.1101/2020.06.18.158196v2>.
- [89] C. J. Mycroft-West, D. Su, I. Pagani et al., "Heparin inhibits cellular invasion by SARS-CoV-2: structural dependence of the interaction of the spike S1 receptor-binding domain with heparin," *Thrombosis and Haemostasis*, vol. 120, no. 12, pp. 1700–1715, 2020.
- [90] A. Nayariseri, R. Khandelwal, M. Madhavi et al., "Shape-based machine learning models for the potential novel COVID-19 protease inhibitors assisted by molecular dynamics simulation," *Current Topics in Medicinal Chemistry*, vol. 20, no. 24, pp. 2146–2167, 2020.
- [91] H. A. Odhar, S. W. Ahjel, A. A. M. A. Albeer, A. F. Hashim, A. M. Rayshan, and S. S. Humadi, "Molecular docking and dynamics simulation of FDA approved drugs with the main protease from 2019 novel coronavirus," *Bioinformation*, vol. 16, no. 3, pp. 236–244, 2020.
- [92] J. T. Ortega, M. L. Serrano, and B. Jastrzebska, "Class A G protein-coupled receptor antagonist famotidine as a therapeutic alternative against SARS-CoV2: an in silico analysis," *Biomolecules*, vol. 10, no. 6, p. 954, 2020.
- [93] P. Pandey, F. Khan, A. K. Rana, Y. Srivastava, S. K. Jha, and N. K. Jha, "A drug repurposing approach towards elucidating the potential of flavonoids as COVID-19 spike protein inhibitors," *Biointerface Research in Applied Chemistry*, vol. 11, no. 1, pp. 8482–8501, 2021.
- [94] S. Parveen and R. B. Alnoman, "Potential exploration of recent FDA-approved anticancer drugs against models of SARS-CoV-2's main protease and spike glycoprotein: a computational study," *Biointerface Research in Applied Chemistry*, vol. 11, no. 3, pp. 10059–10073, 2021.
- [95] K. A. Peele, C. Potla Durthi, T. Srihansa et al., "Molecular docking and dynamic simulations for antiviral compounds against SARS-CoV-2: a computational study," *Informatics in Medicine Unlocked*, vol. 19, Article ID 100345, 2020.
- [96] L. Pinzi, A. Tinivella, F. Caporuscio, and G. Rastelli, "Drug repurposing and polypharmacology to fight SARS-CoV-2 through inhibition of the main protease," *Frontiers in Pharmacology*, vol. 12, Article ID 636989, 2021.
- [97] R. Pokhrel, P. Chapagain, and J. Siltberg-Liberles, "Potential RNA-dependent RNA polymerase inhibitors as prospective therapeutics against SARS-CoV-2," *Journal of Medical Microbiology*, vol. 69, no. 6, pp. 864–873, 2020.
- [98] A. K. Ray, P. S. S. Gupta, S. K. Panda, S. Biswal, and M. K. Rana, "Repurposing of FDA approved drugs for the identification of potential inhibitors of SARS-CoV-2 main protease," 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c74b60842e659a0edb3136>.
- [99] C. Sachdeva, A. Wadhwa, A. Kumari, F. Hussain, P. Jha, and N. K. Kaushik, "In silico potential of approved antimalarial drugs for repurposing against COVID-19," *OMICS: A*

- Journal of Integrative Biology*, vol. 24, no. 10, pp. 568–580, 2020.
- [100] P. Sang, S.-H. Tian, Z.-H. Meng, and L.-Q. Yang, “Anti-HIV drug repurposing against SARS-CoV-2,” *RSC Advances*, vol. 10, no. 27, pp. 15775–15783, 2020.
 - [101] S. Saxena, K. Meher, M. Rotella et al., “Identification of SGLT2 inhibitor ertugliflozin as a treatment for COVID-19 using computational and experimental paradigm,” 2021, <https://www.biorxiv.org/content/10.1101/2021.06.18.448921v1>.
 - [102] H. Setianingsih, E. Yatmasari, M. F. Ilmawan, H. Zahroh, and E. D. P. Lestari, “Targeting multiple SARS-CoV-2 and human proteins: in silico approach for COVID-19 drug repurposing,” *International Journal of Pharmaceutical Research*, vol. 12, no. 2, pp. 3184–3192, 2020.
 - [103] B. Shah, P. Modi, and S. R. Sagar, “In silico studies on therapeutic agents for COVID-19: drug repurposing approach,” *Life Sciences*, vol. 252, Article ID 117652, 2020.
 - [104] S. K. Sharma and A. K. Mishra, “Analysis of therapeutic targets for COVID 19-nCoV/SARS-CoV-2 and discovery by virtual screening methods,” *TANG (Humanitas Medicine)*, vol. 10, no. 2, p. 14, 2020.
 - [105] N. Shekhar, P. Sarma, M. Prajapat et al., “In silico structure-based repositioning of approved drugs for spike glycoprotein S2 domain fusion peptide of SARS-CoV-2: rationale from molecular dynamics and binding free energy calculations,” *mSystems*, vol. 5, no. 5, 2020.
 - [106] S. K. Singh, A. K. Upadhyay, and M. S. Reddy, “Screening of potent drug inhibitors against SARS-CoV-2 RNA polymerase: an in silico approach,” *3 Biotech*, vol. 11, no. 2, pp. 1–13, 2021.
 - [107] M. Sinha, A. Gupta, S. Gupta et al., “Analogue discovery of safer alternatives to HCQ and CQ drugs for SAR-CoV-2 by computational design,” *Computers in Biology and Medicine*, vol. 130, Article ID 104222, 2021.
 - [108] H. Soni, D. Gautam, S. Sharma, and J. Malik, “Rifampicin as potent inhibitor of COVID-19 main protease: in-silico docking approach,” *Saudi Journal of Medical and Pharmaceutical Sciences*, vol. 6, no. 9, pp. 588–593, 2020.
 - [109] A. Tariq, R. M. Mateen, M. S. Afzal, and M. Saleem, “Paromomycin: a potential dual targeted drug effectively inhibits both spike (S1) and main protease of COVID-19,” *International Journal of Infectious Diseases*, vol. 98, pp. 166–175, 2020.
 - [110] G. Tatar, E. Ozyurt, and K. Turhan, “Computational drug repurposing study of the RNA binding domain of SARS-CoV-2 nucleocapsid protein with antiviral agents,” *Bio-technology Progress*, vol. 37, no. 2, Article ID e3110, 2021.
 - [111] E. Tejera, C. R. Munteanu, A. López-Cortés, A. Cabrera-Andrade, and Y. Pérez-Castillo, “Drugs repurposing using QSAR, docking and molecular dynamics for possible inhibitors of the SARS-CoV-2 Mpro protease,” *Molecules*, vol. 25, no. 21, 2020.
 - [112] K. Terali, B. Baddal, and H. O. Gülcan, “Prioritizing potential ACE2 inhibitors in the COVID-19 pandemic: insights from a molecular mechanics-assisted structure-based virtual screening experiment,” *Journal of Molecular Graphics and Modelling*, vol. 100, Article ID 107697, 2020.
 - [113] A. Trezza, D. Iovinelli, A. Santucci, F. Prisch, and O. Spiga, “An integrated drug repurposing strategy for the rapid identification of potential SARS-CoV-2 viral inhibitors,” *Scientific Reports*, vol. 10, no. 1, Article ID 13866, 2020.
 - [114] O. M. Ugurel, O. Mutlu, E. Sariyer et al., “Evaluation of the potency of FDA-approved drugs on wild type and mutant SARS-CoV-2 helicase (NSP13),” *International Journal of Biological Macromolecules*, vol. 163, pp. 1687–1696, 2020.
 - [115] S. Unni, S. Aouti, S. Thiagarajan, and B. Padmanabhan, “Identification of a repurposed drug as an inhibitor of spike protein of human coronavirus SARS-CoV-2 by computational methods,” *Journal of Biosciences*, vol. 45, no. 1, 2020.
 - [116] C. Vaishali, P. Prakash, S. Sharma, B. Raj, and D. Rath, “Structure based drug repurposing through targeting NSP9 replicase and spike proteins of SARS-CoV-2,” 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c74b26337d6c3f76e27a44>.
 - [117] D. Verma, S. Kapoor, S. Das, and K. Thakur, “Potential inhibitors of SARS-CoV-2 main protease (M Pro) identified from the library of FDA approved drugs using molecular docking studies,” 2020, <https://www.preprints.org/manuscript/202004.0149/v1>.
 - [118] T. Z. Wei, H. Wang, X. Q. Wu et al., “In silico screening of potential spike glycoprotein inhibitors of SARS-CoV-2 with drug repurposing strategy,” *Chinese Journal of Integrative Medicine*, vol. 26, no. 9, pp. 663–669, 2020.
 - [119] H. Xu, B. Liu, Z. Xiao et al., “Computational and experimental studies reveal that thymoquinone blocks the entry of coronaviruses into in vitro cells,” *Infectious Diseases and Therapy*, vol. 10, no. 1, pp. 483–494, 2021.
 - [120] I. Abdelli, F. Hassani, S. Bekkel Brikci, and S. Ghalem, “In silico study the inhibition of angiotensin converting enzyme 2 receptor of COVID-19 by Ammoides verticillata components harvested from western Algeria,” *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 9, pp. 3263–3276, 2021.
 - [121] M. E. Abouelela, H. K. Assaf, R. A. Abdelhamid et al., “Identification of potential SARS-CoV-2 main protease and spike protein inhibitors from the genus aloe: an in silico study for drug development,” *Molecules*, vol. 26, no. 6, p. 1767, 2021.
 - [122] S. Adem, V. Eyupoglu, I. Sarfraz, A. Rasul, and M. Ali, “Identification of potent COVID-19 main protease (Mpro) inhibitors from natural polyphenols: an in silico strategy unveils a hope against CORONA,” 2020, <https://www.preprints.org/manuscript/202003.0333/v1>.
 - [123] A. E. Allam, H. K. Assaf, H. A. Hassan, K. Shimizu, and Y. A. M. M. Elshaier, “Anin silicoperception for newly isolated flavonoids from peach fruit as privileged avenue for a countermeasure outbreak of COVID-19,” *RSC Advances*, vol. 10, no. 50, pp. 29983–29998, 2020.
 - [124] A. G. Al-Sehemi, F. A. Olotu, D. Sanal Dev et al., “Natural products database screening for the discovery of naturally occurring SARS-cov-2 spike glycoprotein blockers,” *ChemistrySelect*, vol. 5, no. 42, Article ID 13309, 2020.
 - [125] G. H. Attia, Y. S. Moemen, M. Youns, A. M. Ibrahim, R. Abdou, and M. A. El Raey, “Antiviral zinc oxide nanoparticles mediated by hesperidin and in silico comparison study between antiviral phenolics as anti-SARS-CoV-2,” *Colloids and Surfaces B: Biointerfaces*, vol. 203, Article ID 111724, 2021.
 - [126] K. F. Azim, S. R. Ahmed, A. Banik, M. M. R. Khan, A. Deb, and S. R. Somana, “Screening and druggability analysis of some plant metabolites against SARS-CoV-2: an integrative computational approach,” *Informatics in Medicine Unlocked*, vol. 20, Article ID 100367, 2020.
 - [127] B. Çakır, B. Okuyan, G. Şener, and T. Tunali-Akbay, “Investigation of beta-lactoglobulin derived bioactive peptides against SARS-CoV-2 (COVID-19): in silico analysis,” *European Journal of Pharmacology*, vol. 891, 2021.

- [128] S. Chatterjee, S. Maiti, A. Banerjee, and M. Kanwar, "In-silico study of hesperidin, epigallocatechin (EGCG), kaempferol and quercetin," 2021, https://assets.researchsquare.com/files/rs-525091/v1_covered.pdf?c=1631867322.
- [129] R. V. Chikhale, S. S. Gurav, R. B. Patil et al., "SARS-CoV-2 host entry and replication inhibitors from Indian ginseng: an in-silico approach," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 4510–4521, 2020.
- [130] R. Chikhale, S. K. Sinha, M. Wanjari et al., "Computational assessment of saikosaponins as adjuvant treatment for COVID-19: molecular docking, dynamics, and network pharmacology analysis," *Molecular Diversity*, vol. 25, pp. 1889–1904, 2021.
- [131] R. V. Chikhale, S. K. Sinha, R. B. Patil et al., "In-silico investigation of phytochemicals from *Asparagus racemosus* as plausible antiviral agent in COVID-19," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 14, 2020.
- [132] P. Chowdhury, "In silico investigation of phytoconstituents from Indian medicinal herb "*Tinospora cordifolia* (giloy)" against SARS-CoV-2 (COVID-19) by molecular dynamics approach," *Journal of Biomolecular Structure and Dynamics*, vol. 39, pp. 6792–6809, 2020.
- [133] M. A. Dahab, M. M. Hegazy, and H. S. Abbass, "Hordatines as a potential inhibitor of COVID-19 main protease and RNA polymerase: an in-silico approach," *Natural Products and Bioprospecting*, vol. 10, no. 6, pp. 453–462, 2020.
- [134] S. Alhumaid, A. Al Mutair, Z. Al Alawi, N. Alhameed, A. R. Z. Zaidi, and M. Tobaiqy, "Efficacy and safety of lopinavir/ritonavir for treatment of COVID-19: a systematic review and meta-analysis," *Tropical Medicine and Infectious Disease*, vol. 5, no. 4, p. 180, 2020.
- [135] L. Ang, E. Song, H. W. Lee, and M. S. Lee, "Herbal medicine for the treatment of coronavirus disease 2019 (COVID-19): a systematic review and meta-analysis of randomized controlled trials," *Journal of Clinical Medicine*, vol. 9, no. 5, p. 1583, 2020.
- [136] P. Das, R. Majumder, M. Mandal, and P. Basak, "In-silico approach for identification of effective and stable inhibitors for COVID-19 main protease (Mpro) from flavonoid based phytochemical constituents of *Calendula officinalis*," *Journal of Biomolecular Structure & Dynamics*, vol. 39, no. 16, pp. 1–16, 2020.
- [137] P. Debnath, B. Debnath, S. Bhaumik, and S. Debnath, "In silico identification of potential inhibitors of ADP-ribose phosphatase of SARS-CoV-2 NSP3 by combining E-pharmacophore- and receptor-based virtual screening of database," *ChemistrySelect*, vol. 5, no. 30, pp. 9388–9398, 2020.
- [138] S. A. Dev and I. Kaur, "Bioactive molecules from eucalyptus essential oil as potential inhibitors of COVID 19 corona virus infection by molecular docking studies," *Kragujevac Journal of Science*, vol. 42, pp. 29–43, 2020.
- [139] C. E. Duru, I. A. Duru, and A. E. Adegboyega, "In silico identification of compounds from *Nigella sativa* seed oil as potential inhibitors of SARS-CoV-2 targets," *Bulletin of the National Research Centre*, vol. 45, no. 1, pp. 1–13, 2021.
- [140] A. El-Demerdash, A. M. Metwaly, A. Hassan et al., "Comprehensive virtual screening of the antiviral potentialities of marine polycyclic guanidine alkaloids against sars-cov-2 (Covid-19)," *Biomolecules*, vol. 11, no. 3, pp. 1–25, 2021.
- [141] O. O. Elekofehinti, O. Iwaloye, S. S. Josiah, A. O. Lawal, M. O. Akinjiyan, and E. O. Ariyo, "Molecular docking studies, molecular dynamics and ADME/tox reveal therapeutic potentials of STOCKIN-69160 against papain-like protease of SARS-CoV-2," *Molecular Diversity*, vol. 25, pp. 1761–1773, 2020.
- [142] S. S. El-Hawary, R. Mohammed, H. S. Bahr et al., "Soybean-associated endophytic fungi as potential source for anti-COVID-19 metabolites supported by docking analysis," *Journal of Applied Microbiology*, vol. 131, no. 3, pp. 1193–1211, 2021.
- [143] M. Emirik, "Potential therapeutic effect of turmeric contents against SARS-CoV-2 compared with experimental COVID-19 therapies: in silico study," *Journal of Biomolecular Structure & Dynamics*, vol. 40, pp. 2024–2037, 2020.
- [144] Z. Fakhar, B. Faramarzi, S. Pacifico, and S. Faramarzi, "Anthocyanin derivatives as potent inhibitors of SARS-CoV-2 main protease: an in-silico perspective of therapeutic targets against COVID-19 pandemic," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 6171–6183, 2020.
- [145] V. A. Falade, T. I. Adelusi, I. O. Adedotun, M. Abdul-Hammed, T. A. Lawal, and S. A. Agboluaje, "In silico investigation of saponins and tannins as potential inhibitors of SARS-CoV-2 main protease (Mpro)," *Silico Pharmacology*, vol. 9, no. 1, 2021.
- [146] I. N. Fitriani, W. Utami, A. T. Zikri, and P. Santoso, "In silico approach of potential phytochemical inhibitor from *Moringa oleifera*, *Cocos nucifera*, *Allium cepa*, *Psidium guajava*, and *Eucalyptus globulus* for the treatment of COVID-19 by molecular docking," 2020, <https://www.scienceopen.com/document?vid=45020f08-2651-475b-b2bc-ee16a177515>.
- [147] S. Gangadevi, V. N. Badavath, A. Thakur et al., "Kobophenol A inhibits binding of host ACE2 receptor with spike rbd domain of SARS-CoV-2, a lead compound for blocking COVID-19," *The journal of physical chemistry letters*, vol. 12, no. 7, pp. 1793–1802, 2021.
- [148] K. Gangarapu, P. Sathiyarajeswaran, K. Kanakavalli et al., "In silico computational screening of kabasura kudineer-official siddha formulation and JACOM-novel herbal coded formulation against SARS-CoV-2 spike protein," 2020, https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3574412.
- [149] R. Ghosh, A. Chakraborty, A. Biswas, and S. Chowdhuri, "Evaluation of green tea polyphenols as novel corona virus (SARS CoV-2) main protease (Mpro) inhibitors—an in silico docking and molecular dynamics simulation study," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 12, 2020.
- [150] R. Ghosh, A. Chakraborty, A. Biswas, and S. Chowdhuri, "Identification of alkaloids from *Justicia adhatoda* as potent SARS CoV-2 main protease inhibitors: an in silico perspective," *Journal of Molecular Structure*, vol. 1229, Article ID 129489, 2021.
- [151] U. S. Gorla, G. K. Rao, U. S. Kulandaivelu, R. R. Alavala, and S. P. Panda, "Lead finding from selected flavonoids with antiviral (SARS-CoV-2) potentials against COVID-19: an in-silico evaluation," *Combinatorial Chemistry & High Throughput Screening*, vol. 24, 2020.
- [152] A. B. Gurung, M. A. Ali, J. Lee, M. A. Farah, and K. M. Al-Anazi, "Unravelling lead antiviral phytochemicals for the inhibition of SARS-CoV-2 Mpro enzyme through in silico approach," *Life Sciences*, vol. 255, Article ID 117831, 2020.
- [153] G. A. Gyebi, O. B. Ogunro, A. P. Adegunloye, O. M. Ogunyemi, and S. O. Afolabi, "Potential inhibitors of coronavirus 3-chymotrypsin-like protease (3CLpro): an in silico screening of alkaloids and terpenoids from African medicinal plants," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 9, pp. 3396–3408, 2021.

- [154] B. Elwakil, M. M. Shaaban, A. A. Bekhit, M. Y. El-Naggar, and Z. A. Olama, "Potential anti-COVID-19 activity of Egyptian propolis using computational modeling," *Future Virology*, vol. 16, no. 2, pp. 107–116, 2021.
- [155] A. Hasan, R. Al Mahamud, K. Jannat et al., "Phytochemicals from *Solanum surattense* Burm. f. have high binding affinities for C-3 like main protease of COVID-19 (SARS-CoV-2)," *Journal of Medicinal Plants Studies*, vol. 8, no. 4, pp. 20–26, 2020.
- [156] H. Hashem, "In silico approach of some selected honey constituents as SARS-CoV-2 main protease (COVID-19) inhibitors," 2020, <https://chemrxiv.org/engage/chemrxiv/article-details/60c749e1842e651579db2e34>.
- [157] M. A. A. Ibrahim, K. A. A. Abdeljawaad, A. H. M. Abdelrahman, and M. E. F. Hegazy, "Natural-like products as potential SARS-CoV-2 Mpro inhibitors: in-silico drug discovery," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 5722–5734, 2020.
- [158] M. A. A. Ibrahim, A. H. M. Abdelrahman, T. A. Hussien et al., "In silico drug discovery of major metabolites from spices as SARS-CoV-2 main protease inhibitors," *Computers in Biology and Medicine*, vol. 126, Article ID 104046, 2020.
- [159] M. A. Isa, A. Mustapha, S. Qazi et al., "In silico molecular docking and molecular dynamic simulation of potential inhibitors of 3C-like main proteinase (3CLpro) from severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) using selected African medicinal plants," *Advances in Traditional Medicine*, vol. 22, 2020.
- [160] E. S. Istifli, P. A. Netz, A. Sihoglu Tepe, M. T. Husunet, C. Sarikurkcu, and B. Tepe, "In silico analysis of the interactions of certain flavonoids with the receptor-binding domain of 2019 novel coronavirus and cellular proteases and their pharmacokinetic properties," *Journal of Biomolecular Structure and Dynamics*, vol. 40, 2020.
- [161] J. T. Jan, T. R. Cheng, Y. P. Juang et al., "Identification of existing pharmaceuticals and herbal medicines as inhibitors of SARS-CoV-2 infection," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 118, no. 5, 2021.
- [162] S. Jo, S. Kim, D. Y. Kim, M. S. Kim, and D. H. Shin, "Flavonoids with inhibitory activity against SARS-CoV-2 3CLpro," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 35, no. 1, pp. 1539–1544, 2020.
- [163] R. S. Joshi, S. S. Jagdale, S. B. Bansode et al., "Discovery of potential multi-target-directed ligands by targeting host-specific SARS-CoV-2 structurally conserved main protease," *Journal of Biomolecular Structure & Dynamics*, vol. 39, no. 9, pp. 3099–3114, 2021.
- [164] N. N. Junior, I. A. Santos, B. A. Meireles et al., "In silico evaluation of lapachol derivatives binding to the NSP9 of SARS-CoV-2," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2021.
- [165] P. Kar, V. Kumar, B. Vellingiri et al., "Anisotine and amarogentin as promising inhibitory candidates against SARS-CoV-2 proteins: a computational investigation," *Journal of Biomolecular Structure & Dynamics*, vol. 40, pp. 4532–4542, 2020.
- [166] I. Khalifa, A. Nawaz, R. Sobhy, S. A. Althawb, and H. Barakat, "Polyacylated anthocyanins constructively network with catalytic dyad residues of 3CLpro of 2019-nCoV than monomeric anthocyanins: a structural-relationship activity study with 10 anthocyanins using in-silico approaches," *Journal of Molecular Graphics and Modelling*, vol. 100, 2020.
- [167] I. Khalifa, W. Zhu, H. H. H. Mohammed, K. Dutta, and C. M. Li, "Tannins inhibit SARS-CoV-2 through binding with catalytic dyad residues of 3CL (pro): an in silico approach with 19 structural different hydrolysable tannins," *Journal of Food Biochemistry*, vol. 44, no. 10, 2020.
- [168] M. T. Khan, A. Ali, Q. Wang et al., "Marine natural compounds as potents inhibitors against the main protease of SARS-CoV-2—a molecular dynamic study," *Journal of Biomolecular Structure and Dynamics*, vol. 39, pp. 3627–3637, 2020.
- [169] T. Kiran Raj, R. Ranjithkumar, B. M. Kanthesh, and T. S. Gopenath, "C-phycocyanin of spirulina plantesis inhibits NSP12 required for replication of sars-COV-2: a novel finding in-silico," *International Journal of Pharmaceutical Sciences and Research*, vol. 11, no. 9, pp. 4271–4278, 2020.
- [170] S. Krupanidhi, K. Abraham Peele, T. Venkateswarulu et al., "Screening of phytochemical compounds of *Tinospora cordifolia* for their inhibitory activity on SARS-CoV-2: an in silico study," *Journal of Biomolecular Structure & Dynamics*, vol. 39, no. 15, pp. 1–5, 2020.
- [171] A. Kumar, G. Choudhir, S. K. Shukla et al., "Identification of phytochemical inhibitors against main protease of COVID-19 using molecular modeling approaches," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 3760–3770, 2020.
- [172] N. Kumar, A. Singh, H. K. Gulati et al., "Phytoconstituents from ten natural herbs as potent inhibitors of main protease enzyme of SARS-COV-2: in silico study," *Phytomedicine*, vol. 1, no. 4, Article ID 100083, 2021.
- [173] B. Kumar, P. Parasuraman, T. P. K. Murthy, M. Murahari, and V. Chandramohan, "In silico screening of therapeutic potentials from *Strychnos nux-vomica* against the dimeric main protease (M^{Pro}) structure of SARS-CoV-2," *Journal of Biomolecular Structure and Dynamics*, vol. 40, pp. 1–19, 2021.
- [174] Y. Li, F. Chu, P. Li et al., "Potential effect of Maxing Shigan decoction against coronavirus disease 2019 (COVID-19) revealed by network pharmacology and experimental verification," *Journal of Ethnopharmacology*, vol. 271, Article ID 113854, 2021.
- [175] S. Maiti and A. Banerjee, "Epigallocatechin gallate and theaflavin gallate interaction in SARS-CoV-2 spike-protein central channel with reference to the hydroxychloroquine interaction: bioinformatics and molecular docking study," *Drug Development Research*, vol. 82, no. 1, pp. 86–96, 2021.
- [176] S. Mahmud, M. A. R. Uddin, G. K. Paul et al., "Virtual screening and molecular dynamics simulation study of plant-derived compounds to identify potential inhibitors of main protease from SARS-CoV-2," *Briefings in Bioinformatics*, vol. 22, no. 2, pp. 1402–1414, 2021.
- [177] S. Mahmud, M. A. R. Uddin, M. Zaman et al., "Molecular docking and dynamics study of natural compound for potential inhibition of main protease of SARS-CoV-2," *Journal of Biomolecular Structure & Dynamics*, vol. 39, no. 16, pp. 1–9, 2020.
- [178] F. Mesli, M. Ghalem, I. Daoud, and S. Ghalem, "Potential inhibitors of angiotensin converting enzyme 2 receptor of COVID-19 by *Corchorus olitorius* Linn using docking, molecular dynamics, conceptual DFT investigation and pharmacophore mapping," *Journal of Biomolecular Structure and Dynamics*, vol. 40, no. 16, 2021.
- [179] M. Mohammadi, S. Rajabi, A. Piroozmand, and S. A. Mirhosseini, "In silico study of pacific oyster antiviral polypeptides as potential inhibitory compounds for SARS-

- CoV-2 main protease," *Jentashapir Journal of Cellular and Molecular Biology*, vol. 11, no. 3, 2020.
- [180] N. A. Murugan, C. J. Pandian, and J. Jeyakanthan, "Computational investigation on *Andrographis paniculata* phytochemicals to evaluate their potency against SARS-CoV-2 in comparison to known antiviral compounds in drug trials," *Journal of biomolecular structure & dynamics*, vol. 39, pp. 4415–4426, 2020.
- [181] B. Naik, N. Gupta, R. Ojha, S. Singh, V. K. Prajapati, and D. Prusty, "High throughput virtual screening reveals SARS-CoV-2 multi-target binding natural compounds to lead instant therapy for COVID-19 treatment," *International Journal of Biological Macromolecules*, vol. 40, pp. 1–17, 2020.
- [182] R. R. Narkhede, A. V. Pise, R. S. Cheke, and S. D. Shinde, "Recognition of natural products as potential inhibitors of COVID-19 main protease (Mpro): in-silico evidences," *Natural Products and Bioprospecting*, vol. 10, no. 5, pp. 297–306, 2020.
- [183] R. Nivetha, S. Bhuvargavan, T. Muthu Kumar, K. Ramanathan, and S. Janarthanan, "Inhibition of multiple SARS-CoV-2 proteins by an antiviral biomolecule, seselin from *Aegle marmelos* deciphered using molecular docking analysis," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2020.
- [184] O. M. Ogunyemi, G. A. Gyebi, A. A. Elfiky et al., "Alkaloids and flavonoids from African phytochemicals as potential inhibitors of SARS-Cov-2 RNA-dependent RNA polymerase: an in silico perspective," *Antiviral Chemistry and Chemotherapy*, vol. 28, 2020.
- [185] S. Padhi, M. Masi, R. Chourasia, Y. Rajashekar, A. K. Rai, and A. Evidente, "ADMET profile and virtual screening of plant and microbial natural metabolites as SARS-CoV-2 S1 glycoprotein receptor binding domain and main protease inhibitors," *European Journal of Pharmacology*, vol. 890, 2021.
- [186] P. Pandey, J. S. Rane, A. Chatterjee et al., "Targeting SARS-CoV-2 spike protein of COVID-19 with naturally occurring phytochemicals: an in silico study for drug development," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 6306–6316, 2020.
- [187] M. Kumar, K. Sundaram, and M. S. Ramasamy, "Coronavirus spike (S) glycoprotein (2019-Ncov) targeted siddha medicines kabasura kudineer and Thonthasura kudineer â€" in silico evidence for corona viral drug," *Asian Journal of Pharmaceutical Research and Health Care*, vol. 12, no. 1, pp. 20–27, 2020.
- [188] F. Rahman, S. Tabrez, R. Ali, A. S. Alqahtani, M. Z. Ahmed, and A. Rub, "Molecular docking analysis of rutin reveals possible inhibition of SARS-CoV-2 vital proteins," *Journal of Traditional and Complementary Medicine*, vol. 11, no. 2, pp. 173–179, 2021.
- [189] N. Rahman, Z. Basharat, M. Yousuf, G. Castaldo, L. Rastrelli, and H. Khan, "Virtual screening of natural products against type II transmembrane serine protease (TMPRSS2), the priming agent of coronavirus 2 (SARS-CoV-2)," *Molecules*, vol. 25, no. 10, p. 2271, 2020.
- [190] A. Rakib, A. Paul, M. Chy et al., "Biochemical and computational approach of selected phytocompounds from *Tinospora crispa* in the management of COVID-19," *Molecules*, vol. 25, no. 17, p. 3936, 2020.
- [191] D. S. F. Ramadhan, T. M. Fakhri, and A. Arfan, "Activity prediction of bioactive compounds contained in *etlingera elatior* against the SARS-CoV-2 main protease: an in silico approach," *Borneo Journal of Pharmacy*, vol. 3, no. 4, pp. 235–242, 2020.
- [192] P. Rangsinth, C. Sillapachaiyaporn, S. Nilkhet, T. Tencomnao, A. T. Ung, and S. Chuchawankul, "Mushroom-derived bioactive compounds potentially serve as the inhibitors of SARS-CoV-2 main protease: an in silico approach," *Journal of Traditional and Complementary Medicine*, vol. 11, no. 2, pp. 158–172, 2021.
- [193] N. Rivero-Segura and J. Gomez-Verjan, "In silico screening of natural products isolated from Mexican herbal medicines against COVID-19," *Biomolecules*, vol. 11, no. 2, p. 216, 2021.
- [194] C. Selvaraj, D. C. Dinesh, U. Panwar, R. Abhirami, E. Boura, and S. K. Singh, "Structure-based virtual screening and molecular dynamics simulation of SARS-CoV-2 guanine-N7 methyltransferase (NSP14) for identifying antiviral inhibitors against COVID-19," *Journal of biomolecular structure & dynamics*, vol. 39, pp. 4582–4593, 2020.
- [195] A. D. Sharma and I. Kaur, "Eucalyptus essential oil bioactive molecules from against SARS-CoV-2 spike protein: insights from computational studies," 2021, <https://europepmc.org/article/ppr/ppr260910>.
- [196] A. D. Sharma, "Eucalyptol (1, 8 cineole) from eucalyptus essential oil a potential inhibitor of COVID 19 corona virus infection by molecular docking studies," 2020, <https://www.preprints.org/manuscript/202003.0455/v1>.
- [197] A. D. Sharma and I. Kaur, "Molecular docking studies on jensenone from eucalyptus essential oil as a potential inhibitor of COVID 19 corona virus infection," 2020, <https://arxiv.org/abs/2004.00217>.
- [198] M. M. A. K. Shawan, S. K. Halder, and M. A. Hasan, "Luteolin and abyssinone II as potential inhibitors of SARS-CoV-2: an in silico molecular modeling approach in battling the COVID-19 outbreak," *Bulletin of the National Research Centre*, vol. 45, no. 1, pp. 27–21, 2021.
- [199] T. Sindhu, K. Arathi, A. Akhilesh et al., "Antiviral screening of Clerodol derivatives as COV 2 main protease inhibitor in novel corona virus disease: in silico approaches," *Asian Journal of Pharmacy and Technology*, vol. 10, no. 2, pp. 60–64, 2020.
- [200] R. Singh, V. K. Bhardwaj, J. Sharma, R. Purohit, and S. Kumar, "In-silico evaluation of bioactive compounds from tea as potential SARS-CoV-2 nonstructural protein 16 inhibitors," *Journal of Traditional and Complementary Medicine*, vol. 12, 2021.
- [201] S. Singh, M. Fulbabu, A. Sonawane, P. Kar, and S. Sadhukhan, "Plant-derived natural polyphenols as potential antiviral drugs against SARS-CoV-2 via RNA-dependent RNA polymerase (RdRp) inhibition: an in-silico analysis," *Journal of Biomolecular Structure and Dynamics*, vol. 39, 2020.
- [202] R. Srimathi, M. K. Mohan Maruga Raja, and M. K. Kathiravan, "In silico screening of traditional herbal medicine derived chemical constituents for possible potential inhibition against sars-cov-2," *Journal of Natural Remedies*, vol. 20, no. 2, pp. 79–88, 2020.
- [203] A. Subbaiyan, K. Ravichandran, S. V. Singh et al., "In silico molecular docking analysis targeting SARS-CoV-2 Spike protein and selected herbal constituents," *Journal of Pure and Applied Microbiology*, vol. 14, no. 1, pp. 989–998, 2020.
- [204] M. Surti, M. Patel, M. Adnan et al., "Ilimaquinone (marine sponge metabolite) as a novel inhibitor of SARS-CoV-2 key target proteins in comparison with suggested COVID-19 drugs: designing, docking and molecular dynamics simulation study," *RSC Advances*, vol. 10, no. 62, pp. 37707–37720, 2020.

- [205] Q. Tao, J. Du, X. Li et al., "Network pharmacology and molecular docking analysis on molecular targets and mechanisms of Huashi Baidu formula in the treatment of COVID-19," *Drug Development and Industrial Pharmacy*, vol. 46, no. 8, pp. 1345–1353, 2020.
- [206] H. I. Umar, S. S. Josiah, T. P. Saliu, T. O. Jimoh, A. Ajayi, and J. B. Danjuma, "In-silico analysis of the inhibition of the SARS-CoV-2 main protease by some active compounds from selected African plants," *Journal of Taibah University Medical Sciences*, vol. 16, no. 2, pp. 162–176, 2021.
- [207] K. D. Umesh, C. Selvaraj, S. K. Singh, and V. K. Dubey, "Identification of new anti-nCoV drug chemical compounds from Indian spices exploiting SARS-CoV-2 main protease as target," *Journal of Biomolecular Structure & Dynamics*, vol. 39, no. 9, pp. 3428–3434, 2021.
- [208] J. S. Yang, J. H. Chiang, S. Tsai et al., "In silico de novo curcuminoid derivatives from the compound library of natural products research laboratories inhibit COVID-19 3CLpro activity," *Natural Product Communications*, vol. 15, no. 9, 2020.
- [209] J.-w. Yu, L. Wang, and L.-d. Bao, "Exploring the active compounds of traditional Mongolian medicine in intervention of novel coronavirus (COVID-19) based on molecular docking method," *Journal of Functional Foods*, vol. 71, Article ID 104016, 2020.
- [210] D.-h. Zhang, K.-l. Wu, X. Zhang, S.-q. Deng, and B. Peng, "In silico screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus," *Journal of Integrative Medicine*, vol. 18, no. 2, pp. 152–158, 2020.
- [211] S. Abu-Melha, M. M. Edrees, S. M. Riyadh, M. R. Abdelaziz, A. A. Elfiky, and S. M. Gomha, "Clean grinding technique: a facile synthesis and in silico antiviral activity of hydrazones, pyrazoles, and pyrazines bearing Thiazole moiety against SARS-CoV-2 main protease (Mpro)," *Molecules*, vol. 25, no. 19, 2020.
- [212] E. Aghaee, M. Ghodrati, and J. B. Ghasemi, "In silico exploration of novel protease inhibitors against coronavirus 2019 (COVID-19)," *Informatics in Medicine Unlocked*, vol. 23, 2021.
- [213] S. Ahmad, Y. Waheed, S. Ismail, M. H. Najmi, and J. K. Ansari, "Rational design of potent anti-COVID-19 main protease drugs: an extensive multi-spectrum in silico approach," *Journal of Molecular Liquids*, vol. 330, 2021.
- [214] S. Ahmed, R. Mahtarin, S. S. Ahmed et al., "Investigating the binding affinity, interaction, and structure-activity-relationship of 76 prescription antiviral drugs targeting RdRp and Mpro of SARS-CoV-2," *Journal of Biomolecular Structure and Dynamics*, vol. 39, 2020.
- [215] M. Alabboud and A. Javadmanesh, "In silico study of various antiviral drugs, vitamins, and natural substances as potential binding compounds with SARS-CoV-2 main protease," *DYSONA-Life Science*, vol. 1, no. 2, pp. 44–63, 2020.
- [216] M. A. Alamri, M. Tahir ul Qamar, M. U. Mirza, S. M. Alqahtani, M. Froeyen, and L. L. Chen, "Discovery of human coronaviruses pan-papain-like protease inhibitors using computational approaches," *Journal of Pharmaceutical Analysis*, vol. 10, no. 6, pp. 546–559, 2020.
- [217] R. Alexpandi, J. F. De Mesquita, S. K. Pandian, and A. V. Ravi, "Quinolines-based SARS-CoV-2 3CLpro and RdRp inhibitors and Spike-RBD-ACE2 inhibitor for drug-repurposing against COVID-19: an in silico analysis," *Frontiers in Microbiology*, vol. 11, p. 1796, 2020.
- [218] A. G. Al-Sehemi, M. Pannipara, R. S. Parulekar et al., "Potential of NO donor furoxan as SARS-CoV-2 main protease (Mpro) inhibitors: in silico analysis," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 15, pp. 1–15, 2020.
- [219] N. A. Al-Shar'i, "Tackling COVID-19: identification of potential main protease inhibitors via structural analysis, virtual screening, molecular docking and MM-PBSA calculations," *Journal of Biomolecular Structure and Dynamics*, vol. 39, pp. 6689–6704, 2020.
- [220] V. N. Badavath, A. Kumar, P. K. Samanta et al., "Determination of potential inhibitors based on isatin derivatives against SARS-CoV-2 main protease (m^{pro}): a molecular docking, molecular dynamics and structure-activity relationship studies," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2020.
- [221] R. O. Barros, F. L. C. C. Junior, W. S. Pereira, N. M. N. Oliveira, and R. M. Ramos, "Interaction of drug candidates with various SARS-CoV-2 receptors: an in silico study to combat COVID-19," *Journal of Proteome Research*, vol. 19, no. 11, pp. 4567–4575, 2020.
- [222] A. Basit, T. Ali, and S. U. Rehman, "Truncated human angiotensin converting enzyme 2; a potential inhibitor of SARS-CoV-2 spike glycoprotein and potent COVID-19 therapeutic agent," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 3605–3614, 2020.
- [223] V. Battisti, O. Wieder, A. Garon, T. Seidel, E. Urban, and T. Langer, "A computational approach to identify potential novel inhibitors against the coronavirus SARS-CoV-2," *Molecular Informatics*, vol. 39, no. 10, Article ID e2000090, 2020.
- [224] C. G. Benítez-Cardoza and J. L. Vique-Sánchez, "Potential inhibitors of the interaction between ACE2 and SARS-CoV-2 (RBD), to develop a drug," *Life Sciences*, vol. 256, 2020.
- [225] C. Cava, G. Bertoli, and I. Castiglioni, "In silico discovery of candidate drugs against COVID-19," *Viruses*, vol. 12, no. 4, 2020.
- [226] Y. W. Chen, C.-P. B. Yiu, and K.-Y. Wong, "Prediction of the SARS-CoV-2 (2019-nCoV) 3C-like protease (3CLpro) structure: virtual screening reveals velpatasvir, ledipasvir, and other drug repurposing candidates," *F1000Research*, vol. 9, p. 129, 2020.
- [227] M. I. Choudhary, M. Shaikh, W. Tul-Wahab, and R. Ur-Rahman, "In silico identification of potential inhibitors of key SARS-CoV-2 3CL hydrolase (Mpro) via molecular docking, MMGBSA predictive binding energy calculations, and molecular dynamics simulation," *PLoS One*, vol. 15, no. 7, Article ID e0235030, 2020.
- [228] K. Chunduru, R. Sankhe, F. Begum et al., "In silico study to evaluate the antiviral activity of novel structures against 3c-like protease of novel coronavirus (COVID-19) and SARS-CoV," *Medicinal Chemistry*, vol. 17, no. 4, pp. 380–395, 2021.
- [229] C. Coelho, G. Gallo, C. B. Campos, L. Hardy, and M. Würtele, "Biochemical screening for SARS-CoV-2 main protease inhibitors," *PLoS One*, vol. 15, no. 10, Article ID e0240079, 2020.
- [230] W. Dai, B. Zhang, X. M. Jiang et al., "Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease," *Science*, vol. 368, no. 6497, pp. 1331–1335, 2020.
- [231] G. de Lima Menezes and R. A. da Silva, "Identification of potential drugs against SARS-CoV-2 non-structural protein 1 (NSP1)," *Journal of Biomolecular Structure and Dynamics*, vol. 39, no. 15, pp. 1–11, 2020.
- [232] B. Debnath, A. K. Saha, S. Bhaumik, and S. Debnath, "In silico identification of potential inhibitors of the main protease of SARS-CoV-2 using combined ligand-based and

- structure-based drug design approach," *Eurasian Journal of Medicine and Oncology*, vol. 4, no. 4, pp. 336–348, 2020.
- [233] S. Di Micco, S. Musella, M. C. Scala et al., "In silico analysis revealed potential anti-SARS-CoV-2 main protease activity by the Zonulin inhibitor larazotide acetate," *Frontiers in Chemistry*, vol. 8, p. 1271, 2021.
- [234] M. A. El Hassab, T. M. Ibrahim, S. T. Al-Rashood, A. Alharbi, R. O. Eskandrani, and W. M. Eldehna, "In silico identification of novel SARS-CoV-2 2'-O-methyltransferase (NSP16) inhibitors: structure-based virtual screening, molecular dynamics simulation and MM-PBSA approaches," *Journal of Enzyme Inhibition and Medicinal Chemistry*, vol. 36, no. 1, pp. 727–736, 2021.
- [235] M. Elmessaoudi-Idrissi, K. Tsukiyama-Kohara, J. Nourlil et al., "Structure-guided discovery approach identifies potential lead compounds targeting Mpro of SARS-CoV-2," *VirusDisease*, vol. 31, no. 4, pp. 549–553, 2020.
- [236] E. L. Feitosa, F. T. D. S. S. Júnior, J. A. O. N. Nery Neto et al., "Covid-19: rational discovery of the therapeutic potential of melatonin as a sars-cov-2 main protease inhibitor," *International Journal of Medical Sciences*, vol. 17, no. 14, pp. 2133–2146, 2020.
- [237] A. B. Gurung, M. A. Ali, J. Lee, M. A. Farah, and K. M. Al-Anazi, "Identification of potential SARS-CoV-2 entry inhibitors by targeting the interface region between the spike RBD and human ACE2," *Journal of Infection and Public Health*, vol. 14, no. 2, pp. 227–237, 2021.
- [238] Z. Haider, M. M. Subhani, M. A. Farooq et al., "In-silico pharmacophoric and molecular docking-based drug discovery against the main protease (Mpro) of SARS-CoV-2, a causative agent COVID-19," *Pakistan Journal of Pharmaceutical Sciences*, vol. 33, no. 6, pp. 2697–2705, 2020.
- [239] D. C. Hall and H. F. Ji, "A search for medications to treat COVID-19 via in silico molecular docking models of the SARS-CoV-2 spike glycoprotein and 3CL protease," *Travel Medicine and Infectious Disease*, vol. 35, Article ID 101646, 2020.
- [240] B. Havranek and S. M. Islam, "An in silico approach for identification of novel inhibitors as potential therapeutics targeting COVID-19 main protease," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 4304–4315, 2020.
- [241] M. A. Ibrahim, A. H. Abdelrahman, K. S. Allemailem, A. Almatroudi, M. F. Moustafa, and M.-E. F. Hegazy, "In silico evaluation of prospective anti-COVID-19 drug candidates as potential SARS-CoV-2 main protease inhibitors," *The Protein Journal*, vol. 40, pp. 296–309, 2021.
- [242] G. Jaiswal and V. Kumar, "In-silico design of a potential inhibitor of SARS-CoV-2 S protein," *PLoS One*, vol. 15, no. 10, Article ID e0240004, 2020.
- [243] M. Jamal, E. Barzegari, and F. Gholami-Borujeni, "Structure-based screening to discover new inhibitors for papain-like proteinase of SARS-CoV-2: an in silico study," *Journal of Proteome Research*, vol. 20, no. 1, pp. 1015–1026, 2020.
- [244] Z. Jin, X. Du, Y. Xu et al., "Structure of Mpro from SARS-CoV-2 and discovery of its inhibitors," *Nature*, vol. 582, no. 7811, pp. 289–293, 2020.
- [245] A. M. Kanhed, D. V. Patel, D. M. Teli, N. R. Patel, M. T. Chhabria, and M. R. Yadav, "Identification of potential Mpro inhibitors for the treatment of COVID-19 by using systematic virtual screening approach," *Molecular Diversity*, vol. 25, no. 1, pp. 383–401, 2021.
- [246] K. Kavitha, S. Sivakumar, and B. Ramesh, "1, 2, 4 triazolo[1, 5-a] pyrimidin-7-ones as novel SARS-CoV-2 main protease inhibitors: in silico screening and molecular dynamics simulation of potential COVID-19 drug candidates," *Biophysical Chemistry*, vol. 267, Article ID 106478, 2020.
- [247] D. A. Krishnan, G. Sangeetha, S. Vajravijayan, N. Nandhagopal, and K. Gunasekaran, "Structure-based drug designing towards the identification of potential anti-viral for COVID-19 by targeting endoribonuclease NSP15," *Informatics in Medicine Unlocked*, vol. 20, Article ID 100392, 2020.
- [248] R. Kumar, R. Kumar, and P. Tanwar, "In silico drug designing for COVID-19: an approach of high-throughput virtual screening, molecular, and essential dynamics simulations," *Journal of Biomolecular Structure and Dynamics*, vol. 40, no. 16, 2021.
- [249] S. Kumar, P. P. Sharma, U. Shankar et al., "Discovery of new hydroxyethylamine analogs against 3CLpro protein target of SARS-CoV-2: molecular docking, molecular dynamics simulation, and structure-activity relationship studies," *Journal of Chemical Information and Modeling*, vol. 60, no. 12, pp. 5754–5770, 2020.
- [250] A. Kwarteng, E. Asiedu, S. A. Sakyi, and S. O. Asiedu, "Targeting the SARS-CoV2 nucleocapsid protein for potential therapeutics using immuno-informatics and structure-based drug discovery techniques," *Biomedicine and Pharmacotherapy*, vol. 132, 2020.
- [251] D. Li, J. Luan, and L. Zhang, "Molecular docking of potential SARS-CoV-2 papain-like protease inhibitors," *Biochemical and Biophysical Research Communications*, vol. 538, pp. 72–79, 2021.
- [252] S. Mathpal, T. Joshi, P. Sharma et al., "A dynamic simulation study of FDA drug from zinc database against COVID-19 main protease receptor," *Journal of Biomolecular Structure and Dynamics*, vol. 40, 2020.
- [253] S. K. Maurya, A. K. Maurya, N. Mishra, and H. R. Siddique, "Virtual screening, ADME/T, and binding free energy analysis of anti-viral, anti-protease, and anti-infectious compounds against NSP10/NSP16 methyltransferase and main protease of SARS CoV-2," *Journal of Receptors and Signal Transduction*, vol. 40, no. 6, pp. 605–612, 2020.
- [254] M. F. A. Mohamed, G. E. D. A. Abu-Rahma, A. M. Hayallah, M. A. Aziz, A. Nafady, and E. Samir, "Molecular docking study reveals the potential repurposing of histone deacetylase inhibitors against Covid-19," *International Journal of Pharmaceutical Sciences and Research*, vol. 11, no. 9, pp. 4261–4270, 2020.
- [255] H. Monajemi and S. M. Zain, "Strong inhibition of M-protease activity of coronavirus by using PX-12 inhibitor based on ab initio ONIOM calculations," *Journal of Chemical Research*, vol. 45, no. 1-2, pp. 136–140, 2021.
- [256] M. Motiwale, N. S. Yadav, S. Kumar et al., "Finding potent inhibitors for COVID-19 main protease (M-pro): an in silico approach using SARS-CoV-3CL protease inhibitors for combating CORONA," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2020.
- [257] O. Mutlu, O. M. Ugurel, E. Sariyer et al., "Targeting SARS-CoV-2 NSP12/NSP8 interaction interface with approved and investigational drugs: an in silico structure-based approach," *Journal of Biomolecular Structure and Dynamics*, vol. 40, 2020.
- [258] D. Naidoo, A. Roy, P. Kar, T. Mutanda, and A. Anandraj, "Cyanobacterial metabolites as promising drug leads against the Mpro and PLpro of SARS-CoV-2: an in silico analysis," *Journal of Biomolecular Structure & Dynamics*, vol. 39, pp. 6218–6230, 2020.

- [259] O. O. Olubiyi, M. Olagunju, M. Keutmann, J. Loschwitz, and B. Strodel, "High throughput virtual screening to discover inhibitors of the main protease of the coronavirus SARS-CoV-2," *Molecules*, vol. 25, no. 14, 2020.
- [260] M. Özdemir, B. Köksoy, D. Ceyhan, M. Bulut, and B. Yalçın, "In silico, 6lu7 protein inhibition using dihydroxy-3-phenyl coumarin derivatives for SARS-CoV-2," *Journal of the Turkish Chemical Society, Section A: Chemistry*, vol. 7, no. 3, pp. 691–712, 2020.
- [261] M. Ozdemir, B. Koksoy, D. Ceyhan et al., "Design and in silico study of the novel coumarin derivatives against SARS-CoV-2 main enzymes," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2020.
- [262] D. V. Patel, D. M. Patel, A. M. Patel et al., "Carbazole-based semicarbazones and hydrazones as multifunctional anti-Alzheimer agents," *Journal of Biomolecular Structure and Dynamics*, vol. 6, no. 2, pp. 1–22, 2021.
- [263] L. Peng, X. Tian, L. Shen et al., "Identifying effective antiviral drugs against SARS-CoV-2 by drug repositioning through virus-drug association prediction," *Frontiers in Genetics*, vol. 11, p. 1072, 2020.
- [264] M. R. F. Pratama, H. Poerwono, and S. Siswodiardjo, "Molecular docking of novel 5-O-benzoylpinostrubin derivatives as SARS-CoV-2 main protease inhibitors," *Pharmaceutical Sciences*, vol. 26, pp. S63–S77, 2020.
- [265] H. Pundir, T. Joshi, T. Joshi et al., "Using Chou's 5-steps rule to study pharmacophore-based virtual screening of SARS-CoV-2 Mpro inhibitors," *Molecular Diversity*, vol. 25, 2020.
- [266] M. T. J. Quimque, K. I. R. Notarte, R. A. T. Fernandez et al., "Virtual screening-driven drug discovery of SARS-CoV2 enzyme inhibitors targeting viral attachment, replication, post-translational modification and host immunity evasion infection mechanisms," *Journal of Biomolecular Structure and Dynamics*, vol. 39, 2020.
- [267] A. Rakib, Z. Nain, S. A. Sami et al., "A molecular modelling approach for identifying antiviral selenium-containing heterocyclic compounds that inhibit the main protease of SARS-CoV-2: an in silico investigation," *Briefings in Bioinformatics*, vol. 22, no. 2, pp. 1476–1498, 2021.
- [268] J. S. Rane, P. Pandey, A. Chatterjee et al., "Targeting virus-host interaction by novel pyrimidine derivative: an in silico approach towards discovery of potential drug against COVID-19," *Journal of Biomolecular Structure and Dynamics*, vol. 39, pp. 5768–5778, 2020.
- [269] P. Rao, A. Shukla, P. Parmar et al., "Reckoning a fungal metabolite, pyranonigrin A as a potential Main protease (M^{Pro}) inhibitor of novel SARS-CoV-2 virus identified using docking and molecular dynamics simulation," *Biophysical Chemistry*, vol. 264, 2020.
- [270] S. Salman, F. H. Shah, J. Idrees et al., "Virtual screening of immunomodulatory medicinal compounds as promising anti-SARS-COV-2 inhibitors," *Future Virology*, vol. 15, no. 5, pp. 267–275, 2020.
- [271] P. Sarma, N. Shekhar, M. Prajapat et al., "In-silico homology assisted identification of inhibitor of RNA binding against 2019-nCoV N-protein (N terminal domain)," *Journal of Biomolecular Structure and Dynamics*, vol. 39, 2020.
- [272] N. Sepay, N. Sepay, A. Al Hoque, R. Mondal, U. C. Halder, and M. Muddassir, "In silico fight against novel coronavirus by finding chromone derivatives as inhibitor of coronavirus main proteases enzyme," *Structural Chemistry*, vol. 31, no. 5, pp. 1831–1840, 2020.
- [273] M. Shehroz, T. Zaheer, and T. Hussain, "Computer-aided drug design against spike glycoprotein of SARS-CoV-2 to aid COVID-19 treatment," *Heliyon*, vol. 6, no. 10, Article ID e05278, 2020.
- [274] B. Singh and A. Das, "In silico drug repositioning of hydroxychloroquine (HCQ) for the treatment of viral diseases," *International Journal of Progressive Research in Science and Engineering*, vol. 2, no. 3, pp. 13–15, 2021.
- [275] A. Stefaniu, L. Pirvu, B. Albu, and L. Pintilie, "Molecular docking study on several benzoic acid derivatives against SARS-CoV-2," *Molecules*, vol. 25, no. 24, 2020.
- [276] W. Tachoua, M. Kabrine, M. Mushtaq, and Z. Ul-Haq, "An in-silico evaluation of COVID-19 main protease with clinically approved drugs," *Journal of Molecular Graphics and Modelling*, vol. 101, Article ID 107758, 2020.
- [277] A. Uniyal, M. K. Mahapatra, V. Tiwari, R. Sandhir, and R. Kumar, "Targeting SARS-CoV-2 main protease: structure based virtual screening, in silico ADMET studies and molecular dynamics simulation for identification of potential inhibitors," *Journal of Biomolecular Structure & Dynamics*, vol. 40, 2020.
- [278] A. Welker, C. Kersten, C. Müller et al., "Structure-activity relationships of benzamides and isoindolines designed as SARS-CoV protease inhibitors effective against SARS-CoV-2," *ChemMedChem*, vol. 16, 2020.
- [279] L. Wen, K. Tang, K. K.-H. Chik et al., "In silico structure-based discovery of a SARS-CoV-2 main protease inhibitor," *International Journal of Biological Sciences*, vol. 17, no. 6, pp. 1555–1564, 2021.
- [280] M. A. White, W. Lin, and X. Cheng, "Discovery of COVID-19 inhibitors targeting the SARS-CoV-2 NSP13 helicase," *The Journal of Physical Chemistry Letters*, vol. 11, no. 21, pp. 9144–9151, 2020.
- [281] Y. Wu, K. Y. Chang, L. Lou, L. G. Edwards, B. K. Doma, and Z. R. Xie, "In silico identification of drug candidates against COVID-19," *Informatics in Medicine Unlocked*, vol. 21, Article ID 100461, 2020.
- [282] N. H. Zaher, M. I. Mostafa, and A. Y. Altaher, "Design, synthesis and molecular docking of novel triazole derivatives as potential CoV helicase inhibitors," *Acta Pharmaceutica*, vol. 70, no. 2, pp. 145–159, 2020.
- [283] V. Zareade, H. Rezaei, G. Shakerinezhad et al., "The identification of novel inhibitors of human angiotensin-converting enzyme 2 and main protease of SARS-CoV-2: a combination of in silico methods for treatment of COVID-19," *Journal of Molecular Structure*, vol. 1237, Article ID 130409, 2021.
- [284] S. Agarwal, J. P. Pethani, H. A. Shah et al., "Identification of a novel orally bioavailable NLRP3 inflammasome inhibitor," *Bioorganic & Medicinal Chemistry Letters*, vol. 30, no. 21, Article ID 127571, 2020.
- [285] D. Akaberi, J. Krambrich, J. Ling et al., "Mitigation of the replication of SARS-CoV-2 by nitric oxide in vitro," *Redox Biology*, vol. 37, Article ID 101734, 2020.
- [286] L. R. Bernstein and L. Zhang, "Gallium maltolate has in vitro antiviral activity against SARS-CoV-2 and is a potential treatment for COVID-19," *Antiviral Chemistry & Chemotherapy*, vol. 28, Article ID 2040206620983780, 2020.
- [287] G. Bocci, S. B. Bradfute, C. Ye et al., "Virtual and in vitro antiviral screening revive therapeutic drugs for COVID-19," *ACS Pharmacology & Translational Science*, vol. 3, no. 6, pp. 1278–1292, 2020.
- [288] R. Cao, H. Hu, Y. Li et al., "Anti-SARS-CoV-2 potential of artemisinins in vitro," *ACS Infectious Diseases*, vol. 6, no. 9, pp. 2524–2531, 2020.

- [289] Z. Chen, Q. Cui, L. Cooper et al., "Ginkgolic acid and anacardic acid are specific covalent inhibitors of SARS-CoV-2 cysteine proteases," *Cell and Bioscience*, vol. 11, no. 1, 2021.
- [290] M. Gendrot, J. Andreani, P. Jardot et al., "In vitro antiviral activity of doxycycline against SARS-CoV-2," *Molecules*, vol. 25, no. 21, 2020.
- [291] Y. Gupta, D. Maciorowski, S. E. Zak et al., "Bisindolylmaleimide IX: a novel anti-SARS-CoV2 agent targeting viral main protease 3CLpro demonstrated by virtual screening pipeline and in-vitro validation assays," *Methods*, vol. 195, 2021.
- [292] F. Hahn, C. Wangen, S. Häge et al., "IMU-838, a developmental DHODH inhibitor in phase II for autoimmune disease, shows anti-SARS-CoV-2 and broad-spectrum antiviral efficacy in vitro," *Viruses*, vol. 12, no. 12, 2020.
- [293] S. I. Hattori, N. Higashi-Kuwata, H. Hayashi et al., "A small molecule compound with an indole moiety inhibits the main protease of SARS-CoV-2 and blocks virus replication," *Nature Communications*, vol. 12, no. 1, p. 668, 2021.
- [294] Y. Jang, J. S. Shin, M. K. Lee et al., "Comparison of antiviral activity of gemcitabine with 2'-fluoro-2'-deoxycytidine and combination therapy with remdesivir against SARS-CoV-2," *International Journal of Molecular Sciences*, vol. 22, no. 4, 2021.
- [295] S. Jeon, M. Ko, J. Lee et al., "Identification of antiviral drug candidates against SARS-CoV-2 from FDA-approved drugs," *Antimicrobial Agents and Chemotherapy*, vol. 64, no. 7, Article ID e00819, 2020.
- [296] K. Liu, R. Zou, W. Cui et al., "Clinical HDAC inhibitors are effective drugs to prevent the entry of SARS-CoV2," *ACS Pharmacology & Translational Science*, vol. 3, no. 6, pp. 1361–1370, 2020.
- [297] Q. Ma, R. Li, W. Pan et al., "Phillyrin (KD-1) exerts anti-viral and anti-inflammatory activities against novel coronavirus (SARS-CoV-2) and human coronavirus 229E (HCoV-229E) by suppressing the nuclear factor kappa B (NF- κ B) signaling pathway," *Phytomedicine*, vol. 78, Article ID 153296, 2020.
- [298] A. Masih, A. K. Agnihotri, J. K. Srivastava, N. Pandey, H. R. Bhat, and U. P. Singh, "Discovery of novel pyrazole derivatives as a potent anti-inflammatory agent in RAW264.7 cells via inhibition of NF- κ B for possible benefit against SARS-CoV-2," *Journal of Biochemical and Molecular Toxicology*, vol. 35, 2020.
- [299] I. O. Omotuyi, O. Nash, B. O. Ajiboye et al., "Aframomum melegueta secondary metabolites exhibit polypharmacology against SARS-CoV-2 drug targets: in vitro validation of furin inhibition," *Phytotherapy Research*, vol. 35, 2020.
- [300] V. K. Outlaw, F. T. Bovier, M. C. Mears et al., "Inhibition of coronavirus entry in vitro and ex vivo by a lipid-conjugated peptide derived from the SARS-CoV-2 spike glycoprotein HRC domain," *mBio*, vol. 11, no. 5, 2020.
- [301] E. Pitsillou, J. Liang, K. Ververis, A. Hung, and T. C. Karagiannis, "Interaction of small molecules with the SARS-CoV-2 papain-like protease: in silico studies and in vitro validation of protease activity inhibition using an enzymatic inhibition assay," *Journal of Molecular Graphics and Modelling*, vol. 104, Article ID 107851, 2021.
- [302] A. Pizzorno, B. Padey, J. Dubois et al., "In vitro evaluation of antiviral activity of single and combined repurposable drugs against SARS-CoV-2," *Antiviral Research*, vol. 181, Article ID 104878, 2020.
- [303] M. Plaze, D. Attali, M. Prot et al., "Inhibition of the replication of SARS-CoV-2 in human cells by the FDA-approved drug chlorpromazine," *International Journal of Antimicrobial Agents*, vol. 57, no. 3, Article ID 106274, 2021.
- [304] L. R. Reznikov, M. H. Norris, R. Vashisht et al., "Identification of antiviral antihistamines for COVID-19 repurposing," *Biochemical and biophysical research communications*, vol. 538, pp. 173–179, 2021.
- [305] Y. Sakurai, M. M. Ngwe Tun, Y. Kurosaki et al., "5-amino levulinic acid inhibits SARS-CoV-2 infection in vitro," *Biochemical and Biophysical Research Communications*, vol. 545, pp. 203–207, 2021.
- [306] N. E. Stone, S. A. Jaramillo, A. N. Jones et al., "Stenoparib, an inhibitor of cellular poly(ADP-ribose) polymerase, blocks replication of the SARS-CoV-2 and HCoV-NL63 human coronaviruses in vitro," *mBio*, vol. 12, no. 1, 2021.
- [307] H. X. Su, S. Yao, W. F. Zhao et al., "Anti-SARS-CoV-2 activities in vitro of Shuanghuanglian preparations and bioactive ingredients," *Acta Pharmacologica Sinica*, vol. 41, no. 9, pp. 1167–1177, 2020.
- [308] F. Touret, M. Gilles, K. Barral et al., "In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2 replication," *Scientific Reports*, vol. 10, no. 1, Article ID 13093, 2020.
- [309] J. A. Tree, J. E. Turnbull, K. R. Buttigieg et al., "Unfractionated heparin inhibits live wild type SARS-CoV-2 cell infectivity at therapeutically relevant concentrations," *British Journal of Pharmacology*, vol. 178, no. 3, pp. 626–635, 2021.
- [310] E. C. Vatansever, K. S. Yang, A. K. Drelich et al., "Bepridil is potent against SARS-CoV-2 in vitro," *Proceedings of the National Academy of Sciences*, vol. 118, no. 10, 2021.
- [311] S. C. Wang, Y. Chen, Y. C. Wang et al., "Tannic acid suppresses SARS-CoV-2 as a dual inhibitor of the viral main protease and the cellular TMPRSS2 protease," *American Journal of Cancer Research*, vol. 10, no. 12, pp. 4538–4546, 2020.
- [312] K. Watashi, "Identifying and repurposing antiviral drugs against severe acute respiratory syndrome coronavirus 2 with in silico and in vitro approaches," *Biochemical and Biophysical Research Communications*, vol. 538, pp. 137–144, 2021.
- [313] S. Yu, Y. Zhu, J. Xu et al., "Glycyrrhizic acid exerts inhibitory activity against the spike protein of SARS-CoV-2," *Phyto-medicine*, vol. 85, Article ID 153364, 2021.
- [314] K. Zandi, F. Amblard, K. Musall et al., "Repurposing nucleoside analogs for human coronaviruses," *Antimicrobial Agents and Chemotherapy*, vol. 65, no. 1, 2021.
- [315] H. Zhang, Y. Yang, J. Li et al., "A novel virtual screening procedure identifies pralatrexate as inhibitor of SARS-CoV-2 RdRp and it reduces viral replication in vitro," *PLoS Computational Biology*, vol. 16, no. 12, Article ID e1008489, 2020.
- [316] L. Zhang, J. Liu, R. Cao et al., "Comparative antiviral efficacy of viral protease inhibitors against the novel SARS-CoV-2 in vitro," *Virologica Sinica*, vol. 35, no. 6, pp. 776–784, 2020.
- [317] Q. Zhu, Y. Zhang, L. Wang et al., "Inhibition of coronavirus infection by a synthetic STING agonist in primary human airway system," *Antiviral Research*, vol. 187, 2021.
- [318] C. Lei, K. Qian, T. Li et al., "Neutralization of SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig," *Nature Communications*, vol. 11, no. 1, p. 2070, 2020.
- [319] A. D. Rathnayake, J. Zheng, Y. Kim et al., "3C-like protease inhibitors block coronavirus replication in vitro and improve survival in MERS-CoV-infected mice," *Science Translational Medicine*, vol. 12, no. 557, 2020.
- [320] A. Schäfer, F. Muecksch, J. C. C. Lorenzi et al., "Antibody potency, effector function, and combinations in protection

- and therapy for SARS-cov-2 infection in vivo,” *Journal of Experimental Medicine*, vol. 218, no. 3, 2020.
- [321] S. Schumann, A. Kaiser, F. Nicoletti et al., “Immune-modulating drug MP1032 with SARS-CoV-2 antiviral activity in vitro: a potential multi-target approach for prevention and early intervention treatment of COVID-19,” *International Journal of Molecular Sciences*, vol. 21, no. 22, 2020.
 - [322] C. Shang, X. Zhuang, H. Zhang et al., “Inhibitors of endosomal acidification suppress SARS-CoV-2 replication and relieve viral pneumonia in hACE2 transgenic mice,” *Virology Journal*, vol. 18, no. 1, p. 46, 2021.
 - [323] T. P. Sheahan, A. C. Sims, S. Zhou et al., “An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice,” *Science Translational Medicine*, vol. 12, no. 541, 2020.
 - [324] T. T. Tai, T. J. Wu, H. D. Wu et al., “A strategy to treat COVID-19 disease with targeted delivery of inhalable liposomal hydroxychloroquine: a preclinical pharmacokinetic study,” *Clinical and Translational Science*, vol. 14, no. 1, pp. 132–136, 2021.
 - [325] M. A. Tortorici, M. Beltramello, F. A. Lempp et al., “Ultrapotent human antibodies protect against SARS-CoV-2 challenge via multiple mechanisms,” *Science*, vol. 370, no. 6519, pp. 950–957, 2020.
 - [326] A. Wahl, L. E. Gralinski, C. E. Johnson et al., “SARS-CoV-2 infection is effectively treated and prevented by EIDD-2801,” *Nature*, vol. 591, 2021.
 - [327] G. Wang, M. L. Yang, Z. L. Duan et al., “Dalbavancin binds ACE2 to block its interaction with SARS-CoV-2 spike protein and is effective in inhibiting SARS-CoV-2 infection in animal models,” *Cell Research*, vol. 31, no. 1, pp. 17–24, 2021.
 - [328] S. Weston, C. M. Coleman, R. Haupt et al., “Broad anti-coronavirus activity of food and drug administration-approved drugs against SARS-CoV-2 in vitro and SARS-CoV in vivo,” *Journal of Virology*, vol. 94, no. 21, 2020.
 - [329] K. M. White, R. Rosales, S. Yildiz et al., “Plitidepsin has potent preclinical efficacy against SARS-CoV-2 by targeting the host protein eEF1A,” *Science*, vol. 371, no. 6532, pp. 926–931, 2021.
 - [330] C. Y. Wu, Y. S. Lin, Y. H. Yang, L. H. Shu, Y. C. Cheng, and H. T. Liu, “GB-2 inhibits ACE2 and TMPRSS2 expression: in vivo and in vitro studies,” *Biomedicine and Pharmacotherapy*, vol. 132, 2020.
 - [331] C. K. Yuen, W. M. Wong, L. F. Mak et al., “Suppression of SARS-CoV-2 infection in ex-vivo human lung tissues by targeting class III phosphoinositide 3-kinase,” *Journal of Medical Virology*, vol. 93, no. 4, pp. 2076–2083, 2021.
 - [332] C. Zhang, Y. Wang, Y. Zhu et al., “Development and structural basis of a two-MAb cocktail for treating SARS-CoV-2 infections,” *Nature Communications*, vol. 12, no. 1, p. 264, 2021.