

Xgrip109: A γ Tubulin–Associated Protein with an Essential Role in γ Tubulin Ring Complex (γ TuRC) Assembly and Centrosome Function

Ona C. Martin,* Ruwanthi N. Gunawardane,* Akihiro Iwamatsu,[‡] and Yixian Zheng*

*Department of Embryology, Carnegie Institution of Washington, Baltimore, Maryland 21210; and [‡]Central Laboratories for Key Technology, Kirin Brewery Company, Ltd., Yokohama 236, Japan

Abstract. Previous studies indicate that γ tubulin ring complex (γ TuRC) can nucleate microtubule assembly and may be important in centrosome formation. γ TuRC contains approximately eight subunits, which we refer to as *Xenopus* gamma ring proteins (Xgrrips), in addition to γ tubulin. We found that one γ TuRC subunit, Xgrip109, is a highly conserved protein, with homologues present in yeast, rice, flies, zebrafish, mice, and humans. The yeast Xgrip109 homologue, Spc98, is a spindle–pole body component that interacts with γ tubulin. In vertebrates, Xgrip109 identifies two families

of related proteins. Xgrip109 and Spc98 have more homology to one family than the other. We show that Xgrip109 is a centrosomal protein that directly interacts with γ tubulin. We have developed a complementation assay for centrosome formation using demembrated *Xenopus* sperm and *Xenopus* egg extract. Using this assay, we show that Xgrip109 is necessary for the reassembly of salt-disrupted γ TuRC and for the recruitment of γ tubulin to the centrosome. Xgrip109, therefore, is essential for the formation of a functional centrosome.

PROPERLY organized microtubule arrays are critical for many cellular functions including mitosis, cytokinesis, and axonal and intracellular transport (for review see Kellogg et al., 1994). Microtubules are dynamic polymers that assemble from α -/ β -tubulin heterodimers (Weisenberg, 1972; Mitchison and Kirschner, 1984). In vivo, microtubule dynamics are coordinately regulated by many cellular factors (for review see Desai and Mitchison, 1998). There are proteins that stabilize (for reviews see Vallee et al., 1984; Olmstead, 1986), destabilize (Endow et al., 1994; Belmont and Mitchison, 1996), or sever (McNally et al., 1996) microtubule polymers, as well as proteins that sort microtubules into different arrays. In addition, microtubule nucleation is temporally and spatially controlled within the cell, occurring primarily at structures called microtubule organizing centers (MTOCs)¹ (Kellogg et al., 1994). The major MTOC in animal cells is the centrosome that consists of a pair of centrioles surrounded by

an electron-dense cloud of pericentriolar material (PCM). The PCM is responsible for microtubule nucleation (Kellogg et al., 1994).

The discovery of γ tubulin as a suppressor of a β -tubulin mutation in *Aspergillus nidulans* (Weil et al., 1986; Oakley and Oakley, 1989) was a major breakthrough in the study of microtubule nucleation at a molecular level. γ Tubulin is highly conserved and has been found in all eukaryotes examined (for review see Oakley, 1992). Most γ tubulins share over 60% amino acid identity, with the exception of *Saccharomyces cerevisiae* γ tubulin, which is only ~40% identical to the other γ tubulins (Sobel and Synder, 1995; Marschall et al., 1996; Spang et al., 1996). γ Tubulin is localized to all MTOCs such as the spindle–pole body (the major fungal MTOC) and the centrosome (Stearns et al., 1991; Zheng et al., 1991).

Genetic studies in *Aspergillus* (Oakley et al., 1990), *Saccharomyces pombe* (Horió et al., 1991), *Saccharomyces cerevisiae* (Sobel and Synder, 1995; Marschall et al., 1996; Spang et al., 1996), and *Drosophila melanogaster* (Sunkel et al., 1995; Tavosanis et al., 1997) have demonstrated that γ tubulin is an essential gene required for the assembly of a functional mitotic spindle. Antibody inhibition or depletion experiments performed in animal cells (Joshi et al., 1992) and in *Xenopus* egg extracts (Felix et al., 1994), respectively, further show the critical role of γ tubulin in microtubule nucleation at the centrosome.

Biochemical studies were also initiated to study how

Address all correspondence to Yixian Zheng, Department of Embryology, Carnegie Institution of Washington, 115 West University Parkway, Baltimore, MD 21210. Tel.: (410) 554-1232. Fax: (410) 243-6311. E-mail: zheng@mail1.ciwemb.edu

¹ Abbreviations used in this paper: γ TuRC, γ tubulin ring complex; Xgrip, *Xenopus* gamma ring protein; Asp, ammonium sulfate pellet; CSF, cytosolic factor; DAPI, 4',6-diamidino-2-phenylindole; EST, expressed sequence tag; GST, glutathione S-transferase; MTOC, microtubule organizing center; ORF, open reading frame; PCM, pericentriolar material; PEG, polyethylene glycol.

γ tubulin is involved in microtubule nucleation at the MTOCs. Human γ tubulin translated in vitro is monomeric (Melki et al., 1993), and binds to microtubules in an end-specific manner (Li and Joshi, 1995). On the other hand, in animal cells, the γ tubulin that is not associated with the centrosome is found in large cytoplasmic complexes (Raff et al., 1993; Stearns and Kirschner, 1994). The purified *Xenopus* γ -tubulin-containing complex has an estimated molecular mass of over 2,000 kD. This complex, the γ tubulin ring complex (γ TuRC), has an open ring structure and can nucleate microtubules in vitro. In addition to multiple γ -tubulin molecules, the γ TuRC contains approximately eight additional polypeptides (Zheng et al., 1995). The *S. cerevisiae* γ tubulin also appears to form a complex with at least two other proteins: the spindle-pole body components Spc98 and Spc97 (Geissler et al., 1996; Knop et al., 1997). However, since this γ -tubulin complex has a much smaller S value (6 S) (Geissler et al., 1996; Knop et al., 1997) than that of the γ TuRC (>25 S), it is not clear what the functional relationship is between these complexes or if the yeast γ -tubulin complex is able to nucleate microtubules in vitro. Based on the structure and function of the γ TuRC, two models were proposed to explain how the γ TuRC may nucleate microtubule assembly (Zheng et al., 1995; Erickson and Stoffer, 1996). One model suggests that the γ TuRC acts as a seed, similar to the plus end of a microtubule, to nucleate microtubule assembly (Zheng et al., 1995). The other model proposes that the γ TuRC unrolls into a filament, similar to a tubulin protofilament, to initiate microtubule polymerization (Erickson and Stoffer, 1996). Further biochemical and structural analyses are needed to study the mechanism of microtubule nucleation by the γ TuRC.

In parallel structural studies, EM tomography has revealed hundreds of γ TuRC-like rings embedded in the PCM of *Drosophila* and of the surf clam *Spisula* (Moritz et al., 1995b; Vogel et al., 1997); in *Drosophila* these rings are known to contain γ tubulin (Moritz et al., 1995a). The combined structural and biochemical studies of γ tubulin led to the hypothesis that the γ TuRC is anchored within the PCM where it acts to nucleate microtubules (Zheng et al., 1995).

To better understand the relationship between the centrosomal γ -tubulin-containing rings and the cytosolic γ TuRC, we have begun to characterize the γ TuRC subunits. Here we report the identity and function of one of the subunits, Xgrip109. We show that Xgrip109 is a highly conserved centrosomal protein that is essential for γ -tubulin function.

Materials and Methods

Buffers

Buffers used were: Hepes 100 (50 mM Hepes, pH 8, 1 mM MgCl₂, 1 mM EGTA, 100 mM KCl); Hepes 300, Hepes 500, and Hepes 1 M are the same as Hepes 100 except that the concentration of KCl is 300 mM, 500 mM, and 1 M, respectively; cytostatic factor (CSF)-XB (10 mM potassium Hepes, pH 7.7, 100 mM KCl, 2 mM MgCl₂, 0.1 mM CaCl₂, 50 mM sucrose, 5 mM EGTA, pH 7.7); BRB80 (80 mM potassium Pipes, pH 6.8, 1 mM MgCl₂, 1 mM EGTA); and OR2 (2.5 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 1 mM NaHPO₄, 5 mM Hepes, final pH adjusted to 8.3).

Generating Mouse Polyclonal Ascites against the Xgrips

To obtain quantities of the Xgrips sufficient for mouse immunization, we

performed a large-scale immunoprecipitation of the γ TuRC from a 15–25% ammonium sulfate cut of a concentrated *Xenopus* CSF-arrested egg extract (Murray, 1991). Briefly, ammonium sulfate was added to 15% (from a 100% ammonium sulfate stock solution) to ~100 ml of concentrated extract that had been clarified by centrifugation at 35,000 rpm in a rotor (model SW55; Beckman Instruments, Inc., Palo Alto, CA) for 2 h. The extract was then centrifuged at 10,000 rpm in another rotor (model SS34; Sorvall Inc., Newtown, CT) for 15 min. The supernatant from this spin was adjusted to 25% ammonium sulfate and then the pellet was collected by centrifugation as before. The 25% ammonium sulfate pellet (Asp), which contained over 90% of γ TuRC, was resuspended in 75 ml of Hepes 100 and then clarified by centrifugation at 30,000 rpm in a rotor (model SW55; Beckman Instruments, Inc.) for 1 h. The supernatant was collected and 2.6 ml of anti- γ -tubulin antibodies (XenC) (Zheng et al., 1995) at a concentration of 2 mg/ml were added. After a 1-h incubation at 4°C, 2.7 ml of settled protein A-agarose beads (Life Technologies, Inc., Gaithersburg, MD) was added and then the incubation was continued for another hour. The protein A-agarose beads were then washed batchwise three times with 10 vol each of Hepes 100, Hepes 300, and Hepes 100. The γ TuRC was eluted batchwise using 1.4 M MgCl₂. The protein sample was precipitated by addition of trichloroacetic acid to 10%, neutralized, dissolved in SDS sample buffer, and then separated using preparative 10% SDS-PAGE. Gel slices containing either Xgrip195, 133, a mixture of 110 and 109, or a mixture of 75 kD (Zheng et al., 1995) proteins were homogenized and used to immunize mice according to Harlow and Lane (1988). We estimate that between 1 and 10 μ g of the total protein was injected into each animal. Ascites fluid was induced after the fifth injection in mice that gave a positive response by Western analysis.

Internal Peptide Sequencing of the Xgrips

To obtain internal peptide sequences, the Xgrips were prepared as described above and separated on preparative 10% SDS-PAGE. After blotting to polyvinylidene difluoride membranes (Applied Biosystems, Inc., Foster City, CA), proteins were visualized by staining with Ponceau S. The membrane-immobilized proteins were reduced, S-carboxymethylated, and then digested in situ with *Achromobacter* protease I and endoprotease Asp-N (Iwamatsu and Yoshida-Kubomura, 1996). Digested peptides were separated by reverse-phase HPLC using an \AA column (model Wakosil-II AR C18 300, 2.0 \times 150 mm; Wako Pure Chemical Industries, Ltd., Osaka, Japan). Amino acid sequencing was carried out with a gas-phase sequencer (model PPSQ-10; Shimadzu Corp., Tokyo, Japan).

DNA Cloning and Sequencing

The mouse polyclonal ascites that recognized Xgrip109 by Western blot analysis were used to screen a λ ZAP^c cDNA library of *Xenopus* oocytes (Stratagene, La Jolla, CA) according to Sambrook et al. (1989), with modifications as described (Hirano and Mitchison, 1994). We screened 7×10^5 plaques and isolated two positive clones (clone 3-1 and clone 3-4) that had overlapping sequences. Because neither clone had a complete 5' coding region, we rescreened the library using clone 3-4 as probe. 20 positives were obtained. One clone (p109-14) was longer than clone 3-4, but it still lacked a complete 5' coding region.

To obtain the missing 5' end, we carried out 5' rapid amplification of cDNA ends (RACE) using a 5'/3' RACE Kit (Boehringer Mannheim Biochemicals, Indianapolis, IN). Three partially overlapping primers corresponding to the 5' region of the known sequence of p109-14 were made: (a) gsp1 (5'-ACTAATCCCACTGCTGCCGAT-3'), (b) gsp2 (5'-CGG-AATTTCGATGCTTGCAGCATTCTGTGC-3'), and (c) gsp3 (5'-CGG-AATTCTGCACACTGCACATTCCGATC-3') for the initial reverse transcription of *Xenopus* mRNA (gsp1) and two rounds of PCR (gsp2 and gsp3). Pfu DNA polymerase (Stratagene) was used in both rounds of PCR reactions. A 650-bp-amplified DNA fragment was cloned, sequenced, and contained the complete 5' coding region of Xgrip109 since the open reading frame (ORF) began with a methionine, which was preceded by an in-frame stop codon 48-bp upstream. The race product contained the 131 amino acids that were missing from the 5' end of p109-14.

Expression of Recombinant Fusion Proteins, Production of Rabbit Antibodies, and Immunoblotting

We produced six fusion proteins between glutathione S-transferase (GST) and six different fragments of the Xgrip109. Only the fusion between GST and the 134–244-amino acid fragment of Xgrip109, p109-2, elicited rabbit

antibodies that worked for Western analysis, immunofluorescence, and immunodepletion. The primers 5'-CGGGATCCCACGAGGCCAGGC-CACAGAGC-3' and 5'-CGGAATTCCTCAACAGATCCACTTGA-GTC-3' were used to PCR-amplify this fragment of Xgrip109 using p10-4 as template. The resulting fragment was subcloned into the BamHI and EcoRI sites of pGEX-2TK (Pharmacia Biotech., Inc., Piscataway, NJ). To purify the p109-2 fusion protein, 1 liter of bacterial culture (BL21 lys.S) expressing p109-2 was pelleted and lysed in ~20 ml of Hepes 100 containing 0.1 mM of PMSF. After centrifugation, the clarified extract was incubated with ~5 ml of prewashed glutathione agarose (Sigma Chemical Co.) for 1 h at 4°C. The glutathione agarose was then packed into a PD-10 column (Pharmacia Biotech., Inc.), washed sequentially with 10 column volumes each of Hepes 100, Hepes 500, and Hepes 100, and then eluted with 10 mM of reduced glutathione (Sigma Chemical Co.) in Hepes 100. Each of these wash and elution buffers contained 0.1 mM PMSF.

The p109-2 fusion protein and a synthetic peptide (C)RLRVSMGTR-GRRSFHV, corresponding to the COOH-terminal 16 amino acids of the Xgrip109 were used to immunize separate rabbits (Spring Valley Laboratories, Inc., Sykesville, MD). An NH₂-terminal cysteine was added to the peptide for sulfhydryl coupling. Peptide conjugation and antibody affinity purification were performed as described (Harlow and Lane, 1988). The antibodies against p109-2 fusion protein and the synthetic peptide are referred to as 109-2 and 109-c, respectively.

The XenC antibodies used in this paper were raised against a synthetic peptide corresponding to the last 15 amino acids of the *Xenopus* γ tubulin (Zheng et al., 1995). Immunoblotting was carried out using affinity-purified antibodies at a concentration of ~1 μ g/ml using the enhanced chemiluminescence detection system (Amersham Pharmacia Biotech., Inc., Piscataway, NJ).

Preparation of *Xenopus* Egg Extracts and Demembrated *Xenopus* Sperm

Concentrated extracts made from CSF-arrested *Xenopus* eggs according to Murray (1991) were further clarified in a rotor (model TLS55; Beckman Instruments, Inc.) at 50,000 rpm, 4°C for 1 h. 20 \times energy mix (150 mM creatine phosphate, 20 mM ATP, 2 mM EGTA, pH 7.7, 20 mM MgCl₂) was added to the supernatant to a final concentration of 1 \times . This extract was frozen in liquid N₂ and then stored at -80°C as 100- μ l aliquots. All centrosome assembly assays were performed using the clarified and frozen extract. Demembrated *Xenopus* sperm was prepared as described (Sawin and Mitchison, 1991).

Centrosome Assembly Assay and Quantitation

1- μ l of *Xenopus* sperm (~1.5 \times 10⁴ sperm/ml) that had been diluted 10-fold in CSF-XB was added to 10 μ l of thawed and clarified *Xenopus* egg extract. After addition of 1 μ l of 4 mg/ml rhodamine-tubulin (Hyman et al., 1991), the mixture was incubated at room temperature for 10 min. The reaction was stopped by diluting with 1 ml of BRB80 containing 30% glycerol (vol/vol) and then layered in a Corex (Corning GlassWorks, Corning, NY) tube onto 2 ml of BRB80 containing 30% glycerol, followed by spinning in a rotor (model HB-6; Sorvall Inc.) at 10,000 rpm for 20–30 min at 20°C onto a glass coverslip placed at the bottom of the tube (Evans et al., 1985). The sperm nuclei and microtubule asters were fixed in -20°C methanol for 5 min followed by hydration in BRB80 containing 0.1% Triton X-100 for 5 min. The samples were then stained with 0.2 μ g/ml of 4',6-diamidino-2-phenylindole (DAPI) dissolved in BRB80 containing 0.1% Triton X-100 for 2 min followed by mounting in antifade (1 mg/ml *p*-phenylenediamine, 0.5 \times PBS, pH 9, 50% glycerol, 0.02% NaN₃).

A total of 100 sperm nuclei were scored in random fields. These 100 sperm nuclei were divided into three groups: (a) sperm nuclei with a microtubule aster attached at the tip, "centrosome"; (b) sperm nuclei with a few disorganized microtubules attached at the tip, "disorganized centrosome"; or (c) sperm nuclei alone. Because the microtubule asters break away from the tips of the sperm nuclei at a low frequency, we also counted the free microtubule asters while scoring the sperm nuclei. The number of the free microtubule asters was subtracted from the sperm nuclei alone group and added to the sperm nuclei with a microtubule aster centrosome group.

30% Ammonium Sulfate Fractionation, Immunodepletion, and Complementation Assays

To fractionate the clarified egg extract with 30% ammonium sulfate, 43 μ l

of 100% ammonium sulfate was added to 100 μ l of extract and then incubated on ice for 15 min, followed by centrifugation in a rotor (model SS34; Sorvall Inc.) at 10,000 rpm for 10 min at 4°C. The pellet, which contained over 90% of the total γ TuRC and only ~20% of the total extract protein, was resuspended in 100 μ l of either Hepes 100 or Hepes 1 M containing 1 mM GTP and then incubated on ice for 80 min. This mixture was either analyzed by sucrose gradient sedimentation (see below) directly or after desalting using a 1-ml P6 Bio-Spin column (Bio-Rad Laboratories, Hercules, CA), equilibrated in Hepes 100 containing 1 mM of GTP. The eluate was either analyzed by sucrose gradient sedimentation or concentrated ~20-fold by placing in a collodion bag (Sartorius, Göttingen, Germany) that was kept on a bed of Sephadex G-50 resin (Pharmacia Biotech., Inc.) at 4°C. The concentrated eluate was then used in complementation assays (see below).

To immunodeplete Xgrip109 and γ tubulin from the resuspended 30% Asp in either Hepes 100 or Hepes 1 M, 10 μ l of 109-2 IgG (~0.25 mg/ml), 4.5 μ l of XenC IgG (~1.6 mg/ml), or 2.5 μ l of random IgG (~3 mg/ml) were bound to 10–20 μ l of settled protein A beads. The beads were washed with either Hepes 1 M or Hepes 100 and then added to the appropriate resuspended pellets that were incubated on ice for 20 min. After rotating for 1 h at 4°C, the protein A beads were collected by pelleting and then washed. The immunoprecipitated proteins were analyzed by SDS-PAGE followed by Coomassie blue staining or Western analysis, probing with XenC and 109-2. To estimate the percentage of total γ tubulin that remained bound to Xgrip109 in high salt, the amount of γ tubulin immunoprecipitated using XenC in low-salt buffer (this represents the total γ tubulin) was compared with the amount of γ tubulin immunoprecipitated using 109-2 in high salt buffer by densitometry scanning. The supernatants from the 109-2 and random IgG immunodepletion reactions were desalted and concentrated ~20-fold as described above and then used for complementation assays (see below). To assess whether γ tubulin and Xgrip109 interact with each other, the immunoprecipitation was carried out with an excess of extract. Under this condition, the amount of γ tubulin present in the extract is in excess of the added γ -tubulin antibodies and results in cleaner immunoprecipitation.

To prepare γ TuRC-depleted egg extract, either 4.5 μ l of XenC antibodies (~1.6 mg/ml), or 2.5 μ l of control random IgG (~3 mg/ml) were bound to 10–20 μ l of settled Affi-Prep Protein A Support (Bio-Rad Laboratories), 100 μ l of the thawed *Xenopus* egg extract was added to the CSF-XB-washed protein A beads and then incubated with rotation at 4°C for 1 h. The protein A beads were then pelleted and the supernatant was used for centrosome assembly assays.

To determine whether the random IgG- or XenC IgG-depleted extracts were functional in centrosome assembly, 7.5 μ l of each extract was combined with 2.5 μ l of Hepes 100 containing 1 mM of GTP, 1 μ l of rhodamine-tubulin, and 1 μ l of *Xenopus* sperm (see above). The centrosome formation assay was carried out as described above. To complement the XenC-depleted extract, 2.5 μ l of the concentrated Asp that was resuspended in Hepes 100, Hepes 1 M, Hepes 1 M depleted with random IgG, or Hepes 1 M depleted with 109-2 IgG was added to 7.5 μ l of XenC-depleted extract instead of buffer. Centrosome formation was quantitated as described above.

Cell Culture and Cell Extract

Cell line XLK-WG, derived from a primary culture of *Xenopus* kidney cells, was provided by Z. Wu and J.G. Gall (both from Carnegie Institution of Washington, Baltimore, MD). The cells were grown at 32°C in a water-saturated atmosphere of 5% CO₂ in air in medium containing 60% RPMI medium 1640 (Life Technologies, Inc.), 20% heat-inactivated FBS (Life Technologies, Inc.), 100 U/ml each of penicillin and streptomycin, and 2 g/liter of NaHCO₃.

To make XLK-WG cell extract, 5 plates (100-mm-diam) of 80% confluent XLK-WG cells were first rinsed with Hepes 100 and then harvested by scraping. The cell pellet was resuspended in 1 ml Hepes 100 containing 0.1% Triton X-100, 1 mM 2-mercaptoethanol, 1 mM PMSF, 0.01 mM benzamidine HCl, 0.001 mg/ml phenanthroline, and 0.01 mg/ml each of aprotinin, leupeptin, and pepstatin A. The cells were homogenized in a Dounce homogenizer, frozen in liquid N₂, and then stored at -80°C.

Immunofluorescence Staining

For indirect immunofluorescence staining, XLK-WG cells were rinsed with OR2, fixed in -20°C methanol for 5 min, followed by hydrating in OR2 for 5 min. After permeabilizing with OR2 containing 0.1% Triton

X-100 for 1 min, the cells were rinsed with OR2, blocked in OR2 containing 2% BSA for 30 min, and then incubated for 1 h with monoclonal anti- γ -tubulin antibody diluted 200-fold (clone number GTU-88; Sigma Chemical Co.) and 109-2 or 109-c diluted 400-fold. After three 5-min washes, the cells were incubated for 1 h with FITC-conjugated goat anti-mouse and Cy3-conjugated goat anti-rabbit antibodies (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) diluted 400-fold. After three 5-min washes, the cells were stained with DAPI and then mounted in antifade.

To localize γ tubulin and Xgrip109 in the sperm nuclei and microtubule asters, we carried out the centrosome assembly assay as described above but in the absence of rhodamine tubulin. The sperm nuclei and microtubule asters were centrifuged onto round glass coverslips (refer to above) and the same immunofluorescence staining steps as for the XLK-WG cells were performed. Photomicrographs were obtained using a cooled charge-coupled device camera (Princeton Scientific Instruments, Inc., Princeton, NJ) on a microscope (model E800; Nikon Corp., Tokyo, Japan). Images were handled digitally using Adobe Photoshop (Adobe Systems Inc., Mountain View, CA).

Sucrose Gradient Sedimentation

To analyze the sedimentation behavior of γ tubulin and Xgrip109 in XLK-WG cell extracts and *Xenopus* egg extracts, 5 ml of 5–40% sucrose step gradients were poured and allowed to diffuse overnight. Each step (950 μ l) contained 5, 10, 20, 30, or 40% sucrose in Hepes 100. The thawed XLK-WG cell extract was clarified at top speed in a microfuge (Eppendorf 5417C; Hamburg, Germany) for 10 min, and 300 μ l of this supernatant or 300 μ l of clarified egg extract was directly loaded onto the sucrose gradients. Another 300 μ l of the supernatant or clarified egg extract was precipitated with 2.5% polyethylene glycol (PEG), mol wt of 8,000 (Sigma Chemical Co.). The pellet, which contained over 90% of the total cellular γ TuRC, was resuspended in 300 μ l of Hepes 100 and then loaded onto a separate gradient. Centrifugation was performed in an SW55 rotor (model SW55; Beckman Instruments, Inc.) at 50,000 rpm for 4.5 h at 4°C. 300- μ l fractions were collected from the top of each gradient. The fractions were separated by 10% SDS-PAGE and then analyzed by Western analysis, probing with anti- γ -tubulin and anti-Xgrip109 antibodies.

We used 2 ml of 5–40% sucrose gradients to analyze 100 μ l of the Asp that was resuspended in either Hepes 100 or Hepes 1 M (see above). The gradients were poured as described above except that each of the five steps was 400 μ l. The standards, BSA (4.4 S), bovine liver catalase (11.3 S), and porcine thyroglobulin (19.4 S), were dissolved in either Hepes 100 or Hepes 1 M and run on identical sucrose gradients as the samples. The gradients were centrifuged in a rotor (model TLS55; Beckman Instruments, Inc.) at 55,000 rpm for 2 h at 4°C. 130- μ l fractions were collected from the top of each gradient and analyzed by SDS-PAGE followed by Western analysis, probing with anti- γ -tubulin and anti-Xgrip109 antibodies. Because the Hepes 1 M resuspended samples were heavier than the 5% sucrose at the top of the gradient, standards dissolved in Hepes 1 M appeared to sediment faster than the same standards dissolved in Hepes 100.

Results

Xgrip109 Is a Conserved Protein

We previously reported the purification and biochemical characterization of γ TuRC (Zheng et al., 1995). γ TuRC consists of approximately eight uncharacterized polypeptides in addition to γ tubulin. We refer to each polypeptide as Xgrip followed by the respective apparent molecular mass of that polypeptide (see Fig. 3 B). We took two approaches to identify the Xgrips. First, we raised mouse polyclonal antibodies against gel-purified Xgrips. Xgrip109 and Xgrip195 both elicited antibody responses in mice. Second, we sequenced the purified Xgrips that were immobilized on polyvinylidene difluoride membrane. We obtained internal peptide sequences for Xgrip109, Xgrip110, and the Xgrip75s. Since we could not resolve Xgrip109 and Xgrip110 well enough on the gel to sequence them separately, the peptide sequences obtained are derived from

both proteins, and the same is true for the Xgrip75 group of proteins.

We screened a λ ZAP *Xenopus* oocyte cDNA library using the mouse polyclonal antibodies against Xgrip109 and cloned overlapping cDNAs that contained a single ORF (refer to Materials and Methods). We found that half of the peptide sequences we obtained had perfect matches in the ORF, confirming that we cloned the correct Xgrip109 cDNA. We believed that the peptide sequences that do not match Xgrip109 are derived from Xgrip110. The longest cDNA clone has a stop codon followed by a poly A tail at its 3' end, but it lacks a complete 5' coding sequence. Using 5' RACE, we cloned the missing 5' end.

The complete Xgrip109 cDNA (GenBank/EMBL/DDBJ accession number AF052663) encodes a protein of 906 amino acids with a predicted molecular mass of 103.6 kD. Searches of the protein databases reveal that Xgrip109 is most closely related to Spc98, a *S. cerevisiae* spindle-pole body component (Geissler et al., 1996). Although overall Xgrip109 is only ~21% identical and 46% similar to Spc98, the stretch of amino acids (~180) in the middle of both proteins share 28% amino acid identity (Fig. 1). When we searched the expressed sequence tag (EST) databases using the Xgrip109 sequence, we found two groups of conserved human, mouse, and zebrafish ESTs that both share homology with a region of Xgrip109 between amino acids 512–684. The first group of ESTs (Fig. 2 A) are over 85% identical to Xgrip109 over the entire region between amino acids 512–684. The second group of ESTs, consisting of one human EST and one mouse EST that are over 85% identical to each other, shares more limited homology (~30% identity) with Xgrip109 over a smaller region between amino acids 534–612. Furthermore, database searches with the entire Xgrip109 sequence also identified homologous ESTs in rice (~52% amino acid identity, accession number C26482) and *Drosophila* (36% amino acid identity, accession number AA246343). These sequence analyses suggest that Xgrip109 is a conserved protein that, at least in vertebrates, may share homology with two families of related proteins.

Xgrip109 Is a Component of the γ TuRC

To study the function of Xgrip109, we raised antibodies against a GST fusion protein with amino acids 134–244 of Xgrip109, as well as a synthetic peptide corresponding to the COOH terminus of Xgrip109 (refer to Materials and Methods). The two affinity-purified antibodies, which we refer to as 109-2 and 109-c, respectively, specifically recognize a protein of 109 kD when used to probe Western blots of *Xenopus* egg extracts (Fig. 3 A).

Two approaches were used to confirm that Xgrip109 is indeed a component of the γ TuRC. First, we compared the patterns of proteins coimmunoprecipitated by antibodies against Xgrip109 (109-2) and γ tubulin (XenC, Fig. 3 B). Both XenC (an antibody raised against the COOH terminus of γ tubulin (Zheng et al., 1995) and 109-2 antibodies immunoprecipitated the same set of proteins. Moreover, this protein profile is identical to that of the γ TuRC that we identified previously (Zheng et al., 1995). Although immunoprecipitations from clarified egg extracts gave a much higher background than immunopre-

Xgrip109 1MAVPDQKSPNVLLQNLCCRLGKGEA DVAQQFQYAVRVIGS NF 43
 Spe98 1 MELEPTLPGIIEALAPQLLSQSH...LQTFVSDVNVNLRSSSTK.SATQL 45
Xgrip109 44 APTVERDEFIVTETIKKIKKFEVQRREA.DGALFSELHRKLSQGVKLNRRWS 92
 Spe98 46 GPLIDFYKLSQSLDSPETTMMHKIEKFLDALPG....TQNTDDMVKYLS 90
Xgrip109 93 ILVLLLSLSEDPKQPKNTSSFA...ALFAQALPR DAHSTFYIYARPOS 138
 Spe98 91 VFQSLLP.SNYRAKIVQKSSGLNMENLANHEHLLSPVRAPSIYEASPEN 139
Xgrip109 139 LPLSYQDRNVCAQNAASIGSSGISSIGMYALNGPTPQSIQQGQSNQTPN 188
 Spe98 140 MDRFSERRSMVSPNRY.VPSSSTYSSVTLRQLSNPYVNTIPE..... 181
Xgrip109 189 MC DALRQOLGSR LAWTLAAGQQPSQSTTTKGLPNTVSRNVPRTRREGDS 238
 Spe98 182EDILKVVSYTLA.....TTSALFPFDHBIQI 209
Xgrip109 239 SGSVEITETSLVRDLL...YVFGIDGKFKMKNSENCKYVGVKVAVSKS 285
 Spe98 210 PSKIPNPFESGLHLFIEAGLLYQSLGVYKVEF.....RMLNISPM 249
Xgrip109 286 LKDIKLSKELSGWLHNKIKKYTDQRSLDRAFLGVQSFCAALHQELKEYY 335
 Spe98 250 KKALIIIEISE.....ELQNYT 265
Xgrip109 336 RLLSVLHSQQLVEDDQGVNLGVESLTLRRLVWTFDPKIRL KFLAALVD 385
 Spe98 266 AFVNNL.....VSSGVVSLKSLYREIYENIIRLRYCRFTE 302
Xgrip109 386 HCQGR GGELASAVHA TKTGDPMYRSLVQHILGLVAYPIILNPLYRWIYD 435
 Spe98 303 HLEELSGDPTLELNIFKSHGDLIRKIATNLFNMSILSYVEYIMMMLTK 352
Xgrip109 436 GELEDYVH EFFVAS DP...VVKTDRLWHDKYSLRKSMPSPMTMQSRKV 482
 Spe98 353 GLLRATYGEFFIAENTDTNGTDDDFIYHIPNQEVRPAFIPKELAYKI 402
Xgrip109 483 LLIGKSNINFLHQVCHDQTPASKAMAVGKSAESPKDAEL....FTDLLEN 527
 Spe98 403 FMIGKSYIFLEKYCKE.....VQWTNEFSKHYVLYQNSYRGIST 443
Xgrip109 528 AFQT KIDAAYFDT KYLLDVLNR NYNLLLEHMQRMRVLLLGQDFIRHLM 577
 Spe98 444 NPFETIINDQSEIVNHTNQLNQRFHYRDVVFALKNILLMGKSDFDMDALI 493
Xgrip109 578 DLLKPELVRRPATTLYQHNLTGILETAVRATNAQF....DNPEILKRLDV 622
 Spe98 494 EKANDILATPDSLSLNYKTRVLQEAVALQSSLRHLMNSPRNSSVINGLDA 543
Xgrip109 623 RLELVSPGDTCWDFVSLDYHVDGPIATVFTRE...CMSHYLRVFNFLWRA 669
 Spe98 544 RVLDLGHGSGVGDVFTLDYLYPPLSLVNLVNRPFGRKBYLRIFNPLWRF 593
Xgrip109 670 KRMEYIL TD IKKGMHMCNAKLKLG...MPLESGVLHQCHILASEMVH 712
 Spe98 594 KKNVYFYQKEMLSNNDIIRSPFKIRGYNPLRDIINKLSRISILRTOQFO 643
Xgrip109 713 FIHQMQYIITPEVLECSWDELWNKVLKQAQ....DLDHITAAHADVFLD 755
 Spe98 644 FNSKMSYYLNCITIEENFKEMTRKLRQRTENKSNQVDFLRLLNNGTIELNG 693
Xgrip109 756 TIIISRCLLDSESRALLNQLRAVFD...QIIEFQNAQDALYRAALEELQORL 803
 Spe98 694 ILTPKRAEVLTKSSSKPKQKHAIEKTLNIDELESVHNTPLTNIL..... 736
Xgrip109 804 QPEERKKERESSEGEWGVTAEEEDVEN KRIQEFOES IPKMRSQLRILTHFY 853
 Spe98 737SHKLFATNTSEISVGDYSGQPYPTSLVLLLSNVYEVFKVY 776
Xgrip109 854 ...QGIVQQLVLLTSTDESRLRFLSRL DPN EHYFA REPRL 892
 Spe98 777 CNLNDIGYEIIFKMNLNDEHASNGLGKFNTELKIVSQYKRFKDRLYIF 826
Xgrip109 893 RVSMGTRGRRSFHV..... 906
 Spe98 827 RADLKNDDGDEEFLLSKSLR 846

Figure 1. Xgrip109 is homologous to the yeast γ -tubulin-interacting protein Spe98. Sequence comparison of Xgrip109 and Spe98. Vertical lines, identical amino acids; two dots, conserved amino acid changes. The overall amino acid identity shared between the two sequences is ~21%. The underlined region shares ~28% amino acid identity. The sequences in Xgrip109 that match the internal peptide are boxed.

precipitations from resuspended, clarified 30% Asps, the γ TuRC components are still readily identifiable (Fig. 3 B). The presence of γ tubulin and Xgrip109 in the immunoprecipitates was confirmed by Western analysis using XenC and 109-2, respectively (data not shown).

Second, we compared the sedimentation behavior of Xgrip109 and γ tubulin on sucrose gradients. *Xenopus* egg extracts or cell extracts made from a *Xenopus* kidney cell line were either directly sedimented (data not shown) on 5–40% sucrose gradients, or first precipitated with 2.5%

A

1 N----- mouse, AA152700
 1 RESFPAAPDLFDLENAGQKIDAPVPEISKVLEULNKVSLLEHMQARVLLLGQDFIHLMDLLK zebrafish, AA495279
 1 KDSFKAELFDLESTQSKIDWVPEISKVLEULNKVSLLEHMQARVLLLGQDFIHLMDLLK Xgrip109 (512-684)
 1 RESFKDARELETQLENAFQTKIDAVFEDTSKALLDULNKNVSLLEHMQARVLLLGQDFIHLMDLLK
 2 -----RLTGLTETAPRATNAQDFSPSELARLDARLEUSPGOTQDFSLDYHVDGPIATVF human, T55505
 71 PELVPPATTLYQHNLTGILETAVRATNAQDFSEILKALDUALLEUSPGOTQDFSLDYHVDGPIATVF mouse, AA152700
 71 PELVPPATTLYQHNLTGILETAVRATNAQDFSEILKALDUALLEUSPGOTQDFSLDYHVDGPIATVF zebrafish, AA495279
 71 PELVPPATTLYQHNLTGILETAVRATNAQDFSEILKALDUALLEUSPGOTQDFSLDYHVDGPIATVF Xgrip109 (512-684)
 59 TRECHSHLVRFNPLIAR----- human, T55505
 141 TRECHSHLVRFNPLIARARMEVILTDIARKGM mouse, AA152700
 141 TRECHSHLVRFNPLIARARMEVILTDIARKGM zebrafish, AA495279
 141 TRECHSHLVRFNPLIARARMEVILTDIARKGM Xgrip109 (512-684)

B

1 EKAFNYASKULLDFLMEEKELVAHLRSIKRVFLMDQGGDFVHFMDLREELAKPVEDIIP human, R13714
 1 EKAFNYASKULLDFLMEEKELVAHLRSIKRVFLMDQGGDFVHFMDLREELAKPVEDIIL mouse, AA543491
 1 DRAVFDTSKVLDLVNLKNVSLLEHMQARVLLLGQDFIHLMDLLKPELVRRPATTLYQ Xgrip109 (534-612)
 61 PALEALLELALRMTASTANTDFF human, R13714
 61 TRLEALLELALRMTASTANTDFF mouse, AA543491
 61 HNLGTILETAVRATNAQDF Xgrip109 (534-612)

Figure 2. There may be two protein families in vertebrates that share homology with Xgrip109. (A) Sequence comparisons among one group of human (GenBank/EMBL/DBJ accession number T55505), mouse (accession number AA152700), zebrafish (accession number AA495279) ESTs, and Xgrip109 (from amino acid 512 to 684). These sequences share over 85% amino acid identity. (B) Sequence comparisons among another group of human (accession number R13714) and mouse (accession number AA543491) ESTs, and Xgrip109 (from amino acid 534 to 612). Although the two EST sequences share over 85% amino acid identity with each other, Xgrip109 is only ~38% identical to the two EST sequences.

PEG and then sedimented (Fig. 3 C). The fractions from these gradients were analyzed by Western blotting, probing with antibodies against Xgrip109 and γ tubulin. We found that Xgrip109 and γ tubulin cosediment (Fig. 3 C). These results show that Xgrip109 is a component of the γ TuRC.

Xgrip109 Is Localized to the Centrosomes

Since Xgrip109 is a component of the γ TuRC, we expected Xgrip109, like γ tubulin, to localize to centrosomes. Double label-immunofluorescence staining for γ tubulin and Xgrip109 in *Xenopus* kidney tissue culture cells (XLK-WG) revealed that Xgrip109, like γ tubulin, does indeed localize to centrosomes (Fig. 4 A).

We next assembled centrosomes in vitro by incubating demembrated *Xenopus* sperm with *Xenopus* egg extracts. In the presence of an ATP-containing energy regenerating system, the pair of centrioles located at the tips of the sperm nuclei can recruit proteins to form a functional centrosome that nucleates a microtubule aster (Felix et al., 1994; Stearns and Kirschner, 1994). Using immunofluorescence staining, we found that, like γ tubulin, Xgrip109 also localized to these in vitro assembled centrosomes (Fig. 4 B). To make sure that the γ tubulin and Xgrip109 staining would not bleed through due to microtubule staining, we omitted the microtubule label and visualized aster formation by phase-contrast microscopy. The extensive microtubule arrays appear clearly in the phase image (Fig. 4 B, arrows).

Xgrip109 Directly Interacts with γ Tubulin in the γ TuRC

To determine if the interaction between Xgrip109 and γ




Figure 3. Xgrip109 is a component of γ TuRC. (A) Xgrip109 antibodies. Lanes 1 and 2 show Western analysis of *Xenopus* egg extracts separated by 10% SDS-PAGE followed by immunoblotting with affinity-purified antibodies 109-2 and 109-c, respectively. Both antibodies specifically recognize a protein of 109 kD in the *Xenopus* egg extract. (B) Antibodies against Xgrip109 (109-2) and γ tubulin (XenC) both immunoprecipitated γ TuRC components. Lanes 1 and 3 are immunoprecipitations carried out from the clarified egg extracts. Lanes 2 and 4 are immunoprecipitations carried out from 30% ammonium sulfate precipitates that were resuspended in low-salt buffer (refer to Materials and Methods). The immunoprecipitated proteins were separated by 10% SDS-PAGE and stained with Coomassie blue. Each of the γ TuRC components (Xgrips) is indicated by its respective molecular mass. Immunoprecipitating antibodies are indicated at the bottom of the lanes. (C) Xgrip109 cosediments with γ tubulin. *Xenopus* egg extracts or extracts made from a *Xenopus* kidney cell line (XKL-WG cell line) were precipitated with 2.5% PEG, a treatment that precipitates more than 90% of the γ tubulin (data not shown). The pellet was resuspended in HEPES 100 and then sedimented on a 5–40% sucrose density gradient. Fractions were collected from the top of the gradients and then analyzed by Western blotting, probing with XenC and 109-2. The last fraction (fraction 17) in each sedimentation experiment included the pellet.

tubulin is direct, we disrupted the γ TuRC with high salt and examined the sedimentation behavior of Xgrip109 and γ tubulin by sucrose gradient sedimentation. The γ TuRC was precipitated from the *Xenopus* egg extract with 30% ammonium sulfate and resuspended in either low-salt or high-salt buffers and then incubated on ice for 60–80 min to allow complete dissociation of γ TuRC in the presence of high salt. Both samples were then separated on 5–40% sucrose gradients followed by Western analysis, probing with antibodies against Xgrip109 and γ tubulin. Protein standards BSA, catalase, and thyroglobulin were dissolved in either low-salt or high-salt buffers and then run on identical gradients. Fig. 5 A shows that high salt caused both γ tubulin and Xgrip109 to migrate as a smaller complex (\sim 11 S). On the other hand, the γ TuRC remained intact ($>$ 19.4 S) when the 30% Asp was resuspended in the low-salt buffer (Fig. 5 A).

To determine whether the salt-dissociated Xgrip109 remained associated with any other γ TuRC components, we immunoprecipitated Xgrip109 from the 30% Asp resuspended in either low-salt or high-salt buffers. We found that γ tubulin was the only γ TuRC subunit that coimmunoprecipitated with Xgrip109 in high salt (Fig. 5 B, lane 2). The presence of Xgrip109 and γ tubulin in the immunoprecipitates was confirmed by Western analysis (data not shown). These results suggest that Xgrip109 directly binds to γ tubulin in γ TuRC. We estimated, based on densitometry scanning, that less than half of the total γ tubulin in the γ TuRC remained associated with Xgrip109 in the high-salt buffer (refer to Materials and Methods). As a control, γ tubulin was also immunoprecipitated in parallel using XenC. Since the XenC antibody did not bind to γ tubulin well in high-salt buffers (Fig. 5 B, lane 1), we could not determine whether the remaining γ tubulin binds to any other Xgrips under high salt conditions.

Immunodepletion of Xgrip109 Blocks γ TuRC Reassembly

Since we were able to disrupt the γ TuRC with high salt, we wanted to test whether removing the salt would allow the γ TuRC to reform. Furthermore, if the γ TuRC could reassemble, we wanted to examine whether Xgrip109 is needed for γ TuRC reformation. We used sucrose gradient sedimentation to analyze what happened to the γ TuRC after salt treatment and immunodepletion with either random IgG or Xgrip109 IgG. A 30% Asp was resuspended in high-salt buffer and then immunodepleted with either random IgG or Xgrip109 antibodies (109-2). The immunodepletion supernatants were either directly loaded on 5–40% sucrose gradients, or desalted, and then loaded on sucrose gradients. We analyzed the sucrose gradient fractions by Western, probing with anti- γ -tubulin and anti-Xgrip109 antibodies. The control, Fig. 6 E, shows that under low-salt conditions, the intact γ TuRC sedimented as a large particle ($>$ 19.4 S). On the other hand, the γ TuRC was completely dissociated when the 30% Asp was resuspended in high-salt buffer and then immunodepleted with either random IgG (Fig. 6 A) or Xgrip109 IgG (Fig. 6 B). When the salt was removed from the random IgG-depleted sample, at least 50% of γ tubulin and Xgrip109 were assembled into a complex that had the same S value as that





Figure 4. Xgrip109 is localized to centrosomes. (A) Xgrip109 and γ tubulin colocalized to the centrosomes in XLK-WG cells. γ tubulin, γ -tubulin localization revealed by fluorescein secondary antibody; Xgrip109, Xgrip109 localization revealed by Cy3 secondary antibody; DAPI, DNA staining with DAPI; Nom, Nomarski images of the cells. (B) Xgrip109 is localized to the in vitro-assembled centrosomes. The in vitro-assembled centrosomes were spun onto glass coverslips and indirect immunofluorescence staining was carried out using anti- γ -tubulin and anti-Xgrip109 antibodies (refer to Materials and Methods). γ tubulin, γ tubulin was localized to the tip of the sperm nucleus; Xgrip109, Xgrip109 was also localized to the tip of the sperm nucleus; DAPI, the two-sperm nuclei that were stained with either anti- γ -tubulin antibody (GTU-88) or anti-Xgrip109 antibodies (109-2) were stained with DAPI. The images are a combination of fluorescence and phase images. Arrows, microtubule asters at the tips of the two-sperm nuclei. Bars: (A) 20 μ m; (B) 10 μ m.



of the endogenous γ TuRC (Fig. 6 C). When the salt was removed from the Xgrip109-depleted sample, the remaining γ tubulin had a similar S value to that in high-salt (Fig. 6 D). We analyzed the proteins that were removed by 109-2 IgG and random IgG by both Coomassie blue staining and Western, probing with 109-2 and XenC (Fig. 6, F and G). It is clear that, compared with random IgG, 109-2 IgG specifically removed Xgrip109 and γ tubulin. These results suggest that 50% of γ TuRC reassembled after desalting. Furthermore, removing Xgrip109 and a small amount of γ tubulin from the resuspended 30% Asp before lowering the salt concentration blocked the reassembly.

A Complementation Assay for Centrosome Assembly

To study the role of Xgrip109 (or any other Xgrips) in the recruitment of γ tubulin to the centrosome and/or the formation of a functional centrosome, we needed to develop a centrosome formation assay where the activity is dependent on a source of γ TuRC that we can biochemically manipulate to selectively remove specific Xgrips. Since we found that the γ TuRC present in a 30% Asp can be dissociated with salt and reformed by desalting, we hope to use this as a source of γ TuRC.

We took advantage of the existing centrosome formation assay using *Xenopus* egg extracts and sperm (Felix et al.,




Figure 5. Xgrip109 interacts directly with γ tubulin. (A) γ TuRC is dissociated in high salt. Clarified *Xenopus* egg extracts were first precipitated with 30% ammonium sulfate. The pellet (30% Asp) was resuspended in either Hepes 1 M (refer to Materials and Methods) or Hepes 100. The resuspended proteins were fractionated on 5–40% sucrose gradients. The sucrose gradient standards used were bovine serum albumin (4.4S), bovine liver catalase (11.3S), and bovine thyroglobulin (19.4S). The protein standards used (indicated at the bottom of each panel in A) were dissolved in either Hepes 1 M or Hepes 100 and then fractionated under identical conditions. Gradient fractions were collected from the top and each fraction was analyzed by SDS-PAGE followed by Western blotting with XenC and 109-2 antibodies. (B) A fraction of the γ tubulin remains associated with Xgrip109 in high salt.

Immunoprecipitations with XenC (lanes 1 and 3) or 109-2 antibodies (lanes 2 and 4) were carried out using 30% Asp resuspended in either Hepes 1 M (lanes 1 and 2) or Hepes 100 (lanes 3 and 4). The precipitated proteins were separated by SDS-PAGE and then stained by Coomassie blue. The γ TuRC components (*Xgrips*) are indicated by their respective molecular masses.

1994; Stearns and Kirschner, 1994). This centrosome formation assay was based on the original observation of Lohka and Masui (1983) who showed that the demembrated sperm and *Xenopus* egg extract can be used to study various cellular processes such as DNA replication, chromosome condensation, and centrosome and spindle formation in vitro. The centrosome formation assay is es-

entially an assay for PCM assembly around the sperm centrioles located at the tip of the sperm nucleus. When the sperm is incubated with the egg extract, PCM assembles around the centriole pair, resulting in the formation of a functional centrosome that can nucleate microtubule asters. Using this assay, Felix et al. (1994) showed that immunodepleting γ TuRC from the egg extract blocked the




Figure 6. Xgrip109 is required for the reformation of salt-disrupted γ TuRC. Clarified *Xenopus* egg extracts were precipitated with 30% ammonium sulfate. The pellet fraction (30% Asp) was resuspended in either Hepes 100 as control or in Hepes 1 M. Sucrose gradients in A–E are all 5–40%. (A) 30% Asp resuspended in Hepes 1M was immunoprecipitated with random IgG and then analyzed on a sucrose gradient. (B) The same as in A, except that the anti-Xgrip109 antibody, 109-2, was used in the immunoprecipitation. (C) The same as in A, except that after immunoprecipitation, a desalting step was included before the sucrose gradient sedimentation. (D) The same as in B, except that after immunoprecipitation, a desalting step was included before the sucrose gradient sedimentation. (E) Control, 30% Asp resuspended in Hepes 100, desalted, and fractionated on a sucrose gradient. (F) SDS-PAGE separation followed by Coomassie blue staining of proteins that were immunoprecipitated with random IgG (lane 1) and 109-2 IgG (lane 2). (G) The same protein samples in F were analyzed by Western probing with XenC and 109-2. Because samples in A and B contained higher amounts of salt than that of the samples in C, D, and E (refer to Materials and Methods), the sucrose gradient fractions of A and B cannot be compared directly to that of C, D, and E. The molecular weight standards for A and B are indicated at the bottom of A; C, D, and E are indicated at the bottom of E. Standards used are BSA (4.4S), bovine liver catalase (11.3S), and bovine thyroglobulin (19.4S).




Figure 7. The complementation assay for centrosome formation. (A) Western analysis of the immunodepleted extracts, 30% ammonium supernatant, the pellet was probed with 109-2 and XenC. Lanes 1–4: clarified extract; clarified extract immunodepleted using random IgG; clarified extract immunodepleted using XenC; 30% ammonium sulfate supernatant (there is no detectable γ tubulin or Xgrip109), respectively. Lanes 5 and 6: 30% Asp resuspended in Hepes 100 (lane 5) or Hepes 1 M (lane 6), desalted, and then concentrated \sim 20-fold. (B) Quantitation of the complementation assays. *Red columns*, percentage of sperm with centrosomes that nucleated astral microtubules; *blue columns*, percentage of sperm without any microtubule nucleation from the tip of the sperm; *green columns*, percentage of sperm with centrosomes that nucleated a few disorganized microtubules; *Ran. IgG*, depletion with random IgG allowed over 80% of sperm centrioles to assemble into centrosomes; *XenC IgG*, depletion with XenC IgG completely abolished the centrosome assembly activity in the extract; *+ASP.100mM*, addition of 30% Asp resuspended in Hepes 100 to the γ TuRC-depleted extract resulted in over 80% of sperm centrioles to assemble into centrosomes; *+ASP.1M*, addition of 1 M salt-treated and desalted 30% Asp to the γ TuRC-depleted extract resulted in over 70% of sperm centrioles to assemble into centrosomes. The error bars were determined from four independent assays in each case. (C) Representative sperm nuclei with or without a microtubule aster from the assays in B are shown. The microtubules were labeled by the addition of a small amount of rhodamine-tubulin in the assays. Xgrip109 was detected using 109-c antibodies and a fluorescein conjugated goat anti-rabbit secondary antibody. The sperm DNA was stained by DAPI. Bar, 10 μ m.

formation of a functional centrosome. We wanted to test whether the γ TuRC present in the 30% Asp can complement the γ TuRC-depleted extract to form a functional centrosome. We have previously found that the pellet alone did not support centrosome formation (data not shown).

Felix et al. (1994) used crude *Xenopus* egg extract for their centrosome formation assays. We found that clarifying the crude egg extract by centrifugation did not affect centrosome formation activity. Using XenC, we immunodepleted the γ TuRC from the clarified extract (Fig. 7 A, lane 3). Consistent with previous results (Felix et al., 1994), this γ TuRC-depleted extract was not able to assemble a functional centrosome judging by the absence of a microtubule aster (Fig. 7 B). As expected, Xgrip109 was also absent from the centrosome (Fig. 7 C). The control extract that was depleted with random IgG, on the other hand, assembled a functional centrosome around the sperm centrioles, and Xgrip109 was present at the centrosome (Fig. 7, B and C).

To test whether the γ TuRC present in the 30% Asp can complement the γ TuRC-depleted extract to form a cen-

trosome, we resuspended this pellet in low-salt buffer, desalted and concentrated it \sim 20-fold. We found that this γ TuRC-containing fraction restored the centrosome assembly activity of the γ TuRC-depleted extract (Fig. 7). We next tested whether an Asp that was resuspended in high salt still retains the complementing activity after salt removal. The pellet was resuspended in high-salt buffer and incubated on ice to allow complete dissociation of γ TuRC. The fraction was desalted, concentrated \sim 20-fold, and then tested in the complementation assay. We found that this salt-treated 30% Asp also complemented the γ TuRC-depleted extract to form a functional centrosome (Fig. 7). Since we know that the γ TuRC in this pellet is dissociated by the high-salt incubation (see above), this complementation assay allows us to study the effect of selective immunodepletion of Xgrrips on centrosome assembly.

Immunodepletion of Xgrip109 Abolishes the Complementing Activity

To study the role of Xgrip109 in γ -tubulin recruitment and centrosome formation, we wanted to deplete the Xgrip109

from the salt-dissociated γ TuRC present in the resuspended Asp, and ask whether the remaining γ TuRC components still function in the complementation assay described above. Xgrip109 was immunodepleted from the pellet resuspended in high-salt buffers (Fig. 8 A, lanes 4–8). Control reactions were immunodepleted with random IgG. The two antibody-depleted samples were desalted into low-salt buffers and then concentrated \sim 20-fold (Fig. 8 A, lanes 5–8 for the extent of immunodepletion). We found that immunodepleting Xgrip109 abolished the complementing activity present in the resuspended pellet (Fig. 8, B and C), whereas the random IgG control retained the complementing activity (Fig. 8, B and C). Although immunodepleting Xgrip109 removed a fraction of γ tubulin (refer to Fig. 5 B, lane 2), we estimated that more than half of the total amount of γ tubulin remained (Fig. 8 A, lanes 4–8). Interestingly, this remaining γ tubulin did not appear to be recruited to the centrosome (Fig. 8 C, γ -tubulin staining of the centrosomes). These results suggest that Xgrip109 is necessary for the recruitment of γ tubulin to the centrosome, and for the formation of a functional centrosome.

Discussion

Xgrip109 Is a Conserved γ Tubulin–Interacting Protein

The γ TuRC consists of approximately eight polypeptides (Xgrips) in addition to γ tubulin. In an effort to further understand the assembly and function of the γ TuRC, we have begun to characterize the γ TuRC components. We cloned and sequenced one of the Xgrips, Xgrip109. Sequence analysis revealed that Xgrip109 is homologous to the yeast γ tubulin–interacting protein Spc98 (Geissler et al., 1996) (refer to Fig. 1). Like Spc98, Xgrip109 also interacts with γ tubulin (refer to Fig. 5). This suggests that Xgrip109 and Spc98 may have similar cellular functions. In *Xenopus*, Xgrip109 is a component of the γ TuRC. The *S. cerevisiae* Spc98 is also a component of a γ -tubulin–containing complex, although the yeast γ -tubulin complex has a much smaller S value than that of the γ TuRC (Geissler et al., 1996; Knop et al., 1997). It will be interesting to study whether the smaller yeast γ -tubulin complex, like the γ TuRC, can nucleate microtubules in vitro.

EST database searches shows that Xgrip109 is highly conserved among humans, mice, fish, rice, and flies. The Xgrip109 EST homologues in vertebrates can be divided into two groups that share sequence identities with a similar region of Xgrip109 (refer to Fig. 2). One group of ESTs shares over 85% amino acid identity with Xgrip109 and with each other in the 140-amino acid overlapping sequences. The other group of ESTs shares over 30% amino acid identity with Xgrip109, and over 85% amino acid identity with each other in the 80-amino acid overlapping sequences (refer to Fig. 2). Approximately 30% of amino acid identity is shared between the two groups of ESTs. This suggests that Xgrip109 is highly conserved in vertebrates and that it belongs to one of the two families of related proteins. We have recently identified two related γ -tubulin–interacting proteins that are components of *Drosophila* γ TuRC and found that each shares sequence homology with one of the two families of proteins (our unpublished data). We suggest that there are two related

families of conserved γ -tubulin interacting proteins that participate in the formation of the γ TuRC.

Xgrip109, γ Tubulin, and γ TuRC Assembly

We found that the salt-dissociated γ TuRC present in a resuspended 30% Asp can reassemble after desalting (refer to Fig. 6). The reassembly was judged by the shifting in the S value of γ tubulin and Xgrip109 to that of the γ TuRC in our experiments. Because only \sim 50% of the total γ TuRC can reassemble, we did not attempt to purify the reassembled γ TuRC due to the limited amount of starting material. Therefore, we did not carry out structural analysis to compare the reassembled γ TuRC with that of the endogenous γ TuRC. However, we believe that the reassembled γ TuRC is at least similar to the endogenous γ TuRC for two reasons. First, immunoprecipitation using anti-Xgrip109 and γ -tubulin antibodies showed that the reassembled γ TuRC contained the same set of proteins as that of the endogenous γ TuRC (data not shown). Second, the reassembled γ TuRC can functionally replace the endogenous γ TuRC in the centrosome formation assay (see below).

Using the γ TuRC reformation assay, we found that removing Xgrip109 and a small fraction of γ tubulin before desalting blocks the ability of the remaining γ tubulin to assemble into a γ TuRC-sized complex after salt removal. In fact, the remaining γ tubulin has a similar S value before and after desalting. This suggests that Xgrip109 is required for γ tubulin to assemble into larger complexes.

We only looked at γ TuRC dissociation and reassembly using 30% ammonium sulfate–precipitated γ TuRC. It will be interesting to test whether the purified γ TuRC can also reassemble after dissociation with high salt, or whether accessory factors present in the 30% Asp fraction are required. Unfortunately, since we can only purify a few micrograms of γ TuRC, this type of experiment is currently impractical. Nevertheless, the fact that the dissociated γ TuRC can be reassembled offers a useful assay to study how γ TuRC is assembled from its subunits.

Xgrip109 Is a Centrosome Component

γ -Tubulin–containing rings that have similar dimensions to that of the purified γ TuRC were found in the PCM of purified centrosomes (Moritz et al., 1995a,b). One important question is whether the γ TuRC or only γ tubulin in the γ TuRC is recruited to the PCM to act as a microtubule nucleator. Because the purified γ TuRC can nucleate microtubules in vitro, we propose that the γ TuRC-like rings in the centrosome contain at least some of the Xgrips present in the γ TuRC (Zheng et al., 1995). The finding that Xgrip109 is a centrosomal protein that interacts directly with γ tubulin suggests that Xgrip109 is likely a component of the centrosomal rings.

The Complementation Assay for Centrosome Assembly

To study how each of the Xgrips is involved in the assembly of a functional centrosome, we developed a complementation assay for centrosome formation. In this assay the centrosome formation activity depends on the combination of two fractions: a γ TuRC-depleted extract and a 30% Asp fraction that contains the γ TuRC. Since the




Figure 8. Xgrip109 is essential for the formation of a functional centrosome. (A) Western analysis of the immunodepleted extract, 30% Asp, and 30% ammonium sulfate supernatant probed with XenC and 109-2. Lanes 1 and 2: clarified extract immunodepleted with either random IgG (lane 1) or XenC (lane 2). Judging by the absence of γ tubulin and Xgrip109, XenC depleted the γ TuRC (lane 2). Lane 3, 30% ammonium sulfate supernatant that does not contain detectable γ TuRC as expected. Lane 4, 30% Asp resuspended in Hepes 1 M. Lanes 5 and 6, 30% Asp resuspended in Hepes 1 M and then immunodepleted using either random IgG (lane 5) or 109-2 (lane 6). Lanes 7 and 8 are the same as lanes 5 and 6, respectively, except that the proteins were desalted into Hepes 100, and concentrated \sim 20-fold. Lanes 5–8 show that immunodepletion of Xgrip109 in 1 M KCl removed only a fraction of γ tubulin (compare γ -tubulin signals in lanes 5 and 6), whereas Xgrip109 is completely depleted. (B) Quantitation of the complementation assay. Red columns, percentages of sperm with microtubule asters nucleated from the assembled centrosomes; blue columns, percentages of sperm without microtubule asters; white columns, percentages of sperm with assembled centrosomes that nucleated only a few disorganized microtubules; Ran. IgG, centrosome formation assays carried out with clarified extracts that were immunodepleted with random IgG. Over 80% of the sperm centrioles assembled into centrosomes. XenC. IgG, centrosome formation assays carried out with clarified extracts that were immunodepleted of γ TuRC using XenC IgG. The centrosome assembly activity was abolished. +Asp Ran. IgG, 30% Asp that was resuspended in Hepes 1 M and immunodepleted with random IgG complemented the γ TuRC-depleted extract to assemble centrosomes. +Asp Xgrip109, 30% Asp that was resuspended in Hepes 1M and immunodepleted of Xgrip109 did not complement the γ TuRC-depleted extract to assemble centrosomes.

(C) Representative sperm nuclei with or without a microtubule aster from the assays in B are shown. The microtubules were labeled by the addition of a small amount of rhodamine-tubulin in the assays. γ Tubulin was detected using an anti- γ -tubulin monoclonal antibody GTU-88 (Sigma Chemical Co.) and a fluorescein-conjugated goat anti-mouse secondary antibody. γ Tubulin is not recruited to the centrosome in the absence of Xgrip109. The sperm DNA was stained by DAPI. Bar, 10 μ m.

γ TuRC in the pellet can be dissociated with high salt and reassembled by removing the salt, we can remove specific subunit(s) by immunodepletion in high salt, remove the salt, and then study whether the remaining components reform a complex that can function in centrosome assembly. We believe that this type of assay will also be useful in studying other centrosomal components.

We know that the γ TuRC in the Asp is the key activity that complements the γ TuRC-depleted extract, because depleting Xgrip109 (Fig. 8) or γ TuRC (our unpublished observation) abolishes the activity. An obvious question is whether the purified γ TuRC is sufficient for the complementation. To address this question in a meaningful way, we are currently improving our purification methods to achieve sufficiently high γ TuRC concentrations for the complementation assay.

γ Tubulin, Xgrip109, γ TuRC, and Centrosome Assembly

This study and previous work by Felix et al. (1994) strongly suggest that γ TuRC is essential for centrosome assembly. An interesting question is how the γ TuRC is recognized by the centrosome assembly machinery. One model is that the intact γ TuRC is recognized by a recruiting factor(s) that assembles the whole γ TuRC to the centrosome (Fig. 9 A). In this case, the observed centrosomal γ -tubulin-containing rings (Moritz et al., 1995a,b) would be similar to the γ TuRC. In an alternative model, γ TuRC is merely a cytoplasmic storage form of the centrosomal ring components. Upon centrosome assembly, γ TuRC is disassembled and only γ tubulin, and possibly some of the Xgrips, such as Xgrip109, are recognized and recruited to the centrosome to form the centrosomal ring (Fig. 9 B).




Figure 9. Recruitment models. (A) Direct recruitment model. γ TuRC directly binds to a hypothetical recruitment factor(s) and assembles to the centrosome to act as a microtubule nucleator. (B) Subunit recruitment model. γ TuRC is first disassembled to subunits. Only some subunits are recruited to the centrosome by the recruitment factor.

Although we cannot yet differentiate between these two models, our findings suggest that the centrosome assembly machinery cannot recognize and recruit γ tubulin unless it is in the γ TuRC or associated with Xgrip109. When all of the Xgrip109 and a small amount of γ tubulin was removed, the remaining γ tubulin did not reassemble into the γ TuRC and was not recruited to the centrosome (refer to Figs. 7 and 8). Since only $\sim 50\%$ of the total γ tubulin remained upon Xgrip 109 depletion, it is formally possible that the failure of recruiting this remaining γ tubulin is merely due to insufficient γ -tubulin concentration. However, we believe that this is an unlikely possibility for the following reasons. First, in our centrosome formation assays, the extract was always diluted to 60% of the original concentration (refer to Materials and Methods) and the centrosome formation and γ TuRC recruiting were not significantly affected. Furthermore, Felix et al. (1994) showed that the extract can be diluted threefold without drastically affecting the centrosome formation activity. These observations suggest that the recruitment of γ tubulin to the centrosome should not be affected when the amount of γ tubulin is only diluted onefold. We are currently testing the role of Xgrip109 in γ -tubulin recruitment.

In conclusion, we have initiated molecular characterizations of the γ TuRC components. With the assays presented in this paper, we hope to determine the role of each of the γ TuRC subunits in centrosome formation and microtubule nucleation.

The cloning of Xgrip109 was initiated in the Alberts lab (University of California, San Francisco, CA). We thank T. Hirano (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY) for his advice on the library screen and Z. Wu and J. Gall (both from Carnegie Institution of Washington, Baltimore, MD) for the *Xenopus* tissue culture cell line. We are grateful to T. Mitchison (Harvard University, Cambridge, MA) for his advice and help with the initial γ TuRC purification and characterization. We thank K. Oogema (Harvard University, Cambridge, MA) C. Wiese, C.-M. Fan, O. Cohen-Fix, and D. Koshland (all four from Carnegie Institution of Washington) for their critical comments on the manuscript.

This work was supported by grants from National Institutes of Health (RO1- GM56312-01) and the Pew Scholar's Award to Y. Zheng.

Received for publication 2 February 1998 and in revised form 11 March 1998.

References

- Belmont, L., and T. Mitchison. 1996. Identification of a protein that interacts with tubulin dimers and increases the catastrophe rate of microtubules. *Cell* 84:623–631.
- Desai, A., and T.J. Mitchison. 1998. Microtubule polymerization dynamics. *Annu. Rev. Cell Dev. Biol.* 13:83–117.
- Endow, S., S. Kang, L. Satterwhite, M. Rose, V. Skeen, and E. Salmon. 1994. Yeast Kar3 is a minus-end microtubule motor protein that destabilizes microtubules preferentially at the minus ends. *EMBO (Eur. Mol. Biol. Organ.) J.* 13:2708–2713.
- Erickson, H., and D. Stoffer. 1996. Tubulin rings are universal polymers of the tubulin family – alpha/beta, gamma, and FtsZ. *J. Cell Biol.* 135:5–8.
- Evans, L., T. Mitchison, and M. Kirschner. 1985. Influence of the centrosome on the structure of nucleated microtubules. *J. Cell Biol.* 100:1185–1191.
- Felix, M.-A., C. Antony, M. Wright, and B. Maro. 1994. Centrosome assembly in vitro: Role of γ -tubulin recruitment in *Xenopus* sperm aster formation. *J. Cell Biol.* 124:19–31.
- Geissler, S., G. Pereira, A. Spang, M. Knop, S. Soues, J. Kilmartin, and E. Schiebel. 1996. The spindle pole body component Spc98p interacts with the gamma-tubulin-like Tub4p of *Saccharomyces cerevisiae* at the sites of microtubule attachment. *EMBO (Eur. Mol. Biol. Organ.) J.* 15:3899–3911.
- Harlow, E., and D. Lane. 1988. *In Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 726 pp.
- Hirano, T., and T. Mitchison. 1994. A heterodimeric coiled-coil protein required for mitotic chromosome condensation in vitro. *Cell* 79:449–458.
- Horio, T., S. Uzawa, M.K. Jung, B.R. Oakley, K. Tanaka, and M. Yanagida. 1991. The fission yeast gamma-tubulin is essential for mitosis and is localized at microtubule organizing centers. *J. Cell Sci.* 99:693–700.
- Hyman, A., D. Drechsel, D. Kellogg, S. Salser, K. Sawin, P. Steffen, L. Wordeman, and T. Mitchison. 1991. Preparation of modified tubulins. *Methods Enzymol.* 196:478–487.
- Iwamatsu, A., and N. Yoshida-Kubomura. 1996. Systematic peptide fragmentation of polyvinylidene difluoride(PVDF)-immobilized proteins prior to microsequencing. *J. Biochem. (Tokyo)* 120:29–34.
- Joshi, H.C., M.J. Palacios, L. McNamara, and D.W. Cleveland. 1992. Gamma-tubulin is a centrosomal protein required for cell cycle-dependent microtubule nucleation. *Nature* 356:80–83.
- Kellogg, D.R., M. Moritz, and B.M. Alberts. 1994. The centrosome and cellular organization. *Annu. Rev. Biochem.* 63:639–674.
- Knop, M., G. Pereira, S. Geissler, K. Brein, and E. Schiebel. 1997. The spindle pole body component Spc97p interacts with the gamma tubulin of *Saccharomyces cerevisiae* and functions in microtubule organization and spindle pole body duplication. *EMBO (Eur. Mol. Biol. Organ.) J.* 16:1550–1564.
- Li, Q., and H. Joshi. 1995. Gamma-tubulin is a minus-end-specific microtubule binding protein. *J. Cell Biol.* 131:207–214.
- Lohka, M.J., and Y. Masui. 1983. Formation in vitro of sperm pronuclei and mitotic chromosomes induced by amphibian ooplasmic components. *Science* 220:719–721.
- Marschall, L., R. Jeng, J. Mulholland, and T. Stearns. 1996. Analysis of Tub4p, a yeast gamma-tubulin-like protein: Implications for microtubule organizing center function. *J. Cell Biol.* 134:443–454.
- McNally, F., K. Okawa, A. Iwamatsu, and R. Vale. 1996. Katanin, the microtubule-severing ATPase, is concentrated at centrosomes. *J. Cell Sci.* 109:561–567.
- Melki, R., I. Vainberg, R. Chow, and N. Cowan. 1993. Chaperonin-mediated folding of vertebrate actin-related protein and gamma-tubulin. *J. Cell Biol.* 122:1301–1310.
- Mitchison, T.J., and M.W. Kirschner. 1984. Dynamic instability of microtubule growth. *Nature* 312:237–242.
- Moritz, M., M. Braunfeld, J. Sedat, B. Alberts, and D. Agard. 1995a. Gamma-tubulin-containing rings in the centrosome. *Nature* 378:638–640.
- Moritz, M., M.B. Braunfeld, J.C. Fung, J.W. Sedat, B.M. Alberts, and D.A. Agard. 1995b. 3D structural characterization of centrosomes from early *Drosophila* embryos. *J. Cell Biol.* 130:1149–1159.
- Murray, A.W. 1991. Cell cycle extracts. *Methods Cell Biol.* 36:58–605.
- Oakley, B.R. 1992. Gamma-tubulin: The microtubule organizer? *Trends Cell Biol.* 2:1–5.
- Oakley, C.E., and B.R. Oakley. 1989. Identification of gamma-tubulin, a new member of the tubulin superfamily encoded by mipA gene of *Aspergillus nidulans*. *Nature* 338:662–664.
- Oakley, B.R., C.E. Oakley, Y. Yoon, and M.K. Jung. 1990. Gamma-tubulin is a component of the spindle pole body that is essential for microtubule function in *Aspergillus nidulans*. *Cell* 61:1289–1301.
- Olmstead, J. 1986. Microtubule-associated proteins. *Annu. Rev. Cell Biol.* 2:421–457.
- Raff, J.W., D.R. Kellogg, and B.M. Alberts. 1993. *Drosophila* gamma tubulin is part of a complex containing two previously identified centrosomal MAPs. *J.*

- Cell Biol.* 121:823–835.
- Sambrook, J., E. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*. C. Nolan, editor. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 545 pp.
- Sawin, K., and T. Mitchison. 1991. Mitotic spindle assembly by two different pathways in vitro. *J. Cell Biol.* 112:925–940.
- Sobel, S., and M. Synder. 1995. A highly divergent gamma-tubulin gene is essential for cell growth and proper microtubule organization in *Saccharomyces cerevisiae*. *J. Cell Biol.* 131:1775–1788.
- Spang, A., S. Geissler, K. Grein, and E. Schiebel. 1996. Gamma-tubulin-like Tub4p of *Saccharomyces cerevisiae* is associated with the spindle-pole body substructures that organize microtubules and is required for mitotic spindle formation. *J. Cell Biol.* 134:429–441.
- Stearns, T., and M. Kirschner. 1994. In vitro reconstitution of centrosome assembly and function: The role of γ -tubulin. *Cell.* 76:623–637.
- Stearns, T., L. Evans, and M. Kirschner. 1991. Gamma-tubulin is a highly conserved component of the centrosome. *Cell.* 65:825–836.
- Sunkel, C., R. Gomes, P. Sampaio, J. Perdigo, and C. Gonzalez. 1995. Gamma-tubulin is required for the structure and function of the microtubule organizing centre in *Drosophila* neuroblasts. *EMBO (Eur. Mol. Biol. Organ.) J.* 14: 28–36.
- Tavosanis, G., S. Llamazares, G. Goulielmos, and C. Gonzalez. 1997. Essential role for gamma-tubulin in the acentriolar female meiotic spindle of *Drosophila*. *EMBO (Eur. Mol. Biol. Organ.) J.* 16:1809–1819.
- Vallee, R., G. Bloom, and W. Theurkauf. 1984. Microtubule-associated proteins: Subunits of the cytomatrix. *J. Cell Biol.* 99:38s–44s.
- Vogel, J., T. Stearns, C. Rieder, and R. Palazzo. 1997. Centrosomes isolated from *Spisula solidissima* oocytes contain rings and an unusual stoichiometric ratio of alpha/beta tubulin. *J. Cell Biol.* 137:193–202.
- Weil, C., C. Oakley, and B. Oakley. 1986. Isolation of mip (microtubule-interacting protein) mutations of *Aspergillus nidulans*. *Mol. Cell. Biol.* 6:2963–2968.
- Weisenberg, R. 1972. Microtubule formation in vitro in solutions containing low calcium concentrations. *Science.* 177:1104–1105.
- Zheng, Y., M.K. Jung, and B.R. Oakley. 1991. Gamma-tubulin is present in *Drosophila melanogaster* and *Homo sapiens* and is associated with the centrosome. *Cell.* 65:817–823.
- Zheng, Y., M. Wong, B. Alberts, and T. Mitchison. 1995. Nucleation of microtubule assembly by a gamma-tubulin-containing ring complex. *Nature.* 378: 578–583.