DOI: 10.1002/jmv.27424

RESEARCH ARTICLE

Comparative safety of mRNA COVID-19 vaccines to influenza vaccines: A pharmacovigilance analysis using WHO international database

Min Seo Kim ^{1,2} <a>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>
Yehuda Shoenfeld ^{6,7} Andreas Kronbichler ⁸ Ai Koyanagi ^{9,10,11}
Elena Dragioti ¹² Kalthoum Tizaoui ¹³ I Sung Hwi Hong ¹⁴ Louis Jacob ^{9,15}
Joe-Elie Salem ¹⁶ Dong Keon Yon ¹⁷ 💿 Seung Won Lee ¹⁸ 💿
Shuji Ogino ^{19,20,21,22} B Hanna Kim ²³ Jerome H. Kim ²⁴ Jean-Louis Excler ²⁴
Florian Marks ^{24,25,26} John D. Clemens ^{24,27,28} Michael Eisenhut ²⁹
Yvonne Barnett 30 Laurie Butler 30 Cristian Petre Ilie 31 Eui-Cheol Shin 32,33 💿
Jae II Shin ³ 💿 📋 Lee Smith ³⁰

¹College of Medicine, Korea University, Seoul, Republic of Korea

³Department of Pediatrics, Yonsei University College of Medicine, Seoul, Republic of Korea

⁵Department of Pediatrics, Eulji University School of Medicine, Daejeon, Republic of Korea

⁶Laboratory of the Mosaics of Autoimmunity, Saint Petersburg State University, Saint-Petersburg, Russian Federation

⁷Zabludowicz Center for Autoimmune Diseases, Sheba Medical Center, Affiliated to Tel-Aviv University School of Medicine, Tel-Hashomer, Israel

⁸Department of Internal Medicine IV, Medical University Innsbruck, Innsbruck, Austria

⁹Research and Development Unit, Parc Sanitari Sant Joan de Déu, Universitat de Barcelona, Fundació Sant Joan de Déu, CIBERSAM, Barcelona, Spain ¹⁰ICREA, Pg. Lluis Companys 23, Barcelona, Spain

¹¹Instituto de Salud Carlos III, Centro de Investigación Biomédica en Red de Salud Mental, CIBERSAM, Madrid, Spain

¹²Pain and Rehabilitation Centre, and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

¹³Laboratory of Microorganismes and Active Biomolecules, Sciences Faculty of Tunis, University Tunis El Manar, Tunis, Tunisia

¹⁴Yonsei University College of Medicine, Seoul, Republic of Korea

¹⁵Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines, Montigny-le-Bretonneux, France

¹⁶Sorbonne Université, INSERM, CIC-1901 Paris-Est, CLIP² Galilée, UNICO-GRECO Cardio-oncology Program, and Department of Pharmacology, Pitié-Salpêtrière Hospital, Assistance Publique–Hôpitaux de Paris, Paris, France

¹⁷Department of Pediatrics, Seoul National University Children's Hospital, Seoul National University College of Medicine, Seoul, Republic of Korea

¹⁸Department of Data Science, Sejong University College of Software Convergence, Seoul, Republic of Korea

¹⁹Cancer Immunology and Cancer Epidemiology Programs, Dana-Farber Harvard Cancer Center, Boston, Massachusetts, USA

²⁰Department of Epidemiology, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA

²¹Program in MPE Molecular Pathological Epidemiology, Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

²²Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, Massachusetts, USA

²³College of Medicine, Ewha Womans University, Seoul, Republic of Korea

²⁴International Vaccine Institute, Seoul, Republic of Korea

²⁵Cambridge Institute of Therapeutic Immunology and Infectious Disease, University of Cambridge School of Clinical Medicine, Cambridge, UK

Min Seo Kim and Se Yong Jung contributed equally to this study.

²Samsung Advanced Institute for Health Sciences and Technology (SAIHST), Sungkyunkwan University, Samsung Medical Center, Seoul, Republic of Korea

⁴Department of Pediatrics, Severance Children's Hospital, Yonsei University College of Medicine, South Korea

EY-MEDICAL VIROLOGY

²⁶University of Antananarivo, Antananarivo, Madagascar

²⁷International Centre for Diarrheal Diseases Research, Dhaka, Dhaka, Bangladesh

²⁸UCLA Fielding School of Public Health, Los Angeles, California, USA

²⁹Luton & Dunstable University Hospital, Luton, United Kingdom

³⁰Centre for Health, Performance, and Wellbeing, Anglia Ruskin University, Cambridge, UK

³¹Queen Elizabeth Hospital Foundation Trust, King's Lynn, Norfolk, England

³²Laboratory of Immunology and Infectious Diseases, Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science and Technology, Daejeon, Republic of Korea

³³The Center for Epidemic Preparedness, Korea Advanced Institute of Science and Technology, Daejeon, Republic of Korea

Correspondence

Jae II Shin, Department of Pediatrics, Yonsei University College of Medicine, 50-1 Yonsei-ro, Seodaemun-gu, C. P. O. Box 8044, Seoul 03722, Republic of Korea. Email: shinji@yuhs.ac

Funding information

New faculty research seed money grant of Yonsei University College of Medicine for 2021 (2021-32-0049).

Abstract

Two messenger RNA (mRNA) vaccines developed by Pfizer-BioNTech and Moderna are being rolled out. Despite the high volume of emerging evidence regarding adverse events (AEs) associated with the COVID-19 mRNA vaccines, previous studies have thus far been largely based on the comparison between vaccinated and unvaccinated control, possibly highlighting the AE risks with COVID-19 mRNA vaccination. Comparing the safety profile of mRNA vaccinated individuals with otherwise vaccinated individuals would enable a more relevant assessment for the safety of mRNA vaccination. We designed a comparative safety study between 18755 and 27895 individuals who reported to VigiBase for adverse events following immunization (AEFI) with mRNA COVID-19 and influenza vaccines, respectively, from January 1, 2020, to January 17, 2021. We employed disproportionality analysis to rapidly detect relevant safety signals and compared comparative risks of a diverse span of AEFIs for the vaccines. The safety profile of novel mRNA vaccines was divergent from that of influenza vaccines. The overall pattern suggested that systematic reactions like chill, myalgia, fatigue were more noticeable with the mRNA COVID-19 vaccine, while injection site reactogenicity events were more prevalent with the influenza vaccine. Compared to the influenza vaccine, mRNA COVID-19 vaccines demonstrated a significantly higher risk for a few manageable cardiovascular complications, such as hypertensive crisis (adjusted reporting odds ratio [ROR], 12.72; 95% confidence interval [CI], 2.47-65.54), and supraventricular tachycardia (adjusted ROR, 7.94; 95% CI, 2.62-24.00), but lower risk of neurological complications such as syncope, neuralgia, loss of consciousness, Guillain-Barre syndrome, gait disturbance, visual impairment, and dyskinesia. This study has not identified significant safety concerns regarding mRNA vaccination in real-world settings. The overall safety profile patterned a lower risk of serious AEFI following mRNA vaccines compared to influenza vaccines.

KEYWORDS

COVID-19, influenza vaccine, mRNA vaccine, post-implementation surveillance, safety, VigiBase

1 | INTRODUCTION

In May 2020, the 42nd Global Advisory Committee on Vaccine Safety (GACVS) addressed pharmacovigilance preparedness for the launch of the future COVID-19 vaccines¹; experts have voiced that achieving herd immunity at the population level through mass vaccination is a potential strategy to control coronavirus disease (COVID-19).² Two vaccines, the Pfizer-BioNTech messenger RNA (mRNA) and the Moderna mRNA vaccine, have completed phase 3 trials,²⁻⁵ and are being actively rolled out. These mRNA vaccines are based on new technologies that have not been deployed to the general population, and as such, concerns about their safety in real-world settings intersect with optimism for their extraordinarily encouraging efficacy in clinical trials.^{2,3,6}

Although the safety profiles of mRNA vaccines have been evaluated in serial clinical trials,^{4,5,7} concerns remain as the safety evaluations in clinical trials were limited to relatively healthy people, excluding vulnerable populations such as children, pregnant women, and individuals with severe underlying illnesses.^{2,3,7} However, due to vaccine shortages,^{3,8,9} vulnerable patients at high risk for severe courses of COVID-19 are prioritized for vaccination.¹⁰ Therefore, the safety results from these trials may be unrepresentative of the populations that are prioritized to receive them.¹¹ This discrepancy between the trial settings and real-world roll-out strategy warrants urgent interim post-implementation surveillance.³

Despite the high volume of emerging evidence regarding adverse events (AEs) associated with the COVID-19 mRNA vaccines, the previous studies have thus far been largely based on the comparison between vaccinated and unvaccinated control, possibly standing out the AE risks with COVID-19 mRNA vaccination. Comparing the safety profile of mRNA vaccinated individuals with otherwise vaccinated individuals would enable a more relevant assessment for the safety of mRNA vaccination.

This study aimed to conduct post-implementation pharmacovigilance analysis for the Pfizer-BioNTech and Moderna mRNA vaccines by investigating vaccinated individuals who were reported for AEFIs to VigiBase, the global database of individual case safety reports (ICSRs) provided by the WHO. To the best of our knowledge, this study is the first to report the comparative safety of the mRNA COVID-19 vaccine against conventional influenza vaccines.

2 | METHODS

2.1 | Study design and data source

The large post-implementation pharmacovigilance study was conducted using VigiBase, a WHO global deduplicated individual case safety reports (ICSR) database,¹² which has collected adverse event (AE) reports from over 130 countries and 23 million ICSRs since inception in 1967. VigiBase is managed by the Uppsala Monitoring Center (UMC, Sweden). For the database, reported adverse reactions were coded into the Medical Dictionary for Regulatory Activities (MedDRA) Preferred Terms (PTs).¹³

MEDICAL VIROLOGY-WILEY

AE following immunization (AEFI) is defined as any untoward medical event that follows immunization and that does not necessarily have a causal relationship with the usage of the vaccine.¹⁴ AEFIs were reported from various sources, including healthcare professionals, pharmaceutical companies, and patients, and the sources are generally provided with post-market notifications. We extracted AEFI cases from VigiBase reported with two novel mRNA COVID-19 vaccines, Pfizer-BioNTech and Moderna mRNA vaccines, and influenza vaccines from the beginning of 2020 to January 17, 2021. AEFI were reported from America, Europe, and Asia with COVID-19 vaccines. The Ethics Committee of Yonsei University Severance Hospital, Seoul, Republic of Korea, approved this study and granted a waiver of review from the formal Institutional Review Board (no. 4-2020-1379) for the use of deidentified data.

2.2 | Baseline characteristics

The baseline characteristics of individuals reported to VigiBase for any AEFI after mRNA COVID-19 and influenza vaccination are described in Table 1. The VigiBase provides data on demographics (age, sex, and regions), drug history (components, dosage, regimen, indications, and duration of administration), AEs (MedDRA PT classification terms, time to onset, seriousness of AEs, fetal outcomes, and death), and general administrative information (date of report, reports from clinical trials, and reporter type).

Common AEFI was defined as AEFI with a frequency ≥1% of all COVID-19 vaccinated individuals reported for any adverse reaction to VigiBase. A serious AEFI is defined as an AEFI that is associated with death, is life-threatening, involves hospitalization or its prolongation, results in chronic damage/disability, and requires interventions to prevent permanent impairment.¹⁴ The selection process of common and serious AEFI is presented in Figures S1–2.

2.3 Removal of potentially false reports

Potentially false reports are partially prevented at an early data collection stage as most national centers review case reports before they are sent to UMC, and incoming reports to the VigiBase are systematically checked according to pre-defined quality criteria; unmet reports are flagged and subsequently inspected by UMC for reprocessing.¹² Despite the effort, the noise safety signals may still exist, and we triaged to select validated safety signals using two approaches. First, we incorporated information component (IC), an indicator value for disproportionate reporting, that has been proven to be effective in avoiding false positive¹⁵ and thus suitable for conducting pharmacovigilance studies using spontaneous adverse reaction reporting databases.¹⁶⁻¹⁸ Second, we triaged to remove potentially false reports of adverse drug reactions (ADRs) using disproportionality analysis and clinical appraisal. Given that false reports by chance are less likely to survive in stringent association tests, we

	COVID-19 vaccine (n = 18 755)	Influenza vaccine (n = 27 895)
Regions reporting		
Americas	6947/18 755 (37.0)	17 730/27 895 (63.6)
Europe	11 787/18 755 (62.9)	8380/27 895 (30.0)
Australia	0/18 755 (0.0)	1377/27 895 (4.8)
Asia	21/18 755 (0.1)	327/27 895 (1.2)
Africa	0/18 755 (0.0)	81/27 895 (0.4)
Report from clinical trials	94/18 755 (0.5)	1326/28 750 (4.8)
Reporting months		
2020.01-2020.10	0/18 755 (0.0)	16 338/27 895 (58.6)
2020.11	1/18 755 (0.0)	2302/27 895 (8.2)
2020.12	2087/18 755 (11.1)	9217/27 895 (32.0)
2021.01	16 667/18 755 (88.9)	898/27 895 (3.2)
Reporter		
Health care professional	8459/18 755 (45.1)	4054/27 895 (14.5)
Non-health care professional	3364/18 755 (17.9)	6009/27 895 (21.5)
Unreported	6942/18 755 (37.0)	17 832/27 895 (64.0)
Age groups		
<45 years	9389/18 755 (50.1)	10 703/27 895 (38.3)
45-64 years	6422/18 755 (34.2)	6504/27 895 (23.3)
65-74 years	449/18 755 (2.4)	5132/27 895 (18.4)
≥75 years	1282/18 755 (6.8)	2777/27 895 (10.0)
Unreported	1213/18 755 (6.5)	2779/27 895 (10.0)
Sex		
Male	3838/18 755 (20.5)	9263/27 895 (33.2)
Female	14 514/18 755 (77.4)	18 262/27 895 (65.5)
Unreported	403/18 755 (2.1)	370/27 895 (1.3)
Serious AEFIs	3737/18 755 (19.9)	3343/27 573 (12.1)
Outcomes	n = 13 058	n = 14 317
Deaths ^a	119/13 058	113/14371
Time to AEFIs onset	n = 10 876	n = 14 925
Median days (IQR)	1.0 (0.0-1.0)	0.0 (0.0-0.0)

TABLE 1Baseline characteristics ofparticipants vaccinated against COVID-19and influenza reported to VigiBase for anyadverse event followingimmunization (AEFI)

Abbreviations: AEFIs, adverse events following immunization; IQR, interquartile range.

^aAs denominator, all vaccinated participants with AEFIs reported rather than all vaccinated persons were used; we did not present percentile estimations given that they must be larger than those observed in real-world settings. The AEFIs for the COVID-19 and influenza vaccine were extracted from January 2020 to January 17, 2021. Values are presented as n (%) or n/N (%), unless otherwise indicated. Severe AEFI was defined as AEFI that is life-threatening, causes persistent or significant disability, requires hospitalization (first or prolonged), or results in death.

ran disproportionality analyses for 1980 ADRs and excluded nonsignificant ADRs that were deemed clinically irrelevant with vaccines or potentially containing false reports, leaving 49 ADRs subjected to comparative analysis of mRNA COVID-19 and influenza vaccines. We further excluded ADRs that were unlikely to be associated with vaccination (i.e., chronic diseases) by manual review. Death, anaphylactic reactions, and selected 49 reported ADRs out of 1980 MedDRA PTs are summarized in Table 2 and analyzed for the

TABLE 2 Adverse events for	llowing immunization (AEFIs,) associated with th	ne COVID-19 and the influ	uenza vaccine in the full	database of the Vig	giBase from January, 2020	
	COVID-19 vaccine ^a	IC/IC _{0.25}	ROR (95% CI)	Influenza vaccine ^a	IC/IC _{0.25}	ROR (95% CI)	Full database ^a (since 2020.01)
Total individuals with AEFIs	18755			27895			2 720 221
Common AEFIs							
Vaccination site pain	824	5.04/4.94	45.20 (41.75-48.92)	868	4.55/4.45	31.99 (29.61-34.57)	3568
Lymphadenopathy	685	4.66/4.55	32.27 (29.67–35.09)	287	2.84/2.67	7.83 (6.94-8.84)	3855
Oral paraesthesia	472	4.68/4.54	32.62 (29.50-36.08)	150	2.47/2.23	5.92 (5.02-6.98)	2608
Myalgia	2137	3.58/3.52	14.54 (13.87-15.24)	1443	2.44/2.37	5.97 (5.65–6.30)	25821
Heart rate increased	357	3.00/2.85	8.67 (7.78-9.65)	175	1.41/1.19	2.73 (2.35-3.17)	6393
Pain in extremities	1524	2.92/2.84	8.55 (8.10-9.02)	2664	3.15/3.10	10.61 (10.17-11.06)	29 187
Headache	4974	2.84/2.79	9.68 (9.36-10.01)	2474	1.31/1.25	2.66 (2.56-2.78)	97 345
Fatigue	3123	2.84/2.79	8.79 (8.45-9.14)	1498	1.21/1.14	2.42 (2.30-2.55)	63151
Lethargy	242	2.95/2.76	8.30 (7.28-9.45)	292	2.65/2.48	6.77 (6.01-7.63)	4491
Pyrexia	3577	2.73/2.68	8.30 (7.99-8.61)	3324	2.05/2.00	4.73 (4.56-4.91)	78189
Chills	2476	2.59/2.54	7.06 (6.76-7.37)	1484	1.28/1.21	2,.55 (2.42-2.69)	59451
Arthralgia	1338	2.58/2.50	6.59 (6.22-6.97)	1190	1.84/1.75	3.79 (3.57-4.02)	32 482
Influenza-like illness	359	2.30/2.15	5.19 (4.66-5.77)	494	2.19/2.06	4.84 (4.42-5.30)	10486
Chest discomfort	398	2.01/1.87	4.21 (3.80-4.65)	231	0.66/0.47	1.60 (1.40-1.82)	14247
Dizziness	2022	1.37/1.31	2.81 (2.68-2.95)	1480	0.35/0.27	1.29 (1.23-1.36)	113 320
Flushing	543	1.43/1.30	2.77 (2.55-3.02)	192	-0.64/-0.85	0.63 (0.55-0.73)	29 262
Blood pressure increased	240	1.37/1.18	2.64 (2.30-3.00)	100	-0.46/-0.76	0.72 (0.59-0.88)	13442
Cough	546	1.14/1.02	2.27 (2.08-2.47)	702	0.93/0.83	1.96 (1.81–2.11)	35 788
Palpitations	511	1.04/0.92	2.10 (1.92-2.30)	183	-1.01/-1.23	0.49 (0.42-0.57)	36 033
Nausea	2515	0.94/0.88	2.07 (1.99-2.16)	1578	-0.31/-0.38	0.80 (0.76–0.84)	190 359
Diarrhea	748	0.43/0.32	1.36 (1.27-1.47)	671	-0.30/-0.41	0.80 (0.75–0.87)	80 681
Dyspnea	774	0.38/0.27	1.31 (1.22-1.41)	791	-0.16/-0.27	0.89 (0.83-0.95)	86465

KIM ET AL.

100

(Continues)

	COVID-19 vaccine ^a	IC/IC _{0.25}	ROR (95% CI)	Influenza vaccine ^a	IC/IC _{0.25}	ROR (95% CI)	Full database ^a (since 2020.01)
Death, anaphylactic reaction, and	uncommon but serious AEF	<u>0</u>					
Death	103	-1.37/-1.66	0.38 (0.31-0.46)	104	-1.93/-2.22	0.26 (0.21-0.31)	38 799
Anaphylactic reaction	149	-0.12/-0.36	0.92 (0.78-1.08)	147	-0.71/-0.95	0.61 (0.52-0.71)	23415
Intensive care	36	3.71/3.20	17.49 (12.37-24.73)	54	3.80/3.39	18.71 (13.98-25.05)	333
Facial paralysis	76	3.27/2.92	10.93 (8.66-13.81)	66	3.10/2.80	9.76 (7.94-12.01)	1081
Resuscitation	8	2.82/1.65	12.26 (5.96-25.24)	12	3.02/2.09	12.87 (7.05-23.52)	102
Syncope	180	1.64/1.42	3.20 (2.76-3.71)	735	3.10/2.99	9.55 (8.85-10.31)	8341
Unresponsive to stimuli	41	1.88/1.41	3.89 (2.85-5.31)	117	2.83/2.56	7.85 (6.50-9.49)	1560
Endotracheal intubation	6	2.41/1.32	7.33 (3.75-14.32)	25	3.40/2.78	15.00 (9.84–22.86)	186
Hypertensive crisis	14	1.95/1.10	4.42 (2.59-7.52)	2	-1.09/-3.68	0.41 (0.10–1.65)	471
Obstructive airway disorder	10	2.13/1.10	5.42 (2.88-10.19)	6	1.51/0.42	3.25 (1.67-6.32)	276
Supraventricular tachycardia	22	2.66/2.00	7.53 (4.91-11.57)	4	-0.16/-1.90	0.88 (0.33-2.35)	443
Sensory loss	14	1.89/1.04	4.19 (2.47-7.14)	26	2.25/1.64	5.35 (3.61-7.95)	495
Neuralgia	33	1.16/0.63	2.30 (1.63-3.25)	107	2.29/2.00	5.20 (4.28-6.31)	2101
Aphonia	14	1.17/0.32	2.38 (1.41-4.04)	18	0.99/0.25	2.06 (1.29-3.29)	860
Lacunar infarction	3	2.31/0.26	16.01 (4.86–52.77)	1	0.89/-2.90	3.33 (0.45-24.43)	30
Vestibular neuronitis	З	2.17/0.12	11.68 (3.60-37.89)	7	3.04/1.78	20.48 (9.06–46.30)	40
Loss of consciousness	72	0.46/0.11	1.39 (1.10-1.75)	488	2.65/2.51	6.76 (6.16-7.41)	7565
Visual impairment	58	-0.37/-0.77	0.77 (0.59–1.00)	141	0.33/0.09	1.27 (1.07-1.50)	10905
Aphasia	13	0.01/-0.88	1.01 (0.58-1.74)	31	0.68/0.13	1.63 (1.14-2.32)	1869
Neuralgic amyotrophy	2	1.46/-1.13	5.05 (1.23-20.71)	16	3.90/3.11	35.93 (20.24–63.80)	59
Gait disturbance	43	-0.77/-1.23	0.58 (0.43-0.78)	198	0.85/0.64	1.83 (1.59–2.11)	10 681
Seizure	35	-0.94/-1.46	0.52 (0.37-0.72)	213	1.08/0.88	2.15 (1.88-2.47)	9795
Dyskinesia	13	-0.87/-1.76	0.54 (0.31-0.92)	64	0.82/0.45	1.80 (1.40-2.30)	3506
Sudden hearing loss	2	0.53/-2.06	1.63 (0.40-6.56)	7	1.68/0.42	3.93 (1.85-8.37)	179
Pericarditis	2	-0.19/-2.78	0.85 (0.21-3.41)	12	1.64/0.71	3.52 (1.98-6.27)	341
Myelitis	1	0.28/-3.52	1.36 (0.19-9.74)	12	2.97/2.04	12.20 (6.69–22.24)	107

(Continued) **TABLE 2**

	COVID-19 vaccine ^a	IC/IC _{0.25}	ROR (95% CI)	Influenza vaccine ^ª	IC/IC _{0.25}	ROR (95% CI)	Full database ^a (since 2020.01)
Myocarditis	2	-1.19/-3.78	0.38 (0.10–1.54)	17	1.09/0.32	2.23 (1.38-3.61)	753
Neuritis	1	-0.11/-3.91	0.89 (0.12-6.35)	14	2.74/1.89	9.07 (5.24–15.69)	163
Guillain-Barre syndrome	1	-1.84/-5.63	0.20 (0.03-1.46)	233	4.92/4.73	48.14 (41.13-56.35)	704
Abhreviations: AFEL adverse even	nt following immunization. IC	information compor	nent: NA not annlicable: R	OR renorting odds ratio			

As denominator, all vaccinated participants with AEFIs reported rather than all vaccinated persons were used; we did not present percentile estimations given that they must be larger than those observed in real-world settings.

vaccines are presented in the supplementary material. The first AEFI associated with the COVID-19 vaccine was reported on December 15, 2020. The IC/IC_{0.25} and ROR of AEFIs associated with COVID-19 and influenza vaccines were compared with the entire database of VigiBase from January 01, 2020, to January 17, 2021. A positive ICo25 value (>0) in bold is the traditional threshold used for statistical signal detection. only serious AEFIs that are significantly associated with either COVID-19 or influenza vaccine are listed in this table, while serious AEFIs that were not associated with the ^bDue to the volume,

JOURNAL (MEDICAL VIROLOGY

comparative safety between the vaccines (Figure 1). Our careful approach to using those reports deemed genuine and clinically meaningful for our comparative analyses minimized the risk of false reports driving the misleading results. The detailed triage process for AEFI is demonstrated in Figures S1-2.

Comparative safety between COVID-19 and 2.4 influenza vaccines

We have set influenza vaccines as a control given that they have endured iterative and thorough safety evaluations in the form of continued population-based post-market surveillance.¹⁹ which have deemed them acceptably safe.^{19,20} The most frequently reported AEFIs and death after COVID-19 and influenza vaccination were compared in overall individuals reported to the database for AEFI. For uncommon but serious AEFIs that were identified to be potentially associated with the COVID-19 and influenza vaccine ($IC_{0.25} > 0$), the variable adjusted reporting odds ratio (ROR) between mRNA COVID-19 and influenza vaccines for specific AEFI was calculated as described in a previous study¹⁸ to identify comparative safety. The adjusted ROR was used to quantify the degree of difference in odds of specific AEFI between the COVID-19 and influenza vaccine; since the odds of specific AEFI in the influenza vaccine were used as a control. ROR > 1 indicates the higher risk of the AEFI in COVID-19 vaccines compared to influenza vaccines.

Statistical analysis 2.5

Given that VigiBase is composed of an extensive sample size (23 880 736 reports from inception), the data are eligible for disproportionality analysis (also known as case-non-case analysis), for which large sample size is essential to guarantee applicable power and resolution.²¹ When individuals exposed to a particular drug or vaccine (cases) have higher odds of reporting for certain adverse reactions than those not exposed to the drug or vaccine (non-cases), the association between the intervention and the adverse reaction suggests a possible safety signal. The IC and ROR are indicator parameters used to detect signals from the disproportionate analysis developed by the UMC;>0 for lower 95% credibility interval endpoint of information component ($IC_{0.25}$) and >1 for lower confidence interval (CI) of ROR are deemed significant, respectively. The formula for the calculation of the IC is presented in Table S3.

The IC was calculated by comparing observed and expected adverse reaction values using the Bayesian neural network method developed by the UMC,¹⁵ and AEFIs associated with vaccines were detected. Probabilistic logic in intelligent systems (information theory) has been proven to be useful in controlling both big data and missing data.¹⁵ This sensitive algorithm allowed the detection of early signals of mRNA vaccines and identified any potential risks. Of note, VigiBase was not designed to verify the causal relationship between

	JOURNAL O	1						KIM et al.
	Y MEDIC	AL VIROL	OGY					
COVID-19 Vaccine AEs	N	IC/IC _{0.25}	ROR (95% CI)]	Influenza Vaccine AEs	N	IC/IC _{0.25}	ROR (95% CI)
1. Headache	4974	2.84/2.79	9.68 (9.36-10.01)		1. Pyrexia	3324	2.05/2.00	4.73 (4.56-4.91)
2. Pyrexia	3577	2.73/2.68	8.30 (7.99-8.61)		2. Pain in extremities	2664	3.15/3.10	10.61 (10.17-11.06)
3. Fatigue	3123	2.84/2.79	8.79 (8.45-9.14)	$k \rightarrow l$	3. Injection site erythema	2487	3.50/3.44	13.85 (13.26-14.47)
4. Nausea	2515	0.94/0.88	2.07 (1.99-2.16)	\mathbb{N}	4. Headache	2474	1.31/1.25	2.66 (2.56-2.78)
5. Chills	2476	2.59/2.54	7.06 (6.76-7.37)	$k \gg 1$	5. Injection site swelling	1953	3.60/3.53	14.68 (13.98-15.42)
6. Myalgia	2137	3.58/3.52	14.54 (13.87-15.24)	$k \times \not \sim /$	6. Erythema	1674	1.65/1.58	3.36 (3.20-3.54)
7. Dizziness	2022	1.37/1.31	2.81 (2.68-2.95)	$\times \times \times$	7. Nausea	1578	-0.31/-0.38	0.80 (0.76-0.84)
8. Pain in extremities	1524	2.92/2.84	8.55 (8.10-9.02)		8. Fatigue	1498	1.21/1.14	2.42 (2.30-2.55)
9. Arthralgia	1338	2.58/2.50	6.59 (6.22-6.97)		9. Chills	1484	1.28/1.21	2.55 (2.42-2.69)
10 Vaccination site pain	824	5.04/4.94	45.20 (41.75-48.92)		10. Dizziness	1480	0.35/0.27	1.29 (1.23-1.36)
24. Eryhema				11	11. Myalgia			
43. Injection site erythema			,		12. Arthralgia			
70. Injection site swelling					17. Vaccination site pain			
93 Death	103	-1.37/-1.66	0.38 (0.31.0.46)					
Jo. Doani	105		0.56 (0.51-0.40)		140. Death	104	-1.93/-2.22	0.26 (0.21-0.31)

FIGURE 1 Comparative safety of mRNA vaccines to conventional influenza vaccines: Common adverse events following immunization (AEFIs). The numbers in the first column represent the ranking of AEFIs. Values >0 for the lower 95% credibility interval endpoint of the information component ($IC_{0.25}$) and >1 for the lower confidence interval (CI) of ROR indicate statistical significance. AEFI, adverse event following immunization; IC, information component; mRNA, messenger RNA; N, number; ROR, reporting odds ratio

the vaccine and health problems; instead, they were established to detect uncommon or unexpected patterns of AEFIs that imply possible safety concerns with vaccines.

We used a multivariable logistic regression model to produce age and sex-adjusted ROR to compare ADR reporting between mRNA COVID-19 and influenza vaccines. Categorical variables are described as number count (%), and continuous variables are reported as the median and interguartile range (IQR). The cases reported from COVID-19 and influenza vaccination and full database reports were compared using the χ^2 test or Fisher's exact test. Statistical significance was defined as two-tailed p < 0.05. Comparative analyses were conducted using the IBM statistical package for the social sciences (SPSS) version 25.0 (SPSS Inc.).

3 RESULTS

1092

From January 1, 2020, to January 17, 2021, 18755 and 27895 AEFIs for the COVID-19 and influenza vaccines were reported to VigiBase. The AEFIs were most frequently reported from individuals under 64 years of age for COVID-19 and influenza vaccine (Table 1). Ninety-four individuals out of 18755 (0.5%) and 1326 individuals out of 28 750 (4.8%) were reported from clinical trials for COVID-19 and influenza vaccines, respectively; the remaining reports were collected from spontaneous, nonclinical trial settings. A total of 23 880 736 and 2 720 221 ICSRs have been reported to VigiBase since the inception of the database (1967) and since 2020, respectively; and these reports were used as non-case.

We identified safety signals associated with the vaccines, which are statistically significant (defined as $IC_{0.25} > 0$) compared to non-cases (Tables 2 and S1).

3.1 **Common adverse events**

COVID-19 and influenza vaccines showed numerous statistically significant AEFIs, of which many were related to systemic reaction and injection site reactogenicity (Table 2). The 10 most common AEFIs and deaths for the entire population are shown in Figure 1. A more detailed list of total AEFIs after COVID-19 vaccination and the selection process of common AEFIs are presented in the Supplementary material. In Figure 1, the cross-over pattern suggested that COVID-19-vaccinated individuals are more likely to experience systemic symptoms such as headache, myalgia, pyrexia, and fatigue, while influenza-vaccinated individuals were more likely to experience injection site reactogenicity events.

3.2 Uncommon but serious adverse events

Our analysis detected uncommon but serious AEFIs that were significantly associated with COVID-19 vaccines (Table 2). We assessed the comparative safety between COVID-19 and influenza vaccines for serious AEFIs by calculating the adjusted ROR; cardiovascular AEFIs were more prevalent with COVID-19 vaccines: hypertensive crisis (adjusted ROR, 12.72; 95% CI, 2.47-65.54) and supraventricular

MEDICAL VIROLOGY - WILEY

1093

N cases

Adi POP 05% CI

tachycardia (adjusted ROR, 7.94; 95% Cl, 2.62–24.00). In contrast, neurologic AEFIs, such as syncope, neuralgia, loss of consciousness, Guillain-Barre syndrome, gait disturbance, visual impairment, and dyskinesia were more prevalent with influenza vaccines (Figure 2).

3.3 | Death

COVID-19-vaccinated individuals experienced fewer deaths compared to those not exposed to the vaccines, possibly indicating a protective effect of the vaccine ($IC_{0.25}$, -1.66; ROR, 0.38; 95% CI, 0.31-0.46, Table 2). Influenza-vaccinated individuals also experienced fewer deaths compared to those not exposed to the vaccines ($IC_{0.25}$, -2.22; ROR, 0.26; 95% CI, 0.21-0.31, Table 2).

4 | DISCUSSION

To the best of our knowledge, this is the first post-implementation pharmacovigilance study to investigate a diverse range of adverse reactions and provide comparative views for the COVID-19 mRNA vaccine and influenza vaccine. This study has not identified significant safety concerns regarding mRNA vaccination in real-world settings. We have set influenza vaccines as a control given that they have undergone iterative and thorough safety evaluations in the form of continued population-based post-market surveillance,¹⁹ which have deemed them acceptably safe.^{19,20} This interim safety surveillance data revealed that the safety profiles of novel mRNA vaccines may be divergent from those of influenza vaccines; the overall pattern suggested that systematic reactions like chill, myalgia, fatigue were more noticeable with the mRNA COVID-19 vaccine, while injection site reactogenicity events were more prevalent with the influenza vaccine (Figure 1). The overall safety profile patterned a lower risk of serious AEFI following mRNA vaccines compared to influenza vaccines (Figure 2).

The two novel vaccines contain mRNAs that encode spike proteins of SARS-CoV-2 formulated in a lipid nanoparticle. In principle, mRNA vaccines have a unique mechanism compared to conventional vaccines in terms of immunogenicity. Exogenously administered mRNA can strongly stimulate the innate immune system through RNA-sensing pattern recognition receptors.²² Although mRNA has been structurally modified to reduce innate immune responses in current mRNA vaccines,²³ the safety of mRNA vaccines needs to be carefully evaluated. Further safety concerns were raised from the fact that the safety evaluations in clinical trials were limited to relatively healthy people while vulnerable patients at high risk for severe courses of COVID-19 were prioritized to the vaccination in realworld settings.^{3,8,9} This study was designed to investigate this gap and promptly detect safety signals undiscovered at the trial level but could snowball as vaccine coverage spans across the billions of people worldwide. Of note, this analysis aims to raise hypotheses for further, more definitive studies, not to test hypotheses and inform recommendations.

		Aujusteu	ROK	Auj-NO	K 9070 CI	(COVID/
Serious adverse events afte	r immunizatio	on				muenza
Intensive care			-	1.22	[0.78; 1.91]	36/54
Resuscitation		-+-		1.22	[0.49; 3.02]	8/12
Unresponsive to stimuli		-#-		0.57	[0.39; 0.82]	41/117
Endotracheal intubation		-++		0.62	[0.28; 1.36]	9/25
Obstructive airway disorder		-++		1.35	[0.54; 3.37]	9/1
Hypertensive crisis			+	→ 12.72	[2.47; 65.54]	14/2
Seizures				0.24	[0.05; 1.12]	35/213
Facial paralysis		-		1.16	[0.85; 1.59]	76/99
Syncope		-+-		0.30	[0.26; 0.36]	180/735
Neuralgia				0.43	[0.29; 0.64]	33/107
Loss of consciousness		H		0.20	[0.16; 0.26]	72/855
Guillain-Barre syndrome	~			0.01	[0.00; 0.06]	1/233
Gait disturbance				0.37	[0.27; 0.52]	43/198
Supraventricular tachycardia				- 7.94	[2.62; 24.00]	22/4
Sensory loss		-+-		0.66	[0.34; 1.29]	14/26
Aphonia		-	-	1.01	[0.49; 2.09]	14/18
Lacunar infarction		-+	+	→ 4.58	[0.46; 45.15]	3/1
Vestibular neuronitis		+	_	0.66	[0.16; 2.77]	3/7
Visual impairment				0.61	[0.45; 0.84]	58/141
Aphasia		-+-		0.68	[0.35; 1.33]	13/31
Neuralgic amyotrophy				0.26	[0.06; 1.20]	2/16
Dyskinesia				0.29	[0.16; 0.53]	13/64
Sudden hearing loss	_		_	0.43	[0.08; 2.19]	2/7
Pericarditis				0.34	[0.07; 1.57]	2/12
Myelitis		·		0.12	[0.02; 0.95]	1/12
Myocarditis		<u> </u>		0.18	[0.04; 0.79]	2/17
Neuritis		•	1 1	0.15	[0.02; 1.16]	1/14
	0.0125 0.	1 0.5 1	2 10	40		

A diverte d DOD

COVID-19 vaccines versus influenza vaccines with respect to serious adverse events after immunization (AEFIs). Adj-ROR, adjusted reported odd ratios; 95% CI, 95% confidential interval; mRNA, messenger RNA

FIGURE 2 Comparative safety of mRNA

Higher risk with influenza vaccines

Higher risk with mRNA vaccines

ILEY-MEDICAL VIROLOGY

Our data revealed that COVID-19-vaccinated individuals experienced significantly fewer deaths compared to those not exposed to the vaccines, possibly indicating a protective effect of the vaccine (Table 2). When stratifying death risk by age group, the proportion of death among all AEFI-reported vaccinated individuals in the age group was higher in the >65 years age groups, and the tendency was more prominent for those ≥75 years old (Table S2). This observation could be explained, in part, by the selective roll-out of mRNA vaccines to particularly vulnerable elderly populations, such as those receiving care in long-term care facilities (LTCF), who are frail and at a higher risk of severe courses. Therefore, it is difficult to attribute the higher odds of death in the elderly, especially those >75 years, to mRNA vaccination per se without more data that may help extricate a causal relationship. Further studies should be conducted to elucidate the causal relationship and underlying mechanisms for this association.

It is noteworthy that mRNA vaccines demonstrated a significantly higher risk for a few cardiovascular complications, such as hypertensive crisis and supraventricular tachycardia (SVT) compared with influenza vaccines; however, risks for most other cardiovascular adverse events such as atrial fibrillation, myocardial infarction, cardiogenic shock, and cardiac failure were not increased with mRNA vaccination (Supporting Information Data). Considering hypertensive crisis and SVT are mostly manageable and rarely cause permanent or chronic damages, these cardiovascular signals are less likely to pose a burden to a large population. Moreover, lower risks of other serious complications, especially neurological complications (i.e., neuralgia, Guillain-Barre syndrome, dyskinesia, and gait disturbance), with mRNA vaccines compared to influenza vaccines may further support the comparative safety of mRNA vaccines in real-world settings (Figure 2). The findings and hypotheses raised from this first postimplementation surveillance data may support evidence-based discussions and risk-benefit assessments for ongoing mass vaccination.

The results of this study should be interpreted in the context of known limitations. First, VigiBase relies on spontaneous reports, and therefore the data are subject to reporting biases. To address this, we triaged to remove potentially false reports using disproportionality analysis and clinical appraisal, as demonstrated in the methods. Second, VigiBase was primarily designed to identify unusual or unexpected safety signals that might be associated with vaccines rather than to determine a causal relationship. Therefore, our analysis should be interpreted as confined to the associations, and be aware that this analysis is intended to raise hypotheses for further, more definitive studies. Last, we employed disproportionality analyses between the AEFIs reported with mRNA COVID-19/influenza vaccines and the total number of individual case safety reports for the entire VigiBase database. However, the advantages of VigiBase and the methods used in this study (disproportionate analysis) have been well established through numerous studies^{16,24-26} and may provide sufficient evidence to bring the potential safety signals to the attention of public health professionals and decision-makers.

CONCLUSION

This pharmacovigilance study has not identified significant safety concerns regarding mRNA vaccination in real-world settings. The overall safety profile patterned a lower risk of serious AEFI following mRNA vaccines compared to influenza vaccines.

ACKNOWLEDGMENTS

We appreciate the members of the custom search team at the Uppsala Monitoring Centre (Uppsala, Sweden) research section, who were invaluable to the successful performance of this study. This study was supported by a new faculty research seed money grant of Yonsei University College of Medicine for 2021 (2021-32-0049). The funders had no role in the design, analyses, or interpretation of the study. Hanna Kim is a spouse of Minseo Kim. However, Hanna Kim contributed to this work independently.

CONFLICT OF INTERESTS

The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the World Health Organization (WHO) and the International Vaccine Institute (IVI). The authors declare that they have no known competing financial interests or personal relationships with companies that could have influenced the work reported in this paper. Dr. Jean-Louis Excler reports nonfinancial support from Brighton Collaboration, personal fees from US Military HIV Research Program, personal fees from Johnson & Johnson, outside the submitted work: Dr. Jerome H. Kim reports personal fees from SK bioscience, personal fees from educational companies during the period covered by this manuscript; Min Seo Kim, Se Yong Jung, Jong Gyun Ahn, Se Jin Park, Yehuda Shoenfeld, Andreas Kronbichler, Ai Koyanagi, Elena Dragioti, Kalthoum Tizaoui, Sung Hwi Hong, Louis Jacob, Joe-Elie Salem, Dong Keon Yon, Seung Won Lee, Shuji Ogino, Hanna Kim, Florian Marks, John D. Clemens, Michael Eisenhut, Yvonne Barnett, Laurie Butler, Cristian Petre Ilie, Eui-Cheol Shin, Jae Il Shin, and Lee Smith have no commercial associations that may present a conflict of interest regarding this manuscript.

AUTHOR CONTRIBUTIONS

Min Seo Kim, Se Yong Jung, and Jae II Shin contributed to the study concept and design. Se Yong Jung and Jae II Shin acquired data. Min Seo Kim and Se Yong Jung analyzed the data. Min Seo Kim and Se Yong Jung wrote the first draft of the manuscript. Min Seo Kim finalized the manuscript. Hanna Kim supported organizing influenza vaccine data and supplementary materials. Joe-Elie Salem, Jerome Kim, Jean-Louis Excler, Florian Marks, John D. Clemens supervised the interpretation of vaccine pharmacovigilance results. Jong Gyun Ahn, Se Jin Park, Yehuda Shoenfeld, Andreas Kronbichler, Ai Koyanagi, Elena Dragioti, Kalthoum Tizaoui, Sung Hwi Hong, Louis Jacob, Dong Keon Yon, Seung Won Lee, Shuji Ogino, Michael Eisenhut, Yvonne Barnett, Laurie Butler, Cristian Petre Ilie, Eui-Cheol Shin, and Lee Smith contributed to the intellectual discussion, organization of contents, and critical revision of the manuscript. JI Shin and JE Salem provided statistical advice or supervised the statistical interpretations. All authors saw and approved the final submitted version.

DATA AVAILABILITY STATEMENT

Study protocol and Statistical code are available from Prof. Shin (e-mail, shinji@yuhs.ac). Data set is available from the WHO Program for International Drug Monitoring through a data use agreement.

ORCID

Min Seo Kim b https://orcid.org/0000-0003-2115-7835 Se Yong Jung b https://orcid.org/0000-0003-1337-563X Kalthoum Tizaoui b https://orcid.org/0000-0001-8524-6058 Louis Jacob b https://orcid.org/0000-0003-1071-1239 Dong Keon Yon b https://orcid.org/0000-0003-1628-9948 Seung Won Lee b https://orcid.org/0000-0001-5632-5208 Shuji Ogino b https://orcid.org/0000-0002-3909-2323 Eui-Cheol Shin b https://orcid.org/0000-0002-6308-9503 Jae II Shin b https://orcid.org/0000-0003-2326-1820

REFERENCES

- World Health Orgnaization. Global Advisory Committee on Vaccine Safety; 2020. Accessed May 28, 2020. https://www.who.int/ vaccine_safety/committee/reports/May_2020/en/
- Connors M, Graham BS, Lane HC, Fauci AS. SARS-CoV-2 vaccines: much accomplished, much to learn. *Ann Intern Med.* 2021;174: 687-690.
- Kim JH, Marks F, Clemens JD. Looking beyond COVID-19 vaccine phase 3 trials. Nature Med. 2021:27(2):1-7.
- Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. N Engl J Med. 2020;383(27): 2603-2615. doi:10.1056/NEJMoa2034577
- Baden LR, El Sahly HM, Essink B, et al. Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. N Engl J Med. 2020;384: 403-416. doi:10.1056/NEJMoa2035389
- Castells MC, Phillips EJ. Maintaining safety with SARS-CoV-2 vaccines. N Engl J Med. 2021.384(7):643–649.
- Walsh EE, Frenck RW Jr., Falsey AR, et al. Safety and immunogenicity of two RNA-based Covid-19 vaccine candidates. N Engl J Med. 2020;383(25):2439-2450. doi:10.1056/NEJMoa 2027906
- Barnabas RV, Wald A. A Public Health COVID-19 Vaccination Strategy to Maximize the Health Gains for Every Single Vaccine Dose. American College of Physicians.
- Matrajt L, Eaton J, Leung T, Brown ER. Vaccine optimization for COVID-19: who to vaccinate first? Sci Adv. 2020;7(6):eabf1374. doi:10.1126/sciadv.abf1374
- Omer S, Faden R, Kochhar S, et al. WHO SAGE Roadmap for Prioritizing Uses of COVID-19 Vaccines in the Context of Limited Supply. World Health Organization; 2020.
- Doshi P. Will covid-19 vaccines save lives? Current trials aren't designed to tell us. *BMJ*. 2020;371:m4037.
- 12. Lindquist M. VigiBase, the WHO global ICSR database system: basic facts. *Drug Inf J.* 2008;42(5):409-419.

- Shimabukuro TT, Nguyen M, Martin D, DeStefano F. Safety monitoring in the vaccine adverse event reporting system (VAERS). *Vaccine*. 2015;33(36):4398-4405.
- 14. World Health Organization. COVID-19 Vaccines: Safety Surveillance Manual. World Health Organization; 2020.
- Bate A, Lindquist M, Edwards IR, et al. A Bayesian neural network method for adverse drug reaction signal generation. *Eur J Clin Pharmacol.* 1998;54(4):315-321.
- Salem J-E, Manouchehri A, Moey M, et al. Cardiovascular toxicities associated with immune checkpoint inhibitors: an observational, retrospective, pharmacovigilance study. *Lancet Oncol.* 2018;19(12): 1579-1589.
- Salem JE, Ederhy S, Lebrun-Vignes B, Moslehi JJ. Cardiac events associated with chimeric antigen receptor T-cells (CAR-T): a VigiBase perspective. J Am Coll Cardiol. 2020;75(19):2521-2523. doi:10. 1016/j.jacc.2020.02.070
- Johnson DB, Manouchehri A, Haugh AM, et al. Neurologic toxicity associated with immune checkpoint inhibitors: a pharmacovigilance study. J Immunother Cancer. 2019;7(1):1-9.
- Liang X-F, Li L, Liu D-W, et al. Safety of influenza A (H1N1) vaccine in postmarketing surveillance in China. N Engl J Med. 2011;364(7): 638-647.
- 20. Yamayoshi S, Kawaoka Y. Current and future influenza vaccines. *Nature Med.* 2019;25(2):212-220.
- Caster O, Aoki Y, Gattepaille LM, Grundmark B. Disproportionality analysis for pharmacovigilance signal detection in small databases or subsets: recommendations for limiting false-positive associations. *Drug Saf.* 2020;43(5):479-487.
- 22. Chow KT, Gale M Jr., Loo Y-M. RIG-I and other RNA sensors in antiviral immunity. *Annu Rev Immunol*. 2018;36:667-694.
- 23. Wang Y, Zhang Z, Luo J, Han X, Wei Y, Wei X. mRNA vaccine: a potential therapeutic strategy. *Mol Cancer*. 2021;20(1):1-23.
- Slade BA, Leidel L, Vellozzi C, et al. Postlicensure safety surveillance for quadrivalent human papillomavirus recombinant vaccine. JAMA. 2009;302(7):750-757.
- Dolladille C, Ederhy S, Sassier M, et al. Immune checkpoint inhibitor rechallenge after immune-related adverse events in patients with cancer. JAMA Oncol. 2020;6:865-871.
- Salem JE, Yang T, Moslehi JJ, et al. Androgenic effects on ventricular repolarization: a translational study from the international pharmacovigilance database to iPSC-cardiomyocytes. *Circulation*. 2019; 140(13):1070-1080. doi:10.1161/circulationaha.119.040162

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Kim MS, Jung SY, Ahn JG, et al. Comparative safety of mRNA COVID-19 vaccines to influenza vaccines: A pharmacovigilance analysis using WHO international database. *J Med Virol*. 2022;94:1085-1095. doi:10.1002/jmv.27424

VILEY