

## RESEARCH

# Curcumin exerts a protective effect against premature ovarian failure in mice

Zhengjie Yan<sup>1,2</sup>, Youjin Dai<sup>3</sup>, Heling Fu<sup>3</sup>, Yuan Zheng<sup>3</sup>, Dan Bao<sup>3</sup>, Yuan Yin<sup>3</sup>, Qin Chen<sup>3</sup>, Xiaowei Nie<sup>4</sup>, Qingting Hao<sup>3</sup>, Daorong Hou<sup>3</sup> and Yugui Cui<sup>2</sup>

<sup>1</sup>College of Animal Science and Technology, Yangzhou University, Yangzhou, People's Republic of China

<sup>2</sup>State Key Laboratory of Reproductive Medicine, Center of Clinical Reproductive Medicine, The First Affiliated Hospital of Nanjing Medical University, Nanjing, People's Republic of China

<sup>3</sup>Key Laboratory of the Model Animal Research, Animal Core Facility of Nanjing Medical University, Nanjing Medical University, Nanjing, People's Republic of China

<sup>4</sup>Department of Reproductive Medicine, Affiliated Hospital of Nanjing University of Traditional Chinese Medicine, Nanjing, China

Correspondence should be addressed to D Hou or Y Cui: [houdaorong@njmu.edu.cn](mailto:houdaorong@njmu.edu.cn) or [cuiygnj@njmu.edu.cn](mailto:cuiygnj@njmu.edu.cn)

## Abstract

This study was designed to investigate the protective effect of curcumin against D-galactose (D-gal)-induced premature ovarian failure (POF) in mice. A mouse POF model was induced by subcutaneous injection of D-gal (200 mg/kg/day) daily for 42 days. Mice in the curcumin group received both D-gal treatment and intraperitoneal injection of curcumin (100 mg/kg/day) for 42 days. Ovarian function, oxidative stress and apoptosis were evaluated. The P, E2 and SOD levels were higher, and the FSH, LH and MDA levels were significantly lower in the curcumin group than those in the D-gal group. The proportion of primordial follicles was also significantly higher in the curcumin group than that in the D-gal group. In addition, curcumin treatment after D-gal administration resulted in significantly lower *Sod2*, *Cat*, 8-OHdG, 4-HNE, NTY and senescence-associated protein P16 expression levels, higher *Amh* expression levels and less apoptosis in granulosa cells than was observed in the D-gal group. Moreover, the p-Akt, Nrf2 and HO-1 protein expression levels were significantly higher and the apoptosis-related cleaved caspase-3 and -9 protein expression levels were markedly lower in the curcumin group than in the D-gal group. In conclusion, curcumin effectively inhibited D-gal-induced oxidative stress, apoptosis and ovarian injury via a mechanism involving the Nrf2/HO-1 and PI3K/Akt signaling pathways, suggesting that curcumin is a potential protective agent against POF.

## Key Words

- ▶ premature ovarian failure
- ▶ D-galactose
- ▶ curcumin
- ▶ ovary aging
- ▶ oxidative stress

*Journal of Molecular  
Endocrinology*  
(2018) **60**, 261–271

## Introduction

Premature ovarian failure (POF), also called premature ovarian insufficiency (POI), affects approximately 1% of women in the general population, in whom it causes amenorrhea and hypergonadotropic hypoestrogenism before the age of 40 years (Kaufman *et al.* 1988, Waggoner *et al.* 1990, Guerrero *et al.* 2000, Bandyopadhyay *et al.* 2003).

However, the pathological mechanism underlying POF remains unclear. The associated ovarian pathology is related to the toxic effects of galactose and its metabolites at both the ovarian and extraovarian levels (Campbell *et al.* 2010a,b, Rubio-Gozalbo *et al.* 2010). Moreover, there is no effective etiological treatment for POF. Hormone replacement

therapy (HRT) is available to treat the symptoms of POF, and follicle donation is available for some POF patients seeking to become pregnant. However, HRT has been confirmed to confer a high risk of coronary heart disease, endometrial cancer and breast cancer in women with POF (Deady 2004).

POF is a common clinical feature of galactosemia (Waggoner *et al.* 1990, Guerrero *et al.* 2000), and women with galactosemia eventually develop POF (Kaufman *et al.* 1988). The estrous cycle of female mice is similar to but shorter than that of humans. The mouse D-galactose (D-gal)-induced POF model is used as a model of aging in mice and has been widely used to study the mechanisms underlying ovarian aging, because the accelerated aging observed in this model is very similar to observations in humans (Song *et al.* 1999, Semba *et al.* 2010). Aging is associated with increased deposition of advanced glycation end products (AGEs) and reactive oxygen species (ROS) in the myocardium, brain, liver, eye, red blood cells, kidney, bone, ovary, muscles and tendons (Kimura *et al.* 1996, Nerlich & Schleicher 1999, Schinzel *et al.* 2001, Simm *et al.* 2004, Odetti *et al.* 2005, Haus *et al.* 2007, Kumar *et al.* 2007, Hyogo & Yamagishi 2008). Many studies have indicated that AGEs and ROS exacerbate and accelerate the aging process and contribute to the early phases of age-related diseases, including atherosclerosis, cataracts, neurodegenerative diseases, renal failure, arthritis, ovarian senescence and age-related macular degeneration (Tian *et al.* 2005, Semba *et al.* 2010). D-gal administration can cause excessive ROS formation and AGE accumulation (Song *et al.* 1999, Mao *et al.* 2010, Lin *et al.* 2012). ROS-induced damage and AGE accumulation are widely accepted causes of aging that gradually damage ovarian functions (Semba *et al.* 2010). Therefore, D-gal has been used to induce the POF model (Bandyopadhyay *et al.* 2003). Curcumin (chemical name: 1, 7-bis (4-hydroxy-3-methoxyphenyl)-1, 6-heptadiene-3, 5-dione) (Aggarwal *et al.* 2005, Ak & Gülçin 2008) is an active component of turmeric rhizomes (*Curcuma longa* Linn), which contain 3–5% curcumin (Zhang *et al.* 2013). Curcumin has been scientifically demonstrated to function as an antioxidant (Fujisawa *et al.* 2004, Calabrese *et al.* 2008, Dinkova-Kostova & Talalay 2008, Augustyniak *et al.* 2010), anti-inflammatory (Aggarwal & Harikumar 2009, Wang *et al.* 2012, Ueki *et al.* 2013), anti-apoptotic (Geng *et al.* 2017) and antibacterial (Mun *et al.* 2013) substance. An increasing number of studies have shown that curcumin directly suppresses proliferation and promotes apoptosis in ovarian cancer cells (Shehzad *et al.* 2010, Watson *et al.* 2010, Terlikowska *et al.* 2014, Vallianou *et al.* 2015, Seo *et al.* 2016) and prevents the adverse effects

of ovarian insufficiency (Tiwari-Pandey & Ram Sairam 2009, Voznesens'ka *et al.* 2010, Alekseyeva *et al.* 2011, Aktas *et al.* 2012), ionizing radiation (Aktas *et al.* 2012), ischemia (Eser *et al.* 2015), oxidative stress (Qin *et al.* 2015) and mycotoxins (Qin *et al.* 2015) on ovarian function. Some studies have also demonstrated that curcumin and its analogues exert a stimulatory effect on ovarian functions, because they promote proliferation and reduce apoptosis in murine ovarian cells (Voznesens'ka *et al.* 2010, Aktas *et al.* 2012) while supporting murine ovarian folliculogenesis (Voznesens'ka *et al.* 2010, Alekseyeva *et al.* 2011) and steroidogenesis (Tiwari-Pandey & Ram Sairam 2009). However, the effects of curcumin on experimentally D-gal-induced ovarian aging have not been reported. In this study, we investigate the protective effect of curcumin on ovarian aging in a D-gal-induced mouse model of POF.

## Materials and methods

### Animals and treatment

A total of 60 C57BL/6 female mice aged 7–8 weeks were used. The animals were obtained from the Animal Core Facility of Nanjing Medical University, Nanjing, China and housed under a 12-h darkness/light cycle in an animal facility with a controlled temperature (20–25°C) and humidity (40%–70%). Food and water were provided *ad libitum* throughout the study. The mice were allowed to acclimatize for 1 week. Then, they were randomly divided into the three following groups with 20 mice per group: control group, D-gal group and curcumin group. The mice in the D-gal group were subcutaneously (s.c.) injected daily with D-gal (200 mg/kg/day) for 42 days (Park & Choi 2012, He *et al.* 2017), the mice in the control group received an equal volume of saline (s.c. daily) for 42 days and the mice in the curcumin group received curcumin (100 mg/kg/day) (Chainani-Wu 2003, Aktas *et al.* 2012) via intraperitoneal injection following a daily D-gal injection for 42 days. All experiments involving animals were approved by the Institutional Animal Care and Use Committee (IACUC) of Nanjing Medical University, and the methods were conducted in accordance with the approved guidelines. On the last day of drug administration (day 42), all mice were killed while under general anesthesia (induced using an intraperitoneal injection of pentobarbital sodium (150 mg/kg)). Blood was collected through a heart puncture, the left ovary was immediately excised and stored at –80°C for biochemical analysis and the right ovary was fixed in 4% paraformaldehyde for histological studies.

### Primordial follicle counting

The right ovaries were fixed for 12h in 4% paraformaldehyde and then embedded in paraffin. The tissues were serially sectioned (6- $\mu$ m thick), mounted on glass slides and stained with H&E. The ovarian follicles were counted according to the methods described in a previous study (Tilly 2003, Bernal *et al.* 2010). Briefly, every fifth section was observed under a microscope. To avoid repeated counting of the same follicle, only follicles with a visible oocyte nucleus were included. The numbers of primordial follicles in all serial sections of an ovary were counted. The following follicle classification was used (Hirshfield & Midgley 1978, Wang *et al.* 1991, Borgeest *et al.* 2002): type 1: primordial follicle, one layer of flattened granulosa cells surrounding the oocyte; type 2: primary follicle, one to two complete layers of cuboidal granulosa cells; type 3: secondary follicle, an oocyte surrounded by more than one layer of cuboidal granulosa cells with no visible antrum; type 4: antral follicle, an oocyte surrounded by multiple layers of cuboidal granulosa cells and containing one or more antral spaces, possibly with a cumulus oophorus and thecal layer and type 5: atretic follicle, a follicle that enters a degenerative process without ovulation. The oocyte nuclei in the atretic follicles shrink, the chromosomes and cytoplasm dissolve, the granulosa layer decreases and the follicular membrane cells are hypertrophic (Paulose *et al.* 2012). The primordial follicle ratio refers to the percentage of the primordial follicle number out of the total follicles. The atretic follicles were included in the denominator when calculating the proportion of primordial follicles or total follicles.

### Sample preparation and biochemical assays

All blood samples were collected while the mice were in diestrus and allowed to clot at room temperature. Then, the samples were centrifuged at 900g for 10min to harvest serum. Serum biochemical parameters, including the serum follicle-stimulating hormone (FSH), luteinizing hormone (LH), progesterone (P) and estradiol (E2) levels, were measured spectrophotometrically (Eon, BioTeK, Vermont, UT, USA) using the following commercially available ELISA kits: FSH (KA2330), LH (KA2332) (Novus Biologicals, Littleton, USA), E2 (582251) and P (582601) (Cayman Chemicals, Ann Arbor, MI, USA). The ovaries were washed in ice-cold saline and homogenized in 0.1M Tris-HCl buffer (pH 7.4). The homogenates were centrifuged at 10,000g for 15min, and the supernatants

were centrifuged at 100,000g for 1h. The resulting supernatant (cytosolic fraction) was used to determine the enzymatic activity and lipid peroxidation levels. The biochemical parameters of the ovaries, including the total superoxide dismutase (SOD) activity and malondialdehyde (MDA) level, were measured spectrophotometrically using commercially available kits for SOD (A001-1) and MDA (A003-1) (Jiancheng Bioengineering Institute, Nanjing, China).

### Q-PCR

Total mRNA was extracted from the ovarian samples using the TRIzol reagent (B5704-1, Takara) according to the manufacturer's instructions and then treated with DNase I (2212, Takara) according to the manufacturer's protocol. The quality and quantity of the RNA were determined using a spectrophotometer (NanoDrop 2000c, Thermo Scientific). cDNA was immediately synthesized using the PrimeScript RT Reagent Kit (RR037A, Takara) according to the manual supplied by the manufacturer. Q-PCR was performed using a Light Cycler PCR QC Kit (Roche) and a 7300 Real-Time PCR System (LC96, Roche). The PCR primers are listed in [Supplementary Table 1](#) (see section on [supplementary data](#) given at the end of this article). The housekeeping gene GAPDH was used as the internal reference. Expression of the target gene was normalized to GAPDH and calculated using the comparative quantification method ( $2^{-\Delta\Delta Ct}$ ). Expression of the target genes was corrected to GAPDH prior to normalization. Firstly, the Ct value of each group was subtracted from the Ct value of the internal reference gene, which was named  $\Delta Ct$ , as follows:  $\Delta Ct = Ct(\text{target gene}) - Ct(\text{internal reference gene})$ . Secondly, the  $\Delta Ct$  of the experimental group was subtracted from the control group, and then the inverse of all of the results was taken to obtain  $-\Delta\Delta Ct$ . Finally, the power operation of  $-\Delta\Delta Ct$  was performed in 2. The GraphPad Prism 5 software was used for chart production.

### Immunohistochemical staining

For the immunohistochemical analysis, paraffin-embedded sections were dewaxed and then subjected to heat-mediated antigen retrieval, which was performed by microwaving the sections for 20min in 10mM sodium citrate (pH 6.0). The sections were allowed to cool for 15min, briefly washed in deionized water and then rinsed twice in PBS. The sections were incubated for 30min in 5% goat serum in DPBS containing 0.1%

Tween and 0.5% BSA. The sections were incubated overnight at 4°C with primary antibodies against 8-hydroxyguanosine (ab26842), 4-hydroxynonenal (ab48506), anti-CDKN2A/p16 (ab189034) (Abcam Biotechnology) and nitrotyrosine (sc-71007) (Millipore Biotechnology) at the appropriate dilutions. The secondary antibody in the Dako REAL EnVisio Detection System (K5007) (DAKO) was used to detect labeling. Then, the specimens were counterstained with hematoxylin for 1 min. All sections were incubated under the same conditions and at the same time using the same antibody concentrations. The tissue sections were observed and photographed with a microscope and semi-quantified using the Image Pro Plus 6.0 software. The integrated optical density (IOD) was collected for each photograph. Five fields for each slice (five slides per animal) were randomly selected for blinded measurements ( $n=8$  per group). The images were quantified by the immunoreactive area (IA) in  $\mu\text{m}^2$  and the IOD. The staining intensity (SI) for each image was calculated as  $\text{SI}=\text{IOD}/\text{IA}$ , and the mean with standard deviation was obtained for each series.

#### **In situ TUNEL fluorescence staining assay**

The terminal deoxynucleotidyl transferase (TdT)-mediated deoxyuridine triphosphate (dUTP) nick end-labeling (TUNEL) assay was performed according to the manufacturer's instructions (11684817910, Roche). Ovarian tissues were fixed in 4% paraformaldehyde overnight, dehydrated, embedded in paraffin, cut into 4- $\mu\text{m}$ -thick sections and placed on numbered polylysine-coated glass slides. Deparaffinized tissue sections were incubated with proteinase K (20 mg/mL) in a humidified chamber for 15 min, and endogenous peroxidase activity was blocked by treating the sections with 3%  $\text{H}_2\text{O}_2$  for 10 min. The sections were incubated with TdT labeling buffer at 37°C for 1 h in a moist chamber. Then, the sections were counterstained with DAPI. The TUNEL-positive cells stained green, and the nuclei were stained with DAPI (blue). To eliminate the histological differences between ovarian tissues, five random fields per slide (five slides per animal, eight animals per group,  $n=8$ ) were examined. In total, 200 random fields ( $5 \times 5 \times 8=200$ ) per group were checked. The TUNEL-positive granulosa cells and the total granulosa cells in the antral follicles were counted. The rate of TUNEL-positive granulosa cells (%) in the antral follicles was analyzed using the Image Pro Plus 6.0 software.

#### **Western blotting**

The Western blotting analysis was performed according to the methods described by Banerjee *et al.* (2012). Briefly, 50  $\mu\text{g}$  of total lysate obtained from ovarian tissue was subjected to 15% polyacrylamide gel electrophoresis and then transferred to a cellulose acetate membrane. The membranes were blocked with 1 $\times$  casein solution for approximately 4 h and then incubated with rabbit monoclonal anti-Akt (4685), anti-p-Akt (Ser473) (4058), anti-cleaved caspase-3 (Asp175) (9664) or anti-cleaved caspase-9 (Asp330) (7237) antibodies obtained from Cell Signaling Technology, rabbit polyclonal anti-Nrf2 (sc-722) or anti-GAPDH (H-12) (sc-166574) antibodies or a mouse monoclonal anti-HO-1 (sc-390991) antibody obtained from Santa Cruz Biotechnology in blocking buffer for 2 h at room temperature. The membranes were washed 3 times with TBST and then incubated with a goat anti-rabbit IgG HRP-conjugated secondary antibody (sc-2004, Santa Cruz Biotechnology) or a goat anti-mouse IgG HRP-conjugated secondary antibody (sc-2005, Santa Cruz Biotechnology). Then, the membranes were washed 3 times in TBST, and the blots were imaged using the ChemiDoc XRS+ Molecular Imager (Bio-Rad) with the Pierce ECL Western Blotting Substrate (32209, Thermo Scientific) and analyzed using image analysis software (ImageJ 1.42). The housekeeping protein GAPDH was used as the internal reference. The Western blotting quantification was corrected to GAPDH expression prior to normalization.


#### **Statistical analysis**

All statistical analyses were performed using SPSS v.16. All results are shown as the mean  $\pm$  standard deviation ( $M \pm \text{s.d.}$ ). All statistical comparisons were performed using one-way ANOVA followed by Duncan's multiple range *post hoc* analysis. A  $P$  value  $<0.05$  was considered significant.

## **Results**

#### **Protective effects of curcumin on the HPG axis and ovarian *Amh* mRNA expression**

The D-gal treatment group had significantly increased serum FSH and LH levels ( $P<0.05$  and  $P<0.01$ ) and significantly decreased E2 and P levels ( $P<0.01$ ) compared to those in the control (Fig. 1A, B, C and D). Due to insufficient sensitivity and stability, an enzyme immunoassay (ELISA) was not used to test the serum anti-Müllerian hormone (AMH) level. Instead, *Amh* mRNA expression in the ovarian tissue was tested using q-PCR. Interestingly, *Amh*

**Figure 1**

Protective effects of curcumin on the HPG axis and ovarian AMH. The serum FSH (A), LH (B), E2 (C) and P (D) levels were tested in three groups, and AMH mRNA expression in the ovarian tissues (E) was also tested. Ten mice per group were examined. All data are shown as the mean  $\pm$  s.d. Statistical significance: \* $P$ <0.05 and \*\* $P$ <0.01 vs the control group, # $P$ <0.05 and ## $P$ <0.01 vs the D-gal group.


mRNA expression was significantly decreased in the D-gal treatment group compared with that in the control group (Fig. 1E). In the POF model induced by D-gal, curcumin treatment significantly decreased the serum FSH and LH levels ( $P$ <0.05, Fig. 1A and B) and increased the serum E2 and P levels ( $P$ <0.01, Fig. 1C and D) and the ovarian *Amh* expression level ( $P$ <0.05, Fig. 1E).

### Protective effect of curcumin on follicular development in the POF model


Follicle counting was performed after H&E staining (Fig. 2A, B, C, D, E and F). The follicle classification was based on the characteristics proposed by Hirshfield & Midgley (1978). Counting of the primordial follicles showed that D-gal treatment reduced the proportion of primordial follicles compared to that in the control group ( $P$ <0.01, Fig. 2G and H). Significantly more follicles were counted at the different developmental stages of maturation in the curcumin treatment group than in the D-gal model group ( $P$ <0.05 or  $P$ <0.01, Fig. 2H).

### Effects of D-gal and curcumin on oxidative stress

The total SOD enzyme activity was significantly lower in the ovarian tissues from the D-gal treatment group than in the tissues from the control group ( $P$ <0.01, Fig. 3A), and the MDA level was significantly higher in the treated group than in the control group ( $P$ <0.01, Fig. 3B). Curcumin treatment resulted in markedly higher SOD levels and lower MDA levels than the levels in the D-gal group (all  $P$ <0.01, Fig. 3A and B). *Cat* and *Sod2* mRNA expression was tested using Q-PCR (Fig. 3C and D). Corresponding with the changes in the SOD activity and MDA level, *Cat* expression was significantly increased in the D-gal group compared to that in the control group ( $P$ <0.01), and curcumin treatment partially but significantly rescued the effect of D-gal ( $P$ <0.05, Fig. 3C). However, the change in *Sod2* expression was seemingly abnormal. The *Sod2* mRNA expression level was significantly increased in the D-gal model ( $P$ <0.01, Fig. 3D), whereas curcumin treatment partially and significantly rescued the effect of D-gal ( $P$ <0.01).

**Figure 2**

Effect of curcumin on follicular development in the POF model. Follicles were observed after H&E staining (A, B, C, D, E and F). The follicle classification was based on the characteristics proposed by Hirshfield and Midgley. (G) Primordial follicle ratio counting was performed in the three groups. Ten mice per group were examined. The primordial follicle ratio refers to the percentage of the primordial follicle number to the total follicle number. (H) The numbers of those follicles at different developmental stages of maturation were summarized. \* $P$ <0.05 and \*\* $P$ <0.01 vs the control group, # $P$ <0.05 and ## $P$ <0.01 vs the D-gal group.

**Figure 3**

Effects of D-gal and curcumin on oxidative stress. The total SOD enzyme activity (A) and the MDA level (B) were measured in the ovarian tissues. *Cat* (C) and *Sod2* (D) mRNA expression was tested using Q-PCR. Ten mice per group were examined. All data are shown as the mean  $\pm$  s.d. Statistical significance: \* $P < 0.05$  and \*\* $P < 0.01$  vs the control group, # $P < 0.05$  and ## $P < 0.01$  vs the D-gal group.



### Protective effect of curcumin on D-gal-induced ovarian cell apoptosis

In the TUNEL assay, the nuclei of the TUNEL-positive (apoptotic) cells stained green (Fig. 4A). The number of apoptotic granulosa cells in the antral follicles was assessed in the three groups. More TUNEL-positive cells were detected in the D-gal group than in the control group ( $P < 0.01$ , Fig. 4B). Curcumin significantly decreased the number of TUNEL-positive cells compared with the number in the D-gal model group, suggesting a protective effect on D-gal-induced ovarian cell apoptosis ( $P < 0.01$ ).

### Effects of curcumin on 4-HNE, NTY, 8-OHdG and p16 protein expression

The cellular locations of the 4-HNE, NTY, 8-OHdG and p16 proteins were examined using immunohistochemistry (Fig. 5). The 4-HNE, NTY and 8-OHdG proteins were mainly located in the ovarian interstitial cells. The 4-HNE, 8-OHdG and NTY protein expression levels were significantly higher in the D-gal group than in the control group ( $P < 0.01$ , Fig. 5B), whereas the expression levels were partially and significantly decreased in the curcumin treatment group ( $P < 0.01$  and  $P < 0.05$ , respectively).

Interestingly, the senescence-associated p16 protein was mainly located in follicular granulosa cells and oocytes (Fig. 5A), and relatively lower p16 expression was observed in the ovarian interstitial cells. Corresponding to the 4-HNE, 8-OHdG and NTY protein expression levels, D-gal treatment induced a significant increase in p16


**Figure 4**

Effect of curcumin on D-gal-induced apoptosis of ovarian cells. Apoptosis was analyzed using *in situ* TUNEL fluorescence staining (A). In the TUNEL assay, the nuclei of TUNEL-positive (apoptotic) cells stained green. Five random fields per section (five sections per tissue from a mouse) were examined in each experiment. The TUNEL-positive granulosa cells and total granulosa cells in the antral follicles were counted. The number of TUNEL-positive granulosa cells (%) was compared among the three groups ( $n = 8$ ) (B). Statistical significance: \* $P < 0.01$  and \*\* $P < 0.01$  vs the control group; # $P < 0.01$  and ## $P < 0.01$  vs the D-gal group.

protein expression, whereas curcumin treatment rescued the effect of D-gal treatment ( $P < 0.01$ , Fig. 5B).

### Effect of curcumin on related pathways in the ovary

We investigated the potential mechanisms involved in the effects of curcumin on D-gal-induced ovarian aging. The expression levels of apoptosis- and oxidative stress signal pathway-associated markers were assessed by Western blotting (Fig. 6). Total Akt protein expression was the same in each group, whereas the p-Akt level was markedly lower in the D-gal group than in the control group ( $P < 0.01$ ) and was significantly higher in the curcumin group than in the D-gal group ( $P < 0.01$ ). The cleaved caspase-3 and cleaved

**Figure 5**

Effects of curcumin on 4-HNE, NTY, 8-OHdG and p16 protein expression. The cellular locations of these proteins were observed using immunohistochemistry (A), and 4-HNE, NTY, 8-OHdG and p16 expression was quantitatively analyzed ( $n=8$ ) (B). Data are shown as the mean  $\pm$  s.d. Statistical significance: \* $P < 0.01$  and \*\* $P < 0.01$  vs the control group; # $P < 0.05$  and ## $P < 0.01$  vs the d-gal group.

caspase-9 levels were significantly lower in the curcumin group than in the D-gal group ( $P < 0.01$ ); moreover, the Nrf2 and HO-1 protein expression levels were significantly lower in the D-gal group than in the control group ( $P < 0.01$ ), whereas the Nrf2 and HO-1 expression levels were significantly higher in the curcumin group than in the D-gal group ( $P < 0.05$  and  $P < 0.01$ , respectively).

## Discussion


In the present study, a mouse POF model was successfully induced by D-galactose. D-galactose treatment resulted in increased ROS and AGEs, increased granulosa cell apoptosis and damaged follicular development. Curcumin partially rescued the effects of D-gal in the mouse POF model via a mechanism involving the Nrf2/HO-1 and PI3K/Akt signaling pathways.

Many studies have shown that D-gal directly induces oxidative stress *in vivo* and that galactose toxicity attenuates FSH bioactivity and inhibits E2 production from granulosa cells (Banerjee *et al.* 2012). We found that the serum FSH and LH levels were significantly increased and the E2 and P levels were decreased in mice treated with D-gal. The number of atretic follicles was increased by the D-gal treatment; conversely, the decreased number of primordial, primary and secondary follicles resulted in a decreased number of total follicles, which was in line with a previous report (Bandyopadhyay *et al.* 2003). AMH is a very important early marker of ovarian aging (Visser *et al.* 2006, Soto *et al.* 2009, Tolikas *et al.* 2011) and reflects the size of the ovarian follicle pool (Feyereisen *et al.* 2006). Lower *Amh* expression

has been detected long before normal menopause (Sanders *et al.* 2009, Seifer *et al.* 2011) and during the normal course of aging in mice (Kevenaar *et al.* 2006). We also found that ovarian *Amh* expression was significantly decreased by D-gal. Thus, the use of D-gal successfully induced the mouse POF model in the present study.

Curcumin is an antioxidant (Augustyniak *et al.* 2010), anti-inflammatory (Ueki *et al.* 2013), anti-apoptotic (Geng *et al.* 2017) and antibacterial (Mun *et al.* 2013) substance. Curcumin was demonstrated to exert a stimulatory effect on ovarian function, because curcumin promoted folliculogenesis and reduced apoptosis in murine ovarian cells (Voznesens'ka *et al.* 2010, Aktas *et al.* 2012). This study was designed to investigate the protective effects of curcumin on ovarian aging in a D-gal-induced mouse model of POF. We found that curcumin decreased the FSH and LH levels and increased the E2 and P levels in the mouse POF model. Interestingly, ovarian AMH expression was significantly increased by curcumin, whereas the total number of follicles increased with the increased numbers of primordial, primary and secondary follicles. Curcumin partially and significantly rescued the effects of D-gal in the POF model, suggesting that curcumin might regulate the reproductive endocrine function and promote follicular development or the maintenance of primordial follicles.

We found that the total SOD enzyme activity was decreased and the MDA level was increased in the D-galactose-induced POF model, whereas *Cat* and *Sod2* expression was significantly increased. Superoxide dismutase (SOD) is a type of enzyme that alternatively catalyzes dismutation of the superoxide ( $O_2^-$ ) radical



**Figure 6**

Effect of curcumin on the expression of apoptosis- and oxidative stress-associated factors in ovarian tissues. The Akt, p-Akt, cleaved caspase-3, cleaved caspase-9, Nrf2 and HO-1 protein expression levels in ovarian tissues were observed using Western blotting (A). The Akt, p-Akt, Nrf2 and HO-1 protein expression levels were quantitatively analyzed ( $n=8$ ) (B). Data are shown as the mean  $\pm$  s.d. Statistical significance: \* $P<0.01$  and \*\* $P<0.01$  vs the control group; # $P<0.05$  and ## $P<0.01$  vs the D-gal group.

into either ordinary molecular oxygen ( $O_2$ ) or hydrogen peroxide ( $H_2O_2$ ). The increased ROS production induced by D-gal is related to the inhibition of SOD activity. Three forms of superoxide dismutase (SOD1, SOD2 and SOD3) are expressed in all mammals. SOD2, which is the mitochondrial SOD and is also known as manganese-dependent superoxide dismutase (MnSOD), is a member of the iron/manganese superoxide dismutase family (Scandalios 2005). SOD2 and catalase are the two most important antioxidant enzymes in the body, and their expression levels are increased following oxidative stress (Scandalios 2005, Zhang *et al.* 2014). D-galactose, which is a reducing sugar that reacts readily with the free amines of amino acids in proteins and peptides to form AGEs

(Bucala & Cerami 1992, Vlassara *et al.* 1994), produces acetaldehyde and hydrogen peroxide under the action of galactose oxidase and thus increases ROS, resulting in the aging of cells (Ho *et al.* 2003). Several studies have shown that oxidative stress can reduce the number of follicles and oocytes (Tarín 1995, 1996). Miyamoto *et al.* (2010) reported that oxidative stress caused a significant decrease in the number of follicles and the ovulated oocytes during repeated ovulation. We found that curcumin increased the total SOD activity and decreased MDA. Curcumin exerts a protective effect by inhibiting oxidative stress in a D-gal-induced model of ovarian aging, and its effects may be mediated via the suppression of ROS and superoxide anion free radicals (Suckow & Suckow 2006, Sikora *et al.* 2010).

The results showed that 4-HNE, NTY and 8-OHdG expression was mainly located in ovarian interstitial cells, and curcumin decreased the expression of those proteins. The D-gal-induced effects on ROS, AGEs and other factors on cellular aging should be extensive (Song *et al.* 1999). Factors such as 8-OHdG, 4-HNE and NTY, which are mainly located in somatic cells, are involved in the aging process related to ROS and AGEs. Interestingly, the p16 protein was highly expressed in ovarian follicles, including oocytes, granular cells and cumulus cells, and expressed at low levels in interstitial cells. The p16 protein negatively regulates cell proliferation and division and promotes apoptosis and senescence; thus, it affects the cell cycle and G1-S conversion by competitively suppressing the combination between CDK4/6 and cyclin D within G1 phase. We found that p16 expression in follicles was significantly increased in the D-gal-induced model and that curcumin significantly decreased D-gal-induced p16 expression. The p16 expression level was positively related to the follicular number and negatively related to the  $E_2$  and P levels. p16 expression was low in young animal tissues and subsequently increased with age (Kim *et al.* 2002, Attema *et al.* 2009). Excessive p16 expression can induce premature aging in cells (Attema *et al.* 2009). Baker and coworkers found that the aging cells reactivated and aging-associated phenotypes were reduced by silencing p16 protein expression (Baker *et al.* 2011). However, how p16 participates in ovarian aging through oxidative stress is unclear (Kim *et al.* 2002, Attema *et al.* 2009). We visualized an important effect of p16 on oxidative stress in ovarian follicles and found that curcumin also exerted a protective effect against ovarian aging partially through the downregulation of p16.



In the present study, we found that curcumin partially rescued the effects of D-gal via multiple mechanisms, including the inhibition of oxidative stress and granular cell apoptosis, as described in other reports (Suckow & Suckow 2006, Sikora *et al.* 2010). Our results also showed that curcumin significantly increased p-Akt, HO-1 and Nrf2 expression in ovarian tissues and decreased the expression of the apoptosis-related proteins cleaved caspase-3 and -9. PI3K/AKT signaling is widely accepted as a key pathway that regulates cell survival (Atif *et al.* 2015). Upregulation of phosphorylated PI3K and AKT can inhibit both the release of Bad and Bax and the activation of downstream pro-apoptotic proteins, such as caspases, resulting in the suppression of apoptosis (Rong & Distelhorst 2008, Atif *et al.* 2015). Nrf2 can bind to antioxidant response elements (AREs) in the promoter regions of Nrf2 target genes, which remove ROS via sequential enzymatic reactions, such as HO-1 (Gozzelino *et al.* 2010). The induction of HO-1 is an adaptive response to oxidative stress (Paine *et al.* 2010). Curcumin has also been reported to activate Nrf2 to upregulate enzymes involved in antioxidant defenses, including SOD and HO-1 (Balogun *et al.* 2003, Zingg *et al.* 2013). Our results indicated that curcumin attenuated D-gal-induced oxidative stress injury in the ovary partly by activating the PI3K/Akt and Nrf2/HO-1 pathways.

In conclusion, curcumin effectively inhibited D-gal-induced oxidative stress, apoptosis and ovarian injury via multiple mechanisms, including the Nrf2/HO-1 and PI3K/Akt signaling pathways. These results suggest that curcumin as an antioxidant is a potential protective agent against POF.

#### Supplementary data

This is linked to the online version of the paper at <https://doi.org/10.1530/JME-17-0214>.

#### Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the reported research.

#### Funding

This work was supported by the the National Key Research and Development Program of China (2017YFC1001602), Science and Technology Development Key Projects Fund of Nanjing Medical University (2016NJMUZD020), the Nature and Science Foundation of China (81270746), the Jiangsu Province Special Program of Medical Science (BL2012009 and ZX201110) and a project funded by PAPD of the Priority Academic Program Development of Jiangsu High Education Institutions (JX10231802).

## References

- Aggarwal BB & Harikumar KB 2009 Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *International Journal of Biochemistry and Cell Biology* **41** 40–59. (<https://doi.org/10.1016/j.biocel.2008.06.010>)
- Aggarwal BB, Aggarwal MS & Shishodia S 2005 Curcumin derived from turmeric (*Curcuma longa*): a spice for all seasons. *Phytopharmaceuticals in Cancer Chemoprevention* **23** 349–387. (<https://doi.org/10.1201/9780203506707.ch23>)
- Ak T & Gülçin I 2008 Antioxidant and radical scavenging properties of curcumin. *Chemico-biological Interactions* **174** 27–37. (<https://doi.org/10.1016/j.cbi.2008.05.003>)
- Aktas C, Kanter M & Kocak Z 2012 Antiapoptotic and proliferative activity of curcumin on ovarian follicles in mice exposed to whole body ionizing radiation. *Toxicology and Industrial Health* **28** 852–863. (<https://doi.org/10.1177/0748233711425080>)
- Alekseyeva IN, Makogon NV, Bryzgina TM, Voznesenskaya TY & Sukhina VS 2011 Effects of NF- $\kappa$ B blocker curcumin on oogenesis and immunocompetent organ cells in immune ovarian injury in mice. *Bulletin of Experimental Biology and Medicine* **151** 432–435. (<https://doi.org/10.1007/s10517-011-1349-1>)
- Atif F, Yousuf S & Stein DG 2015 Anti-tumor effects of progesterone in human glioblastoma multiforme: role of PI3K/Akt/mTOR signaling. *Journal of Steroid Biochemistry and Molecular Biology* **146** 62–73. (<https://doi.org/10.1016/j.jsbmb.2014.04.007>)
- Attema JL, Pronk CJ, Norddahl GL, Nygren JM & Bryder D 2009 Hematopoietic stem cell ageing is uncoupled from p16 INK4A-mediated senescence. *Oncogene* **28** 2238–2243. (<https://doi.org/10.1038/onc.2009.94>)
- Augustyniak A, Bartosz G, Cipak A, Duburs G, Horáková L, Luczaj W, Majekova M, Odysseus AD, Rackova L, Skrzydlewska E, *et al.* 2010 Natural and synthetic antioxidants: an updated overview. *Free Radical Research* **44** 1216–1262. (<https://doi.org/10.3109/10715762.2010.508495>)
- Baker DJ, Wijshake T, Tchkonja T, LeBrasseur NK, Childs BG, van de Sluis B, Kirkland JL & van Deursen JM 2011 Clearance of p16<sup>INK4a</sup>-positive senescent cells delays ageing-associated disorders. *Nature* **479** 232–236. (<https://doi.org/10.1038/nature10600>)
- Balogun E, Hoque M, Gong P, Killeen E, Green CJ, Foresti R, Alam J & Motterlini R 2003 Curcumin activates the haem oxygenase-1 gene via regulation of Nrf2 and the antioxidant-responsive element. *Biochemical Journal* **371** 887–895. (<https://doi.org/10.1042/bj20021619>)
- Bandyopadhyay S, Chakrabarti J, Banerjee S, Pal AK, Goswami SK, Chakravarty BN & Kabir SN 2003 Galactose toxicity in the rat as a model for premature ovarian failure: an experimental approach readdressed. *Human Reproduction* **18** 2031–2038. (<https://doi.org/10.1093/humrep/deg414>)
- Banerjee S, Chakraborty P, Saha P, Bandyopadhyay SA, Banerjee S & Kabir SN 2012 Ovotoxic effects of galactose involve attenuation of follicle-stimulating hormone bioactivity and up-regulation of granulosa cell p53 expression. *PLoS ONE* **7** e30709. (<https://doi.org/10.1371/journal.pone.0030709>)
- Bernal AB, Vickers MH, Hampton MB, Poynton RA & Sloboda DM 2010 Maternal undernutrition significantly impacts ovarian follicle number and increases ovarian oxidative stress in adult rat offspring. *PLoS ONE* **5** e15558. (<https://doi.org/10.1371/journal.pone.0015558>)
- Borgeest C, Symonds D, Mayer LP, Hoyer PB & Flaws JA 2002 Methoxychlor may cause ovarian follicular atresia and proliferation of the ovarian epithelium in the mouse. *Toxicological Sciences* **68** 473–478. (<https://doi.org/10.1093/toxsci/68.2.473>)
- Bucala R & Cerami A 1992 Advanced glycosylation: chemistry, biology, and implications for diabetes and aging. *Advances in Pharmacology* **23** 1–34. ([https://doi.org/10.1016/S1054-3589\(08\)60961-8](https://doi.org/10.1016/S1054-3589(08)60961-8))

- Calabrese V, Bates TE, Mancuso C, Cornelius C, Ventimiglia B, Cambria MT, Di Renzo L, De Lorenzo A & Dinkova-Kostova AT 2008 Curcumin and the cellular stress response in free radical-related diseases. *Molecular Nutrition and Food Research* **52** 1062–1073. (<https://doi.org/10.1002/mnfr.200700316>)
- Campbell BK, Kendall NR, Onions V & Scaramuzzi RJ 2010a The effect of systemic and ovarian infusion of glucose, galactose and fructose on ovarian function in sheep. *Reproduction* **140** 721–732. (<https://doi.org/10.1530/REP-10-0185>)
- Campbell BK, Onions V, Kendall NR, Guo L & Scaramuzzi RJ 2010b The effect of monosaccharide sugars and pyruvate on the differentiation and metabolism of sheep granulosa cells *in vitro*. *Reproduction* **140** 541–550. (<https://doi.org/10.1530/REP-10-0146>)
- Chainani-Wu N 2003 Safety and anti-inflammatory activity of curcumin: a component of tumeric (*Curcuma longa*). *Journal of Alternative and Complementary Medicine* **9** 161–168. (<https://doi.org/10.1089/107555303321223035>)
- Deady J 2004 Clinical monograph: hormone replacement therapy. *Journal of Managed Care Pharmacy* **10** 33–47. (<https://doi.org/10.18553/jmcp.2004.10.1.33>)
- Dinkova-Kostova AT & Talalay P 2008 Direct and indirect antioxidant properties of inducers of cytoprotective proteins. *Molecular Nutrition and Food Research* **52** S128–S138. (<https://doi.org/10.1002/mnfr.200700195>)
- Eser A, Hizli D, Haltas H, Namuslu M, Kosus A, Kosus N & Kafali H 2015 Effects of curcumin on ovarian ischemia-reperfusion injury in a rat model. *Biomedical Reports* **3** 807–813. (<https://doi.org/10.3892/br.2015.515>)
- Feyereisen E, Méndez Lozano DH, Taiab J, Hesters L, Frydman R & Fanchin R 2006 Anti-Müllerian hormone: clinical insights into a promising biomarker of ovarian follicular status. *Reproductive Biomedicine Online* **12** 695–703. ([https://doi.org/10.1016/S1472-6483\(10\)61081-4](https://doi.org/10.1016/S1472-6483(10)61081-4))
- Fujisawa S, Atsumi T, Ishihara M & Kadoma Y 2004 Cytotoxicity, ROS-generation activity and radical-scavenging activity of curcumin and related compounds. *Anticancer Research* **24** 563–569.
- Geng X, Hong Q, Wang W, Zheng W, Li O, Cai G, Chen X & Wu D 2017 Biological membrane-packed mesenchymal stem cells treat acute kidney disease by ameliorating mitochondrial-related apoptosis. *Scientific Reports* **7** 41136. (<https://doi.org/10.1038/srep41136>)
- Gozzelino R, Jeney V & Soares MP 2010 Mechanisms of cell protection by heme oxygenase-1. *Annual Review of Pharmacology and Toxicology* **50** 323–354. (<https://doi.org/10.1146/annurev.pharmtox.010909.105600>)
- Guerrero NV, Singh RH, Manatunga A, Berry GT, Steiner RD & Elsas LJ 2000 Risk factors for premature ovarian failure in females with galactosemia. *Journal of Pediatrics* **137** 833–841. (<https://doi.org/10.1067/mpd.2000.109148>)
- Haus JM, Carrithers JA, Trappe SW & Trappe TA 2007 Collagen, cross-linking, and advanced glycation end products in aging human skeletal muscle. *Journal of Applied Physiology* **103** 2068–2076. (<https://doi.org/10.1152/jappphysiol.00670.2007>)
- He L, Ling L, Wei T, Wang Y & Xiong Z 2017 Ginsenoside Rg1 improves fertility and reduces ovarian pathological damages in premature ovarian failure model of mice. *Experimental Biology and Medicine* **242** 683–691. (<https://doi.org/10.1177/1535370217693323>)
- Hirshfield AN & Midgley AR Jr 1978 Morphometric analysis of follicular development in the rat. *Biology of Reproduction* **19** 597–605. (<https://doi.org/10.1095/biolreprod19.3.597>)
- Ho SC, Liu JH & Wu RY 2003 Establishment of the mimetic aging effect in mice caused by D-galactose. *Biogerontology* **4** 15–18. (<https://doi.org/10.1023/A:1022417102206>)
- Hyogo H & Yamagishi S 2008 Advance glycation end products (AGEs) and their involvement in liver disease. *Current Pharmaceutical Design* **14** 969–972. (<https://doi.org/10.2174/138161208784139701>)
- Kaufman FR, Xu YK, Ng WG & Donnell GN 1988 Correlation of ovarian function with galactose-1-phosphate uridylyl transferase levels in galactosemia. *Journal of Pediatrics* **112** 754–756. ([https://doi.org/10.1016/S0022-3476\(88\)80697-8](https://doi.org/10.1016/S0022-3476(88)80697-8))
- Kevenaar ME, Meerasahib MF, Kramer P, van de Lang-Born BM, de Jong FH, Groome NP, Themmen AP & Visser JA 2006 Serum anti-müllerian hormone levels reflect the size of the primordial follicle pool in mice. *Endocrinology* **147** 3228–3234. (<https://doi.org/10.1210/en.2005-1588>)
- Kim H, You S, Farris J, Kong BW, Christman SA, Foster LK & Foster DN 2002 Expression profiles of p53-, p16(INK4a)-, and telomere-regulating genes in replicative senescent primary human, mouse, and chicken fibroblast cells. *Experimental Cell Research* **272** 199–208. (<https://doi.org/10.1006/excr.2001.5420>)
- Kimura T, Takamatsu J, Ikeda K, Kondo A, Miyakawa T & Horiuchi S 1996 Accumulation of advanced glycation end products of the Maillard reaction with age in human hippocampal neurons. *Neuroscience Letters* **208** 53–56. ([https://doi.org/10.1016/0304-3940\(96\)12537-4](https://doi.org/10.1016/0304-3940(96)12537-4))
- Kumar PA, Kumar MS & Reddy GB 2007 Effect of glycation on alpha-crystallin structure and chaperone-like function. *Biochemical Journal* **408** 251–258. (<https://doi.org/10.1042/BJ20070989>)
- Lin N, Zhang H & Su Q 2012 Advanced glycation end-products induce injury to pancreatic beta cells through oxidative stress. *Diabetes and Metabolism* **38** 250–257. (<https://doi.org/10.1016/j.diabet.2012.01.003>)
- Mao GX, Deng HB, Yuan LG, Li DD, Li YY & Wang Z 2010 Protective role of solidroside against aging in a mouse model induced by D-galactose. *Biomedical and Environmental Sciences* **23** 161–166. ([https://doi.org/10.1016/S0895-3988\(10\)60047-5](https://doi.org/10.1016/S0895-3988(10)60047-5))
- Miyamoto K, Sato EF, Kasahara E, Jikumaru M, Hiramoto K, Tabata H, Katsuragi M, Odo S, Utsumi K, Inoue M, *et al.* 2010 Effect of oxidative stress during repeated ovulation on the structure and functions of the ovary, oocytes, and their mitochondria. *Free Radical Biology and Medicine* **49** 674–681. (<https://doi.org/10.1016/j.freeradbiomed.2010.05.025>)
- Mun SH, Joung DK, Kim YS, Kang OH, Kim SB, Seo YS, Kim YC, Lee DS, Shin DW, Kweon KT, *et al.* 2013 Synergistic antibacterial effect of curcumin against methicillin-resistant *Staphylococcus aureus*. *Phytomedicine* **20** 714–718. (<https://doi.org/10.1016/j.phymed.2013.02.006>)
- Nerlich AG & Schleicher ED 1999 N(epsilon)-(carboxymethyl)lysine in atherosclerotic vascular lesions as a marker for local oxidative stress. *Atherosclerosis* **144** 41–47. ([https://doi.org/10.1016/S0021-9150\(99\)00038-6](https://doi.org/10.1016/S0021-9150(99)00038-6))
- Odetti P, Rossi S, Monacelli F, Poggi A, Ciriigliaro M, Federici M & Federici A 2005 Advanced glycation end products and bone loss during aging. *Annals of the New York Academy of Sciences* **1043** 710–717. (<https://doi.org/10.1196/annals.1333.082>)
- Paine A, Eiz-Vesper B, Blasczyk R & Immenschuh S 2010 Signaling to heme oxygenase-1 and its anti-inflammatory therapeutic potential. *Biochemical Pharmacology* **80** 1895–1903. (<https://doi.org/10.1016/j.bcp.2010.07.014>)
- Park JH & Choi TS 2012 Polycystic ovary syndrome (PCOS)-like phenotypes in the D-galactose-induced aging mouse model. *Biochemical and Biophysical Research Communications* **427** 701–704. (<https://doi.org/10.1016/j.bbrc.2012.09.099>)
- Paulose T, Hannon PR, Peretz J, Craig ZR & Flaws JA 2012 Estrogen receptor alpha overexpressing mouse antral follicles are sensitive to atresia induced by methoxychlor and its metabolites. *Reproductive Toxicology* **33** 353–360. (<https://doi.org/10.1016/j.reprotox.2012.01.007>)
- Qin X, Cao M, Lai F, Yang F, Ge W, Zhang X, Cheng S, Sun X, Qin G, Shen W, *et al.* 2015 Oxidative stress induced by zearalenone in porcine granulosa cells and its rescue by curcumin *in vitro*. *PLoS ONE* **10** e0127551. (<https://doi.org/10.1371/journal.pone.0127551>)
- Rong Y & Distelhorst CW 2008 Bcl-2 protein family members: versatile regulators of calcium signaling in cell survival and apoptosis. *Annual Review of Physiology* **70** 73–91. (<https://doi.org/10.1146/annurev.physiol.70.021507.105852>)
- Rubio-Gozalbo ME, Gubbels CS, Bakker JA, Menheere PP, Wodzig WK & Land JA 2010 Gonadal function in male and female patients with classic galactosemia. *Human Reproduction Update* **16** 177–188. (<https://doi.org/10.1093/humupd/dmp038>)

- Sanders RD, Spencer JB, Epstein MP, Pollak SV, Vardhana PA, Lustbader JW & Fridovich-Keil JL 2009 Biomarkers of ovarian function in girls and women with classic galactosemia. *Fertility and Sterility* **92** 344–351. (<https://doi.org/10.1016/j.fertnstert.2008.04.060>)
- Scandalios JG 2005 Oxidative stress: molecular perception and transduction of signals triggering antioxidant gene defenses. *Brazilian Journal of Medical and Biological Research* **38** 995–1014. (<https://doi.org/10.1590/S0100-879X2005000700003>)
- Schinzl R, Münch G, Heidland A & Sebekova K 2001 Advanced glycation end products in end-stage renal disease and their removal. *Nephron* **87** 295–303. (<https://doi.org/10.1159/000045934>)
- Seifer DB, Baker VL & Leader B 2011 Age-specific serum anti-Müllerian hormone values for 17, 120 women presenting to fertility centers within the United States. *Fertility and Sterility* **95** 747–750. (<https://doi.org/10.1016/j.fertnstert.2010.10.011>)
- Semba RD, Nicklett EJ & Ferrucci L 2010 Does accumulation of advanced glycation end products contribute to the aging phenotype? *Journals of Gerontology: Series A, Biological Sciences and Medical Sciences* **65** 963–975. (<https://doi.org/10.1093/gerona/gdq074>)
- Seo JA, Kim B, Dhanasekaran DN, Tsang BK & Song YS 2016 Curcumin induces apoptosis by inhibiting sarco/endoplasmic reticulum Ca<sup>2+</sup>-ATPase activity in ovarian cancer cells. *Cancer Letters* **371** 30–37. (<https://doi.org/10.1016/j.canlet.2015.11.021>)
- Shehzad A, Wahid F & Lee YS 2010 Curcumin in cancer chemoprevention: molecular targets, pharmacokinetics, bioavailability, and clinical trials. *Archiv Der Pharmazie* **343** 489–499. (<https://doi.org/10.1002/ardp.200900319>)
- Sikora E, Bielak-Zmijewska A, Mosieniak G & Piwocka K 2010 The promise of slow down ageing may come from curcumin. *Current Pharmaceutical Design* **16** 884–892. (<https://doi.org/10.2174/138161210790883507>)
- Simm A, Casselmann C, Schubert A, Hofmann S, Reimann A & Silber RE 2004 Age associated changes of AGE-receptor expression: RAGE upregulation is associated with human heart dysfunction. *Experimental Gerontology* **39** 407–413. (<https://doi.org/10.1016/j.exger.2003.12.006>)
- Song X, Bao M, Li D & Li YM 1999 Advanced glycation in D-galactose induced mouse aging model. *Mechanisms of Ageing and Development* **108** 239–251. ([https://doi.org/10.1016/S0047-6374\(99\)00022-6](https://doi.org/10.1016/S0047-6374(99)00022-6))
- Soto N, Iñiguez G, López P, Larenas G, Mujica V, Rey RA & Codner E 2009 Anti-Müllerian hormone and inhibin B levels as markers of premature ovarian aging and transition to menopause in type 1 diabetes mellitus. *Human Reproduction* **24** 2838–2844. (<https://doi.org/10.1093/humrep/dep276>)
- Suckow BK & Suckow MA 2006 Lifespan extension by the antioxidant curcumin in *Drosophila melanogaster*. *International Journal of Biomedical Science* **2** 402–405.
- Tarín JJ 1995 Aetiology of age-associated aneuploidy: a mechanism based on the 'free radical theory of ageing'. *Human Reproduction* **10** 1563–1565. (<https://doi.org/10.1093/HUMREP/10.6.1563>)
- Tarín JJ 1996 Potential effects of age-associated oxidative stress on mammalian oocytes/embryos. *Molecular Human Reproduction* **2** 717–724. (<https://doi.org/10.1093/molehr/2.10.717>)
- Terlikowska KM, Witkowska AM, Zujko ME, Dobrzycka B & Terlikowski SJ 2014 Potential application of curcumin and its analogues in the treatment strategy of patients with primary epithelial ovarian cancer. *International Journal of Molecular Sciences* **15** 21703–21722. (<https://doi.org/10.3390/ijms151221703>)
- Tian J, Ishibashi K, Ishibashi K, Reiser K, Grebe R, Biswal S, Gehlbach P & Handa JT 2005 Advanced glycation endproduct-induced aging of the retinal pigment epithelium and choroid: a comprehensive transcriptional response. *PNAS* **102** 11846–11851. (<https://doi.org/10.1073/pnas.0504759102>)
- Tilly JL 2003 Ovarian follicle counts – not as simple as 1, 2, 3. *Reproductive Biology and Endocrinology* **1** 11. (<https://doi.org/10.1186/1477-7827-1-11>)
- Tiwari-Pandey R & Ram Sairam M 2009 Modulation of ovarian structure and abdominal obesity in curcumin- and flutamide-treated aging FSH-R haploinsufficient mice. *Reproductive Sciences* **16** 539–550. (<https://doi.org/10.1177/1933719109332822>)
- Tolikas A, Tsakos E, Gerou S, Prapas Y & Loufopoulos A 2011 Anti-Müllerian hormone (AMH) levels in serum and follicular fluid as predictors of ovarian response in stimulated (IVF and ICSI) cycles. *Human Fertility* **14** 246–253. (<https://doi.org/10.3109/14647273.2011.608464>)
- Ueki M, Ueno M, Morishita J & Maekawa N 2013 Curcumin ameliorates cisplatin-induced nephrotoxicity by inhibiting renal inflammation in mice. *Journal of Bioscience and Bioengineering* **115** 547–551. (<https://doi.org/10.1016/j.jbiosc.2012.11.007>)
- Vallianou NG, Evangelopoulos A, Schizas N & Kazakis C 2015 Potential anticancer properties and mechanisms of action of curcumin. *Anticancer Research* **35** 645–651.
- Visser JA, de Jong FH, Laven JS & Themmen AP 2006 Anti-Müllerian hormone: a new marker for ovarian function. *Reproduction* **131** 1–9. (<https://doi.org/10.1530/rep.1.00529>)
- Vlassara H, Bucala R & Striker L 1994 Pathogenic effects of advanced glycosylation: biochemical, biologic, and clinical implications for diabetes and aging. *Laboratory Investigation* **70** 138–151.
- Voznesens'ka T, Bryzhina TM, Sukhina VS, Makohon NV & Aleksieieva IM 2010 Effect of NF-kappaB activation inhibitor curcumin on the oogenesis and follicular cell death in immune ovarian failure in mice. *Fiziolohichnyi Zhurnal* **56** 96–101.
- Waggoner DD, Buist NR & Donnell GN 1990 Long-term prognosis in galactosaemia: results of a survey of 350 cases. *Journal of Inherited Metabolic Disease* **13** 802–818. (<https://doi.org/10.1007/BF01800204>)
- Wang XN, Roy SK & Greenwald GS 1991 *In vitro* DNA synthesis by isolated preantral to preovulatory follicles from the cyclic mouse. *Biology of Reproduction* **44** 857–863. (<https://doi.org/10.1095/biolreprod44.5.857>)
- Wang ME, Chen YC, Chen IS, Hsieh SC, Chen SS & Chiu CH 2012 Curcumin protects against thioacetamide-induced hepatic fibrosis by attenuating the inflammatory response and inducing apoptosis of damaged hepatocytes. *Journal of Nutritional Biochemistry* **23** 1352–1366. (<https://doi.org/10.1016/j.jnutbio.2011.08.004>)
- Watson JL, Greenshields A, Hill R, Hilchie A, Lee PW, Giacomantonio CA & Hoskin DW 2010 Curcumin-induced apoptosis in ovarian carcinoma cells is p53-independent and involves p38 mitogen-activated protein kinase activation and downregulation of Bcl-2 and survivin expression and Akt signaling. *Molecular Carcinogenesis* **49** 13–24. (<https://doi.org/10.1002/mc.20571>)
- Zhang DW, Fu M, Gao SH & Liu JL 2013 Curcumin and diabetes: a systematic review. *Evidence-based Complementary and Alternative Medicine* **2013** 636053. (<https://doi.org/10.1155/2013/636053>)
- Zhang JQ, Shen M, Zhu CC, Yu FX, Liu ZQ, Ally N, Sun SC, Li K & Liu HL 2014 3-Nitropropionic acid induces ovarian oxidative stress and impairs follicle in mouse. *PLoS ONE* **9** e86589. (<https://doi.org/10.1371/journal.pone.0086589>)
- Zingg JM, Hasan ST & Meydani M 2013 Molecular mechanisms of hypolipidemic effects of curcumin. *BioFactors* **39** 101–121. (<https://doi.org/10.1002/biof.1072>)

Received in final form 2 February 2018

Accepted 7 February 2018

Accepted Preprint published online 7 February 2018