

Original Article



Factors influencing decision-making around opportunistic salpingectomy: a nationwide survey

Miranda P. Steenbeek ¹, Laura A.M. van Lieshout ^{1,2}, Johanna W.M. Aarts ¹,
Jurgen M.J. Piek ², Sjors F.P.J. Coppus ^{1,3}, Leon F.A.G. Massuger ¹,
Rosella P.M.G. Hermens ⁴, Joanne A. de Hullu ¹

¹Radboud Institute for Health Sciences, Department of Obstetrics and Gynecology, Radboud University Medical Center, Nijmegen, The Netherlands

²Department of Obstetrics and Gynecology, Catharina Hospital, Eindhoven, The Netherlands

³Máxima Medical Centre, Veldhoven, The Netherlands

⁴Radboud Institute for Health Sciences, Scientific Institute for Quality of Healthcare, Radboud University Medical Center, Nijmegen, The Netherlands

OPEN ACCESS

Received: Jul 24, 2018

Revised: Aug 24, 2018

Accepted: Aug 28, 2018

Correspondence to

Miranda P. Steenbeek

Radboud Institute for Health Sciences,
Department of Obstetrics and Gynecology,
Radboud University Medical Center, P.O. box
9101, 6500HB, Nijmegen, The Netherlands.
E-mail: Miranda.Steenbeek@radboudumc.nl

Copyright © 2019. Asian Society of
Gynecologic Oncology, Korean Society of
Gynecologic Oncology

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Miranda P. Steenbeek
<https://orcid.org/0000-0001-6097-1579>
Laura A.M. van Lieshout
<https://orcid.org/0000-0003-0613-5267>
Johanna W.M. Aarts
<https://orcid.org/0000-0002-0674-9405>
Jurgen M.J. Piek
<https://orcid.org/0000-0002-0487-0631>
Sjors F.P.J. Coppus
<https://orcid.org/0000-0001-6786-5824>
Leon F.A.G. Massuger
<https://orcid.org/0000-0002-1539-8985>

ABSTRACT

Objective: To explore current practice and influencing factors on adoption of the opportunistic salpingectomy (OS), particularly regarding the decision making, to eventually enhance the development and implementation of clear guidelines.

Methods: This nationwide cross-sectional survey study was conducted in all hospitals in the Netherlands. An anonymous online survey was sent to gynecologists with special interest in gynecological oncology, gynecological endoscopy or urogynecology and all Dutch gynecology trainees. The survey mainly focused on current practice regarding OS and identification of influencing factors on the level of innovation, organization, healthcare professional and individual patient.



Results: The response rate was 348 out of 597 gynecologists (58.3%) and 142 out of 340 trainees (41.8%). Current practice of discussing and performing the OS varied widely, with ovarian cancer (OC) risk reduction as most important supportive factor on innovation level. Supportive factors on the level of organization and healthcare provider were; working in a non-training hospital, knowledge of current literature and extensive work experience (in years and annual number of hysterectomies). On individual patient level, a vaginal approach of hysterectomy, negative family history for OC and the presence of firm adhesions were suppressive factors for the OS.

Conclusion: In this study we evaluated the current practice regarding the opportunistic salpingectomy in the Netherlands and identified influencing factors on different levels to raise awareness and attribute to development of a targeted implementation strategy, on both national and international level.

Keywords: Salpingectomy; Ovarian Neoplasms; Primary Prevention; Risk Reduction

INTRODUCTION

Ovarian cancer (OC) is often detected at an advanced stage of disease, limiting 5-year survival to 30%–40%. The lifetime risk to develop OC varies from 1.3% in the general population up to 17%–44% in *BRCA1/2* germline mutation carriers [1]. High grade serous ovarian cancer (HGSC) accounts for 70% of all epithelial OCs [2]. As curative therapies are limited and

Rosella P.M.G. Hermens 
<https://orcid.org/0000-0001-7624-7120>
 Joanne A. de Hullu 
<https://orcid.org/0000-0001-5578-092X>

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Author Contributions

Conceptualization: S.M., H.R.P.M.G., D.H.J.A.;
 Formal analysis: S.M., V.L.L.A.M., H.R.P.M.G.,
 D.H.J.A.; Data curation: S.M., V.L.L.A.M.,
 A.J.W.M., P.J.M.J., C.S.F.P.J., M.L.F.A.G.,
 H.R.P.M.G., D.H.J.A.; Methodology: S.M.,
 H.R.P.M.G., C.S.F.P.J., D.H.J.A.; Software:
 S.M., H.R.P.M.G.; Supervision: A.J.W.M.,
 P.J.M.J., C.S.F.P.J., M.L.F.A.G., H.R.P.M.G.,
 D.H.J.A.; Validation: S.M., H.R.P.M.G., D.H.J.A.;
 Visualization: S.M., V.L.L.A.M.; Writing
 - original draft: S.M., V.L.L.A.M.; Writing -
 review & editing: A.J.W.M., P.J.M.J., C.S.F.P.J.,
 M.L.F.A.G., H.R.P.M.G., D.H.J.A.

effective screening methods are lacking, emphasis should be directed towards preventive measures. Therefore, *BRCA 1/2* germline mutation carriers are currently recommended to remove both ovaries and fallopian tubes (salpingo-oophorectomy) around the age of 40, resulting in premature menopause.

In the wake of the discovery of the Serous Tubal Intraepithelial Carcinoma (STIC) in 2001, the fallopian tube has been shown to play an important role in HGSC pathogenesis [3]. This insight has provided new opportunities for OC risk reduction by performing a salpingectomy. In *BRCA 1/2* germline mutation carriers this finding led to the initiation of several studies (NCT02321228) [4]. These studies compare early salpingectomy (after completion of childbearing) and delayed oophorectomy with the standard salpingo-oophorectomy to delay menopause and increase quality of life.

For the general population, opportunistic salpingectomy (OS) is defined as an additional salpingectomy during (benign) pelvic surgery and is currently a point of discussion among gynecologists [5]. According to a recent meta-analysis, opportunistic removal of the Fallopian tubes might decrease the overall incidence of OC up to 50% [6]. Some researchers suggest a risk reduction up to 70%, based on the assumption that OS will prevent all OCs of the serous subtype (HGSC) [5]. However, an OS has potential downsides; the Fallopian tube and ovary partially share their blood supply and thus, salpingectomy could diminish ovarian reserve. Several studies investigated this issue, with no difference in ovarian reserve after salpingectomy [7-9]. Furthermore, there is no evidence for an increased complication rate after OS [6,8] and literature on surgical time suggests a minor increase of 0–13 minutes, depending on surgical approach [8,10-12].

Currently there is no consensus regarding OS in the Netherlands, resulting in a large variety of practice, probably due to a lack of clear guidelines. According to previous research in other countries, 43%–77% of gynecologists discuss the possibility of an OS [13-15]. It is likely that the adoption rate is influenced by several factors, which can be found on different levels, such as the innovation itself (OS), the organization of care, healthcare professionals attitudes towards OS and characteristics of the individual patient [16].

In this study we aim to explore current practice and the influencing factors on OS adoption, particularly regarding the decision making, to eventually enhance the development of clear guidelines and subsequent implementation.

MATERIALS AND METHODS

1. Study design and population

This observational cross-sectional study among gynecologists and obstetrics and gynecology trainees was conducted between October and December 2017. The nationwide study sample consisted of 605 gynecologists (members of the Dutch Society of Obstetrics and Gynecology [NVOG]) with special interest in gynecological oncology, gynecological endoscopy or urogynecology and all 364 current Dutch gynecology trainees.

2. Setting

All gynecologists in the Netherlands are affiliated to the NVOG and benign pelvic surgery is solely performed in community hospitals with affiliations to all Dutch healthcare insurance

companies. The surgeries investigated in this survey are performed in all 92 Dutch hospitals and is fully covered by the health insurance after paying a flat rate each year when using healthcare in general. The costs of all surgeries exceed this flat rate, which means an additional salpingectomy does not influence the individual patient's costs.

3. Development survey

The survey was developed by a team of researchers and clinicians (S.M., H.R.P.M.G., C.S.F.P.J. and D.H.J.A.) based on current literature and subsequently pilot-tested by A.J.W.M., V.L.L.A.M. and H.D. The survey consisted of four subsections with 44 questions in total. It included questions about demographics, current practice of OS and subsections to identify influencing factors on different levels as formulated by Grol et al. [16]. The survey focused particularly on the level of the innovation (OS), organization, healthcare professional and individual patient [16];

- The questions about baseline demographics for gynecologists contained information about the years of work experience, type of institution, special interest, number of (vaginal) hysterectomies performed annually and the knowledge of current literature. For trainees, the questions were about the year of residency, special interest, number of (vaginal) hysterectomies performed and also the knowledge of current literature. Knowledge of current literature was based on self-assessment of the respondents with three possible answers; 1) knowledge of current literature on the suggested influence of OS for OC risk reduction (hypothesis), 2) knowledge of the hypothesis but not the literature itself, or 3) no knowledge of the hypothesis at all.
- The current practice domain (12 questions) covered both how frequent patients were informed on the possibility of OS and how frequent the procedure was carried out.
- Potential supportive and suppressive influencing factors were identified from literature and could be stratified according to perceived importance. Most of the included potential influencing factors were on the level of the innovation itself. Respondents were asked to allocate a total of 100 points to different influencing factors, allocating the highest amount of points to the most important factor. Influencing factors not covered by the questionnaire could be reported in a free text field.
- To identify and stratify influencing factors on patient level, case-vignettes were designed. The vignettes consisted of a fictive case of a 40-year old women intending to undergo a hysterectomy for menstrual complaints with varying patient-specific factors such as surgical modality (abdominal, laparoscopic or vaginal approach), family history (no or one family member with OC) and presence of intra-abdominal adhesions (mild or firm). Prior to presentation of the case-vignette, respondents received a short overview of background information, consisting of the lifetime risk of OC in the general population (1.3%), in patients with one family member with OC (2.5%) and the estimated risk reduction attributed to OS (50%–70%). As this study focused on the general population, there was no vignette with more than one family member with OC. The section started out with a vignette of a patient in optimal circumstances to advise salpingectomy (abdominal approach, one family member with OC and no intra-abdominal adhesions). Further on, patient-specific factors got more complicated and the number of respondents willing and reluctant to advise OS due to a combination of factors were identified (15 questions).

4. Data collection

A nationwide email-survey was sent out with an online computer programme using anonymous tokens so no identifiable information was collected. Due to the nature of the survey and anonymous collection of data, institutional review board approval was not required for this study. All fields in the questionnaire were set to be obligatory to answer in order to minimize missing data. It took approximately 10 minutes to complete the survey. A maximum of two reminders were sent out to non-responding participants. Surveys were eligible for inclusion if at least one subsection after baseline characteristics was completed. In the baseline characteristics section, gynecologists were asked to estimate the number of hysterectomies performed annually. If this number was below ten, the questionnaire could not be completed and ended after baseline characteristics. The Checklist for Reporting Results of Internet E-Surveys (CHERRIES) was followed (**Appendix 1**) [17].

5. Data analysis

Data was analyzed in SPSS (released 2013, IBM SPSS Statistics, version 22.0; IBM Corp., Chicago, IL, USA). Baseline characteristics of respondents, results on the current practice domain and the influencing factors on innovation and individual level were descriptively analyzed (median with interquartile range [IQR] and percentages). Influencing factors on organization and professional level were analyzed by the use of multivariate logistic regression analysis. As dependent variable, the current practice was converted to a bivariate variable ('never,' 'rarely,' 'about half the time' versus 'usually,' 'always') per type of surgery. The following baseline characteristics were used as independent variables among gynecologists; years of work experience, type of institution, special interest, number of (vaginal) hysterectomies performed annually and knowledge of current literature. Among trainees the imputed independent variables were; year of residency, number of (vaginal) hysterectomies performed with or without supervision and the knowledge of current literature. At first, univariate logistic regression analysis was performed and independent variables with a p-value ≤ 0.20 were analyzed in multivariate logistic regression analysis. The multivariate analysis was performed using backward-stepwise method and a p-value ≤ 0.05 was considered statistically significant.

RESULTS

1. Study population

In total, we received 510 responses, 20 of which were excluded because respondents did not fill out one complete domain, apart from baseline characteristics. The response rate was 348 out of 597 gynecologists (58.3%) and 142 out of 340 trainees (41.8%). A total of 47 gynecologists reported to perform less than ten hysterectomies per year, meaning that these surveys were not eligible for analysis. Ultimately, 443 surveys were included in the analysis, consisting of 68% gynecologists and 32% trainees.

The median duration of work experience among gynecologists was 12 years (IQR 7–19) and was over 25 years for 33 out of 301 (11%) gynecologists. Approximately two-thirds of the gynecologists (67.1%) worked in a (academic) training-hospital. Most respondents had special interest in benign gynecology/endoscopy and gynecologic oncology (70.2%). Out of 142 trainees, 49 were in the final two years of their training and started a subject of special interest (e.g., urogynecology, gynecologic oncology or benign gynecology). The most

Table 1. Baseline characteristics of respondents

Baseline characteristics	Value
Gynecologists	301 (67.9)
Work experience in years	12 (7–19)
Type of institution	
(Academic) Training hospital	202 (67.1)
Non-academic non-training hospital	99 (32.9)
Special interest	
Urogynecology	102 (23.0)
Benign gynecology/endoscopy	191 (43.1)
Gynecologic oncology	120 (27.1)
Other	6 (2.0)
No. of hysterectomies per year	30 (20–40)
No. of vaginal hysterectomies per year	5 (0–12)
Knowledge of current literature	
Knows current literature	182 (60.5)
Knows hypothesis, not the literature	119 (39.5)
Never heard of hypothesis	0 (0.0)
Gynecology trainees	142 (32.1)
Year of residency	
1 or 2	39 (27.5)
3 or 4	54 (38.0)
5 or 6	49 (34.5)
Special interest (residents in 5th or 6th year only)	
Urogynecology	2 (4.1)
Benign gynecology/endoscopy	15 (30.6)
Gynecologic oncology	9 (18.4)
Other/unknown	23 (53.1)
No. of hysterectomies performed with or without supervision	25 (10–50)
No. of vaginal hysterectomies performed with or without supervision	8 (3–15)
Knowledge of current literature	
Knows current literature	57 (40.1)
Knows hypothesis, not the literature	85 (59.9)
Never heard of hypothesis	0 (0.0)

Values are presented as number (%) or median (interquartile range).

common subject of special interest was partially the same as among gynecologists (benign gynecology/endoscopy) (**Table 1**).

2. Current practice of the opportunistic salpingectomy

Out of 443 respondents, 71.3% said to always or in the majority of cases discuss OS when planning an abdominal hysterectomy; 58.7% said to always or in the majority of the cases perform an OS. In case of laparoscopic hysterectomy, the results were 71.3% and 56.2%, respectively. In the discussion of vaginal hysterectomy, the respondents indicated OS to be feasible in a median of five out of ten vaginal hysterectomies (IQR 2–7). Gynecologists who performed over 30 vaginal hysterectomies a year expected feasibility in eight out of ten vaginal hysterectomies (range 4–9). Furthermore, 6.7% reported to always or usually inform patients about OS prior to vaginal hysterectomy, 26.9% never discussed this possibility with patients. The actual performance rate was reported as always or usually by 6.3% of respondents. Approximately half of the respondents indicated to rarely or never discuss OS in case of a laparotomy (46.0%) or laparoscopy (48.5%), other than hysterectomy. In terms of permanent contraception, 47.2% said to never or rarely inform patients on the option of salpingectomy. Approximately one third of the respondents (33.0%) did perform salpingectomies as sterilization technique in at least half of the patients. The abovementioned results are summarized in **Fig. 1**.




Fig. 1. Current practice of gynecologists and trainees in The Netherlands.

3. Influencing factors related to the OS

Domain 1: influencing factors at innovation level

A total of 408 respondents completed this section (Fig. 2). Among the supportive factors, reduction of OC risk was the most important and heavy weighing (median of 80 points, IQR 60–90), followed by the explicit request from a patient (median of 10 points, IQR 0–20). Supportive factors mentioned in the free text field were the removal of sterilization devices (e.g., clips or rings) or a positive family history of OC.




Fig. 2. A forest plot of potential influencing factors and the distribution of allocated points.

The stratification of suppressive factors was more diverse, with an approximately equal rank for possible increased surgical risk (median of 20 points, IQR 0–50) and possible negative effect on ovarian reserve (median of 10 points, IQR 0–40). Other factors (suggested in free text fields) were complicated or difficult situations (for example due to adhesions), no consent from the patient and the lack of a Dutch guideline on OS.

Respondents were reluctant to recommend OS as definitive contraception technique because no options for re-fertilization are available (37.7%), insufficient proof of potential benefits (31.4%), possibly increased risk of complications (30.9%), possibly deleterious effects on ovarian reserve (25.7%) and prolonged duration of surgery (22.6%). Furthermore, 22 respondents commented that lack of consensus in the Netherlands formed the most important reason not to advise OS as method of sterilization.

Domain 2: influencing factors at organization and healthcare professional level

When counseling a patient for surgery, gynecologists with more knowledge of current literature were more likely to usually or always discuss OS, just as gynecologists working in a non-training hospital. Work experience, either in years or number of procedures a year, was also a factor in whether or not gynecologists discuss OS, especially in vaginal hysterectomies.

A higher performance rate was found among gynecologists from non-training hospitals. Another supportive factor was knowledge of current literature and not just the hypothesis. Furthermore, the amount of work experience in years was positively correlated with higher performance rate of OS. During laparotomy for benign gynecological disease or tubal sterilization, the subgroup of gynecologists with a special interest in oncology was more likely to perform OS compared to gynecologists with other interests.

For trainees, only the knowledge of current literature was a significant supportive factor for OS in laparotomy for benign gynecological disease. An overview of all statistically significant results is presented in **Table 2**.

Domain 3: influencing factors at individual patient level

The recommendation on whether or not to perform OS based on patient characteristics, is represented in **Fig. 3**. In the optimal situation (e.g., abdominal or laparoscopic approach,

Table 2. Multivariate analysis of influencing factors on organization and professional level

Variables	Never, rarely, half the time	Usually, always	Univariate analysis		Multivariate analysis		
			OR (95% CI)	p-value	OR (95% CI)	p-value	
Gynecologists (n=301)							
Discuss the OS in case of:							
Abdominal hysterectomy	14 52 33 33	83 144 83 144	Non-training hospital Training hospital No Yes	1 0.47 (0.24-0.89) 1 1.74 (1.00-3.02)	0.02 0.05	1 0.43 (0.23-0.83) 1 1.89 (1.08-3.33)	0.01 0.03
Laparoscopic hysterectomy	18 76	77 119	Non-training hospital Training hospital	1 0.37 (0.20-0.66)	0.00	1 0.37 (0.20-0.66)	0.00
Vaginal hysterectomy	90 96	19 46	No Yes	1 2.27 (1.24-4.17)	0.01	1 2.07 (1.10-3.88)	0.02
	10 (6-16) 8 (3-15)	17 (9-24) 7 (2-10)	Work experience (yr) Vaginal hysterectomies a year	1.07 (1.03-1.11) 0.97 (0.94-1.01)	0.00 0.12	1.08 (1.04-1.12) 0.96 (0.93-0.99)	0.00 0.04
Benign gynecological laparoscopy	30 (20-40)	30 (25-40)	No. of hysterectomies a year	1.01 (1.00-1.02)	0.15	1.01 (1.00-1.03)	0.04
Sterilization	94 98	21 60	No Yes	1 2.74 (1.55-4.86)	0.00	1 2.74 (1.54-4.86)	0.00
Perform the OS in case of:							
Abdominal hysterectomy	18 76	77 119	Non-training hospital Training hospital	1 0.37 (0.20-0.66)	0.00	1 0.37 (0.20-0.66)	0.00
Laparoscopic hysterectomy	15 66	72 112	Non-training hospital Training hospital	1 0.35 (0.19-0.67)	0.00	1 0.35 (0.19-0.67)	0.00
Vaginal hysterectomy	90 96	19 46	No Yes	1 4.86 (1.61-14.64)	0.01	1 4.30 (1.39-13.33)	0.01
	11 (6-18) 11 (6-17)	20 (13-27) 14.5 (8-24)	Work experience (yr) Work experience (yr)	1.12 (1.06-1.18) 1.12 (1.06-1.18)	0.00 0.00	1.11 (1.05-1.17) 1.05 (1.01-1.10)	0.00 0.01
Benign gynecological laparoscopy	129 71	45 33	No Yes	1 2.40 (1.04-5.56)	0.00	1 2.90 (1.52-5.54)	0.00
	11 (6-18)	14 (9-24)	Work experience (yr)	1.05 (1.01-1.09)	0.01	1.05 (1.01-1.09)	0.02
Sterilization	102 132	10 34	No Yes	1 2.55 (1.24-5.25)	0.01	1 2.48 (1.16-5.28)	0.02
	79 160	16 18	Non-training hospital Training hospital	1 0.56 (0.27-1.15)	0.11	1 0.41 (0.19-0.89)	0.02
	10 (6-18)	16 (8-25)	Work experience (yr)	1.07 (1.02-1.12)	0.00	1.08 (1.03-1.12)	0.00
	161 78	15 19	No Yes	1 2.62 (1.26-5.42)	0.01	1 2.87 (1.34-6.17)	0.01
Gynecology trainees (n=142)							
Discuss the OS in case of:							
Benign gynecological laparoscopy	59 33	6 12	No Yes	1 3.58 (1.23-10.41)	0.02	1 3.58 (1.23-10.41)	0.02

Values are presented as number or median (interquartile range).
CI, confidence interval; OR odds ratio; OS, opportunistic salpingectomy.

Opportunistic salpingectomy: a nationwide survey




Fig. 3. Influencing factors on individual patient level.

FH+, one first-degree family member with ovarian cancer; FH-, no first-degree family member with ovarian cancer.

*Salpingectomy advised: no; †Salpingectomy advised: yes; ‡Advise to perform laparoscopic hysterectomy instead of the vaginal approach; §Mild adhesions are defined as easily to be removed, with minimal risk of damaging adjacent organs; ¶Firm adhesions are defined as difficult to remove, with risk of damaging adjacent organs.

one family member with OC and no intra-abdominal adhesions) 94% of the respondents would advise OS. For patients with a negative family history of OC, approximately 73% of the respondents would advise OS. Mild adhesions lead to a decline of approximately 3% while firm adhesions were a suppressive factor for an extra 50%–60% of the respondents, resulting in approximately 11% of respondents advising OS when combined with a negative family history. In case of one family member with OC, respondents were more willing to perform OS, with an additional 25.6%–31.6% of the respondents advising OS.

In case of vaginal hysterectomy, 56.4% of the respondents reported to recommend OS to patients with a positive family history. Of these respondents, 54.6% advised a strategic conversion to laparoscopy to perform OS. In case of a negative family history, OS was recommended by 22.6% of the respondents and 3.8% advised the strategic conversion. The most important reason to strategically convert a vaginal hysterectomy to laparoscopy was to maximize the chances of a successful OS (76.6%). Other reasons were a possibly increased complication rate in case of a vaginal approach (50.8%) and technical difficulties (41.1%). An important consideration to persevere in the vaginal route was the preference for the vaginal approach according to the Dutch guideline (76.2%).

Respondents were also asked about their recommendations on OS as sterilization technique. A total of 84 (20.7%) respondents would advise OS to all patients, 101 (24.9%) respondents would recommend OS to patients with one family member with OC and a total of 221 (54.4%) would not recommend OS at all, but would advise other techniques instead (i.e., clips, rings).

DISCUSSION

In this nationwide study performed in the Netherlands, we assessed both the current practice and influencing factors on the level of innovation, organization, healthcare professional and individual patient characteristics. The diverse results in the current practice domain represent the lack of consensus in the Netherlands regarding OS. On the level of innovation, OC risk-reduction was considered to be the most influential factor. On organizational level, OS appeared to be more routinely implemented in non-training hospitals compared to training hospitals. On healthcare professional level, implementation was mainly dependent on the knowledge of current literature, years of work experience and annual number of (vaginal) hysterectomies performed. Among trainees, solely knowledge of current literature was associated with higher performance rate when discussing OS preceding laparotomy for benign gynecological disease. Strong influencing factors on individual patient level were vaginal approach of hysterectomy, negative family history of OC and presence of firm adhesions during surgery.

On innovation level, OC risk reduction was by far the most influential factor in favor of OS. Intervention-related factors not to perform OS were possible effect on surgical risk and ovarian reserve. In addition, the absence of a Dutch guideline was perceived as a suppressive factor for OS. Previous research however, has not found evidence for an increased complication risk, estimated blood loss or duration of hospital stay [8,10,18]. Furthermore, studies on ovarian reserve after OS found no difference in ovarian function after hysterectomy with or without OS [8,10].

On organizational level, OS appeared to be better implemented in non-training hospitals when compared to training hospitals. This finding is in contrast with our expectations. The number of hysterectomies a gynecologist performs a year would have been a plausible explanation, yet in our cohort the number of hysterectomies per gynecologist was higher in training hospitals. Although unlikely, the lower implementation rate in training hospitals might be the result of reluctance to 'complicate' the procedure for trainees with an additional surgical step. Furthermore, several respondents commented on the absence of a univocal guideline in the Netherlands. Over the past few years several countries published statements and guidelines regarding OS. The general tendency of these papers is that in patients

at population risk of OC, OS should be considered and discussed prior to hysterectomy (American College of Obstetricians and Gynecologists [ACOG], Arbeitsgemeinschaft Gynäkologische Onkologie [AGO], The Society of Gynecologic Oncology of Canada [GOC], Royal Australian and New Zealand College of Obstetricians and Gynaecologists [RANZCOG]) and other pelvic surgery (Royal College of Obstetricians and Gynecologists [RCOG], Society of Gynecologic Oncology [SGO]) [19-22]. Yet, the implementation of new innovations goes beyond guideline development and is in need of a specific implementation strategy to achieve the aimed change [16]. In Canada, efforts to increase the implementation of OS have increased the number of hysterectomies with OS from 5% to 35% in a few years' time [23].

On healthcare professional level, OS was more likely to be adopted by experienced gynecologists (either in years of work experience or amount of hysterectomies performed annually) and in case of knowledge of current literature. These insights create opportunities to improve OS adoption as increased surgical risk and possible decrease of ovarian reserve were most frequently mentioned as suppressive factor. Thus, improvement in knowledge of current literature might increase implementation rate. Despite our findings, Jones et al. [15] investigated the implementation of OS and found no association between practice patterns and physicians characteristics. This discrepancy could be explained by differences in the subset of analyzed characteristics or study population. Aside from our findings regarding professional experience, gynecological surgeries performed by high-volume surgeons are associated with lower conversion rates, decreased operative complication rates and shorter duration of hospital stay [24]. Moreover, for gynecological surgeons with advanced surgical skills, the addition of a surgical step is probably less challenging, with easier implementation of OS consequently.

On individual patient level, the most influencing factors were a vaginal approach of hysterectomy, negative family history of OC and presence of firm adhesions. The realization of OS is more challenging during vaginal hysterectomy when compared to a laparoscopic or abdominal approach. Therefore, a tailored approach to OS during this type of surgery is comprehensible. Previous research demonstrated OS to be feasible in 81%–88% of vaginal hysterectomies [12,25]. However, respondents in our cohort estimated OS to be feasible in approximately half of the cases. This difference could be explained by the type of gynecologists performing the hysterectomies in the feasibility studies. Furthermore, decreasing number of vaginal hysterectomies per gynecologist might lead to a loss of expertise and less gynecologists with sufficient experience to safely perform the intervention. In the study of Robert et al. [25] all hysterectomies were performed or supervised by two experienced vaginal surgeons and in the study of Antosh et al. [12] the type of gynecologists performing the surgeries were not specified. The more experienced gynecologists in the present study (performing over 30 vaginal hysterectomies annually) estimated the feasibility to be 80% [12,25].

Aside from feasibility of OS during vaginal hysterectomy, very few patient-specific factors have been investigated prior to this survey study on the overall implementation of OS. However, patient-specific factors are essential when aiming for a more widespread adoption of OS. A higher performance rate will result in OS implementation among an increased variety of circumstances, including difficult surgeries (in the case of firm adhesions for example). However, as OS is a preventive measure, it should not be carried out at all costs. Generally, the costs should always be weighed against the (potential) benefits; our respondents agree that firm adhesions appear to be a strong influencing factor.

There are several strengths and limitations in the present study. The nationwide character gives a representative view on different influencing factors. The influencing factors among OS have not been investigated before and therefore attribute to specific implementation strategies on both a national and international basis. Additionally, the use of a case-vignette provides results that give insight in daily practice and increases the accuracy of self-assessment of the respondents. On the contrary, the self-assessment of respondents is one of the possible limitations of the study as self-assessment is susceptible to recall bias. Moreover, less than 50% of the trainees (against almost 60% of the gynecologists) filled in the questionnaire and possibly our respondents are more prone to complete the questionnaire when interested in the OS. Therefore, non-responders might have an even lower rate of OS in the current practice. This is a potential source of selection bias in the results and therefore the results might not be applicable to all gynecologists and trainees in general. However, the representation of gynecologists from the various types of organizations in our study was similar to the current situation in the Netherlands, with approximately one-third of the gynecologists working in a non-training hospital [26]. Although extrapolation of results to gynecologists and trainees in other countries is complicated.

In this study we objectified the current care regarding OS in the Netherlands and identified influencing factors on different levels to raise awareness and attribute to development of a specific implementation strategy, on both national and international level. Future research should be focused on the presence of factors identified in this study in the actual clinical practice and the effect of specific implementation strategies.

ACKNOWLEDGEMENTS

We would like to thank Dana Huppelschoten (D.H.) for extensive pilot-testing of the questionnaire prior to this study and Steven Teerenstra for statistical support.

REFERENCES

1. Kuchenbaecker KB, Hopper JL, Barnes DR, Phillips KA, Mooij TM, Roos-Blom MJ, et al. Risks of breast, ovarian, and contralateral breast cancer for *BRCA1* and *BRCA2* mutation carriers. *JAMA* 2017;317:2402-16. [PUBMED](#) | [CROSSREF](#)
2. McCluggage WG. Morphological subtypes of ovarian carcinoma: a review with emphasis on new developments and pathogenesis. *Pathology* 2011;43:420-32. [PUBMED](#) | [CROSSREF](#)
3. Piek JM, van Diest PJ, Zweemer RP, Jansen JW, Poort-Keesom RJ, Menko FH, et al. Dysplastic changes in prophylactically removed fallopian tubes of women predisposed to developing ovarian cancer. *J Pathol* 2001;195:451-6. [PUBMED](#) | [CROSSREF](#)
4. Harmsen MG, Arts-de Jong M, Hoogerbrugge N, Maas AH, Prins JB, Bulten J, et al. Early salpingectomy (Tubectomy) with delayed oophorectomy to improve quality of life as alternative for risk-reducing salpingo-oophorectomy in *BRCA1/2* mutation carriers (TUBA study): a prospective non-randomised multicentre study. *BMC Cancer* 2015;15:593. [PUBMED](#) | [CROSSREF](#)
5. Dilley SE, Straughn JM, Leath CA 3rd. The evolution of and evidence for opportunistic salpingectomy. *Obstet Gynecol* 2017;130:814-24. [PUBMED](#) | [CROSSREF](#)
6. Yoon SH, Kim SN, Shim SH, Kang SB, Lee SJ. Bilateral salpingectomy can reduce the risk of ovarian cancer in the general population: a meta-analysis. *Eur J Cancer* 2016;55:38-46. [PUBMED](#) | [CROSSREF](#)

7. Venturella R, Lico D, Borelli M, Imbrogno MG, Cevenini G, Zupi E, et al. 3 to 5 years later: long-term effects of prophylactic bilateral salpingectomy on ovarian function. *J Minim Invasive Gynecol* 2017;24:145-50.
[PUBMED](#) | [CROSSREF](#)
8. Van Lieshout LA, Pijlman B, Vos MC, de Groot MJ, Houterman S, Coppus SF, et al. Opportunistic salpingectomy in women undergoing hysterectomy: results from the HYSTUB randomised controlled trial. *Maturitas* 2018;107:1-6.
[PUBMED](#) | [CROSSREF](#)
9. Song T, Kim MK, Kim ML, Jung YW, Yun BS, Seong SJ, et al. Impact of opportunistic salpingectomy on anti-Müllerian hormone in patients undergoing laparoscopic hysterectomy: a multicentre randomised controlled trial. *BJOG* 2017;124:314-20.
[PUBMED](#) | [CROSSREF](#)
10. Song T, Lee SH, Kim WY, Heo EJ, Kim TJ. Opportunistic salpingectomy does not affect ovarian reserve or surgical outcomes in patients undergoing laparoscopic myomectomy. *Gynecol Obstet Invest* 2017;82:468-74.
[PUBMED](#) | [CROSSREF](#)
11. Ganer Herman H, Gluck O, Keidar R, Kerner R, Kovo M, Levran D, et al. Ovarian reserve following cesarean section with salpingectomy vs tubal ligation: a randomized trial. *Am J Obstet Gynecol* 2017;217:472.e1-6.
[PUBMED](#) | [CROSSREF](#)
12. Antosh DD, High R, Brown HW, Oliphant SS, Abed H, Philip N, et al. Feasibility of prophylactic salpingectomy during vaginal hysterectomy. *Am J Obstet Gynecol* 2017;217:605.e1-5.
[PUBMED](#) | [CROSSREF](#)
13. Gill SE, Mills BB. Physician opinions regarding elective bilateral salpingectomy with hysterectomy and for sterilization. *J Minim Invasive Gynecol* 2013;20:517-21.
[PUBMED](#) | [CROSSREF](#)
14. Chene G, de Rochambeau B, Le Bail-Carval K, Beaufile E, Chabert P, Mellier G, et al. Current surgical practice of prophylactic and opportunistic salpingectomy in France. *Gynecol Obstet Fertil* 2016;44:377-84.
[PUBMED](#) | [CROSSREF](#)
15. Jones NL, Schulkin J, Urban RR, Wright JD, Burke WM, Hou JY, et al. Physicians' perspectives and practice patterns toward opportunistic salpingectomy in high- and low-risk women. *Cancer Invest* 2017;35:51-61.
[PUBMED](#) | [CROSSREF](#)
16. Grol R, Wensing M, Eccles M. Improving patient care: the implementation of change in clinical practice. Edinburgh/New York: Elsevier Butterworth Heinemann; 2005.
17. Eysenbach G. Improving the quality of web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2004;6:e34.
[PUBMED](#) | [CROSSREF](#)
18. Ghezzi F, Cromi A, Siesto G, Bergamini V, Zefiro F, Bolis P. Infectious morbidity after total laparoscopic hysterectomy: does concomitant salpingectomy make a difference? *BJOG* 2009;116:589-93.
[PUBMED](#) | [CROSSREF](#)
19. Pölcher M, Hauptmann S, Fotopoulou C, Schmalfeldt B, Meinhold-Heerlein I, Mustea A, et al. Opportunistic salpingectomies for the prevention of a high-grade serous carcinoma: a statement by the Kommission Ovar of the AGO. *Arch Gynecol Obstet* 2015;292:231-4.
[PUBMED](#) | [CROSSREF](#)
20. Committee on Gynecologic Practice. Committee opinion No. 620: salpingectomy for ovarian cancer prevention. *Obstet Gynecol* 2015;125:279-81.
[PUBMED](#) | [CROSSREF](#)
21. Salvador S, Scott S, Francis JA, Agrawal A, Giede C. No. 344-opportunistic salpingectomy and other methods of risk reduction for ovarian/fallopian tube/peritoneal cancer in the general population. *J Obstet Gynaecol Can* 2017;39:480-93.
[PUBMED](#) | [CROSSREF](#)
22. Royal College of Obstetricians & Gynaecologists. The distal fallopian tube as the origin of non-uterine pelvic high-grade serous carcinoma. Scientific impact paper No. 44. London: Royal College of Obstetricians & Gynaecologists; 2014.
23. McAlpine JN, Hanley GE, Woo MM, Tone AA, Rozenberg N, Swenerton KD, et al. Opportunistic salpingectomy: uptake, risks, and complications of a regional initiative for ovarian cancer prevention. *Am J Obstet Gynecol* 2014;210:471.e1-11.
[PUBMED](#) | [CROSSREF](#)
24. Worley MJ Jr, Anwandter C, Sun CC, dos Reis R, Nick AM, Frumovitz M, et al. Impact of surgeon volume on patient safety in laparoscopic gynecologic surgery. *Gynecol Oncol* 2012;125:241-4.
[PUBMED](#) | [CROSSREF](#)

25. Robert M, Cenaiko D, Sepandj J, Iwanicki S. Success and complications of salpingectomy at the time of vaginal hysterectomy. *J Minim Invasive Gynecol* 2015;22:864-9.
[PUBMED](#) | [CROSSREF](#)
26. van der Velden LFJ, Hingstman L. *Monitor arbeidsmarkt gynaecologen 2006*. Utrecht: NIVEL; 2006.

Appendix 1. CHERRIES checklist

Design	Describe survey design	Nationwide cross-sectional survey among all gynaecologists with special interest in gynaecological oncology, gynaecological endoscopy or urogynaecology and all Dutch gynaecology trainees.
Institutional Review Board (IRB) approval and informed consent process	IRB approval	IRB approval was not applicable for this study.
	Informed consent	The participants were invited to complete the survey by email. The email contained information about the content of the survey, including aim, identity of the investigators, estimated time investment to complete the survey and anonymous data handling.
	Data protection	For data collection the web-based tool LimeSurvey was used. All data was automatically converted to anonymous tokens, therefore it was impossible to link e-mail addresses to the corresponding surveys. In this way total anonymity of participants was secured.
Development and pre-testing	Development and testing	The survey was developed by a team of researchers and clinicians based on current literature, subsequently it was extensively pilot-tested to assess the usability and technical functionality.
Recruitment process and description of the sample having access to the questionnaire	Open survey versus closed survey	Participants were only able to fill in the survey by the use of a personal ink sent by email, therefore it was a closed survey.
	Contact mode	The (potential) participants were contacted by e-mail. A maximum of two reminders were sent out to participants who had not responded.
	Advertising the survey	Not applicable, no advertising was used.
Survey administration	Web/E-mail	The survey was web-based (LimeSurvey) and data was automatically transferred to a database.
	Context	Not applicable.
	Mandatory/voluntary	It was a voluntary survey.
	Incentives	No incentives were offered.
	Time/date	Participants were invited to complete the survey between October and December 2017.
	Randomization of items or questionnaires	Not applicable.
	Adaptive questioning	The survey could only be completed when participants performed at least ten hysterectomies a year.
	Number of items	The survey contained a total of 44 questions.
	Number of screens (pages)	The questionnaire was distributed across six pages (screens) with a minimum of two and a maximum of 16 questions per page (screen).
	Completeness check	The questions in the survey were all mandatory to complete before entering the next page (screen) with questions (an automatic option in LimeSurvey).
Review step	Respondents were not able to use the 'back button' because information on the last page (screen) of the survey could influence the answers given on the earlier pages.	
Response rates	Unique site visitor	
	View rate (ratio unique of unique survey visitors/unique site visitors)	Not applicable.
	Participation rate (ratio unique survey page visitors/agreed to participate)	In total, 937 participants received the invitation E-mail and 510 replied/visited the first page of the survey (54%).
	Completion rate (ratio agreed to participate/finished survey)	The completion rate was 96%.
Preventing multiple entries from the same individual	Cookies used	Not applicable because of the unique tokens.
	IP check	Not applicable because a unique token was sent to each E-mail address which could only be used once.
	Log file analysis	Not applicable because a unique token was sent to each E-mail address which could only be used once.
Analysis	Registration	Not applicable because a unique token was sent to each E-mail address which could only be used once
	Handling of incomplete questionnaires	Surveys were eligible for inclusion if at least one subsection (besides the baseline characteristics) was completed.
	Questionnaires submitted with an atypical timestamp	Not applicable.
	Statistical correction	Not applicable.

REFERENCE

Eysenbach G. Improving the quality of web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2014;6:e34.