




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Designing a virtual mental health consultation program to support and strengthen childcare centers impacted by COVID-19: A randomized controlled trial protocol

Ruby Natale, Yaray Agosto, Rebecca J. Bulotsky-Shearer, Sara M. St. George, Jason Jent



PII: S1551-7144(22)00348-2
DOI: <https://doi.org/10.1016/j.cct.2022.107022>
Reference: CONCLI 107022
To appear in: *Contemporary Clinical Trials*
Received date: 29 July 2022
Revised date: 16 November 2022
Accepted date: 21 November 2022

Please cite this article as: R. Natale, Y. Agosto, R.J. Bulotsky-Shearer, et al., Designing a virtual mental health consultation program to support and strengthen childcare centers impacted by COVID-19: A randomized controlled trial protocol, *Contemporary Clinical Trials* (2022), <https://doi.org/10.1016/j.cct.2022.107022>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Designing a Virtual Mental Health Consultation Program to Support and Strengthen Childcare Centers Impacted by COVID-19: A Randomized Controlled Trial Protocol

Ruby Natale, PhD, PsyD^a, Yaray Agosto, MPH^a, Rebecca J. Bulotsky-Shearer, PhD^b, Sara M. St. George, PhD^c, Jason Jent, PhD^a

^a University of Miami Miller School of Medicine, Department of Pediatrics, 1601 NW 12th Ave, Miami, Florida 33136, USA

^b University of Miami, Department of Psychology, 5665 Ponce de Leon Blvd, Coral Gables, Florida 33146, USA

^c University of Miami Miller School of Medicine, Department of Public Health Sciences, 1120 NW 14th St, Miami, Florida 33136, USA

Email Addresses:

Ruby Natale: rnatale@med.miami.edu

Yaray Agosto: yagosto@med.miami.edu

Rebecca J. Bulotsky-Shearer: rshearer@med.miami.edu

Sara M. St. George: s.stgeorge@med.miami.edu

Jason Jent: jjent@med.miami.edu

Address correspondence to Ruby Natale, University of Miami Miller School of Medicine, Department of Pediatrics, 1601 NW 12th Ave. Miami, FL 33136. Phone: (954) 552-0446. Email: rnatale@med.miami.edu

Abstract

Background: The COVID-19 pandemic has had negative psychosocial impacts on young children; teachers in childcare centers continue to be overwhelmed by how to address the downstream psychological effects children are experiencing. This randomized controlled trial

will study the role of a community-based, childcare center-support system in improving resilience and mitigating the long-term impacts of COVID-19 on children's development.

Methods: This study will be modeled on a successful Early Childhood Mental Health Consultation (ECMHC) intervention which utilizes mental health consultants to deliver a Jump Start+: COVID 19 Support (JS+CS) virtual toolkit to childcare centers in Miami-Dade County via a Kubi robot. The toolkit comprises four strength-based strategies likely to be effective in improving resiliency following disasters: Safety Planning, Effective Communication, Adult Self-Care, and Trauma-Informed Behavior Support. Our first aim will utilize a cluster randomized trial to examine the effectiveness of JS+CS on improving the psychosocial functioning of young children, as compared to an obesity prevention intervention control group. Children will be followed at 6, 12, 18, and 24 months. The second aim will examine the mechanisms that contribute to effective uptake by teachers of the JS+CS support strategies on child outcomes. The third aim will explore implementation barriers/facilitators as well as potential societal contextual factors (e.g., vaccine uptake) to help centers serving disproportionately affected minority communities recover from and prepare for future crises.

Conclusion: This design will inform the refinement and scaling of JS+CS and generalize impacts to other childcare center interventions in the context of disasters.

ClinicalTrials.gov Identifier: NCT05445518

Keywords: early childhood, childcare center, COVID-19, virtual consultation, resiliency

1. Introduction

Childcare centers are essential businesses that have been negatively impacted by the COVID-19 pandemic and other public health crises, with downstream negative effects on teachers, children, and families; yet little is known about effective public health interventions that can address centers' pressing short- and long-term needs.

The pandemic has had devastating impacts on children's psychosocial health, ranging from long-term social isolation to heightened family stressors.¹ Children as young as two are experiencing anxiety, irritability, anger, and stress.^{2,3} The incidence and severity of behavior problems are increasing in children with and without a prior history of behavior problems,⁴⁻⁶ and once through the acute phase of the pandemic, "sleeper" effects on children's mental health are predicted.⁷ Early data found that children ages 2-5 years had elevated behavior problems, which were higher in parents reporting lower-income and more COVID-19 stress.^{8,9} School closures and cumulative exposure to adverse events can negatively impact social-emotional, behavioral, and cognitive development during the critical period (0-5 years) when the foundation of learning is built.¹⁰⁻¹⁴ Enrollment in high-quality childcare centers with nurturing caregivers can buffer these negative effects by building resiliency and by strengthening self-regulation and executive functioning skills.¹⁵⁻¹⁷ Unfortunately, our Childcare Worker Job Stress Inventory showed that early childhood teachers felt overwhelmed with additional burdens of maintaining a safe environment during the pandemic while delivering quality classroom practices.^{18,19} The result has been staff turnover and poorly functioning classrooms.^{20,21} Therefore, it is essential to intervene at multiple levels to increase support for teachers, build child resilience, and foster a center environment anchored in policies that promote positive growth during stressful times.

COVID-19 has also provided a strategic opportunity to deliver virtual intervention support to childcare centers. Safety standards at childcare centers have changed, including

limiting visitors, even parents or consultants. With decreased rates of infectious disease transmission, including flu, many centers are maintaining these visitation restrictions and prefer teleconsultations. In addition to reducing disease transmission, virtual supports such as mental health teleconsultations reduce travel time and driving fatigue for consultants. One novel way to increase service accessibility and feasibility is by using a Kubi, an affordable telepresence robot that holds a tablet that can connect to Zoom or other video conferencing software. The remote user connects to the tablet using the Kubi Connect App, which pairs the tablet to the telepresence robot via Bluetooth. Teleconsultation via the Kubi is superior to other methods because it allows for a broader rotating view to watch and listen as teachers practice strategies (teacher-child interactions), and it aids in observations by providing a more real-life, face-to-face experience of how the classroom and center functions. In order to reduce classroom distractions by the Kubi during remote teleconsultations, the video camera of the remote consultant is turned off. While preliminary evidence supports the use of Kubi telepresence robots in hard-to-access schools, literature assessing their use in childcare settings is limited.^{22,23}

In response to the urgent need to support childcare centers during the pandemic, our team was one of the first in the country to develop a mental health toolkit, Jump Start+: COVID Support (JS+CS), to prepare them to handle public health crises.¹⁸ The program uses the JS+CS toolkit so that mental health consultants (MHCs), who are considered “trusted messengers,” provide centers with psychosocial coping strategies to boost resilience and easy-to-access safety information to address infection-related concerns. The toolkit is organized around four program pillars: two are psychosocial (Self-Care, Trauma-Informed Behavior Support) and two are safety-related (Safety, Communication). Importantly, and given evolving safety standards that

limited consultant visitation, we quickly pivoted to providing JS+CS consultations virtually so that we could continue to support teachers during the pandemic.

While our preliminary data show that JS+CS helps reduce teacher stress and improve centers' capacity to implement emergency preparedness guidelines, the short- and longer-term impact of JS+CS on child, teacher, and center outcomes has not yet been rigorously tested.¹⁹ As such, there are three aims of this study. Aim 1 works to improve the psychosocial functioning of children during and after COVID-19. Specifically, we hypothesize that relative to the control group, children in the intervention group will exhibit improved psychosocial outcomes as demonstrated by reduced behavior problems and improved resiliency (executive functioning and prosocial behaviors). Aim 2 works to improve teacher psychosocial functioning, so that teachers can be more resilient and provide better quality classroom practices. We hypothesize that JS+CS will improve teacher psychosocial functioning as evidenced by changes in attitudes (self-efficacy), beliefs (stress), and classroom practices (teacher-child interactions). Aim 3 will help strengthen the capacity of childcare centers to improve physical safety and psychosocial support standards by understanding barriers and facilitators to effectively implementing JS+CS. We hypothesize that centers implementing JS+CS policies will improve the physical safety and psychosocial practices that lead to better quality childcare.

1.1 Theoretical Model

The Early Childhood Mental Health Consultation (ECMHC) theory of change serves as the conceptual framework (Figure 1). ECMHC, through mental health consultants, has proven effective in addressing young children's behavioral health by building the capacity of childcare teachers to develop the attitudes and skills necessary to effectively support the development of resilience in young children.²⁴⁻²⁷ The JS+CS Toolkit is organized into four pillars that work

together to build resilience by improving psychosocial coping through *Self-Care and Trauma-informed Behavior Supports* and disaster recovery practices by improving *Safety and Communication practices*. The pillars are based on Caring for Our Children–National Health and Safety Standards²⁸, supplemented with CDC COVID-19 guidelines for Childcare Centers²⁹ and other evidenced-based practices.^{30,31}




Figure 1. Program Model Overview

The JS+CS toolkit is delivered by Mental Health Consultants (MHCs). They provide toolkit implementation strategies at three levels to improve child psychosocial functioning (Aim 1), teacher psychosocial functioning and classroom practices (Aim 2), and the center’s capacity to maintain safety and psychosocial support (Aim 3). MHCs are master’s level clinicians trained in the ECMHC framework which uses the “consultative stance” to build relationships.^{32,33} The multimedia toolkit provides strategies on how to implement the program pillars to achieve outcomes.

Overall, we posit that child psychosocial functioning (prosocial and reduced internalizing/externalizing) and resiliency (children’s self-regulation) will improve by working with teachers

to improve their teaching practices (e.g., use of trauma-focused behavior management), teacher's increased self-efficacy when facing challenging behaviors (attitudes), and their beliefs (support from center directors).^{18,20,34,35} Table 1 provides examples of strategies implemented that are hypothesized to lead to change at 3 levels: center level, teacher level, and child level.

Considering response burden and power, specifically, we will evaluate family moderating factors including stress, low income, vaccine uptake/confidence, and parent safety practices outside of school^{36,37} based on the Halfon et al. risk model.³⁸

Pillar 1: Physical & Psychological Safety	Pillar 2: Communication	Pillar 3: Self-Care	Pillar 4: Trauma-informed Behavior Support
<p>Center Level Goal: Comply with CDC guidelines <i>Strategy: Refine policies in accordance with CDC's disaster recovery guidelines and share JS+CS COVID cartoon videos with parents and children</i> Measure: Health Environment Rating Scale – Program</p>	<p>Center Level Goal: Develop strong partnerships with teachers and parents Strategy: Director/teacher collaborative goal setting and support in staff with parent communication Measure: Key Informant Interviews</p>	<p>Center Level Goal: Promote staff resilience Strategy: Develop wellness policies, resiliency planning to identify self-care practices, staff social support, and scheduled stress check-ins with teachers Measure: Health Environment Rating Scale – Program and Key Informant interview</p>	<p>Center Level Goal: Promote center-wide policy on positive behaviors and transform negative behaviors Strategy: Ongoing staff continuing education on trauma-informed classroom management strategies Measure: Health Environment Rating Scale – Program</p>
<p>Teacher Level Goal: Be a healthy role model Strategy: Practice COVID-19 related safety guidelines and model self-regulation to promote safety and security within the classroom</p>	<p>Teacher Level Goal: Develop strong partnerships with families Strategy: Hold ongoing staff and parent meetings to review updated COVID safety procedures.</p>	<p>Teacher Level Goal: Plan ahead for stress management Strategy: Practice daily mindfulness and additional verbal and non-verbal strategies to model self-regulation techniques for children and self.</p>	<p>Teacher Level Goal: Promote positive classroom climate Strategy: Use 1,2,3 calming and positive reinforcement with specific praise for child prosocial behaviors and engagement</p>

<i>Measure: Health Environment Rating Scale – Classroom</i>	<i>Measure: Child Care Worker Job Stress Inventory</i>	<i>Measure: Child Care Worker Job Stress Inventory</i>	<i>Measure: Health Environment Rating Scale – Classroom</i>
<p>Child Level Goal: Ensure children are emotionally supported through grief and stress</p> <p>Strategy: Use social stories to teach children about overcoming challenges</p> <p>Measure: Devereux Early Childhood Assessment</p>	<p>Child Level Goal: Promote positive behaviors at home and school</p> <p>Strategy: Establish school-home communication notes and shared reward-based system to support positive child behaviors)</p> <p>Measure: Child Behavior Checklist</p>	<p>Child Level Goal: Promote children’s sense of self and attachment security</p> <p>Strategy: Use I Love You Rituals (Becky Bailey) for expressing anger and need for attention</p> <p>Measure: Devereux Early Childhood Assessment</p>	<p>Child Level Goal: Promote children’s self-control/regulation and initiative</p> <p>Strategy: Use trauma-informed practices such as “calm down box” with feather, scents etc. to teach children breathing and relaxation</p> <p>Measure: Strengths and Difficulties Questionnaire</p>

2. Methods

2.1 Participants and Study Sampling

Implementation is guided by Reach, Effectiveness, Adoption, Implementation, and Maintenance (RE-AIM) framework.³⁹ RE-AIM uses strategies to adopt and integrate evidence-based health interventions and change practice patterns within specific settings. The RE-AIM framework has been used in our pilot work and full details can be found elsewhere.⁴⁰ We will use a Type 1 implementation-effectiveness approach to identify implementation facilitators and barriers that will offer critical insights on the uptake of COVID-19 interventions in real time.⁴¹ This Type 1 approach is superior to other effectiveness studies because it will inform the refinement and scaling up of JS+CS so that results can be rapidly used to support other childcare centers in the context of disasters.⁴¹

Within the Type 1 hybrid implementation-effectiveness design⁴², we will utilize a cluster-randomized trial to assess effectiveness and collect data on observed barriers and facilitators to better understand the uptake of implementation for JS+CS. As the intervention is multi-level and center-wide, we will randomly assign centers (N=24) to one of two groups (intervention or

attention control). In cluster-randomized trials, clusters (i.e., childcare centers) rather than individuals are randomized to intervention or control groups and outcomes are measured on individuals within those clusters. The intervention group will receive the JS+CS program (n=12). The attention control group (n=12) is an active, time matched group that will receive the HC2 obesity prevention program, a program unrelated to JS+CS.^{43,44}

Randomization will be balanced via block randomization for ethnicity with half the centers in each group serving predominantly Hispanic families and the other half non-Hispanic Black families, allowing for ethnic group comparisons within balance in each group. In order to have adequate power for our multi-level nesting design⁴⁵ (described below), we will recruit 720 children/families (360 intervention, 360 control), 120 teachers/classrooms (60 intervention, 60 control), and 24 directors (12 each group). To obtain unbiased medium-level effect sizes, previous studies have recommended including above 20 clusters (i.e., centers) and above 5-6 individuals (i.e., children).^{46,47} Thus, we are being conservative in estimating 24 centers, with 120 classrooms and 6 children in each. All centers will be part of the Quality Improvement System, a governing body which enrolls childcare centers and monitors the quality of services provided to 417 centers in the county. Center inclusion criteria: (1) have ≥ 50 children (≥ 30 of whom are 18 month-3 years old); (2) be located in the low-income census tract, with at least 50% of families receiving childcare subsidy; (3) serve at least 60% Hispanic or 60% Non-Hispanic Black families; (4) have directors and teachers who agree to participate; and (5) be a Quality Improvement System center with a star rating of 3 indicating average quality practices (range 1-5) and not previously enrolled in Jump Start.

2.2 Intervention Group

There are 3 steps to ensure that the toolkit is successful, sustainable, and implemented with fidelity. **STEP 1 Toolkit Implementation.** Content: The toolkit includes: (1) infographics with goals and myth-busting facts that pertain to intervention pillars; (2) how-to cartoon videos demonstrating resiliency-based coping strategies for teachers and directors; (3) cartoon videos directed at preschool children to assist in adherence to infectious disease control guidelines that are relevant post-pandemic (e.g., hand washing). To ensure our toolkit is up to date, our Scientific Advisory Committee meets monthly to review the latest CDC guidelines and propose toolkit changes. Structure: The Toolkit Action Plan is a standardized implementation protocol used by MHCs in each teleconsultation to 1) set the goal based on the pillar for the week, 2) provide strategies for achieving the goals using infographics and “how-to” videos located on our website, 3) have teachers practice strategies while MHCs provide real-time feedback. Six weeks are devoted to each pillar before moving on to the next (in the order of Self-Care, Behavior Support, Safety, Communication). Dose: MHCs provide 24 weeks of teleconsultations for 4 hours weekly per center, divided between directors and teachers. Center-level teleconsultations occur with the directors for 1 hour per week to build center-wide capacity by focusing on how they can implement that week’s pillar policies into practice. Classroom-level teleconsultations occur in a small virtual group (2-3 teachers) for 1 hour per week, during “Break and Breathe”, to provide teachers with goals/strategies to adopt pillars into their daily lives. Examples of goals/strategies include helping teachers develop a self-care plan to build resiliency (beliefs) and use coping strategies (self-efficacy) and collaborating/engaging with parents by posting cartoon videos with parent tips through a multimodal platform such as WhatsApp or Facebook (practices). Child-level teleconsultations occur for 2 hours each week (20 minutes individually

for each teacher) to focus on implementing the goals/strategies with the children (teacher practices). See Table 1. Telepresence robot integration: Each school will receive a Kubi Plus with Enhanced Audio telepresence robot that the MHCs will use to deliver the intervention. The robots, which are operated remotely by MHCs, are stationary devices that can tilt and pan around a classroom while video conferencing with individuals in the classroom. It has microphones and large speakers for group classroom sound. MHCs can use the audio to provide live classroom practice strategies and teacher feedback using a single Bluetooth earbud to minimize child distractions. Before initiating services, MHCs will train teachers on how to a) structure the environment for telepresence robot classroom consultation or observation sessions, b) initiate and end sessions, and 3) troubleshoot common problems.

STEP 2 Sustainability and Cultural and Linguistic Relevance. From the first teleconsultation, a sustainability plan is developed as part of the Action Plan. This includes the identification of a “Program Champion,” a lead teacher (paid a \$500 stipend/per annum) who pledges to support the center in maintaining toolkit strategies. In years 2 and 3, the MHCs will provide quarterly 2-hour refresher trainings (booster sessions) with all teachers and the Program Champion to help maintain their skill set in supporting center-wide implementation of toolkit pillars and policy standards. We use the 8 dimensions of the Ecological Validity Model to ensure that the toolkit has the necessary cultural and linguistic adaptations to reduce health disparities.⁴⁸ The Ecological Validity Model is the most widely used approach to culturally adapting interventions.⁴⁹ Specific cultural targeting strategies include peripheral, evidential, linguistic, constituent involving, and socio-cultural strategies.⁵⁰ As an example, all materials will be translated into Spanish (linguistic strategies) and will include photos of ethnic minorities

(peripheral strategies) and information about national and local prevalence data related to teacher practices and children of ethnic minority groups (evidential strategies).

STEP 3 Monitoring Fidelity. Fidelity is based on Dane and Schneider's 5 components: Adherence, Exposure, Participant Responsiveness, Quality of Delivery, and Program Differentiation (described in Measures).⁵¹

2.3 Active Time-Matched Control Group

Centers randomized to the attention control will receive an existing obesity prevention program, Healthy Caregivers-Healthy Children (HC2) by the MCHC. This program was fully developed and implemented through federal funding by the PI and was empirically found to improve obesity prevention practices within centers that were ethnically similar to this study.^{43,44,52} Control centers receive all the same pre- and post- measures and incentives as the intervention arm to ensure retention. The same 3-step process will be followed for the control group and intervention group.

2.4 Procedures

The study has been approved by University's Institutional Review Board and has been registered with ClinicalTrials.gov number NCT05445518. We will use CONSORT guidelines' extension for social and psychological intervention trials to report RCT enrollment and retention data as well as to generate a participant flow diagram. We are recruiting, within the existing childcare network, 24 childcare centers that serve low-income families from ethnic minority backgrounds. Interested center directors will complete a readiness screener that assesses eligibility. Eligible center directors who give informed consent will then complete the baseline survey. Graduate-level research assistants (RAs) will be from commensurate ethnic minority backgrounds and have experience working in their assigned area of the county (north and south).

Centers will be recruited and consented first, followed by teachers, and then families. A virtual kick off with a “Meet and Greet the Robot” will occur weekly for the first 2 weeks for teachers and then 2 weeks later for parents, during which the study will be explained by our program’s MHC. Participants giving verbal agreement are contacted by the RA based on their preference (email or phone) to obtain consent and measures. Teacher self-reports, teacher ratings of children, center director information, and parent ratings all will be collected by electronic administration via REDCap with an option for oral administration of measures. The RAs will utilize the Robot for classroom observational measures via live streaming to avoid video recordings of children. All data will be entered into the REDCap system, a secure, web-based application designed exclusively to support data capture for research studies. Measures will be collected at 6-month intervals: baseline (T1), 6 months (T2), 12 months (T3), 18 months (T4), and 24 months (T5). All the same measures will be administered at all five timepoints.

2.5 Measures

2.5.1 Aim 1

Given that the toolkit is designed to mitigate COVID-19’s effects by improving resilience, we are using a parent and teacher report measure, the Devereux Early Childhood Assessment (DECA) to determine improvements to those child prosocial behaviors that are linked to resilient behaviors in young children.^{53,54} Internal consistency is adequate in a low-income diverse English and Spanish speaking sample.⁵⁵ Children’s externalizing and internalizing measures will be assessed by parent and teacher report via the Strengths and Difficulties Questionnaire. This is a primary outcome measure.

Caregiver life stress will be measured by the Everyday Stressors Index (ESI, moderator) with good reliability, validity, and internal consistency in our samples ($\alpha = .98$), and in other

studies including low-income families with young children.⁵⁶⁻⁵⁹ **Family characteristics** (i.e., English proficiency, gender, race, ethnicity, country of birth, single parent status, primary language spoken in the home, employment, income level, and years of education) will be measured by our JS+CS demographic survey.⁴⁰ This is a secondary outcome measure.

Family COVID-19 precaution behaviors will be measured using a dichotomized checklist of caregiver reported behaviors that follow CDC recommendations and adult and child vaccine uptake and confidence (if emergency authorization is approved for 1–3-year-olds) measured by specific items from the NIH Rapid Acceleration of Diagnostics (RADx) COVID-19 Common Data Elements Survey.⁶⁰⁻⁶² According to ECMHC, child development outcomes are mediated by teachers' practices (measured by implementation of JS+CS strategies assessed through the Health Environment Rating Scale–Classroom (HERS-C) and teachers' self-efficacy and beliefs (both measured by the Childcare Worker Job Stress Inventory described below).²⁴⁻²⁷

2.5.2 Aim 2

Prior ECMHC and JS research suggests that improving classroom practices impacts child development.^{25,40,63} **Classroom practices (psychosocial support safety strategies)** will be assessed through the Health Environment Rating Scale–Classroom (HERS-C) observation. This is a primary outcome measure. The HERS-C is a 30-minute observation developed by the investigators to align with core national standards including NAEYC and Caring for Our Children-3rd edition and has been used for the past 8 years.^{28,31} It is comprised of 5 domains (safety, behavioral supports, communication, self-care, and nutrition/physical activity) and scored via a 7-point Likert scale. It is useful because 1) it maps on to each of the 4 JS+CS program pillars; 2) no other measures exist that align with ECMHC key program implementation factors; and 3) it includes Nutrition and Physical Activity domains that will be used to measure

expected changes in the control centers that are receiving the obesity prevention intervention.^{64,65}

Validity is demonstrated with positive associations with the Classroom Assessment Scoring System (CLASS (r 's ranging from .30-.60)). Adequate interrater agreement has been achieved in prior studies ($\geq .80$).

Teachers' attitudes (sense of self-efficacy related to managing stressors) and teacher's beliefs (job stress, coping, distress) will be measured by the Childcare Worker Job Stress Inventory, which assesses 3 domains of workplace stress, including job demands, resources, and control.⁶⁶ The domains have strong internal consistency and constructive validity and have been used effectively in our JS pilot and other ECMHC models with low-income children.⁶⁷ **This is a secondary outcome measure.**

Teacher and consultant fidelity variables will be included as moderators: 1) Adherence Checklist: coded on proportion of strategies that were delivered compared to the number prescribed in the Action Plan, 2) Exposure Contact Log: coded on the frequency and length of consultation sessions and total minutes spent viewing online toolkit, 3) Quality of Delivery: total score out of 100 on the Fidelity Form, 4) Program Differentiation: based on a dichotomous variable on the Fidelity Form to determine if intervention spillover occurred between JS+:CS and HC2, 5) Responsiveness: total score on the Usefulness and Ease of use domains of the Technology Acceptability Fast form.

Teacher demographic characteristics (e.g., age, gender, race, ethnicity, country of birth, primary language spoken in the home, English proficiency, years of education) will be collected via the JS Demographic survey. We will use specific study questions from the RADx Common Data Elements to measure **vaccine uptake and confidence**.⁶² Given the

disproportionate impact of COVID-19 on Black and Hispanic populations⁶⁸, these factors as well as other demographics and vaccine confidence will be examined as covariates.

2.5.3 Aim 3

Adoption of center-wide policy pillars including COVID-19-related practices will be assessed quantitatively by the HERS-Program-level at all 5 time points. The HERS-P includes the same domains as the HERS-C but is focused on center policies. 1) Safety (physical safety and disaster recovery procedures), 2) Trauma-informed Behavioral Supports (teacher-child relationships, structured environment, social-emotional program practices), 3) Self-Care (staff support and supervision), and 4) Communication (teacher-parent communication and connection). This observational measure will be completed by trained RAs who have established inter-rater reliability. This is a primary outcome measure.

Key informant interviews Semi-structured interview guides will be developed based on available COVID-19 literature⁶⁹ and the results of the HERS to assess implementation barriers and facilitators and to inform necessary toolkit or cultural adaptations. Two graduate-level RAs will conduct interviews by phone or videoconference at T1, T2, T3, and T5 with a subset of directors, teachers, and parents until saturation is achieved in each of the block randomized centers to ensure equal ethnic composition of Hispanic and non-Hispanic Black participants.

Societal contextual variables outside of the centers' control (e.g., population surge in cases by zip code, vaccine access and uptake by staff and children if authorization provides, self-reported infection), will be measured by the COVID-19 Impact Survey, the RADx COVID-19 Common Data Elements.⁶²

2.6 Data Analyses

Quantitative data analysis will test for the minimal detectable effect size for intervention effects for the first two research aims based on an a priori power analysis, for child and teacher/classroom level outcomes measures. Minimum detectable effect sizes were estimated based on prior evidence of child- and teacher-level effects in pilot studies.^{70,71}

Aim 1 will use multilevel and latent growth modeling to examine short- and long-term effects of the intervention versus control condition, on child behavior problems, prosocial behaviors, and executive functioning skills. To account for the nested data structure in the cluster-randomized design, we will use multilevel models.⁷⁰ Latent growth models (LGM) will examine the long-term maintenance of intervention effects on children's outcomes over time (from baseline to 24 months). The Johnson-Neyman technique will assess moderating effects of child and family risk variables (e.g., child behavior problems and parent stress) that could influence children's outcomes.⁷²

Aim 2 will use multiple regression and latent growth modeling to examine differences in trajectories of teachers' classroom practices, attitudes (self-efficacy) and beliefs (job stress, coping, distress) between the 2 conditions (intervention and control groups) and over the 5 assessment time points. The LGM will control for teachers' baseline sociodemographic variables by specifying those as covariates at level 2.

Aim 3. Similar to other hybrid designs, we will assess implementation using an exploratory sequential mixed methods approach or one in which qualitative data are used to explain quantitative findings.^{73,74} This approach has been informative in prior Type 1 implementation-effectiveness studies to explore barriers/facilitators as well as potential societal contextual factors (e.g., vaccine uptake) to help centers serving disproportionately affected

minority communities recover from and prepare for future crises.⁷³ This ensures that the JS+CS toolkit is responsive to the needs of racially and ethnically diverse childcare providers, is perceived as valuable, and is adaptable to the broader community/policy context.

We will first examine data collected from the HERS-Program level to determine a) overall implementation adoption (average total HERS score) and b) implementation adoption of each JS+CS pillar (average HERS score by domain). We will categorize scores as either “meeting implementation standards” (i.e., HERS score ≥ 5) or “not meeting implementation standards” (i.e., HERS score < 5). The focus of our qualitative interviews will be based on quantitative findings. For centers whose scores show they are meeting vs. not meeting implementation standards, interviews will focus on implementation facilitators vs. barriers to the intervention, respectively. Given the evolving nature of COVID-19 guidelines, we will employ a rapid qualitative analysis approach to summarize main points from key informant interviews into matrices and explore relevant themes. Although there are different RQA approaches, we will follow the steps described by Hamilton⁵ which involve developing brief summary documents to summarize key points from interviews, transferring the information into matrices to view data across all participants and all domains of interest, and using a matrix analysis approach to explore relevant themes. This process has been recommended to address rapidly evolving, urgent health equity challenges⁷⁶ and will allow us to analyze qualitative data efficiently and systematically. We will integrate the quantitative and qualitative findings by producing joint displays. Specifically, we will enter HERS-P scores into tables alongside qualitative findings and illustrative quotes, to compare quotes from centers who had scores indicating they were meeting vs. not meeting implementation standards.

3. Discussion

There are few available resources to aid centers in promoting psychosocial support to address the downstream psychological impacts of COVID-19 on teaching staff and children. The purpose of this study is to evaluate a multi-level public health intervention that will promote resilience while mitigating COVID-19's long-term impacts in these high-risk populations. To date, based on PubMed, there has been no rigorously tested research on the long-term impacts of COVID-19 on young children and staff in childcare centers or on interventions to mitigate the negative effects. Childcare centers are a population that have not traditionally been given support yet have been severely impacted by COVID-19.⁷⁷ The contribution of a novel multi-lingual intervention (i.e., English, Spanish, Creole) can have far reaching impacts for diverse communities across the country.

Georgetown University's Early Childhood Mental Health Consultation is an evidence-based multi-level intervention with documented short- and long-term effects.^{63,78,79} At the center level, it provides quality supports for staff and children.⁷⁸ At the teacher level, it improves resilience by helping them develop strategies to overcome teaching challenges, strengthen self-efficacy, reduce stress, and improve classroom practices.^{78,80} At the child level, it improves child resilience by increasing prosocial skills and executive functioning.⁷⁸ JS+CS adapted the Georgetown Model during the COVID-19 pandemic to provide virtual support by promoting self-care and improving coping strategies needed at the child, teacher, and center levels. The effectiveness of this virtual program will be evaluated using this RCT. Evaluation of the telepresence robot consultation approach is vital for understanding how to best support teachers during public health crises. Virtual consultation reduces travel time and allows access to rural and remote centers, which increases consultants' capacity to serve more centers in need. Kubi

technology is an affordable telehealth option that can be adapted to diverse clients and settings beyond childcare centers.

As with all community-based research, there are anticipated barriers, such as treatment noncompliance (withdrawing or moving to a non-participating or control center). In the event of treatment non-compliance, the causal estimates of interest will be the intent to treat (ITT) and the treatment-on-treated effects. In addition, given this is a technology-based intervention, teachers who are not comfortable with the use of technology may have difficulty accessing services. In order to overcome this barrier, we will assemble a step-by-step technology guide on how to troubleshoot various scenarios as well as short video demos for teachers. The consultants will be well-versed and trained in troubleshooting, and technology support specialists on the team will be available for additional assistance. Lastly, measurement tools needed to be considered. Specifically, given the limitations of self-report, we have intentionally included multi-method measures, such as observational measures at the center-, classroom-, and child-level. Barring these challenges, the results of this study will provide insight on how to best prepare childcare centers and build their capacity to respond to subsequent public health crises and provide children with life-long coping skills.

Funding: This work is supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) [grant number R01HD105474]. The authors have no conflicts of interests or financial relationships relevant to this article to disclose.

References

1. Fore HH. A wake-up call: COVID-19 and its impact on children's health and wellbeing. *Lancet Glob Health*. 2020;8(7):e861-e862.
2. Dalton L, Rapa E, Stein A. Protecting the psychological health of children through effective communication about COVID-19. *Lancet Child Adolesc Health*. 2020;4(5):346-347.
3. Mental health and psychological resilience during the COVID-19 pandemic. <https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/mental-health-and-psychological-resilience-during-the-covid-19-pandemic>. Published 2020. Accessed July 5, 2022.
4. Kluger J. The coronavirus seems to spare most kids from illness, but its effect on their mental health is deepening. <https://time.com/5876478/children-mental-health-coronavirus>. Published 2020. Accessed July 5, 2022.
5. How COVID-19 Is Affecting the Mental Health of Children and Adolescents. <https://www.psychcongress.com/multimedia/how-covid-19-affecting-mental-health-children-and-adolescents>. Published 2020. Accessed July 5, 2022.
6. Lee J. Mental health effects of school closures during COVID-19. *Lancet Child Adolesc Health*. 2020;4(6):421.
7. Wade M, Prime H, Browne DT. Why we need longitudinal mental health research with children and youth during (and after) the COVID-19 pandemic. *Psychiatry research*. 2020;290:113143.
8. Davidson B, Schmidt E, Mallar C, et al. Risk and resilience of well-being in caregivers of young children in response to the COVID-19 pandemic. *Transl Behav Med*. 2021;11(2):305-312.
9. Robertson EL, Piscitello J, Schmidt E, Mallar C, Davidson B, Natale R. Longitudinal transactional relationships between caregiver and child mental health during the COVID-19 global pandemic. *Child Adolesc Psychiatry Ment Health*. 2021;15(1):66.
10. Mental Health During COVID-19: Signs Your Child May Need More Support. <https://www.healthychildren.org/English/health-issues/conditions/COVID-19/Pages/Signs-your-Teen-May-Need-More-Support.aspx>. Published 2021. Accessed July 5, 2022.
11. Enlow MB, Devick KL, Brunst KJ, Lipton LR, Coull BA, Wright RJ. Maternal Lifetime Trauma Exposure, Prenatal Cortisol, and Infant Negative Affectivity. *Infancy*. 2017;22(4):492-513.
12. Blair C. Developmental Science and Executive Function. *Curr Dir Psychol Sci*. 2016;25(1):3-7.
13. Raver CC. Placing emotional self-regulation in sociocultural and socioeconomic contexts. *Child Dev*. 2004;75(2):346-353.

14. Raver CC, Blair C, Willoughby M. Poverty as a predictor of 4-year-olds' executive function: new perspectives on models of differential susceptibility. *Dev Psychol.* 2013;49(2):292-304.
15. Bartlett JD, Griffin J, Thomson D. Resources for supporting children's emotional well-being during the COVID-19 pandemic. *Child Trends.* 2020;12.
16. Deborah A. Phillips JPS. From neurons to neighborhoods: The science of early childhood development. *National Academies Press.* 2000.
17. Yoshikawa H, Wuermli AJ, Britto PR, et al. Effects of the Global Coronavirus Disease-2019 Pandemic on Early Childhood Development: Short- and Long-Term Risks and Mitigating Program and Policy Actions. *J Pediatr.* 2020;223:188-193.
18. Natale R, Davidson B, Woody M, et al. A Model of Care for Delivering Supports to Childcare Providers during the COVID-19 Pandemic. *Research Square.* 2021.
19. Natale R, Jent J, Shearer R. Supporting Childcare Centers in a COVID-19 Hotspot: Mitigating Psychosocial Impacts. *Journal of Prevention.* 2022. (In submission)
20. Collie RJ. COVID-19 and Teachers' Somatic Burden, Stress, and Emotional Exhaustion: Examining the Role of Principal Leadership and Workplace Buoyancy. *AERA Open.* 2021;7:2332858420986187.
21. Cipriano C, Rappolt-Schlichtmann G, Brackett M. Supporting school community wellness with social and emotional learning (SEL) during and after a pandemic. *Edna Bennet Pierce Prevention Research Center The Pennsylvania State University.* 2020.
22. Fischer AJ, Bloomfield BS, Clark RR, McNeill AL, Erchul WP. Increasing student compliance with teacher instructions using telepresence robot problem-solving teleconsultation. *International Journal of School & Educational Psychology.* 2019;7(sup1):158-172.
23. King HC, Bloomfield BS, Wu S, Fischer AJ. A Systematic Review of School Teleconsultation: Implications for Research and Practice. *School Psychology Review.* 2022;51(2):237-256.
24. Conners-Burrow NA, Patrick T, Kyzer A, McKelvey L. A preliminary evaluation of REACH: Training early childhood teachers to support children's social and emotional development. *Early Child Educ J.* 2017;45(2):187-199.
25. Gilliam WS, Maupin AN, Reyes CR. Early Childhood Mental Health Consultation: Results of a Statewide Random-Controlled Evaluation. *J Am Acad Child Adolesc Psychiatry.* 2016;55(9):754-761.
26. Heller SS, Booth A, Keyes A, Nagle G, Sidell M, Rice J. Implementation of a mental health consultation model and its impact on early childhood teachers' efficacy and competence. *Infant Ment Health J.* 2011;32(2):143-164.
27. Upshur C, Wenz-Gross M, Reed G. A pilot study of early childhood mental health consultation for children with behavioral problems in preschool. *Early Childhood Research Quarterly.* 2009;24(1):29-45.
28. American Academy of Pediatrics, American Public Health Association, National Resource Center for Health and Safety in Child Care and Early Education. Caring for Our Children, National Health and Safety Performance Standards; Guidelines for Early Care and Education Programs. 4th ed. Itasca, IL. Am Acad Pediatrics; 2019.
29. Centers for Disease Control and Prevention. Guidance for Operating Child Care Programs during COVID-19. <https://www.cdc.gov/coronavirus/2019->

- ncov/community/schools-childcare/guidance-for-shelchildcare.html. Published 2020. Updated March 12, 2021. Accessed July 5, 2022.
30. Hemmeter ML, Snyder PA, Fox L, Algina J. Evaluating the implementation of the Pyramid Model for promoting social-emotional competence in early childhood classrooms. *Topics in Early Childhood Special Education*. 2016;36(3):133-146.
 31. National Association for the Education of Young Children. NAEYC Early Learning Program Accreditation Standards and Assessment Items. In: National Association for the Education of Young Children Washington, DC; 2019.
 32. Johnston K, Brinamen C. Mental Health Consultation in Child Care: Transforming Relationships among Directors, Staff, and Families. *ZERO TO THREE*. 2006.
 33. Hepburn K. Defining the consultative stance in the context of relationship-based work. In: Georgetown University Center for Child and Human Development's Center for Early Childhood Mental Health Consultation.
 34. Kalluri N, Kelly C, Garg A. Child Care During the COVID-19 Pandemic: A Bad Situation Made Worse. *Pediatrics*. 2021;147(3):e2020041525.
 35. Edmondson AC, Higgins M, Singer S, Weiner J. Understanding Psychological Safety in Health Care and Education Organizations: A Comparative Perspective. *Research in Human Development*. 2016;13(1):65-83.
 36. Nguyen LH, Joshi AD, Drew DA, et al. Racial and ethnic differences in COVID-19 vaccine hesitancy and uptake. *medRxiv*. 2021.
 37. McKune SL, Acosta D, Diaz N, et al. Psychosocial health of school-aged children during the initial COVID-19 safer-at-home school mandates in Florida: a cross-sectional study. *BMC Public Health*. 2021;21(1):663.
 38. Halfon N, Larson K, Lu M, Tullis E, Ruess S. Lifecourse health development: past, present and future. *Matern Child Health J*. 2014;18(2):344-365.
 39. Glasgow RE, Harden SM, Gaglio B, et al. RE-AIM Planning and Evaluation Framework: Adapting to New Science and Practice With a 20-Year Review. *Front Public Health*. 2019;7:64. Published 2019 Mar 29. doi:10.3389/fpubh.2019.00064
 40. Author et al., 2020. (Citation blinded for review)
 41. Wolfenden L, Williams CM, Wiggers J, Nathan N, Yoong SL. Improving the translation of health promotion interventions using effectiveness-implementation hybrid designs in program evaluations. *Health Promot J Austr*. 2016;27(3):204-207.
 42. Curran GM, Bauer M, Mittman B, Pyne JM, Stetler C. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care*. 2012;50(3):217-226.
 43. Natale RA, Messiah SE, Asfour L, Uhlhorn SB, Delamater A, Arheart KL. Role modeling as an early childhood obesity prevention strategy: effect of parents and teachers on preschool children's healthy lifestyle habits. *J Dev Behav Pediatr*. 2014;35(6):378-387.
 44. Natale RA, Messiah SE, Asfour LS, Uhlhorn SB, Englebert NE, Arheart KL. Obesity Prevention Program in Childcare Centers: Two-Year Follow-Up. *Am J Health Promot*. 2017;31(6):502-510.
 45. Raudenbush SW, Bryk AS. *Hierarchical linear models: Applications and data analysis methods*. Vol 1: sage; 2002.

46. Scherbaum CA, Ferreter JM. Estimating statistical power and required sample sizes for organizational research using multilevel modeling. *Organizational Research Methods*. 2009;12(2):347-367.
47. McNeish DM, Stapleton LM. The effect of small sample size on two-level model estimates: A review and illustration. *Educational Psychology Review*. 2016;28(2):295-314.
48. Bernal G, Bonilla J, Bellido C. Ecological validity and cultural sensitivity for outcome research: issues for the cultural adaptation and development of psychosocial treatments with Hispanics. *J Abnorm Child Psychol*. 1995;23(1):67-82.
49. Pina AA, Polo AJ, Huey SJ. Evidence-Based Psychosocial Interventions for Ethnic Minority Youth: The 10-Year Update. *J Clin Child Adolesc Psychol*. 2019;48(2):179-202.
50. Kreuter MW, Lukwago SN, Bucholtz RD, Clark EM, Sanders-Thompson V. Achieving cultural appropriateness in health promotion programs: targeted and tailored approaches. *Health Educ Behav*. 2003;30(2):133-146.
51. Dane AV, Schneider BH. Program integrity in primary and early secondary prevention: are implementation effects out of control? *Clin Psychol Rev*. 1998;18(1):23-45.
52. Natale RA, Lopez-Mitnik G, Uhlhorn SB, Asfour L, Messiah SE. Effect of a Child Care Center-Based Obesity Prevention Program on Body Mass Index and Nutrition Practices Among Preschool-Aged Children. *Health Promotion Practice*. 2014;15(5):695-705.
53. LeBuffe PA, Naglieri JA. The Devereux Early Childhood Assessment (DECA): A measure of within-child protective factors in preschool children. *NHSA Dialog: A Research-to-Practice Journal for the Early Intervention Field*. 1999;3(1):75-80.
54. Naglieri JA, LeBuffe PA, Shapiro VB. Assessment of social-emotional competencies related to resilience. In: *Handbook of resilience in children*. Springer; 2013:261-272.
55. Crane J, Mincic MS, Winsler A. Parent-teacher agreement and reliability on the Devereux Early Childhood Assessment (DECA) in English and Spanish for ethnically diverse children living in poverty. *Early Education & Development*. 2011;22(3):520-547.
56. Hall LA. *Social supports, everyday stressors, and maternal mental health*, University of North Carolina at Chapel Hill Chapel Hill; 1983.
57. Hall LA, Gurley DM, Seans B, Kryscio RJ. Psychosocial predictors of maternal depressive symptoms, parenting attitudes, and child behavior in single-parent families. *Nurs Res*. 1991;40(4):214-220.
58. Hall LA, Farel AM. Maternal stresses and depressive symptoms: correlates of behavior problems in young children. *Nursing research*. 1988.
59. Gomez ML, Ashford K, Linares AM, Hall LA. A Pilot Test of the Everyday Stressors Index-Spanish Version in a Sample of Hispanic Women Attending Prenatal Care. *J Nurs Meas*. 2015;23(3):519-531.
60. Gilliam WS, Malik AA, Shafiq M, et al. COVID-19 Transmission in US Child Care Programs. *Pediatrics*. 2021;147(1):e2020031971.
61. Murray TS, Malik AA, Shafiq M, et al. Association of Child Masking With COVID-19-Related Closures in US Childcare Programs. *JAMA Netw Open*. 2022;5(1):e2141227.
62. National Institutes of Health. NIH RADx-UP Common Data Elements. 2021. <https://radx-up.org/learning-resources/cdes>.

63. Watts TW, Gandhi J, Ibrahim DA, Masucci MD, Raver CC. The Chicago School Readiness Project: Examining the long-term impacts of an early childhood intervention. *PLoS One*. 2018;13(7):e0200144.
64. Hollar TL, Cook N, Natale R, Quinn D, Phillips T, DeLuca M. Training early childcare providers in evidence-based nutrition strategies can help improve nutrition policies and practices of early childcare centres serving racially and ethnically diverse children from low-income families. *Public Health Nutr*. 2018;21(7):1212-1221.
65. Natale R, Camejo S, Sanders LM. Communities putting prevention to work: results of an obesity prevention initiative in child care facilities. *Journal of Research in Childhood Education*. 2016;30(3):306-319.
66. Curbow B, Spratt K, Ungaretti A, McDonnell K, Breckler S. Development of the child care worker job stress inventory. *Early Childhood Research Quarterly*. 2000;15(4):515-536.
67. Raver CC, Jones SM, Li-Grining C, Zhai F, Bub K, Presler E. CSRP's impact on low-income preschoolers' preacademic skills: self-regulation as a mediating mechanism. *Child development*. 2011;82(1):362-378.
68. Webb Hooper M, Napoles AM, Perez-Stable EJ. COVID-19 and Racial/Ethnic Disparities. *JAMA*. 2020;323(24):2466-2467.
69. Centers for Disease Control and Prevention. Monitoring and Evaluating Mitigation Strategies in Child Care Programs. <https://www.cdc.gov/coronavirus/2019-ncov/php/monitoring-evaluating-community-mitigation-strategies/resources-by-setting/child-care-evaluation.html>. Published 2020. Updated November 13, 2020. Accessed July 5, 2022.
70. Downer JT, Williford AP, Bulotsky-Shearer RJ, et al. Using Data-Driven, Video-Based Early Childhood Consultation with Teachers to Reduce Children's Challenging Behaviors and Improve Engagement in Preschool Classrooms. *School Mental Health*. 2018;10(3):226-242.
71. Williford AP, Bulotsky-Shearer R, Bichay K, Reilly S, Downer JT. Adapting assessments of child engagement to develop an early childhood consultation model. *Journal of Applied Developmental Psychology*. 2018;56:67-78.
72. Preacher KJ, Rucker LD, Hayes AF. Addressing Moderated Mediation Hypotheses: Theory, Methods, and Prescriptions. *Multivariate Behav Res*. 2007;42(1):185-227.
73. Aarons GA, Fetter DL, Sommerfeld DH, Palinkas LA. Mixed methods for implementation research: application to evidence-based practice implementation and staff turnover in community-based organizations providing child welfare services. *Child Maltreat*. 2012;17(1):67-79.
74. Creswell JW, Clark VLP. *Designing and conducting mixed methods research*. Sage publications; 2017.
75. Hamilton AB, Finley EP. Qualitative methods in implementation research: An introduction. *Psychiatry Research*. 2019;280:112516.
76. St. George SM, Harkness AH, Rodriguez-Diaz CE, et al. Applying rapid qualitative analysis for health equity: Lessons learned using "EARS" with Latino communities. *Int J Qual Methods*. 2022. (Under review).
77. Laurencin CT, McClinton A. The COVID-19 Pandemic: a Call to Action to Identify and Address Racial and Ethnic Disparities. *J Racial Ethn Health Disparities*. 2020;7(3):398-402.

78. Duran FB, Hepburn KS, Kaufmann RK, et al. Research synthesis: Early childhood mental health consultation. Nashville, TN: The Center on the Social and Emotional Foundations for Early Learning at Vanderbilt; 2010.
79. Raver CC, Jones SM, Li-Grining CP, Metzger M, Smallwood K, Sardin L. Improving Preschool Classroom Processes: Preliminary Findings from a Randomized Trial Implemented in Head Start Settings. *Early Child Res Q.* 2008;63(3):253-255.
80. Gu Q, Day C. Teachers resilience: A necessary condition for effectiveness. *Teaching and Teacher education.* 2007;23(8):1302-1316.

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof