



# Answer to the Letter to the Editor Regarding the Comments on the Article “Radiological Evaluation of the Femoral Tunnel Positioning in Anterior Cruciate Ligament Reconstruction”

## *Resposta à carta ao editor referente aos comentários sobre o artigo “Radiological Evaluation of the femoral Tunnel Positioning in Anterior Cruciate Ligament Reconstruction”*

Luciano Rodrigo Peres<sup>1</sup> Matheus Silva Teixeira<sup>1</sup>  Caetano Scalizi Júnior<sup>1</sup> Wolf Akl Filho

<sup>1</sup>Knee Group, Orthopedics and Traumatology Service, Hospital do Servidor Público Estadual de São Paulo, São Paulo, SP, Brazil

Address for correspondence Matheus Silva Teixeira, Rua Silveira Peixoto 380, apto. 601, Água Verde, Curitiba, PR, 80240-120, Brazil (e-mail: dr.matheust@gmail.com).

Rev Bras Ortop 2021;56(1):127.

Dear editor,

First of all, we would like to thank you for your interest in our article. We appreciate your references regarding our study.

The main objective of our study was to evaluate the inclination and length of femoral tunnels and to compare the measurements obtained through computed tomography (CT) and anteroposterior (AP) radiography. Since the femoral tunnel is oblique, there is no supplementary test nor gold standard method to measure its length. Length assessment in an axial CT or CT with three-dimensional (3D) reconstruction seems more reliable. However, as described in our study and reported by the cited authors, the difficulty in comparing different tests is evident, mainly because radiography is subject to a bias regarding knee positioning in addition to the overlap and magnification of anatomical bone landmarks.

The radiological measurements in AP view were based on a line that inferiorly touches the image formed by the overlap of the entire femoral condyle, not considering femoral rotation. Since the femoral tunnel is oblique, a CT standard axial

section does not consider a more proximal position of the anatomical curvature of the condyle. Therefore, any comparison is unfeasible, since these reference lines may not be parallel.

Therefore, we decided to measure the inclination and length of the femoral tunnel at the same tomographic position, considering the lower portion of the femoral condyles. A perfect comparison between tests would require the knowledge of the real anatomical inclination and length of the femoral tunnel, either in a cadaveric evaluation or using an anatomical model made with a 3D printer.

### Conflict of Interests

The authors have no conflict of interests to declare.

### References

- 1 Hensler D, Working ZM, Illingworth KD, Tashman S, Fu FH. Correlation between femoral tunnel length and tunnel position in ACL reconstruction. *J Bone Joint Surg Am* 2013;95(22):2029–2034

received  
July 29, 2020  
accepted  
September 16, 2020

DOI <https://doi.org/10.1055/s-0040-1722595>.  
ISSN 0102-3616.

© 2021. Sociedade Brasileira de Ortopedia e Traumatologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil